

# Derived Representation Theory and the Algebraic $K$ -theory of Fields

Gunnar Carlsson<sup>1</sup>  
 Department of Mathematics  
 Stanford University  
 Stanford, California 94305

## 1 Introduction

Quillen’s higher algebraic  $K$ -theory for fields  $F$  has been the object of intense study since its introduction in 1972 [32]. The main direction of research has been the construction of “descent spectral sequences” whose  $E_2$ -term involved the cohomology of the absolute Galois group  $G_F$  with coefficients in the algebraic  $K$ -theory of an algebraically closed field. The form of such a spectral sequence was conjectured in [34] and [22], but it was soon realized that it could not be expected to converge to algebraic  $K$ -theory exactly. However, it appeared likely that it could converge to algebraic  $K$ -groups in sufficiently high dimensions, i.e. in dimensions greater than the cohomological dimension of  $G_F$ . This observation was formalized into the *Quillen-Lichtenbaum conjecture*. This conjecture has attracted a great deal of attention. Over the years, a number of special cases [42], [18] have been treated, and partial progress has been made [10]. Voevodsky in [47] proved results which led to the verification of the conjecture at  $p = 2$ . More recent work of V. Voevodsky appears to have resolved this long standing conjecture at odd primes as well. Moreover, it also appears to resolve the Beilinson-Lichtenbaum conjecture about the form of a spectral sequence which converges exactly to the algebraic  $K$ -theory of  $F$ .

The existence of this spectral sequence does not, however, provide a homotopy theoretic model for the algebraic  $K$ -theory spectrum of the field  $F$ . It is the goal of this paper to propose such a homotopy theoretic model, and to verify it in some cases. We will construct a model which depends only on the complex representation theory (or twisted versions, when the roots of unity are not present in  $F$ ) of  $G_F$ , without any other explicit arithmetic information about the field  $F$ , and is therefore contravariantly functorial in  $G_F$ . It is understood in algebraic topology that explicit space level models for spaces and spectra are generally preferable to strictly algebraic calculations of homotopy groups for the spaces. Knowledge of homotopy groups alone does not allow one to understand the behavior of various constructions and maps of spectra in the explicit way. In this case, there are specific reasons why one would find such a model desirable.

- It appears likely that one could begin to understand *finite* descent prob-

---

<sup>1</sup>Supported in part by NSF DMS 0104162

lems, since the behavior of restriction and induction maps for subgroups of finite index is relatively well understood.

- It appears that the Milnor  $K$ -groups should be identified with the homotopy groups of the so-called *derived completion* of the complex representation ring of  $G_F$ . This would relate an explicitly arithmetic invariant (Milnor  $K$ -theory) with an explicitly group-theoretic invariant coming out of the representation theory. The Bloch-Kato conjecture provides one such connection, relating Milnor  $K$ -theory with Galois cohomology. These ideas provide another one, arising out of the representation theory of the group rather than from cohomology.
- It is well understood that representation theory of Galois groups plays a key role in many problems in number theory. This construction provides another explicit link between representation theory and arithmetic in fields which it would be very interesting to explore.

The idea of the construction is as follows. We assume for simplicity that  $F$  contains an algebraically closed subfield. A more involved and more general version of the construction is discussed in the body of the paper. Let  $F$  be a field, with separable closure  $\overline{F}$ , and define the category of *descent data* for the extension  $F \subset \overline{F}$  (denoted by  $V^G(\overline{F})$ ) to have objects the finite dimensional  $\overline{F}$ -vector spaces  $V$  with  $G_F$ -action satisfying  $\gamma(\overline{f}v) = \overline{f}'\gamma(v)$ , with equivariant  $\overline{F}$ -linear isomorphisms. It is standard descent theory [17] that this category is equivalent to the category of finite dimensional  $F$ -vector spaces. On the other hand, let  $Rep_F[G_F]$  denote the category of finite dimensional continuous representations of the profinite group  $G_F$ . There is a canonical homomorphism from  $Rep_F[G_F]$  to  $V^G(\overline{F})$  obtained by applying  $\overline{F} \otimes_F -$  and extending the action via the Galois action of  $G_F$  on  $\overline{F}$ . We then have the composite

$$A_F : KRep_k[G_F] \longrightarrow KRep_F[G_F] \longrightarrow K^G(\overline{F})$$

This map is of course very far from being an equivalence, since  $\pi_0 KRep_k[G_F]$  is isomorphic to  $R[G_F]$ , a non-finitely generated abelian group, and  $\pi_0 K^G(\overline{F}) \cong \mathbb{Z}$ . However, we observe that both spectra are *commutative  $S$ -algebras* in the sense of [13], and further that the map  $A_F$  is a homomorphism of commutative  $S$ -algebras. We describe in this paper a derived version of completion, which is applicable to any homomorphism of commutative  $S$ -algebras. For such a homomorphism  $f : A \rightarrow B$ , we denote this derived completion by  $A_B^\wedge$ . We also let  $\mathbb{H}_p$  denote the mod- $p$  Eilenberg-MacLane spectrum.  $\mathbb{H}_p$  is also a commutative  $S$ -algebra, and we have an evident commutative diagram of commutative  $S$ -

algebras

$$\begin{array}{ccc}
 KRep_k[G_F] & \xrightarrow{A_F} & K^G \simeq KF \\
 \varepsilon_p \downarrow & & \varepsilon_p \downarrow \\
 \mathbb{H}_p & \xrightarrow{id} & \mathbb{H}_p
 \end{array}$$

The vertical maps are induced by forgetful functors which forget the vector space structure and only retain the dimension mod  $p$ . The completion construction is natural for such commutative squares, and we obtain a map of spectra  $\alpha_F(p) : KRep_k[G_F]_{\mathbb{H}_p}^{\wedge} \rightarrow KF_{\mathbb{H}_p}^{\wedge}$ . We will refer to  $\alpha_F(p)$  as the “representational assembly (at  $p$ )”. In this paper, we make this construction precise, and then study it, with the following results.

- We prove that  $\alpha_F(p)$  is an equivalence for fields of characteristic prime to  $p$  containing a separably closed subfield with topologically finitely generated abelian absolute Galois group. This would include, for example, iterated power series fields  $k((x_1))((x_2)) \cdots ((x_n))$ . We regard this as a “proof of concept”, in that it shows that the map is highly non-trivial. For example, this result can be extended to prove that for all fields of characteristic prime to  $p$  containing a separably closed subfield,  $\alpha_F(p)$  is surjective on  $\pi_1$ .
- We discuss how generally one would expect  $\alpha_F(p)$  to be an equivalence. Although we have no counterexamples, we suspect that for non-abelian absolute Galois groups, the statement above is inadequate and we would expect instead an equivariant form, involving the derived completions of Green functors, to be correct construction. We discuss some supporting evidence that this should be the case.
- We discuss twisted versions, i.e. versions of  $\alpha_F(p)$  which can be constructed when there is no separably closed subfield. This can even include situations where the roots of unity are not in the ground field. The case for field of positive characteristic is particularly nice, where there is a very clean statement involving the  $K$ -theory of twisted group rings. In this context, the corresponding conjecture for the case of a power series field over a finite field has been proved by G. Lyo in [23].
- We construct a possible geometric model for the derived representation theory, separate from the  $k$ -theory of the field. For non-abelian profinite groups, the representation rings are complicated non-Noetherian rings, and evaluating derived completions is a daunting task. For this reason, it appears sensible to approach the computation indirectly, by comparing it to a more geometric object. One such object is *deformation  $K$ -theory*, a construction on discrete groups which builds a version of equivariant

$K$ -theory which takes into account the presence of continuous families of deformations of representations. T. Lawson in [20] has proved that for a finitely generated nilpotent group  $\Gamma$ , the derived representation theory of the pro- $p$  completion of  $\Gamma$  is equivalent to the  $p$ -adic completion of its deformation  $K$ -theory. This suggests strongly that approaches related to deformation  $K$ -theory may very well be useful models for the derived representation theory.

In future work, we plan to study various indirect methods for evaluation of the derived representation theory, including some which are particularly well suited for comparisons with the motivic methods. We ultimately hope to be able to show, using these methods, that the homotopy groups of the derived completion at  $p$  of the representation ring of the separable Galois group (regarded as a ring spectrum via the Eilenberg-MacLane construction) will be identified with  $p$ -completed Milnor  $K$ -theory, via an explicit homomorphism. We also expect that these methods will increase our geometric understanding of this construction, which will take us in the direction of proving the conjecture as stated. We also hope to formulate a conjecture which applies to all fields, even fields of characteristic zero where the roots of unity are not present in the ground field.

The ultimate hope is that the clarification of the relationship between arithmetically defined descriptions of algebraic  $K$ -theory, such as the motivic spectral sequence, with descriptions which involve the Galois group and its representation theory directly, will shed more light on arithmetic and algebraic geometric questions.

The author wishes to express his thanks to a number of mathematicians for useful conversations, including D. Bump, R. Cohen, W. Dwyer, A. Elmendorf, E. Friedlander, L. Hesselholt, M. Hopkins, W.C. Hsiang, J.F. Jardine, M. Levine, J. Li, I. Madsen, M. Mandell, J.P. May, H. Miller, J. Morava, F. Morel, K. Rubin, C. Schlichtkrull, V. Snaith, R. Thomason, R. Vakil, and C. Weibel.

## 2 Preliminaries

We assume the reader to be familiar with the category of spectra, as developed in [12] and [13] or [19]. In either of these references, the category of spectra is shown to possess a coherently commutative and associative monoidal structure, called the smash product. Of course, on the level of homotopy categories, this monoidal structure is the usual smash product. The presence of such a monoidal structure makes it possible to define *ring spectra* as monoid objects in *Spectra*. It is also possible to define the notion of a commutative ring spectrum, with appropriate higher homotopies encoding the commutativity as well as associativity and distributivity. In line with the terminology of [13], we will refer to these objects as  $S$ -algebras and commutative  $S$ -algebras, respectively.

For the relationship with earlier notions of  $A_\infty$  and  $E_\infty$  ring spectra, see [13]. These same references will also introduce notions we will make use of, such as the Universal coefficient and the Künneth spectral sequences. See [8], section 2.1, for a discussion of the relevant notions as they will be used in the present paper.

We also remind the reader of the construction by Bousfield-Kan of the  $l$ -completion of a space (simplicial set) at a prime  $l$ , denoted by  $X_l^\wedge$ . Bousfield and Kan construct a functorial cosimplicial space  $T_l X$ , and define the  $l$ -completion of  $X$  to be  $\text{Tot}(T_l X)$ . This construction gives rise to a functorial notion of completion at a prime  $l$ . It extends in a straightforward way to spectra by applying completion at  $l$  levelwise. It is extended further in [8] to the notion of completion of a module  $M$  over a commutative  $S$ -algebra  $A$  at a commutative  $A$ -algebra  $B$ , denoted  $M_B^\wedge$ . The Bousfield-Kan construction is the special case of this construction where  $A$  is the sphere spectrum and  $B$  is the mod- $l$  Eilenberg-MacLane spectrum. A number of properties of this construction are established in [8]. We will use them extensively. We will use the special case of a group ring at one point in the paper. The relevant results are the following.

**Proposition 2.1** *Suppose that we have a homomorphism  $\varphi : G. \rightarrow H.$  of abelian simplicial groups, so that  $B\varphi$  induces an isomorphism on mod- $p$  homology groups. Then the evident homomorphism of commutative  $S$ -algebras  $|A[G.]|_{\mathbb{H}_p}^\wedge \rightarrow |A[H.]|_{\mathbb{H}_p}^\wedge$  is a weak equivalence of  $S$ -algebras, where  $\mathbb{H}_p$  is an algebra over the group ring via augmentation.*

**Proof:** It follows immediately from 2.2 below and the Atiyah-Hirzebruch spectral sequence that the map of cosimplicial spectra induced by  $\varphi$  is a levelwise equivalence, since in this case the graded algebras

$$\pi_*(B \underset{A}{\wedge} B \underset{A}{\wedge} \cdots \underset{A}{\wedge} B)$$

are  $\mathbb{F}_p$ -vector spaces. It is standard that levelwise equivalences of cosimplicial spectra induce equivalences of total spectra.  $\square$

**Proposition 2.2** *Suppose that we are given a commutative  $S$ -algebra  $A$  and an abelian simplicial group  $G.$ , and a homomorphism  $A \rightarrow B$  of commutative  $S$ -algebras. Suppose further that  $i : G. \rightarrow E.$  is an inclusion of simplicial groups, and that  $E.$  is contractible. Then the commutative  $S$ -algebra*

$$\underbrace{|A[G.]| \underset{A}{\wedge} |A[G.]| \cdots \underset{A}{\wedge} |A[G.]|}_{k \text{ factors}}$$

*is equivalent to the commutative  $S$ -algebra*

$$\underbrace{|B \underset{A}{\wedge} B \underset{A}{\wedge} \cdots \underset{A}{\wedge} B[E^k./G^{k-1}.]|}_{k \text{ factors}}$$

where  $G^{k-1}$  is included in  $E^k$  via the homomorphism  $(g_1, \dots, g_{k-1}) \rightarrow (i(g_1), i(g_2) - i(g_1), i(g_3) - i(g_2), \dots, i(g_{k-1}) - i(g_{k-2}), i(g_{k-1}))$ . Note that the quotient group is a model for  $BG^{k-1}$ .

**Proof:** Immediate from the definitions. □

We refer the reader to [32] for results concerning  $K$ -theory spectra, notably the localization, devissage, and reduction by resolution theorems. These theorems apply equally well to the completed versions  $KA_l^\wedge$ , where  $l$  is a prime. We will also recall Suslin's theorem about the  $K$ -theory of an algebraically closed field [41].

**Theorem 2.3** *Let  $k \rightarrow F$  be an inclusion of algebraically closed fields of characteristic  $p \neq l$  ( $p$  may be 0). The natural map  $Kk \rightarrow KF$  induces an equivalence  $Kk_l^\wedge \rightarrow KF_l^\wedge$ . In fact, the proof shows that the map of pro-spectra  $Kk_l^\wedge \rightarrow KF_l^\wedge$  is a weak equivalence in the sense that it induces an isomorphism of homotopy pro-groups.*

An important method for constructing spectra from combinatorial data is via *infinite loop space machines* (see [26] or [37]), which are functors from the category of symmetric monoidal categories to spectra. The algebraic  $K$ -theory functor is a prime example of this construction, since it can be obtained by applying an infinite loop space machine to the symmetric monoidal category of finitely generated projective modules over a ring. When the symmetric monoidal category has a coherently associative and distributive second monoidal structure, such as tensor product of modules, the spectrum constructed by an infinite loop space machine will have the structure of an  $S$ -algebra. If in addition the second monoidal structure is coherently commutative, the spectrum will be a commutative  $S$ -algebra. See [11] for these results. Since the tensor product of finitely generated projective modules over a commutative ring  $R$  is coherently commutative, we have

**Proposition 2.4** *For any commutative ring  $A$ , the spectrum  $KA$  is equipped with a commutative  $S$ -algebra structure in a canonical way. The  $S$ -algebra structure is functorial for homomorphisms of commutative rings.*

We also recall the ideas of equivariant stable homotopy theory. See for example [16] or [7] for information about this theory. In summary, for a finite group  $G$ , there is a complete theory of  $G$ -equivariant spectra which includes suspension

maps for one-point compactifications of all orthogonal representations of  $G$ . The proper analogue of homotopy groups takes its values in the abelian category of *Mackey functors*, which is a suitably defined category of diagrams over a category whose objects are finite  $G$ -sets, and whose morphisms include maps of  $G$ -sets, and also transfer maps attached to orbit projections. The category of Mackey functors admits a coherently commutative and associative tensor product, which is denoted by  $\square$ . Consequently, by analogy with the theory of rings, we define a *Green functor* to be a monoid object in the category of Mackey functors. The theory of ring functors has analogues for most of the standard theorems and constructions of ring theory. In particular, it is possible to define ideals, modules, completions, tensor products of modules over a Green functor  $R$ , and modules  $\text{Hom}_R(M, N)$  for any Green functor and modules  $M$  and  $N$  over  $R$ . The category of modules over a Green functor also has enough projectives, so homological algebra can be carried out in this category. It is also easy to verify that by passing to direct limits, it is possible to directly extend the ideas about Mackey and Green functors to profinite groups. See [3] for background material about Mackey and Green functors.

As mentioned above, the Mackey functor valued analogue of homotopy groups plays the same role for equivariant spectra that ordinary homotopy groups play for spectra. For example, it is shown in [21] that there exist Eilenberg-MacLane spectra attached to every Mackey functor. Moreover, a theory of ring and module spectra in the equivariant category has been developed by May and Mandell [24] and [28] in such a way that the Mackey-functor valued homotopy group becomes a Green functor in an evident way, and that the usual spectral sequences (Künneth and Universal coefficient) hold for modules in this category.

### 3 Categories of descent data

Let  $A$  be any commutative ring, equipped with a group action by a profinite group  $G$ . We assume that the action of  $G$  is continuous, in the sense that the stabilizer of any element of  $A$  is an open and closed subgroup of finite index in  $G$ .

**Definition 3.1** *By a linear descent datum for the pair  $(G, A)$ , we will mean a finitely generated  $A$ -vector space  $M$ , together with a continuous action of  $G$  on  $M$  so that  $g(am) = a^g g(m)$  for all  $g \in G$ ,  $a \in A$ , and  $m \in M$ . We define two categories of linear descent data,  $V^G(A)$  and  $V(G, A)$ . The objects of  $V^G(A)$  are all linear descent data for the pair  $(G, A)$ , and the morphisms are all equivariant  $A$ -linear morphisms. The objects of  $V(G, A)$  are also all linear descent data for  $(G, A)$ , but the morphisms are all  $A$ -linear morphisms (without any equivariance requirements). The group  $G$  acts continuously on the category  $V(G, A)$ , by conjugation of maps (so the action is trivial on objects),*

and the fixed point subcategory is clearly  $V^G(A)$ . Note that both categories are symmetric monoidal categories under direct sum.

We note that  $\otimes_A$  provides a coherently associative and commutative monoidal structure on  $V(G, A)$  and  $V^G(A)$ .

**Definition 3.2** We define the spectra  $K^G(A)$  and  $K(G, A)$  to be the spectra obtained by applying an infinite loop space machine ([26] or [37]) to the symmetric monoidal categories of isomorphisms of  $V^G(A)$  and  $V(G, A)$ , respectively.  $K(G, A)$  is a spectrum with  $G$ -action, with fixed point spectrum  $K^G(A)$ . The tensor product described above makes each of these spectra into commutative  $S$ -algebras using the results of [11].

There are various functors relating these categories (and therefore their  $K$ -theory spectra). We have the fixed point functor

$$(-)^G : V^G(A) \longrightarrow V^{\{e\}}(A^G) \cong A^G - mod$$

defined on objects by  $M \rightarrow M^G$ . We also have the induction functor

$$A \otimes_{A^G} - : A^G - mod \cong V^{\{e\}}(A^G) \longrightarrow V^G(A)$$

given on objects by  $M \rightarrow A \otimes_{A^G} M$ . The following is a standard result in descent theory. See [17] for details.

**Proposition 3.3** Let  $F$  be a field, and suppose that we are given a continuous action of a profinite group  $G$  on  $F$ , so that  $F^G \hookrightarrow F$  is a Galois extension with Galois group  $G$ . Suppose further that  $G$  is in fact the inverse limit of groups of order prime to the characteristic of  $F$ . Then both  $(-)^G$  and  $F \otimes_{F^G} -$  are equivalences of categories.

**Proof:** This is standard in the case when  $G$  is finite, and follows by a direct limit argument in the profinite group case.  $\square$

We have an analogue of this result when  $A$  is not a field. We will write  $V^G(A)_{proj}$  for the full subcategory of  $V^G(A)$  on the modules which are finitely generated projective as modules over  $A$ .

**Proposition 3.4** Suppose that  $A$  is an infinite Galois extension of a ring  $A_0 \subseteq A$ , in the sense of [9], and suppose that  $A$  is a union of finite Galois extensions. The group  $\Gamma$  of automorphisms of  $A$  over  $A_0$  is a profinite group, is equipped



with a continuous action on  $A$ , and we have  $A_0 = A^\Gamma$ . Suppose further that  $A_0$  contains a field  $k$ , and that no subgroup of  $\Gamma$  admits a continuous homomorphism to the group  $\mathbb{Z}/p\mathbb{Z}$ , where  $p$  is the characteristic of  $k$ . Of course, this condition is vacuous if  $\text{char}(k) = 0$ . Then the restriction of the functor  $(-)^\Gamma : V^\Gamma(A) \rightarrow A^\Gamma - \text{mod}$  to the subcategory  $V^\Gamma(A)_{\text{proj}}$  factors through the category  $\text{Proj}(A^\Gamma)$  of finitely generated projective modules over  $A^\Gamma$ , and the restriction is an equivalence of categories.

**Proof:** This is direct from Theorem 1.1, Ch. III of [9], and follows from a direct limit argument in the general case.  $\square$

We also have

**Proposition 3.5** *Let  $F$  be a field. The category  $V(G, F)$  is canonically equivalent to the category  $\text{Vect}(F)$  of finitely generated vector spaces over  $F$ .*

**Proof:** The definition of the morphisms has no dependence on the group action, and the result follows immediately.  $\square$

**Remark:** Note that when the group action is trivial, i.e.  $G$  acts by the identity, and  $A$  is a field, then the category  $V^G(F)$  is just the category of finite dimensional continuous  $F$ -linear representations of  $G$ .

**Definition 3.6** *In the case when the  $G$ -action is trivial, and  $A$  is a field  $F$ , we will also write  $\text{Rep}_F[G]$  for  $V^G(F)$ .*

**Proposition 3.7** *The functor*

$$\text{Rep}_{F^G}[G] \cong V^G(F^G) \xrightarrow{F^G \otimes -} V^G(F) \cong V^{\{e\}}(F^G) \cong \text{Vect}(F^G)$$

*respects the tensor product structure, and  $K(F^G)$  becomes an algebra over the  $S$ -algebra  $K\text{Rep}_{F^G}[G]$ .*

## 4 An example

Let  $k$  denote an algebraically closed field of characteristic 0, let  $k[[x]]$  denote its ring of formal power series, and let  $F = k((x))$  denote the field of fractions of  $k[[x]]$ . In this section, we wish to show that the representation theory of its absolute Galois group can be used to create a model for the spectrum  $Kk((x))$ .

The field  $F$  contains the subring  $A = k[[x]]$  of actual power series, and is obtained from it by inverting  $x$ . We begin by analyzing the spectrum  $KF$ . It

follows from the localization sequence (see [32]) that we have a fibration sequence of spectra

$$Kk \longrightarrow KA \longrightarrow KF.$$

Further, the ring  $A$  is a *Henselian local ring* with residue class field  $k$ . It follows from a theorem of O. Gabber (see [14]) that the map of spectra  $KA \rightarrow Kk$  induces an isomorphism on homotopy groups with finite coefficients, and consequently an equivalence on  $l$ -adic completions  $KA_l^\wedge \rightarrow Kk_l^\wedge$ . The fiber sequence above now becomes, up to homotopy equivalence, a fiber sequence

$$Kk_l^\wedge \xrightarrow{i} Kk_l^\wedge \longrightarrow KF_l^\wedge.$$

All three spectra in this sequence become module spectra over the commutative  $S$ -algebra  $Kk_l^\wedge$ , and the sequence consists of maps which are  $Kk_l^\wedge$ -module maps. Since the inclusion of rings  $k \hookrightarrow F$  induces an isomorphism  $\pi_0 Kk_l^\wedge \cong \pi_0 KF_l^\wedge \cong \mathbb{Z}_l$ , we find that the map  $\pi_0(i) : \pi_0 Kk_l^\wedge \rightarrow \pi_0 KF_l^\wedge$  is the zero map. Since the module  $\pi_* Kk_l^\wedge$  is cyclic over the ring  $\pi_* Kk_l^\wedge$ , this means that the inclusion induces the zero map on all homotopy groups. The conclusion is that as a  $\pi_* Kk_l^\wedge$ -module,

$$\pi_* KF_l^\wedge \cong \pi_* Kk_l^\wedge \oplus \pi_* \Sigma Kk_l^\wedge$$

where the second summand is topologically generated by the unit  $x$  viewed as an element of  $K_1(F)$ . In fact, the algebra structure is also determined, since the square of the one dimensional generator is zero. Therefore the homotopy groups are given by

$$\pi_* KF \cong \Lambda_{\mathbb{Z}_l[x]}(\xi)$$

where  $\Lambda$  denotes the Grassmann algebra functor, where the polynomial generator  $x$  is in dimension 2, and where the exterior generator  $\xi$  is in dimension 1.

The following result is proved in [49]. It is derived from the fact that the algebraic closure of  $k((x))$  is the field of *Puiseux series*, i.e. the union of the fields  $k((x^{\frac{1}{n}}))$ .

**Proposition 4.1** *The absolute Galois group  $G$  of  $k((x))$  is the group  $\hat{\mathbb{Z}}$ , the profinite completion of the group of integers.*

Let  $F$  denote the field  $k((x))$ , and let  $E$  denote  $\overline{k((x))}$ . We have observed in 3.7 that the spectrum  $KF$ , which is equivalent to  $K^G(E)$ , becomes an algebra

spectrum over the  $S$ -algebra  $KRep_F[G] \cong K^G F$ . We wish to explore the nature of this algebra structure, and to use the derived completion to demonstrate that the algebraic  $K$ -theory of  $F$  can be constructed directly from the representation theory of  $G$  over an algebraically closed field, e.g.  $\mathbb{C}$ .

We have the map of  $S$ -algebras  $KRep_F[G] \rightarrow K^G(E) \cong KF$ , induced by the functor

$$id_E \otimes_F - : V^G(F) \longrightarrow V^G(E).$$

We may compose this map with the canonical map  $KRep_k[G] \rightarrow KRep_F[G]$  to obtain a map of  $S$ -algebras

$$\hat{\alpha} : KRep_k[G] \rightarrow K^G(E).$$

We note that as it stands this map doesn't seem to carry much structure.

**Proposition 4.2**  $\pi_* KRep_k[G] \cong R[G] \otimes ku_*$ , where  $R[G]$  denotes the complex representation ring. (The complex representation ring of a profinite group is defined to be the direct limit of the representation rings of its finite quotients.)

**Proof:** Since  $k$  is algebraically closed of characteristic zero, the representation theory of  $G$  over  $k$  is identical to that over  $\mathbb{C}$ . This shows that  $\pi_0 KRep_k[G] \cong R[G]$ . In the category  $Rep_k[G]$ , every object has a unique decomposition into irreducibles, each of which has  $\mathbb{C}$  as its endomorphism ring. The result follows directly.  $\square$

For the rest of this section, we will adopt the notational convention that all  $K$ -theory spectra and groups are  $l$ -adically completed, where  $l$  is a prime. So, when we write  $KF$ , it will denote the spectrum  $(KF)_l^\wedge$ , and  $K_*F$  will denote  $\pi_*((KF)_l^\wedge)$ . We will occasionally include the completion notation for emphasis, but not generally.

We also know from the above discussion that  $K_*F \cong K_*k \oplus K_{*-1}k$ . It is now easy to check that the map  $KRep_k[G] \rightarrow K^G E \cong KF$  induces the composite

$$R[G] \otimes K_*k \xrightarrow{\varepsilon \otimes id} K_*k \hookrightarrow K_*k \oplus K_{*-1}k \cong K_*F$$

which does not appear to carry much information about  $KF$ . However, we may use the derived completion as follows. As usual, let  $\mathbb{H}_l$  denote the mod- $l$  Eilenberg-MacLane spectrum. We now have a commutative diagram of  $S$ -algebras

$$\begin{array}{ccc}
KRep_k[G] & \xrightarrow{\hat{\alpha}} & K^G E \\
\downarrow \varepsilon & & \downarrow \varepsilon \\
\mathbb{H}_l & \xrightarrow{id} & \mathbb{H}_l
\end{array}$$

where  $\varepsilon$  in both cases denotes the augmentation map which sends any vector space or representation to its dimension mod  $l$ . The naturality properties of the derived completion construction yield a map

$$\alpha_F(l) : KRep_k[G]_{\mathbb{H}_l}^{\wedge} \rightarrow K^G E_{\mathbb{H}_l}^{\wedge} \xrightarrow{\sim} KF_{\mathbb{H}_l}^{\wedge}$$

which we refer to as the *representational assembly map* for  $F$ . The rightmost equivalence is induced by the functor which carries an object in  $V^G(E)$  to its  $G$ -fixed point subspace. The goal of this section is to show that despite the fact that  $\hat{\alpha}$  carries little information about  $K_*F$ ,  $\alpha_F(l)$  is an equivalence of spectra.

**Proposition 4.3**  $KF_{\mathbb{H}_l}^{\wedge}$  is equivalent to the  $l$ -adic completion of  $KF$ .

**Proof:** We note that the  $KRep_k[G]$ -module structure on  $KF$  extends over the augmentation, i.e. there is a commutative diagram

$$\begin{array}{ccc}
KRep_k[G] \wedge KF & \longrightarrow & KF \\
\downarrow \varepsilon \wedge id & & \downarrow id \\
Kk \wedge KF & \longrightarrow & KF
\end{array}$$

where the horizontal maps are the structure maps for the module structures. Letting  $A = KRep_k[G]$  and  $B = Kk$ , we write  $KF_A$  and  $KF_B$  for  $KF$  regarded as  $A$  and  $B$  module spectra, respectively. We obtain a natural map

$$(KF_A)_{\mathbb{H}_l} \rightarrow (KF_B)_{\mathbb{H}_l}$$

which is an equivalence by Proposition 3.2 (6) of [8]. Next, let  $KF_S$  denote  $KF$  regarded as a module spectrum over the sphere spectrum  $S^0$ . Then we similarly obtain a natural map  $(KF_S)_{\mathbb{H}_l} \rightarrow (KF_B)_{\mathbb{H}_l}$ , which is an equivalence. But, example 8.2 of [8] shows that  $(KF_S)_{\mathbb{H}_l}$  is equivalent to the Bousfield-Kan  $l$ -adic completion of  $KF$ .  $\square$

It remains to determine the structure of  $\pi_* KRep_k[G]_{\mathbb{H}_l}^{\wedge}$ .

**Proposition 4.4** *There is an equivalence of spectra*

$$KRep_k[G]_{\mathbb{H}_l}^\wedge \cong ku_l^\wedge \wedge S_+^1.$$

In particular the homotopy groups are given by  $\pi_i KRep_k[G]_{\mathbb{H}_l}^\wedge \cong \mathbb{Z}_l$  for all  $i \geq 0$ , and  $\pi_i KRep_k[G]_{\mathbb{H}_l}^\wedge \cong 0$  otherwise.

**Proof:** We first show that the  $S$ -algebra  $KRep_k[G]$  may be identified with the group ring  $Kk[\chi(\mathbb{Z}_l)] \cong Kk[\mathbb{Z}/l^\infty\mathbb{Z}]$ , where  $\chi(\mathbb{Z}_l)$  denotes the group of complex characters of  $\mathbb{Z}_l$ , and  $\mathbb{Z}/l^\infty\mathbb{Z}$  denotes the union  $\bigcup_n \mathbb{Z}/l^n\mathbb{Z}$ . We observe that because  $k$  is algebraically closed of characteristic zero, the irreducible representations are in bijective correspondence with the characters of  $\mathbb{Z}_l$ . We let  $C[\chi(\mathbb{Z}_l)]$  denote the category whose objects are finite sets  $X$  equipped with a function  $\varphi_X : X \rightarrow \chi(\mathbb{Z}_l)$ , and whose morphisms are set isomorphisms respecting the character valued functions.  $C[\chi(\mathbb{Z}_l)]$  admits a notion of sum (disjoint union) and of product (the product of objects  $(X, \varphi_X)$  and  $(Y, \varphi_Y)$  is the set  $X \times Y$  equipped with the function  $\varphi_{X \times Y}$  defined by  $\varphi_{X \times Y}(x, y) = \varphi_X(x)\varphi_Y(y)$ ). There is an evident functor from  $C[\chi(\mathbb{Z}_l)]$  to  $Rep_k[\mathbb{Z}_l]$ , which carries disjoint unions to direct sums and products to tensor products, and which sends an object  $(X, \varphi_X)$  in  $C[\chi(\mathbb{Z}_l)]$  to the free  $k$ -vector space on the set  $X$ , with the representation on each element  $x \in X$  specified by the character  $\varphi_X(x)$ . Consequently, using the results of [11], we obtain a homomorphism of commutative  $S$ -algebras

$$S[\chi(\mathbb{Z}_l)] \longrightarrow KRep_k[\mathbb{Z}_l].$$

Since  $KRep_k[\mathbb{Z}_l]$  is an algebra over the commutative  $S$ -algebra  $Kk$ , we obtain a homomorphism  $Kk[\chi(\mathbb{Z}_l)] \rightarrow KRep_k[\mathbb{Z}_l]$ , which is readily verified to be an equivalence of spectra.

There is further a homomorphism  $\chi(\mathbb{Z}_l) \rightarrow S^1$ , where  $S^1$  denotes the singular complex of the circle group, and where the homomorphism is induced by the inclusion of topological groups

$$\chi(\mathbb{Z}_l) \cong \mathbb{Z}/l^\infty\mathbb{Z} \hookrightarrow S^1.$$

as the group of  $l$ -power torsion points. The map of classifying spaces induces isomorphisms on mod- $l$  homology groups, and consequently by 2.1 a weak equivalence on derived completions.  $\square$

We have now shown that the derived completion  $KRep_k[G]_{\mathbb{H}_l}^\wedge$  has the same homotopy groups as the  $K$ -theory spectrum  $KF$ . We must now show that the representational assembly induces an isomorphism on homotopy groups. We recall the localization sequence [32] which allows us to compute  $K_*(F)$ . Consider the category  $\mathcal{M} = Mod(k[[x]])$  of finitely generated modules over  $k[[x]]$ . Let  $\mathcal{T} \subseteq \mathcal{P}$  denote the full subcategory of  $x$ -torsion modules. Then the category  $Vect(F)$  may be identified with the quotient abelian category  $\mathcal{M}/\mathcal{T}$ , and we have the usual localization sequence

$$K\mathcal{T} \rightarrow K\mathcal{M} \rightarrow K\mathcal{P}/\mathcal{T}$$

which is identified with the fiber sequence

$$Kk \rightarrow Kk[[x]] \rightarrow KF.$$

We now observe that there is a twisted version of this construction. Let  $\mathcal{O}_E$  denote the integral closure of  $k[[x]]$  in  $E$ .  $\mathcal{O}_E$  is closed under the action of the group  $G$ . We now define  $\mathcal{E}^G$  to be the category whose objects are finitely generated  $\mathcal{O}_E$  modules equipped with a  $G$ -action so that  $g(em) = e^g g(m)$ . Further, let  $\mathcal{T}^G$  denote the full subcategory of torsion modules.

**Proposition 4.5** *The quotient abelian category  $\mathcal{E}^G/\mathcal{T}^G$  can be identified with the category of linear descent data  $V^G(E)$ .*

**Proof:** Straightforward along the lines of Swan's proof in [44] that  $\mathcal{M}/\mathcal{T} \cong \text{Vect}F$ .  $\square$

**Corollary 4.6** *There is up to homotopy a fiber sequence of spectra*

$$K\mathcal{T}^G \longrightarrow K\mathcal{E}^G \longrightarrow K^G(E) \cong KF.$$

We note that for each of the categories  $\mathcal{T}^G$ ,  $\mathcal{E}^G$ , and  $V^G(E)$ , objects may be tensored with finite dimensional  $k$ -linear representations of  $G$  to give new objects in the category. This construction has sufficient coherence so that the results of [11] imply that each of the  $K$ -theory spectra are module spectra over  $K\text{Rep}_k[G]$ , and that the maps between them are morphisms of module spectra.

**Lemma 4.7** *The homotopy fiber sequence of 4.6 is a homotopy fiber sequence of  $K\text{Rep}_k[G]$ -module spectra.*

Derived completion at  $\mathbb{H}_l$  preserves fiber sequences of module spectra ([8], Proposition 3.2 (3)), so we have a homotopy fiber sequence

$$(K\mathcal{T}^G)_{\mathbb{H}_l}^{\wedge} \longrightarrow (K\mathcal{E}^G)_{\mathbb{H}_l}^{\wedge} \longrightarrow (K^G(E))_{\mathbb{H}_l}^{\wedge} \cong KF.$$

The last equivalence is 4.3. We will now show that

- $(K\mathcal{T}^G)_{\mathbb{H}_l}^{\wedge} \simeq *$
- $(K\mathcal{E}^G)_{\mathbb{H}_l}^{\wedge} \simeq K\text{Rep}_k[G]_{\mathbb{H}_l}^{\wedge}$ .

The result will follow immediately.

**Proposition 4.8**  $(K\mathcal{T}^G)_{\mathbb{H}_l}^\wedge \simeq *$ .

**Proof:** Let  $\mathcal{T}_n^G$  denote the subcategory of finitely generated torsion  $k[[t^{\frac{1}{n}}]]$ -modules  $M$  equipped with  $G$ -action so that  $g(fm) = f^g g(m)$  for  $g \in G$ ,  $f \in k[[t^{\frac{1}{n}}]]$ , and  $m \in M$ . We have obvious functors

$$\tau_n^k = k[[t^{\frac{1}{k}}]] \otimes_{k[[t^{\frac{1}{n}}]]} - : \mathcal{T}_n^G \rightarrow \mathcal{T}_k^G$$

whenever  $n$  divides  $k$ , and an equivalence of categories

$$\lim_{\rightarrow} \mathcal{T}_n^G \longrightarrow \mathcal{T}^G.$$

A straightforward devissage argument now shows that the inclusion of the full subcategory of objects of  $\mathcal{T}_n^G$  on which  $x^{\frac{1}{n}}$  acts by 0 induces an equivalence on  $K$ -theory spectra. It follows that  $K_*\mathcal{T}_n^G \cong R[G] \otimes K_*k$ , and moreover that  $K\mathcal{T}_n^G \simeq KRep_k[G]$  as module spectra over  $KRep_k[G]$ . We will now need to analyze the map of  $K$ -theory spectra induced by the functors  $\tau_n^k$ . Recall that  $R[G] \cong \mathbb{Z}[\mathbb{Q}/\mathbb{Z}]$ . For all positive integers  $n$  and  $s$ , let  $\nu_{n,s}$  denote the element in  $R[G]$  corresponding to the element  $\sum_{i=0}^{s-1} [i/ns] \in \mathbb{Z}[\mathbb{Q}/\mathbb{Z}]$ . We claim that we have a commutative diagram

$$\begin{array}{ccc} K_*\mathcal{T}_n^G & \xrightarrow{\sim} & R[G] \otimes K_*k \\ \tau_n^{n,s} \downarrow & & \downarrow (\cdot \nu_{n,s}) \otimes id \\ K_*\mathcal{T}_{ns}^G & \xrightarrow{\sim} & R[G] \otimes K_*k. \end{array}$$

That we have such a diagram is reduced to a  $\pi_0$  calculation, since the diagram consists of  $R[G] \otimes K_*k$ -modules which are obtained by extension of scalars from  $R[G] \otimes K_0k$ . It therefore suffices to check the commuting of this diagram on the element  $1 \in K_*\mathcal{T}_n^G$ . But 1 under  $\tau_n^{n,s}$  clearly goes to the module  $k[[x^{\frac{1}{ns}}]]/(x^{\frac{1}{n}})$ . It is easily verified that as an element in the representation ring, this element is  $\nu_{n,s}$ .

We are now ready to describe  $\mathbb{H}_l \bigwedge_{KRep[G]} K\mathcal{T}^G$ . Since smash products commute with filtering colimits, we see that

$$\mathbb{H}_l \bigwedge_{KRep[G]} K\mathcal{T}^G \simeq \lim_{\rightarrow} \mathbb{H}_l \bigwedge_{KRep[G]} K\mathcal{T}_n^G.$$

But since  $K_*\mathcal{T}_n^G$  is a free  $R[G] \otimes K_*k$ -module of rank 1,  $\mathbb{H}_l \wedge_{KRep[G]} K\mathcal{T}_n^G \cong \mathbb{H}_l$ .

It is readily checked that the induced map

$$id_{\mathbb{H}_l} \wedge_{KRep[G]} (\cdot \nu_{ns}) \otimes id$$

is multiplication by  $s$  on  $\pi_*\mathbb{H}_l$ . It now clearly follows that  $\pi_*\mathbb{H}_l \wedge_{KRep[G]} K\mathcal{T}^G = 0$ , and consequently that

$$\underbrace{\pi_*\mathbb{H}_l \wedge_{KRep[G]} \cdots \wedge_{KRep[G]} \mathbb{H}_l \wedge_{KRep[G]} K\mathcal{T}^G}_{k \text{ factors}} = 0$$

for all  $k$ . It now follows directly from the definition of the completion, together with the fact that the total spectrum of a cosimplicial spectrum which is level-wise contractible is itself contractible, that  $(K\mathcal{T}^G)_{\mathbb{H}_l}^{\wedge} \simeq *$ .  $\square$

**Corollary 4.9** *The map  $(K\mathcal{E}^G)_{\mathbb{H}_l}^{\wedge} \rightarrow (K^G(E))_{\mathbb{H}_l}^{\wedge}$  is a weak equivalence of spectra.*

**Proof:** Completion preserves fibration sequences of spectra.  $\square$

We next analyze  $K_*\mathcal{E}^G$  in low degrees. Let  $\mathbb{N}$  be the partially ordered set of positive integers, where  $m \leq n$  if and only if  $m|n$ . We define a directed system of  $R[G]$ -modules  $\{A_n\}_{n>0}$  parametrized by  $\mathbb{N}$  by setting  $A_n = R[G]$ , and where whenever  $m|n$ , we define the bonding map from  $A_m$  to  $A_n$  to be multiplication by the element  $\nu_{m, \frac{n}{m}}$ . We let  $QR[G]$  denote the colimit of this module. It follows from the proof of 4.8 that  $K_*\mathcal{T}^G \cong QR[G] \otimes K_*k$ . In particular,  $K_*\mathcal{T}^G = 0$  in odd degrees. On the other hand, we know that  $K_*F \cong \mathbb{Z}_l$  in odd degrees, but that the elements in odd degree map injectively under the connecting homomorphism in the localization sequence. Consequently, we have

**Proposition 4.10**  $K_*\mathcal{E}^G = 0$  in odd degrees. In particular,  $K_1\mathcal{E}^G = 0$ .

**Proposition 4.11**  $K_0\mathcal{E}^G \cong R[G]$ .

**Proof:** Objects of  $\mathcal{E}^G$  are the same thing as modules over the twisted group ring  $\mathcal{O}_E\langle G \rangle$ , and a devissage argument shows that the inclusion

$$Proj(\mathcal{O}_E\langle G \rangle) \hookrightarrow Mod(\mathcal{O}_E\langle G \rangle)$$



induces an isomorphism, so that we may prove the corresponding result for finitely generated projective modules over  $\mathcal{O}_E\langle G \rangle$ . Note that we have a ring homomorphism

$$\pi : \mathcal{O}_E\langle G \rangle \longrightarrow k[G]$$

given by sending all the elements  $x^{\frac{1}{n}}$  to zero.  $\mathcal{O}_E\langle G \rangle$  is complete in the  $I$ -adic topology, where  $I$  is the kernel of  $\pi$ . As in [44], isomorphism classes of projective modules over  $\mathcal{O}_E\langle G \rangle$  are now in bijective correspondence via  $\pi$  with the isomorphism classes of projective modules over  $k[G]$ , which gives the result.  $\square$

**Remark 4.1** There should be a better proof of this result, proceeding via a non-commutative version of the results of O. Gabber [14]. That is, one should prove directly that the reduction map  $\mathcal{O}_E\langle G \rangle \rightarrow k[G]$  is an equivalence by proving that it is sufficiently ‘‘Henselian’’. It seems quite likely that such a proof could be made to work, which would mean that the proof would not require as input any computational information about the field in question.

**Corollary 4.12** *The functor  $\mathcal{O}_E \otimes_k - : \text{Rep}_k[G] \rightarrow \mathcal{E}^G$  induces a weak equivalence on spectra. Consequently, the map  $K\text{Rep}_k[G]_{\mathbb{H}_l}^\wedge \rightarrow (K\mathcal{E}^G)_{\mathbb{H}_l}^\wedge$  is a weak equivalence.*

**Proof:** 4.10 and 4.11 identify the homotopy groups of  $K\mathcal{E}^G$  in degrees 0 and 1. From the localization sequence above, together with the fact that multiplication by the Bott element induces isomorphisms  $K_i\mathcal{T}^G \rightarrow K_{i+2}\mathcal{T}^G$  and  $K_iF \rightarrow K_{i+2}F$ , it follows that it also induces an isomorphism  $K_i\mathcal{E}^G \rightarrow K_{i+2}\mathcal{E}^G$ . So, the map of the statement of the corollary induces an isomorphism on homotopy groups, which is the required result.  $\square$

**Theorem 4.13** *The map  $\alpha_F(l) : K\text{Rep}_k[G]_{\mathbb{H}_l}^\wedge \rightarrow K^G(E)_{\mathbb{H}_l}^\wedge \cong KF_{\mathbb{H}_l}^\wedge$  is a weak equivalence of spectra.*

**Proof:**  $\alpha_F(l)$  is the composite of the maps of Corollary 4.9 and Corollary 4.12, both of which we have shown are weak equivalences.  $\square$

## 5 The case of an abelian absolute Galois group

We again adopt the convention that all  $K$ -theory spectra are assumed to be  $l$ -completed, and that  $K$ -groups are interpreted as homotopy groups of  $l$ -completed

$K$ -theory spectra. We will now show how to extend the result of the previous section to the case of geometric fields (i.e. containing and algebraically closed subfield)  $F$  of characteristic zero with  $G_F$  a free pro- $l$  abelian group. Let  $F$  be geometric, and let  $k \subseteq F$  denote an algebraically closed subfield. Let  $A = k[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_n^{\pm 1}]$ . We let  $B$  denote the ring obtained by adjoining all  $p$ -th power roots of unity of the variables  $t_i$  to  $A$ , so

$$B = \bigcup_s k[t_1^{\pm \frac{1}{l^s}}, t_2^{\pm \frac{1}{l^s}}, \dots, t_n^{\pm \frac{1}{l^s}}]$$

We choose a sequence of elements  $\zeta_n$  in  $k$  so that  $\zeta_n$  is a primitive  $l^n$ -th root of unity, and so that  $\zeta_n^l = \zeta_{n-1}$ . We define an action of the group  $\mathbb{Z}_l^n$  on  $B$  by  $\tau_i(t_i^{\frac{1}{l^n}}) = \zeta_n t_i^{\frac{1}{l^n}}$ , and  $\tau_j(t_i^{\frac{1}{l^n}}) = t_i^{\frac{1}{l^n}}$  for  $i \neq j$ , where  $\{\tau_1, \tau_2, \dots, \tau_n\}$  is a set of topological generators for  $\mathbb{Z}_l^n$ .

**Proposition 5.1** *The  $l$ -completed  $K$ -theory groups of  $A$  and  $B$  are given by*

- $K_*A \cong \Lambda_{K_*k}(\theta_1, \theta_2, \dots, \theta_n)$
- $K_*B \cong K_*k$

**Proof:** The first result is a direct consequence of the formula for the  $K$ -groups of a Laurent polynomial ring. The second also follows from this fact, together with analysis of the behavior of the  $l$ -th power map on a Laurent extension.  $\square$

We now may construct a representational assembly in this case as well, to obtain a representational assembly

$$\alpha_L(l) : KRep_k[G]_{\mathbb{H}_l}^\wedge \longrightarrow K^G(B)_{\mathbb{H}_l}^\wedge \cong KA_{\mathbb{H}_l}^\wedge \cong KA_l^\wedge$$

where the first equivalence follows from 3.4, and the second in an identical fashion to that in the proof of 4.3.

**Proposition 5.2**  $\alpha_L(l)$  *is an equivalence of spectra.*

**Proof:** We first consider the case  $n = 1$ , and the field  $F = k((x))$ , where we have already done the analysis. It is immediate that the inclusion  $A \rightarrow F$  defined by  $t \rightarrow x$  extends to an equivariant homomorphism  $B \rightarrow E$  of  $k$ -algebras, and consequently that we get a commutative diagram

$$\begin{array}{ccc} KRep_k[G] & & \\ \alpha_L(l) \downarrow & \searrow \alpha_F(l) & \\ KA & \xrightarrow{\cong} & KF \end{array}$$

It is readily checked that the inclusion  $KA \rightarrow KF$  is a weak equivalence, and so that the horizontal arrow in the diagram is a weak equivalence. On the other hand, we have already shown that  $\alpha_F(l)$  is a weak equivalence of spectra, which shows that  $\alpha_L(l)$  is an equivalence. For larger values of  $n$ , the result follows by smashing copies of this example together over the coefficient  $S$ -algebra  $K_*k$ .  $\square$

**Remark 5.1** As in Remark 4.1, we believe that there is another proof of this result, this time following from the homotopy property of [32]. The idea would be to observe that the group action preserves the polynomial subrings  $k[t_1^{\frac{1}{l^n}}, \dots, t_n^{\frac{1}{l^n}}]$ , and that a corresponding twisted polynomial ring should have  $K$ -theory computable using Quillen's results.

Now consider any geometric field  $F$  of characteristic zero, with  $G_F \cong \mathbb{Z}_l^n$ , and let  $k$  be an algebraically closed subfield. It is a direct consequence of Kummer theory that there is a family of elements  $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$  so that  $E = \overline{F}$  can be written as

$$E = \bigcup_n F(\sqrt[l^n]{\alpha_1}, \sqrt[l^n]{\alpha_2}, \dots, \sqrt[l^n]{\alpha_n})$$

It is therefore further clear that we may define a  $k$ -algebra homomorphism  $i : B \rightarrow E$  by setting  $i(t_i^{\frac{1}{l^n}}) = \sqrt[l^n]{\alpha_i}$ , where  $\sqrt[l^n]{\alpha_i}$  is chosen as the choice of  $l^n$ -th root so that the action of  $\tau_i$  is via multiplication by  $\zeta_n$ . It follows that we obtain a commutative diagram

$$\begin{array}{ccc} KRep_k[G]_{\mathbb{H}_l}^{\wedge} & & \\ \alpha_L(l) \downarrow & \searrow \alpha_F(l) & \\ KA & \xrightarrow{Ki} & KF \end{array}$$

Since we now know that  $\alpha_L(l)$  is a weak equivalence, it will now suffice to show that  $Ki$  is a weak equivalence of spectra. But the work of Geisser-Levine [15] shows that if the Bloch-Kato conjecture holds, then the  $l$ -completed version of the Bloch-Lichtenbaum spectral sequence converging to the algebraic  $K$ -theory has as its  $E_2$ -term an exterior algebra on generators in degree one, corresponding to the elements  $\alpha_1, \alpha_2, \dots, \alpha_n$ , from which this result follows directly.

**Proposition 5.3** *If the Bloch-Kato conjecture holds, then  $\alpha_F(l)$  is an equivalence of spectra for geometric fields of characteristic prime to  $l$  with finitely generated abelian separable Galois group.*

## 6 Surjectivity on $\pi_1$ in the geometric case

Let  $F$  be any field containing an algebraically closed subfield  $k$ , and suppose  $l$  is prime to the characteristic of  $k$ . We claim that the map  $\alpha_F(l)$  induces a surjection on  $\pi_1$ .

**Proposition 6.1** *For any field  $F$ , we have*

$$\pi_1((KF)_l^\wedge) \cong (F^*)_l^\wedge$$

where the right hand term denotes the usual algebraic  $l$ -adic completion of the group  $F^*$ .

**Proof:** Since  $\pi_0(KF) \cong \mathbb{Z}$  is a finitely generated abelian group, this result follows from Proposition 5.1 in Ch VI of [4].  $\square$

Functoriality of the assembly construction now gives us a density result.

**Proposition 6.2** *Suppose  $F$  contains an algebraically closed subfield  $k$ , and that  $l$  is prime to  $\text{char}(k)$ . Then the image of the homomorphism*

$$\pi_1(\alpha_F(l)) : \pi_1(K\text{Rep}_k[G]_{\mathbb{H}_l}^\wedge) \rightarrow \pi_1((KF)_l^\wedge) \cong (F^*)_l^\wedge$$

is dense in the profinite topology on  $(F^*)_l^\wedge$ . In

**Proof:** It clearly suffices to prove that for any  $n$  and any element  $x \in F^*/l^n F^*$ , there is an element  $\xi \in \pi_1(K\text{Rep}_k[G]_{\mathbb{H}_l}^\wedge)$  so that  $\rho(\pi_1(\alpha_F(l))(\xi)) = x$ , where  $\rho$  is the projection  $F^* \rightarrow F^*/l^n F^*$ . In order to do this, we select  $\bar{x} \in F^*$  so that  $\rho(\bar{x}) = x$ . We next consider the extension  $\bigcup_n F(x^{\frac{1}{l^n}})$  of  $F$ . This extension is preserved under the action of the separable Galois group  $G$ , and therefore yields a particular topologically cyclic quotient of  $G$ , which we denote by  $Q$ . We note that  $Q$  is also a pro- $l$  group, and is therefore isomorphic to  $\mathbb{Z}_l$ . Select a particular identification  $\theta : Q \simeq \mathbb{Z}_l$ . Next, define a ring homomorphism  $A = k[t^{\pm 1}] \rightarrow F$  by sending the variable  $t$  to  $\bar{x}$ . Choosing a sequence of primitive  $l^n$ -th roots of unity  $\zeta_n$  as in the beginning of Section 5, we extend this map to a ring homomorphism

$$i : B = \bigcup_n k[t^{\pm \frac{1}{l^n}}] \rightarrow F$$

via the requirement that  $t^{\pm \frac{1}{l^n}}$  be sent to the unique  $l^n$ -th root of  $x$  on which the topological generator  $\theta^{-1}(1)$  acts by multiplication by  $\zeta_n$ . It is now clear from the construction that the diagram

$$\begin{array}{ccc}
KRep_k[\mathbb{Z}_l]_{\mathbb{H}_l}^{\wedge} & \xrightarrow{\alpha_L(l)} & KA_l^{\wedge} \\
\searrow & & \downarrow \\
& & KF_l^{\wedge} \\
& \swarrow_{\alpha_F(l)} & 
\end{array}$$

where the vertical arrow is induced by the ring homomorphism  $A \rightarrow F$  defined above. It was proved in Section 5 that  $\alpha_L(l)$  is an equivalence, hence induces an isomorphism on  $\pi_1$ . The result now follows immediately.  $\square$

The actual surjectivity now follows from

**Proposition 6.3** *The groups  $\pi_t(KRep_k[G]_{\mathbb{H}_l}^{\wedge})$  are all  $l$ -complete.*

**Proof:** It follows from the definition of the derived completions that the derived completion is the total spectrum of a cosimplicial spectrum which in every level is a mod- $l$  Eilenberg-MacLane spectrum, hence is  $l$ -complete in the sense of [4]. It is standard that the total space of such a cosimplicial spectrum itself has  $l$ -complete homotopy groups.  $\square$

## 7 Representational assembly in the geometric case

Suppose now that we are in the case of a *geometric field*, i.e. a field  $F$  containing an algebraically closed subfield  $k$ . Let  $E$  denote the algebraic closure of  $F$ , and let  $G$  denote the absolute Galois group. As in the case of the example of the previous section, we have a composite functor

$$Rep_k[G] \longrightarrow Rep_F[G] \cong V^G(F) \longrightarrow V^G(E)$$

and the induced map of completed spectra

$$KRep_k[G]_{\mathbb{H}_l}^{\wedge} \longrightarrow K^G(E)_{\mathbb{H}_l}^{\wedge} \cong KF_{\mathbb{H}_l}^{\wedge}$$

We may ask whether this map is an equivalence as occurred in the case of the example in the previous section. In order to discuss the plausibility of this kind of statement, let us examine what occurs in the example, when  $G = \mathbb{Z}_l$ . In this case,  $\pi_0 KRep_k[G] \cong \mathbb{Z}[\mathbb{Q}/\mathbb{Z}]$ . This is a very large, non-Noetherian ring. However, after derived completion, we find that  $\pi_0 KRep_k[G]_{\mathbb{H}_l}^{\wedge} \cong \mathbb{Z}_l$ . On the other hand, we have from Theorem 7.1 of [8] that  $\pi_0 KRep_k[G]_{\mathbb{H}_l}^{\wedge}$  is isomorphic to

the ordinary (not derived) completion of the ring  $R[G]$  at the ideal  $J = (l) + I$ , where  $I$  denotes the augmentation ideal in  $R[G]$ . So the conclusion is that  $\varprojlim R[G]/J^n \cong \mathbb{Z}_l \cong (R[G]/I)^\wedge$ . The reason this happens is the following result about the ideal  $I$ .

**Proposition 7.1** *We have that  $I/I^2$  is a divisible group, and that  $I^k = I^{k+1}$  for all  $k$ .*

**Proof:** It is a standard result for any group  $G$  that in the group ring  $\mathbb{Z}[G]$ , the augmentation ideal  $I[G]$  has the property that  $I[G]/I[G]^2 \cong G^{ab}$ . Since  $\mathbb{Q}/\mathbb{Z}$  is abelian, we find that  $I/I^2 \cong \mathbb{Q}/\mathbb{Z}$ , which is divisible. For the second statement, we note that  $I^k/I^{k+1}$  is a surjective image of  $\underbrace{I/I^2 \otimes \dots \otimes I/I^2}_{k \text{ factors}}$ . But it is clear

that  $\mathbb{Q}/\mathbb{Z} \otimes \mathbb{Q}/\mathbb{Z} = 0$ , so  $I^k/I^{k+1} = 0$ . This gives the result.  $\square$

**Corollary 7.2** *We have that  $((l)+I)^k = (l^k)+I$ . Equivalently, for any element  $\theta \in I$ , and any positive integers  $s$  and  $t$ , there is an  $\eta \in I$  and  $\mu \in I^k$  so that*

$$\theta = l^s \eta + \mu.$$

**Proof:** We have

$$((l) + I)^k = \sum_t l^t I^{k-l} = (l^k) + l^{k-1}I + I^2.$$

But 7.1 implies that  $l^{k-1}I + I^2 = I$ , which gives the result.  $\square$

It now follows that  $R[G]/J^k \cong R[G]/(I + (l^k)) \cong \mathbb{Z}/l^k\mathbb{Z}$ , and therefore that the completion of  $R[G]$  at the ideal  $J$  is just  $\mathbb{Z}_l$ .

This result extends to torsion free abelian profinite groups. That is, any such group also has the property that if  $I$  is the augmentation ideal in the representation ring, then  $I^k = I^2$  for  $k \geq 2$ , and that  $I/I^2$  is divisible. However, for non-abelian Galois groups, there appears to be no obvious reason why such a result should hold, and indeed we believe that it does not. However, there is a modification of this statement which is true for absolute Galois groups and which suggests a conjecture which can be formulated for any geometric field.

Recall that the representation ring of a finite group is not just a ring, but is actually a part of a *Green functor*. This was discussed in section 2, where there is a discussion of Mackey and Green functors. See [3] for a thorough discussion of these objects. A Mackey functor for a group  $G$  is a functor from a category of orbits to abelian groups. In the case of the representation ring, there is a functor  $\mathcal{R}$  given on orbits by  $\mathcal{R}(G/K) = R[K]$ . The maps induced by projections of orbits induce restriction maps on representation rings, and transfers

induce inductions. This functor is actually a commutative *Green functor*, in the sense that there is a multiplication map  $\mathcal{R} \square \mathcal{R} \rightarrow \mathcal{R}$ , which is associative and commutative. Also for any finite group  $G$ , there is another Green functor  $\mathcal{Z}$  given on objects by  $\mathcal{Z}(G/K) = \mathbb{Z}$ , and for which projections of orbits induce the identity map and where transfers associated to projections induce multiplication by the degree of the projection. We may also consider  $\mathcal{Z}/l\mathcal{Z}$ , which is obtained by composing the functor  $\mathcal{Z}$  with the projection  $\mathbb{Z} \rightarrow \mathbb{Z}/l\mathbb{Z}$ . The *augmentation* is the morphism of Green functors  $\mathcal{R} \rightarrow \mathcal{Z}$  which is given on an object  $G/K$  by the augmentation of  $R[K]$ . The mod- $l$  augmentation  $\varepsilon$  is the composite  $\mathcal{R} \rightarrow \mathcal{Z} \rightarrow \mathcal{Z}/l\mathcal{Z}$ . The theory of Green functors is entirely parallel with the theory of rings, modules, and ideals. We may therefore speak of the *augmentation ideal*  $\mathcal{I}$  and the ideal  $\mathcal{J} = (l) + \mathcal{I}$ , as well as the powers of these ideals. We can therefore also speak of completion at a Green functor ideal. We also observe that the theory of Mackey and Green functors extends in an obvious way to profinite groups, by considering the category of finite  $G$ -orbits. We want to identify a class of profinite groups  $G$  for which

$$\mathcal{R}_l^\wedge = \varprojlim \mathcal{R}/\mathcal{J}^n$$

is isomorphic to  $\mathcal{Z}_l^\wedge = \mathbb{Z}_l \otimes \mathcal{Z}$ .

**Definition 7.3** *Let  $G$  be a profinite group. We say  $G$  is totally torsion free if every subgroup of finite index has a torsion free abelianization.*

**Example:** Free profinite groups and free profinite  $l$ -groups are totally torsion free.

**Example:** Free profinite abelian and profinite  $l$ -abelian groups.

**Example:** Let  $\Gamma$  denote the integral Heisenberg group, i.e. the group of upper triangular integer matrices with ones along the diagonal. Then the profinite and pro- $l$  completion of  $\Gamma$  is not totally torsion free.

**Proposition 7.4** *Let  $G$  be the absolute Galois group of a geometric field  $F$ . Let  $G_l$  denote the maximal pro- $l$  quotient of  $G$ . Then  $G_l$  is totally torsion free.*

**Proof:** Consider any subgroup  $K$  of finite index in  $G_l$ . Then let  $E$  denote the extension of  $F$  corresponding to  $K$ . Then the abelianization of  $K$  corresponds to the maximal pro- $l$  Abelian extension of  $E$ . Let  $\mathbb{N}$  denote the partially ordered set of positive integers. Define a functor  $\Phi$  from  $\mathbb{N}$  to abelian groups by  $\Phi(n) = E^*/(E^*)^{l^n}$ , and on morphisms by  $\Phi(m \leq m+n) = E^*/(E^*)^{l^m} \xrightarrow{(-)^{l^n}} E^*/(E^*)^{l^{m+n}}$ . Let  $\mathcal{E}$  denote the direct limit over  $\mathbb{N}$  of  $\Phi$ . Kummer theory then asserts the existence of a perfect pairing

$$K^{ab} \times \mathcal{E} \rightarrow \mathbb{Q}/\mathbb{Z}$$

So  $K^{ab} \cong \text{Hom}(\mathcal{E}, \mathbb{Q}/\mathbb{Z})$ . The group  $\mathcal{E}$  is clearly  $l$ -divisible, and it is easily verified that the  $\mathbb{Q}/\mathbb{Z}$ -dual of an  $l$ -divisible group is  $l$ -torsion free.  $\square$

We have the following result concerning totally torsion free profinite groups.

**Proposition 7.5** *Let  $G$  be a totally torsion free  $l$ -profinite group. Then the natural map  $\mathcal{R}_l^\wedge \rightarrow \mathcal{Z}_l^\wedge$  is an isomorphism. Consequently, for  $G$  the maximal pro- $l$  quotient of the absolute Galois group of a geometric field, we have that  $\mathcal{R}_l^\wedge \cong \mathcal{Z}_l^\wedge$ .*

**Proof:** We will verify that  $\mathcal{R}_l^\wedge(G/G) \rightarrow \mathcal{Z}_l^\wedge(G/G)$  is an isomorphism. The result at any  $G/K$  for any finite index subgroup will follow by using that result for the totally torsion free group  $K$ . It will suffice to show that for every finite dimensional representation  $\rho$  of  $G/N$ , where  $N$  is a normal subgroup of finite index, and every choice of positive integers  $s$  and  $t$ , there are elements  $x \in \mathcal{R}(G/G)$  and  $y \in \mathcal{I}^t(G/G)$  so that  $[\dim \rho] - [\rho] = l^s x + y$ . We recall *Blichfeldt's theorem* [39], which asserts that there is a subgroup  $L$  of  $G/N$  and a one-dimensional representation  $\rho_L$  of  $L$  so that  $\rho$  is isomorphic to the representation of  $G/N$  induced up from  $\rho_L$ . It follows that  $[\dim \rho] - [\rho] = i_L^{G/N}(1 - \rho_L)$ . Let  $\bar{L}$  denote the subgroup  $\pi^{-1}L \subseteq G$ , where  $\pi : G \rightarrow G/N$  is the projection, and let  $\rho_{\bar{L}} = \rho_L \circ \pi$ . Then we clearly also have  $[\dim \rho] - [\rho] = i_{\bar{L}}^G(1 - \rho_{\bar{L}})$ . Now,  $1 - \rho_{\bar{L}}$  is in the image of  $R[\bar{L}^{ab}] \rightarrow R[\bar{L}]$ , and let the corresponding one-dimensional representation of  $\bar{L}^{ab}$  be  $\rho_{\bar{L}^{ab}}$ . Since  $\bar{L}^{ab}$  is abelian and torsion free (by the totally torsion free hypothesis), we may write  $1 - \rho_{\bar{L}^{ab}} = l^s \xi + \eta$ , where  $\eta \in I^t(\bar{L}^{ab})$  and where  $\xi \in R[\bar{L}^{ab}]$ , by 7.2. This means that we may pull  $\xi$  and  $\eta$  back along the homomorphism  $\bar{L} \rightarrow \bar{L}^{ab}$ , to get elements  $\bar{\xi} \in R[\bar{L}]$  and  $\bar{\eta} \in I^t(\bar{L})$  so that  $\rho_{\bar{L}} = l^s \bar{\xi} + \bar{\eta}$ . Since  $\mathcal{I}^t$  is closed under induction, we have that  $i_{\bar{L}}^G(\bar{\eta}) \in \mathcal{I}^t(G/G)$ . Now we have that  $[\dim \rho] - [\rho] = i_{\bar{L}}^G(1 - \rho_{\bar{L}}) = l^s i_{\bar{L}}^G \bar{\xi} + i_{\bar{L}}^G \bar{\eta}$ . The result follows.  $\square$

We recall the relationship between Mackey functor theory and equivariant stable homotopy theory. The natural analogue for homotopy groups in the world of equivariant spectra takes its values in the category of Mackey functors. The Adams spectral sequence and the algebraic to geometric spectral sequence have  $E_2$ -terms which are computed using derived functors of Hom and  $\square$ -product, which we will denote by *Ext* and *Tor* as in the non-equivariant case. We will now observe that the constructions we have discussed are actually part of equivariant spectra.

**Proposition 7.6** *For any profinite group  $G$ , there is a stable category of  $G$ -spectra, which has all the important properties which the stable homotopy theory*



of  $G$ -spectra for finite groups has. In particular, there are fixed point subspectra for every subgroup of finite index as well as transfers for finite  $G$ -coverings. The “tom Dieck” filtration holds as for finite groups. The homotopy groups of a  $G$ -spectrum form a Mackey functor. Moreover, the homotopy groups of a  $G$ - $S$ -algebra are a Green functor, and the homotopy groups of a  $G$ -module spectrum are a module over this Green functor.

**Proof:** For any homomorphism  $f : G \rightarrow H$  of finite groups, there is a pullback functor  $f^*$  from the category of  $H$ -spectra to the category of  $G$ -spectra. Hence, for a profinite group  $g$ , we get a direct limit system of categories parametrized by the normal subgroups of finite index, with the value at  $N$  being the category of  $G/N$ -spectra. A  $G$  spectrum can be defined as a family of spectra  $\mathcal{S}_N$  in the category of  $G/N$ -spectra, together with isomorphisms

$$(G/N_1 \rightarrow G/N_2)^*(\mathcal{S}_{N_2}) \xrightarrow{\sim} \mathcal{S}_{N_1}.$$

This general construction can be made into a theory of  $G$  spectra with the desired properties.  $\square$

**Proposition 7.7** *Let  $F$  be a field, with  $E$  its algebraic closure, and  $G$  the absolute Galois group. Then there is a  $G$ - $S$ -algebra with total spectrum  $K(G, E)$ , with fixed point spectra  $K(G, E)^H \cong K^H(E)$ . The attached Green functor is given by*

$$G/L \rightarrow \pi_* K(G, E)^L \cong \pi_* K^L(E) \cong K_*(E^L).$$

*Similarly, there is a  $G$ - $S$ -algebra with total spectrum  $K(G, F)$ , and with fixed point spectra  $K(G, F)^L \cong K^L(F) \cong K \text{Rep}_F[L]$ . (Note that the  $G$ -action on  $F$  is trivial.) In this case, the associated Green functor is given by*

$$G/L \rightarrow \pi_* K(G, F)^L \cong \pi_* K^L(F) \cong K_* \text{Rep}_F[L]$$

*In the case when  $F$  contains all the  $l$ -th power roots of unity, we find that the Mackey functor attached to  $K(G, F)$  is  $ku_* \otimes \mathcal{R}$ . The functor  $E \otimes_F -$  induces a map of  $G$ - $S$ -algebra  $K(G, F) \rightarrow K(G, E)$ , which induces the functors of Proposition 3.7 on fixed point spectra.*

**Proof:** This result is essentially a consequence of the equivariant infinite loop space recognition principle [40].  $\square$

We also recall the results of [21], where it was shown that in the category of  $G$ -equivariant spectra, there is an Eilenberg-MacLane spectrum for every Mackey functor. We will let  $\mathcal{H}_l$  denote the Eilenberg-MacLane spectrum attached to

the Green functor  $\mathcal{Z}/l\mathcal{Z}$ . Derived completions of homomorphisms of  $S$ -algebras are defined as in the non-equivariant case.

**Proposition 7.8** *Let  $F$  be a geometric field, with the algebraically closed subfield  $k$ . There is a commutative diagram of  $G$ - $S$ -algebras*

$$\begin{array}{ccc} K(G, k) & \xrightarrow[k]{E \otimes} & K(G, E) \\ & \searrow \varepsilon & \downarrow \varepsilon \\ & & \mathcal{H}_l \end{array}$$

where the maps  $\varepsilon$  are given by “dimension mod  $l$ ”. Consequently, there is a map of derived completions

$$\alpha^{rep} : K(G, k)_{\mathcal{H}_l}^{\wedge} \longrightarrow K(G, E)_{\mathcal{H}_l}^{\wedge}$$

It follows from the equivariant algebraic to geometric spectral sequence that  $\pi_0 K(G, k)_{\mathcal{H}_l}^{\wedge} \cong \mathcal{R}_l^{\wedge}$ , and by Proposition 7.5 this is isomorphic to  $\mathcal{Z}_l^{\wedge}$ , which is in turn isomorphic to the Green functor  $\pi_0 K(G, E)$ . This makes plausible the following conjecture.

**Conjecture 7.9** *Let  $F$  be a geometric field. The representational assembly map*

$$\alpha^{rep} : K(G, k)_{\mathcal{H}_l}^{\wedge} \rightarrow K(G, E)_{\mathcal{H}_l}^{\wedge}$$

is an equivalence of  $G$ - $S$ -algebras. On  $G$ -fixed point spectra, we have an equivalence of  $S$ -algebras

$$(\alpha^{rep})^G : (K(G, k)_{\mathcal{H}_l}^{\wedge})^G \longrightarrow (K(G, E)_{\mathcal{H}_l}^{\wedge})^G \cong K(E^G) \cong KF.$$

## 8 Representational assembly in the twisted case

In the case of a field  $F$  which does not contain an algebraically closed subfield, it is not as easy to construct a model for the  $K$ -theory spectrum of the field. As usual, let  $F$  denote a field,  $E$  its algebraic closure, and  $G$  its absolute Galois group. We observe first that for any subfield of  $F' \subseteq E$  which is closed under the action of  $G$ , we obtain as above a diagram

$$\begin{array}{ccc}
K(G, F') & \xrightarrow[E']{E \otimes} & K(G, E) \\
& \searrow \varepsilon & \downarrow \varepsilon \\
& & \mathcal{H}_l
\end{array}$$

and consequently a map

$$K(G, F')_{\mathcal{H}_l}^{\wedge} \longrightarrow K(G, E)_{\mathcal{H}_l}^{\wedge}.$$

We may conjecture (as an extension of Conjecture 7.9 above) that this map is always an equivalence, and indeed we believe this to be true. In the case when  $F'$  is algebraically closed, we believe that the domain of the map has what we would regard as a simple form, i.e. is built out of representation theory of  $G$  over an algebraically closed field. In general, though, the  $K$ -theory of the category  $V^G(F')$  may not be a priori any simpler than the  $K$ -theory of  $F$ . However, when  $F'$  is a field whose  $K$ -theory we already understand well, then we expect to obtain information this way. In addition to algebraically closed fields, we have an understanding of the  $K$ -theory of finite fields from the work of Quillen [33]. So, consider the case where  $F$  has finite characteristic  $p$ , distinct from  $l$ . We may in this case let  $F'$  be the maximal finite subfield contained in  $F$ , which is of the form  $\mathbb{F}_q$ , where  $q = p^n$  for some  $n$ . In this case we have, in parallel with Conjecture 7.9,

**Conjecture 8.1** *For  $F$  of finite characteristic  $p \neq l$ , with  $\mathbb{F}_q$  the maximal finite subfield of  $F$ , the map*

$$K(G, \overline{\mathbb{F}_q})_{\mathcal{H}_l}^{\wedge} \longrightarrow K(G, E)_{\mathcal{H}_l}^{\wedge}$$

*is an equivalence of  $G$ - $S$ -algebras. In particular, we have an equivalence*

$$(K(G, \overline{\mathbb{F}_q})_{\mathcal{H}_l}^{\wedge})^G \simeq K(G, E)_{\mathcal{H}_l}^{\wedge} \cong KF.$$

We now argue that this is a reasonable replacement for Conjecture 7.9 in this case. We note that the fixed point category of  $V(G, \overline{\mathbb{F}_q})$  is the category  $V^G(\overline{\mathbb{F}_q})$  of linear descent data. We have the straightforward observation

**Proposition 8.2** *For any finite group  $G$  acting on  $\mathbb{F}_q$  by automorphisms, the category of linear descent data  $V^G(\mathbb{F}_q)$  is equivalent to the category of left modules over the twisted group ring  $\mathbb{F}_q\langle G \rangle$ . Similarly, for a profinite group  $G$  acting on  $\mathbb{F}_q$ , we find that the category of linear descent data is equivalent to the category of continuous modules over the twisted group ring  $\mathbb{F}_q\langle G \rangle$ .*

Note that for finite  $G$ ,  $\mathbb{F}_q\langle G \rangle$  is a finite dimensional semisimple algebra over  $\mathbb{F}_p$ , so it is a product of matrix rings over field extensions of  $\mathbb{F}_p$ . Consequently, we should view its  $K$ -theory as essentially understood, given Quillen's computations for finite fields. For this reason, we regard the above conjecture as a satisfactory replacement for Conjecture 7.9.

In the characteristic 0 case, though, we have more difficulties, since in this case we do not know the  $K$ -theory of the prime field  $\mathbb{Q}$ . Indeed, we would like to make conjectures about the  $K$ -theory of  $\mathbb{Q}$  involving the representation theory of  $G_{\mathbb{Q}}$ . We do, however, thanks to the work of Suslin [42], understand the  $K$ -theory of the field  $\mathbb{Q}_p$  and  $\mathbb{Z}_p$  where  $p \neq l$ .

**Theorem 8.3 (Suslin; see [42])** *Let  $K$  denote any finite extension of  $\mathbb{Q}_p$ , and let  $\mathcal{O}_K$  denote its ring of integers. Let  $(\pi)$  denote its unique maximal ideal. The quotient homomorphism  $\mathcal{O}_K \rightarrow \mathcal{O}_K/\pi\mathcal{O}_K$  induces an isomorphism on  $l$ -completed  $K$ -theory.*

Now, suppose we have any field  $F$  containing  $\mathbb{Q}_p$  and the  $l$ -th roots of unity, and let  $E$  denote its algebraic closure. Let  $L$  be  $\bigcup_n \mathbb{Q}_p(\zeta_{l^n}) \subseteq E$ , so  $L \subseteq F$ .  $L$  is of course closed under the action of the absolute Galois group  $G = G_F$ . By abuse of notation, we will write  $V(G, \mathcal{O}_L)$  for the category of twisted  $G - \mathcal{O}_L$ -modules over  $\mathcal{O}_L$ , i.e. finitely generated  $\mathcal{O}_L$ -modules  $M$  equipped with a  $G$ -action so that  $g(rm) = r^g g(m)$  for all  $r \in \mathcal{O}_L$ ,  $m \in M$ , and  $g \in G$ . The following is an easy consequence of Theorem 8.3 above.

**Proposition 8.4** *The functor  $V(G, \mathcal{O}_L) \rightarrow V(G, \mathcal{O}_L/\pi\mathcal{O}_L)$  induces an equivalence  $K^G \mathcal{O}_L \rightarrow K^G \mathcal{O}_L/\pi\mathcal{O}_L$ .*

Since  $\mathcal{O}_L/\pi\mathcal{O}_L$  is a semisimple algebra over a finite field, we will regard it as an understood quantity. This means that in this case, we also have a version of the representational assembly via the diagram

$$(K^G \mathcal{O}_L/\pi\mathcal{O}_L)_{\mathcal{H}_l}^{\wedge} \xleftarrow{\sim} (K^G \mathcal{O}_L)_{\mathcal{H}_l}^{\wedge} \xrightarrow{\mathcal{O}_L^{\otimes E}} (K^G E)_{\mathcal{H}_l}^{\wedge} \cong KF$$

This kind of result extends to the case where  $F$  contains a Henselian local ring whose residue class field is algebraic over a finite field.

## 9 The ascent map and assembly for the case $\mu_{l^\infty} \subseteq F$

The constructions of the Section 7 provide explicit maps from derived completions of spectra attached to representation categories of the absolute Galois

group  $G$  of a field  $F$  to  $KF$ . However, they only apply to the geometric case, i.e. where  $F$  contains an algebraically closed subfield. We believe, though, that the statement relating the  $K$ -theory of the field to the derived representation theory of the absolute Galois group should be true for all fields containing the  $l$ -th power roots of unity. The purpose of this section is to develop a criterion which when satisfied will produce an equivalence between these spectra. The criterion will involve a map which we call the *ascent* map. The terminology “ascent” refers to the fact that the method gives a description of  $KE$  coming from information about  $KF$ , rather than describing  $KF$  in terms of  $KE$ , as is done in the descent. In the interest of clarity, we will first describe the approach as it works in the abelian case, where it is not necessary to pass to the generality of Mackey and Green functors, and then indicate the changes necessary when we deal with the more general situation. Finally, we suppose that  $F$  contains all the  $l$ -th power roots of unity, and as usual we let  $E$  denote the algebraic closure of  $F$  and  $G$  denote the absolute Galois group of  $F$ .

Consider the forgetful functor  $V^G(E) \xrightarrow{\phi} V(G, E) \cong Vect(E)$ , which simply forgets the  $G$ -action. It induces a map

$$KF \cong K^G(E) \xrightarrow{K\phi} K(G, E) \cong KE.$$

We now have a commutative diagram

$$\begin{array}{ccc} KRep_F[G] \wedge K^G(E) & \longrightarrow & K^G(E) \\ \varepsilon \wedge id \downarrow & & \downarrow K\phi \\ KF \wedge K^G(E) & \longrightarrow & KE \\ id \wedge K\phi \downarrow & & \downarrow id \\ KF \wedge KE & \longrightarrow & KE \end{array}$$

of spectra, where the horizontal maps are multiplication maps defining algebra structures, and where the vertical maps are built out of the augmentation map  $\varepsilon$  and the forgetful functor  $\phi$ . Note that the middle horizontal map exists because  $\varepsilon$  is induced by the forgetful functor  $Rep_F[G] \rightarrow Vect(F)$  which forgets the  $G$  action. Considering the four corners of this diagram, we obtain a map of spectra

$$asc_F : KF \underset{KRep_F[G]}{\wedge} K^G(E) \longrightarrow K(G, E) \cong KE.$$

It is readily verified that  $asc_F$  is a homomorphism of commutative  $S$ -algebras. We note further that both sides of this map are  $KF$ -algebras, and that  $asc_F$  is

a homomorphism of  $KF$ -algebras. As usual, let  $\mathbb{H}$  denote the mod- $l$  Eilenberg-MacLane spectrum, so we have an obvious morphism  $KF \rightarrow \mathbb{H}$  of  $S$ -algebras. We will now explore the consequences of the hypothesis that the homomorphism

$$id_{\mathbb{H}} \wedge_{KF} KF \wedge_{KRep_F[G]} K^G(E) = \mathbb{H} \wedge_{KRep_F[G]} K^G(E) \rightarrow \mathbb{H} \wedge_{KF} KE$$

is a weak equivalence.

For any homomorphism of  $S$ -algebras  $f : D \rightarrow E$ , we let  $T(D \rightarrow E)$  denote the cosimplicial  $S$ -algebra given by

$$T^k(D \rightarrow E) = \underbrace{E \wedge_D E \wedge_D \cdots \wedge_D E}_{k+1 \text{ factors}}$$

so  $Tot(T(D \rightarrow E)) \cong D_E^\wedge$ . Now suppose that  $D$  and  $E$  are both  $k$ -algebras, where  $k$  is a commutative  $S$ -algebra, and that we are given a homomorphism  $k \rightarrow \mathbb{H}$ . We can now construct a bisimplicial spectrum  $\Sigma^\cdot = \Sigma^\cdot(D \rightarrow E, k \rightarrow \mathbb{H})$  by setting

$$\Sigma^{kl} = \underbrace{\mathbb{H} \wedge_k \mathbb{H} \wedge_k \cdots \wedge_k \mathbb{H} \wedge_k T^l(D \rightarrow E)}_{k+1 \text{ factors}}.$$

**Proposition 9.1** *Suppose that  $k$ ,  $D$ , and  $E$  are all  $(-1)$ -connected, and that  $\pi_0 k \cong \pi_0 E \cong \mathbb{Z}$ . Then  $Tot(\Sigma^\cdot) \cong D_{\mathbb{H} \wedge_k E}^\wedge$ , where  $\varepsilon$  is the composite  $D \rightarrow E \rightarrow \mathbb{H} \wedge_k E$ . If in addition,  $\pi_0 D \cong \mathbb{Z}$ , then  $Tot(\Sigma^\cdot)$  is equivalent to the  $l$ -adic completion of  $D$ .*

**Proof:** The diagonal bisimplicial spectrum is canonically equivalent to the cosimplicial spectrum  $T(D \rightarrow E \wedge_k \mathbb{H})$ . By use of the Künneth spectral sequence, it is clear that  $E \wedge_k \mathbb{H}$  is  $(-1)$  connected and that  $\pi_0 E \wedge_k \mathbb{H} \cong \mathbb{F}_l$ . The first result now follows from Theorem 6.1 of [8]. The second follows immediately from Theorem 3.2, item 4 of [8].  $\square$

Whenever we have a commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ A' & \longrightarrow & B' \end{array}$$

of  $k$ -algebras, we have an induced map

$$\Sigma''(A \rightarrow B) \rightarrow \Sigma''(A' \rightarrow B').$$

Now suppose we have two  $k$ -algebra homomorphisms  $A \rightarrow B$  and  $A \rightarrow C$ , so that each of the composites  $k \rightarrow A \rightarrow B$  and  $k \rightarrow A \rightarrow C$  are weak equivalences. We then obtain a commutative diagram of commutative  $k$ -algebras

$$\begin{array}{ccc} B & \longrightarrow & B \underset{A}{\wedge} C \\ \downarrow & & \downarrow \\ B \underset{A}{\wedge} C \underset{k}{\wedge} A & \longrightarrow & B \underset{A}{\wedge} C. \end{array}$$

The upper horizontal arrow and the left hand vertical arrow are obvious inclusions, the right hand arrow is the identity, and the lower horizontal arrow is the multiplication map obtained by regarding  $B \underset{A}{\wedge} C$  as a right module over the  $k$ -algebra  $A$ . We consequently obtain a map of bicosimplicial  $k$ -algebras

$$\Sigma''(B \rightarrow B \underset{A}{\wedge} C) \longrightarrow \Sigma''(B \underset{A}{\wedge} C \underset{k}{\wedge} A \rightarrow B \underset{A}{\wedge} C).$$

**Proposition 9.2** *Under the hypotheses above, i.e. that  $k \rightarrow B$  and  $k \rightarrow C$  are weak equivalences, this map of bicosimplicial  $k$ -algebras is a levelwise weak equivalence, and hence induces an equivalence of total spectra.*

**Proof:** There is a commutative diagram

$$\begin{array}{ccc} B \underset{A}{\wedge} T(A \rightarrow C) & \xrightarrow{\sim} & T(B \rightarrow B \underset{A}{\wedge} C) \\ \downarrow & & \downarrow \\ B \underset{A}{\wedge} T(C \underset{k}{\wedge} A \rightarrow C) & \xrightarrow{\sim} & T(B \underset{A}{\wedge} C \underset{k}{\wedge} A \rightarrow B \underset{A}{\wedge} C) \end{array}$$

where the horizontal arrows are levelwise weak equivalences. This follows from the standard isomorphism

$$B \underset{A}{\wedge} \underbrace{(C \underset{A}{\wedge} C \underset{A}{\wedge} \cdots \underset{A}{\wedge} C)}_{k \text{ factors}} \rightarrow \underbrace{(B \underset{A}{\wedge} C) \underset{B}{\wedge} (B \underset{A}{\wedge} C) \underset{B}{\wedge} \cdots \underset{B}{\wedge} (B \underset{A}{\wedge} C)}_{k \text{ factors}}.$$

It now follows that the map  $T(A \rightarrow C) \rightarrow T(C \underset{k}{\wedge} A \rightarrow C)$  is a levelwise weak equivalence, since the composite

$$A \cong k \underset{k}{\wedge} A \longrightarrow C \underset{k}{\wedge} A$$

is a weak equivalence. Applying the functors

$$\underbrace{\mathbb{H} \underset{k}{\wedge} \mathbb{H} \underset{k}{\wedge} \cdots \underset{k}{\wedge} \mathbb{H} \underset{k}{\wedge} -}_{k \text{ factors}}$$

to this equivalence, we obtain the result for  $\Sigma''$ .  $\square$

We now examine the consequences of this result in our case. The spectra will be  $k = KF$ ,  $A = KRep_F[G]$ ,  $B = K^G(E)$ , and  $C = KF$ , viewed as an  $A$ -algebra via the augmentation which forgets the action. This data clearly satisfies the hypotheses above. If  $id_{\mathbb{H}} \underset{KF}{\wedge} asc_F$  is a weak equivalence, then the arrow  $B \rightarrow B \underset{A}{\wedge} C$  can be identified with the arrow  $K^G(E) \rightarrow K(G, E)$ , which in turn can be identified canonically with the arrow  $KF \rightarrow KE$ . It follows that

$$Tot(\Sigma''(B \rightarrow B \underset{A}{\wedge} C)) \cong Tot(\Sigma''(KF \rightarrow KE)) \cong KF_l^\wedge$$

On the other hand,

$$\begin{aligned} & \Sigma''(B \underset{A}{\wedge} C \underset{k}{\wedge} A \rightarrow B \underset{A}{\wedge} C) \cong \\ & \Sigma''(KF \underset{KRep_F[G]}{\wedge} K^G(E) \underset{KF}{\wedge} KRep_F[G] \rightarrow KF \underset{KRep_F[G]}{\wedge} K^G(E)) \end{aligned}$$

and  $asc_F$  induces a map of bicosimplicial spectra

$$\begin{aligned} & \Sigma''(KF \underset{KRep_F[G]}{\wedge} K^G(E) \underset{KF}{\wedge} KRep_F[G] \rightarrow KF \underset{KRep_F[G]}{\wedge} K^G(E)) \longrightarrow \\ & \Sigma''(KE \underset{KF}{\wedge} KRep_F[G] \rightarrow KE). \end{aligned}$$

By Proposition 3.2, item 2 of [8], if  $id_{\mathbb{H}} \underset{KF}{\wedge} asc_F$  is a weak equivalence, this map of bicosimplicial spectra is a weak equivalence. The natural homomorphism  $KE \underset{KF}{\wedge} KRep_F[G] \rightarrow KRep_E[G]$  now induces another map of bicosimplicial spectra

$$\Sigma''(KE \underset{KF}{\wedge} KRep_F[G] \rightarrow KE) \longrightarrow \Sigma''(KRep_E[G] \rightarrow KE).$$

We now have

**Proposition 9.3** *When all  $l$ -th power roots of unity are in  $F$ , the evident map  $KRep_F[G] \underset{KF}{\wedge} KE \rightarrow KRep_E[G]$  is a weak equivalence.*



**Proof:** It is a standard fact in representation theory that for any field  $F$  which contains the  $l$ -th power roots of unity, the isomorphism classes of representations of any finite  $l$ -group are in bijective correspondence with the isomorphism classes of representations of the same group in  $\mathbb{C}$ . Moreover, the endomorphism ring of any irreducible representation is a copy of  $F$ . We therefore have isomorphisms

$$\pi_* KRep_F[G] \cong K_*F \otimes R[G]$$

and

$$\pi_* KRep_E[G] \cong K_*E \otimes R[G].$$

where  $R[G]$  denotes the complex representation ring. The  $E_2$ -term of the Künneth spectral sequence for  $\pi_* KRep_F[G] \wedge_{KF} KE$  now has the form

$$Tor_{K_*F}(K_*F \otimes R[G], K_*E) \cong K_*Rep_E[G]$$

which gives the result.  $\square$

We now have the following conclusion.

**Theorem 9.4** *Suppose that  $F$  contains all the  $l$ -th power roots of unity. Suppose further that  $id_{\mathbb{H}} \wedge_{KF} asc_F$  is a weak equivalence. Then we have a weak equivalence*

$$KF_l^\wedge \cong KRep_E[G]_{KE}^\wedge.$$

We observed in Section 7 that we did not expect to have an equivalence

$$KF \cong KRep_k[G]_{\mathbb{H}}^\wedge$$

as stated except in the abelian case, but that there is an analogous “fully equivariant version” involving equivariant spectra and Green functors which we expect to hold in all cases. This suggests that we should not expect the ascent map as formulated above to hold except in the abelian case. However, it is possible to modify the ascent map to extend it to the fully equivariant version of the completion, and we indicate how.

We fix the group  $G = G_F$ , and we define various equivariant  $S$ -algebras. We let  $\underline{KV}$  denote the equivariant spectrum obtained using the category of descent data described above. For any subgroup  $H \subseteq G$ , the fixed point spectrum  $\underline{KV}^H$  is equivalent to the  $K$ -theory spectrum  $K(E^H)$ . For any field  $F$ , we let  $\underline{KRep}_F$  denote the equivariant spectrum corresponding to the trivial action of  $\overline{G}$  on  $F$ , so for any  $H \subseteq G$ , we have that  $\underline{KRep}_F^H \cong KRep_F[H]$ . Also for any field  $F$ , we will write  $\underline{KF}$  for the equivariant spectrum  $KF \wedge S^0$ , where  $S^0$  denotes the  $G$ -equivariant sphere spectrum. We have that  $\underline{KF}^H \cong KF \wedge G/H_+$  for any subgroup of finite index  $H \subseteq G$ . Finally, we let  $\underline{\mathcal{H}}$  denote the  $G$ -spectrum

associated to the Green functor  $\mathcal{Z}/l\mathcal{Z}$  described above in section 7. We note that there is an obvious homomorphism of  $G$ -equivariant  $S$ -algebras  $\underline{KF} \rightarrow \underline{\mathcal{H}}_l$ . We can easily verify that there is a natural analogue of the ascent map in this equivariant setting, i.e. a map of  $G$ -equivariant  $S$ -algebras  $asc_F : \underline{KF} \wedge_{\underline{KRep}_F} \underline{KV} \rightarrow \underline{KE}$ . We now have the following equivariant generalization of 9.4, whose proof is identical to the nonequivariant version given above.

**Theorem 9.5** *Let  $F, E$ , and  $G$  be as above. Suppose further that the map of equivariant spectra*

$$id_{\mathcal{H}_l} \wedge_{\underline{KF}} asc_F : \mathcal{H}_l \wedge_{\underline{KRep}_F} \underline{KV} \rightarrow \mathcal{H}_l \wedge_{\underline{KF}} \underline{KE}$$

*is a weak equivalence of spectra. Then there is a canonically defined equivalence of  $G$ -equivariant  $S$ -algebras*

$$\underline{KV}_{\mathcal{H}_l} \wedge \rightarrow \underline{KRep}_{E_{\mathcal{H}_l}} \wedge.$$

*Note that the fixed point sets of the equivariant spectrum  $\underline{KV}_{\mathcal{H}_l} \wedge$  are just the  $l$ -adic completions of the  $K$ -theory spectra of the corresponding subfields.*

**Remark:** We expect that the equivariant map  $id_{\mathcal{H}_l} \wedge_{\underline{KF}} asc_F$  will be an equivalence for all fields containing all the  $l$ -th power roots of unity.

**Remark:** The above discussion could be a reflection of an interesting convergence theorem within  $\mathbb{A}^1$  homotopy theory (see [30] or [31] for treatments of this construction). We suppose that  $\mathbb{A}^1$  homotopy theory has been extended to an equivariant theory for the action of profinite groups. We further suppose that algebraic  $K$ -theory has been extended to a fully equivariant cohomology theory  $K^G$  on this equivariant category, where by “fully equivariant” we mean a theory which admits deloopings with respect to arbitrary continuous linear representations of a fixed profinite group  $G$ . Let  $F$  be a field,  $E$  its separable closure, and suppose  $G = G_F$  is its separable Galois group. We let the group  $\Gamma = G \times G$  act on  $E$  by projection on the first factor. We let  $G_0 = \{e\} \times G \subseteq \Gamma$ , and let  $\Delta$  denote the diagonal subgroup in  $\Gamma$ . We consider the square

$$\begin{array}{ccc} Spec(E) \times \Gamma & \longrightarrow & Spec(E) \times \Gamma/\Delta \\ \downarrow & & \downarrow \\ Spec(E) \times \Gamma/G_0 & \longrightarrow & Spec(E) \end{array}$$

of pro-schemes, where the sets  $\Gamma$ ,  $\Gamma/\Delta$ , and  $\Gamma/G_0$  are all regarded as discrete pro-schemes. If this square is a homotopy pullback square in the equivariant  $\mathbb{A}^1$  category, and if the Eilenberg-Moore spectral sequence associated to the equivariant cohomology theory  $K^\Gamma$  converges, then this would confirm the hypotheses of the theorems above.

## 10 Derived representation theory and deformation $K$ -theory

So far, our understanding of the derived completion of representation rings is limited to knowledge of spectral sequences for computing them, given information about  $Tor$  or  $Ext$  functors of these rings. In this (entirely speculative) section, we want to suggest that there should be a relationship between derived representation theory and deformations of representations.

We consider first the representation theory of finite  $l$ -groups  $G$ . In this case, the derived representation ring of  $G$  is just the  $l$ -adic completion of  $R[G]$ , as can be verified using the results of Atiyah [1]. In particular, the derived representation ring has no higher homotopy. However, when we pass to a profinite  $l$ -group, such as  $\mathbb{Z}_l$ , we find that the derived representation ring does have higher homotopy, a single copy of  $\mathbb{Z}_l$  in dimension 1. This situation appears to be parallel to the following situation. Consider the infinite discrete group  $\Gamma = \mathbb{Z}$ . We may consider the category  $Rep_{\mathbb{C}}[\Gamma]$  of finite dimensional representations of  $\Gamma$ , and its  $K$ -theory spectrum  $KRep_{\mathbb{C}}[\Gamma]$ . The homotopy of this spectrum is given by

$$\pi_* KRep_{\mathbb{C}}[\Gamma] \cong \mathbb{Z}[S^1] \otimes K_*\mathbb{C}$$

where  $S^1$  is the circle regarded as a discrete group. This isomorphism arises from the existence of Jordan normal form, which (suitably interpreted) shows that every representation of  $\Gamma$  admits a filtration by subrepresentations so that the subquotients are one-dimensional, and are therefore given by multiplication by a uniquely defined non-zero complex number. This construction does not take into account the topology of  $\mathbb{C}$  at all. We might take the topology into account as in the following definition.

**Definition 10.1** *Consider any discrete group  $\Gamma$ . For each  $k$ , we consider the category  $Rep_{\mathbb{C}}^k[\Gamma]$  whose objects are all possible continuous actions of  $\Gamma$  on  $\Delta[k] \times V$  which preserve the projection  $\pi : \Delta[k] \times V \rightarrow \Delta[k]$ , and which are linear on each fiber of  $\pi$ . It is clear that the categories  $Rep_{\mathbb{C}}^k[\Gamma]$  fit together into a simplicial symmetric monoidal category, and we define the **deformation  $K$ -theory spectrum of  $\Gamma$** ,  $K^{def}[\Gamma]$ , as the total spectrum of the simplicial spectrum*

$$k \rightarrow KRep_{\mathbb{C}}^k[\Gamma].$$

**Remark:** The terminology is justified by the observation that, for example, an object in the category of 1-simplices  $Rep_{\mathbb{C}}^1[\Gamma]$  is exactly a path in the space of representations of  $\Gamma$  in  $GL(V)$ , or a deformation of the representation at  $0 \times V$ .

One can easily check that  $\pi_0 K^{def}[\mathbb{Z}] \cong \mathbb{Z}$ , with the isomorphism given by sending a representation to its dimension. This follows from the fact that any two representations of the same dimension of  $\mathbb{Z}$  can be connected by a deformation. In fact, one can apply the functor  $\pi_0$  levelwise to the simplicial

spectrum  $KRep_{\mathbb{C}}[\mathbb{Z}]$ , and attempt to compute the homotopy groups of the simplicial abelian group  $k \rightarrow \pi_0 KRep_{\mathbb{C}}^k[\mathbb{Z}]$ . As above, it is easy to see that  $\pi_0$  of this simplicial abelian group is zero, and it appears likely that

$$\pi_*(\pi_0 KRep_{\mathbb{C}}[\mathbb{Z}]) \cong \pi_*\mathbb{Z}[S^1] \cong H_*(S^1)$$

where  $S^1$  is the circle regarded as a topological group. Note that  $S^1$  is the character group of the original discrete group  $\mathbb{Z}$ .

Now consider the profinite group  $\mathbb{Z}_l$ . We may define the deformation  $K$ -theory of this group as the direct limit of its finite quotients, and in view of the rigidity of complex representations of finite groups we have that

$$KRep_{\mathbb{C}}[\mathbb{Z}_l] \cong K^{def}[\mathbb{Z}_l]$$

We have an inclusion  $\mathbb{Z} \hookrightarrow \mathbb{Z}_l$ , and therefore a map of spectra  $K^{def}[\mathbb{Z}_l] \rightarrow K^{def}[\mathbb{Z}]$ . We now obtain a composite map of derived completions

$$KRep_{\mathbb{C}}[\mathbb{Z}_l]_{\mathbb{H}}^{\wedge} \simeq K^{def}[\mathbb{Z}_l]_{\mathbb{H}}^{\wedge} \longrightarrow K^{def}[\mathbb{Z}]_{\mathbb{H}}^{\wedge}$$

where  $\mathbb{H}$  denotes the mod- $l$  Eilenberg-MacLane spectrum, regarded as a module over the deformation  $K$ -theory via the map which takes any representation to its dimension mod  $l$ . Since the homotopy groups on the two sides appear isomorphic, it appears likely that this composite is a homotopy equivalence. We now observe that for any discrete group  $\Gamma$ , we may consider its pro- $l$  completion  $\Gamma_l^{\wedge}$ , and obtain a map  $KRep_{\mathbb{C}}[\Gamma_l^{\wedge}] \rightarrow K^{def}[\Gamma]$ . This map induces a map  $j$  of derived completions, and we ask the following question.

**Question:** For what discrete groups does the map  $j$  induce a weak equivalence of spectra after derived completion at the augmentation map  $\varepsilon : K^{def}[\Gamma] \rightarrow K\mathbb{C}$ ? Note that the completion may have to be taken in the category of  $\Gamma_l^{\wedge}$ -equivariant spectra.

We can further ask if the effect of derived completion can be computed directly on the pro- $l$  group, rather than by permitting deformations on a discrete subgroup. This cannot be achieved over  $\mathbb{C}$  since representations of finite groups are rigid, but perhaps over some other algebraically closed field.

**Question:** Can one construct a deformation  $K$ -theory of representations of a pro- $l$  group  $G$ , using an algebraic deformation theory like the one discussed in [29], so that it coincides with the derived completion of  $KRep_{\mathbb{C}}[G]$ ?

We conclude by pointing out an analogy between our theory of the representational assembly and other well known constructions. Consider a  $K(\Gamma, 1)$  manifold  $X$ . The category  $VB(X)$  of complex vector bundles over  $X$  is a symmetric monoidal category, and we can construct its  $K$ -theory. On the other hand, we have a functor from the category of complex representations of  $\Gamma$  to  $VB(X)$ , given by

$$\rho \rightarrow (\tilde{X} \times_{\Gamma} \rho \rightarrow X)$$

where  $\tilde{X}$  denotes the universal covering space of  $X$ . This functor produces a map of  $K$ -theory spectra. It is also easy to define a “deformation version” of  $KVB(X)$ , and one obtains a map

$$K^{def}[\Gamma] \rightarrow K^{def}VB(X).$$

This map should be viewed as the analogue in this setting for the representational assembly we discussed in the case of  $K$ -theory of fields, using the point of view that fields are analogues of  $K(G, 1)$  manifolds in the algebraic geometric context.

## References

- [1] M.F. Atiyah, *Characters and cohomology of finite groups*, Inst. Hautes Etudes Scient. Publ. **9** (1961), 23-64.
- [2] S. Bloch and S. Lichtenbaum, *A spectral sequence for motivic cohomology*, preprint, available on the  $K$ -theory server at U. of Illinois, Urbana Champaigne.
- [3] S. Bouc, *Green functors and  $G$ -sets* Lecture Notes in Mathematics, **1671** Springer-Verlag, Berlin, 1997.
- [4] A.K. Bousfield and D.M. Kan, **Homotopy limits, completions and localizations**. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972. v+348 pp.
- [5] G. Carlsson, *Equivariant stable homotopy theory and Segal’s Burnside ring conjecture*, Annals of Mathematics **120** (1984) 189-224.
- [6] G. Carlsson, *Equivariant stable homotopy and Sullivan’s conjecture*, Inventiones Mathematicae **103** (1991) 497-525.
- [7] G. Carlsson, *A survey of equivariant stable homotopy theory*, Topology 31 (1992), no. 1, 1-27.
- [8] G. Carlsson, *Derived completions in stable homotopy theory*, Journal of Pure and Applied Algebra, Volume 212, Issue 3, March 2008, Pages 550-577
- [9] F. DeMeyer and E. Ingraham, **Separable Algebras over Commutative Rings**, Lecture Notes in Mathematics, Vol. 181 Springer-Verlag, Berlin-New York 1971 iv+157 pp.
- [10] W. Dwyer, E. Friedlander, V.P. Snaith, and R.W. Thomason, *Algebraic  $K$ -theory eventually surjects onto topological  $K$ -theory*, Invent. Math. 66 (1982), no. 3, 481-491.

- [11] A. D. Elmendorf and M. A. Mandell, *Rings, modules, and algebras in infinite loop space theory*, Adv. Math. **205** (2006), no. 1, 163–228.
- [12] A.D. Elmendorf; I. Kriz; M.A. Mandell; J.P. May, *Modern foundations for stable homotopy theory*. Handbook of algebraic topology, 213–253, North-Holland, Amsterdam, 1995.
- [13] A. D. Elmendorf, I. Kriz, M.A. Mandell, and J.P. May, **Rings, modules, and algebras in stable homotopy theory. With an appendix by M. Cole**. Mathematical Surveys and Monographs, 47. American Mathematical Society, Providence, RI, 1997.
- [14] O. Gabber, *K-theory of Henselian local rings and Henselian pairs* Algebraic K-theory, commutative algebra, and algebraic geometry (Santa Margherita Ligure, 1989), 59–70, Contemp. Math., 126, Amer. Math. Soc., Providence, RI, 1992.
- [15] T. Geisser and M. Levine, *The Bloch-Kato conjecture and a theorem of Suslin-Voevodsky*. J. Reine Angew. Math. 530 (2001), 55–103.
- [16] J. P. C. Greenlees and J. P. May, Equivariant stable homotopy theory. Handbook of algebraic topology, 277–323, North-Holland, Amsterdam, 1995.
- [17] A. Grothendieck, et al, with M. Artin and J.-L. Verdier, *Theorie des topos et cohomologie etale des schemas*, Lecture Notes in Mathematics **269**, **270**, **305**, Springer Verlag, 1972-3.
- [18] L. Hesselholt and I. Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, Topology **36** (1997), no. 1, 29–101.
- [19] M. Hovey, B. Shipley, and J. Smith, *Symmetric spectra* J. Amer. Math. Soc. 13 (2000), no. 1, 149–208.
- [20] T. Lawson, Completed representation ring spectra of nilpotent groups. Algebr. Geom. Topol. **6** (2006), 253–286
- [21] G. Lewis, J.P. May, and J. McClure, *Ordinary  $RO(G)$ -graded cohomology* Bull. Amer. Math. Soc. (N.S.) **4** (1981), no. 2, 208–212.
- [22] S. Lichtenbaum, *On the values of zeta and L-functions I* Ann. of Math. (2) **96** (1972), 338–360.
- [23] G. Lyo, *Semilinear actions of Galois groups and the algebraic K-theory of fields*, Thesis, University of California at Berkeley, 2007.
- [24] M.A. Mandell and J. P. May, Equivariant orthogonal spectra and S-modules. Mem. Amer. Math. Soc. 159 (2002), no. 755
- [25] J.P. May, **Simplicial Objects in Algebraic Topology** Van Nostrand 1967.

- [26] J.P. May, *Geometry of iterated loop spaces*, Lecture Notes in Mathematics **271**, Springer Verlag, 1972
- [27] J.P. May, *Multiplicative infinite loop space theory*. J. Pure Appl. Algebra 26 (1982), no. 1, 1–69.
- [28] J. P. May, Equivariant and nonequivariant module spectra. J. Pure Appl. Algebra 127 (1998), no. 1, 83–97.
- [29] B. Mazur, *An introduction to the deformation theory of Galois representations*. Modular forms and Fermat’s last theorem (Boston, MA, 1995), 243–311, Springer, New York, 1997.
- [30] F. Morel, *An introduction to  $\mathbb{A}^1$ -homotopy theory*, Contemporary developments in algebraic  $K$ -theory, 357–441 (electronic), ICTP Lect. Notes, XV, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [31] F. Morel and V. Voevodsky,  *$\mathbb{A}^1$ -homotopy theory of schemes*, Inst. Hautes tudes Sci. Publ. Math. No. 90 (1999), 45–143 (2001).
- [32] D.G. Quillen *Higher algebraic K-theory I* Lecture Notes in Mathematics **341**, Springer Verlag 1973, 85-147
- [33] D.G. Quillen, *On the cohomology and K-theory of the general linear groups over a finite field*, Annals of Mathematics **96** (1972), 552-586.
- [34] D.G. Quillen, *Higher algebraic K-theory*, Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974), Vol. 1, pp. 171–176. Canad. Math. Congress, Montreal, Que., 1975.
- [35] D.G. Quillen, *Finite generation of the groups  $K_i$  of rings of algebraic integers*. Algebraic  $K$ -theory, I: Higher  $K$ -theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), pp. 179–198. Lecture Notes in Math., Vol. 341, Springer, Berlin, 1973.
- [36] J. Rognes, and C. Weibel, *Two-primary algebraic K-theory of rings of integers in number fields* Appendix A by Manfred Kolster. J. Amer. Math. Soc. **13** (2000), no. 1, 1–54.
- [37] G.B. Segal, *Configuration spaces and iterated loop spaces*, Inventiones Mathematicae **21** (1973) 213-221.
- [38] G.B. Segal, *Equivariant stable homotopy theory* Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 2, pp. 59–63. Gauthier-Villars, Paris, 1971.
- [39] J.P. Serre, **Linear representations of finite groups**. Translated from the second French edition by Leonard L. Scott. Graduate Texts in Mathematics, Vol. 42. Springer-Verlag, New York-Heidelberg, 1977.

- [40] K. Shimakawa, *Infinite loop  $G$ -spaces associated to monoidal categories*, Publ. Res. Inst. Math. Sci **25**, (no.2), 239-262 (1989)
- [41] A.A. Suslin, *On the  $K$ -theory of algebraically closed fields*, Inventiones Mathematicae **73**, 1983, 241-245.
- [42] A.A. Suslin, *On the  $K$ -theory of local fields* Proceedings of the Luminy conference on algebraic  $K$ -theory (Luminy, 1983). J. Pure Appl. Algebra 34 (1984), no. 2-3, 301–318.
- [43] A. Suslin, Andrei and V. Voevodsky, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*. The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), 117–189, NATO Sci. Ser. C Math. Phys. Sci., 548, Kluwer Acad. Publ., Dordrecht, 2000.
- [44] R.G. Swan,  *$K$ -theory of finite groups and orders*, Lecture Notes in Mathematics **76**, Springer Verlag, 1968.
- [45] R. W. Thomason, *Algebraic  $K$ -theory and tale cohomology* Ann. Sci. cole Norm. Sup. (4) **18** (1985), no. 3, 437–552.
- [46] V. Voevodsky, A. Suslin, and E.M. Friedlander, **Cycles, transfers, and motivic homology theories** Annals of Mathematics Studies, **143** Princeton University Press, Princeton, NJ, 2000.
- [47] V. Voevodsky, *The Milnor conjecture*, preprint, available at the  $K$ -theory server, Univ. of Illinois, Urbana-Champaign.
- [48] V. Voevodsky, *Voevodsky's Seattle lectures:  $K$ -theory and motivic cohomology*. Algebraic  $K$ -theory (Seattle, WA, 1997), 283–303, Proc. Sympos. Pure Math., 67, Amer. Math. Soc., Providence, RI, 1999.
- [49] H. Völklein, **Groups as Galois groups. An introduction**. Cambridge Studies in Advanced Mathematics, 53. Cambridge University Press, Cambridge, 1996