

Experimental Evidence for Quantum Structure in Cognition

Diederik Aerts

*Center Leo Apostel for Interdisciplinary Studies
Department of Mathematics and Department of Psychology
Vrije Universiteit Brussel, 1160 Brussels, Belgium
E-Mail: diraerts@vub.ac.be*

Sven Aerts

*Center Leo Apostel for Interdisciplinary Studies
Department of Mathematics
Vrije Universiteit Brussel, 1160 Brussels, Belgium
E-Mail: saerts@vub.ac.be*

Liane Gabora

*Psychology and Computer Science
University of British Columbia
Kelowna, British Columbia, Canada
E-Mail: liane.gabora@ubc.ca*

Abstract

We prove a theorem that shows that a collection of experimental data of membership weights of items with respect to a pair of concepts and its conjunction cannot be modeled within a classical measure theoretic weight structure in case the experimental data contain the effect called overextension. Since the effect of overextension, analogue to the well-known guppy effect for concept combinations, is abundant in all experiments testing weights of items with respect to pairs of concepts and their conjunctions, our theorem constitutes a no-go theorem for classical measure structure for common data of membership weights of items with respect to concepts and their combinations. We put forward a simple geometric criterion that reveals the non classicality of the membership weight structure and use experimentally measured membership weights estimated by subjects in experiments from [29] to illustrate our geometrical criterion. The violation of the classical weight structure is similar to the violation of the well-known Bell inequalities studied in quantum mechanics, and hence suggests that the quantum formalism and hence the modeling by quantum membership weights, as for example in [18], can accomplish what classical membership weights cannot do.

1 Introduction

Many branches of mathematics, such as geometry, complexity theory, and even number theory, were originally conceived not as domains of mathematics, but as describing a particular domain of physical reality. It was only much later that they were conceived more abstractly, and their applicability to a wide range of phenomena was realized. We believe this is also proving to be the case for the

mathematical formalisms originally developed to describe events observed in the microworld: quantum mechanics.

Meanwhile the mathematical formalism of quantum mechanics has indeed been used successfully to model situations pertaining to domains different from the micro-world, for example, in economics [1, 2, 3], operations research and management sciences [4, 5], psychology and cognition [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18], game theory [19, 20], and language and artificial intelligence [21, 22, 23, 24, 25, 26].

More specifically, in [18] a quantum mechanical representation for experimental data corresponding to membership weights of items with respect to pairs of concepts and their conjunctions was elaborated. It was also proven that these data cannot be modelled by a classical theory of membership weights, i.e. a theory where membership weights are represented within a measure theoretic structure (see theorems 3 and 6).

In the present paper we introduce a very simple geometrical criterion that allows the identification of the classical or non classical quantum-like nature of membership weight data gathered on pairs of concepts and their conjunctions, or more generally on collections of concepts and conjunctions of some of the pairs in these collections. More specifically we determine for such a collection of concepts and some of the conjunctions of these concepts a geometrical figure called a polytopes in a real vector space, and a geometrical way of representing the measured membership weights of this collection and its conjunctions by means of a vector in this real vector space called a correlation vector. We prove that ‘if this correlation vector is located inside the polytope a classical measure theoretic model exists for these data, while if the correlation vector is located outside of the polytope there does not exist such a model’.

2 Membership weights on pairs of concepts and their conjunctions

It has been shown that *Guppy* is neither a very typical example of *Pet* nor *Fish* but is a very typical example of *Pet-Fish* [27]. Hence, the typicality of a specific item with respect to the conjunction of concepts can behave in an unexpected way. The problem is often referred to as the ‘pet-fish problem’ and the effect is usually called the ‘guppy effect’. The guppy effect is abundant; it appears almost in every situation where concepts combine. Meanwhile many experiments and analyses of this effect and related to the problem of combining concepts have been conducted [27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41].

The guppy effect was not only identified for the typicality of items with respect to concepts and their conjunctions but also for the membership weights of items with respect to concepts and their conjunction [29]. For example, subjects rate *Cuckoo* a better member of the conjunction ‘*Bird and Pet*’ than of the concept *Pet* on its own. This is a strange effect; if the conjunction of concepts behaved like a conjunction of logical propositions the second should be at least as great as the first. This deviation from what one would expect of a standard classical interpretation of conjunctions of concepts is referred to as ‘overextension’ [29]. Table 1 gives the list of six pairs of concepts and their conjunction for which in [29] the membership weights were measured with respect to different items, and in Table 2 the outcomes of these measurements are given for each of the items.

3 Classical and non classical membership weights

This behavior of a standard classical weight for a conjunction is described mathematically for the case of one pair of concepts and their conjunction in section 3 of [18]. Consider weights $\mu(A_1)$, $\mu(A_2)$ and

$\mu(A_1$ and $A_2)$ of an item X with respect to a pair of concepts A_1 and A_2 and their conjunction ‘ A_1 and A_2 ’. We say that they are ‘classical membership weights’ if and only if there exists a normed measure space $(\Omega, \sigma(\Omega), P)$ and events $E_{A_1}, E_{A_2} \in \sigma(\Omega)$ of the events algebra $\sigma(\Omega)$ such that

$$P(E_{A_1}) = \mu(A_1) \quad P(E_{A_2}) = \mu(A_2) \quad \text{and} \quad P(E_{A_1} \cap E_{A_2}) = \mu(A_1 \text{ and } A_2) \quad (1)$$

A normed measure P is a function defined on a σ -algebra $\sigma(\Omega)$ over a set Ω and taking values in the interval $[0, 1]$ such that the following properties are satisfied: (i) The empty set has measure zero, i.e. $P(\emptyset) = 0$; (ii) Countable additivity or σ -additivity: if E_1, E_2, E_3, \dots is a countable sequence of pairwise disjoint sets in $\sigma(\Omega)$, the measure of the union of all the E_i is equal to the sum of the measures of each E_i , i.e. $P(\bigcup_{i=1}^{\infty} E_i) = \sum_{i=1}^{\infty} P(E_i)$; (iii) The total measure is one, i.e. $P(\Omega) = 1$. The triple $(\Omega, \sigma(\Omega), P)$ is called a normed measure space, and the members of $\sigma(\Omega)$ are called measurable sets. A σ -algebra over a set Ω is a nonempty collection $\sigma(\Omega)$ of subsets of Ω that is closed under complementation and countable unions of its members. Measure spaces are the most general structures devised by mathematicians and physicists to represent weights.

We generalize this definition to the case of n concepts A_1, A_2, \dots, A_n with weights $\mu(A_i)$ for each concept A_i , and weights $\mu(A_i$ and $A_j)$ for the conjunction of concepts A_i and A_j . It is not necessary that weights are measured with respect to each one of the possible pairs of concepts. Hence, to describe this situation formally, we consider a set S of pairs of indices $S \subseteq \{(i, j) \mid i < j; i, j = 1, 2, \dots, n\}$ corresponding to those pairs of concepts for which the weights have been measured with respect to the conjunction of these pairs. As a consequence, the following set of weights have been experimentally determined

$$p_i = \mu(A_i) \quad i = 1, 2, \dots, n \quad p_{ij} = \mu(A_i \text{ and } A_j) \quad (i, j) \in S$$

We say that the set of weights in (2) is a ‘classical set of membership weights’ if it has a normed measure representation, hence if there exists a normed measure space $(\Omega, \sigma(\Omega), P)$ with $E_{A_1}, E_{A_2}, \dots, E_{A_n} \in \sigma(\Omega)$ elements of the event algebra, such that

$$p_i = P(E_{A_i}) \quad i = 1, 2, \dots, n \quad p_{ij} = P(E_{A_i} \cap E_{A_j}) \quad (i, j) \in S \quad (2)$$

4 Geometrical characterization of membership weights

We now introduce a geometric language that makes it possible to verify the existence of a normed measure representation for the set of weights in (2), inspired by the characterization of Kolmogorovian probability models in [42]. First we introduce a $n + |S|$ -dimensional correlation vector

$$\vec{p} = (p_1, p_2, \dots, p_n, \dots, p_{ij}, \dots) \quad (3)$$

where $|S|$ is the cardinality of S . Denote $R(n, S) = \mathbb{R}^{n+|S|}$ the $n + |S|$ dimensional vector space over the real numbers. Let $\epsilon \in \{0, 1\}^n$ be an arbitrary n -dimensional vector consisting of 0’s and 1’s. For each ϵ we construct the following vector $\vec{u}^\epsilon \in R(n, S)$

$$u_i^\epsilon = \epsilon_i \quad i = 1, 2, \dots, n \quad u_{ij}^\epsilon = \epsilon_i \epsilon_j \quad (i, j) \in S \quad (4)$$

The set of convex linear combinations of the u ’s is called the classical correlation polytope

$$c(n, S) = \{ \vec{f} \in R(n, S) \mid \vec{f} = \sum_{\epsilon \in \{0,1\}^n} \lambda_\epsilon \vec{u}^\epsilon; \lambda_\epsilon \geq 0; \sum_{\epsilon \in \{0,1\}^n} \lambda_\epsilon = 1 \} \quad (5)$$

The following theorem can now be proven similar to what was done in [42] for the case of Kolmogorovian probabilities

Theorem. *The set of weights*

$$p_i = \mu(A_i) \quad i = 1, 2, \dots, n \quad p_{ij} = \mu(A_i \text{ and } A_j) \quad (i, j) \in S$$

admits a normed measure space, and hence is a classical set of membership weights, if and only if its correlation vector \vec{p} belongs to the correlation polytope $c(n, S)$

Proof. Suppose that (6) is a classical set of weights, and hence we have a normed measure space $(\Omega, \sigma(\Omega), P)$ and events $E_{A_i} \in \sigma(\Omega)$ such that (2) are satisfied. Let us show that in this case $\vec{p} \in c(n, S)$. For an arbitrary subset $X \subset \Omega$ we define $X^1 = X$ and $X^0 = \Omega \setminus X$. Consider $\epsilon = (\epsilon_1, \dots, \epsilon_n) \in \{0, 1\}^n$ and define $A(\epsilon) = \bigcap_{\epsilon_i=1} A_i^{\epsilon_i}$. Then we have that $A(\epsilon) \cap A(\epsilon') = \emptyset$ for $\epsilon \neq \epsilon'$, $\bigcup_{\epsilon} A(\epsilon) = \Omega$, and $\bigcup_{\epsilon, \epsilon_j=1} A(\epsilon) = A_j$. We put now $\lambda_{\epsilon} = P(A(\epsilon))$. Then we have $\lambda_{\epsilon} \geq 0$ and $\sum_{\epsilon} \lambda_{\epsilon} = 1$, and $p_i = P(A_i) = \sum_{\epsilon, \epsilon_i=1} \lambda_{\epsilon} = \sum_{\epsilon} \lambda_{\epsilon} \epsilon_i$. We also have $p_{ij} = P(A_i \cap A_j) = \sum_{\epsilon, \epsilon_i=1, \epsilon_j=1} \lambda_{\epsilon} = \sum_{\epsilon} \lambda_{\epsilon} \epsilon_i \epsilon_j$. This means that $\vec{p} = \sum_{\epsilon} \lambda_{\epsilon} u^{\epsilon}$, which shows that $\vec{p} \in c(n, S)$. Conversely, suppose that $\vec{p} \in c(n, S)$. Then there exists numbers $\lambda_{\epsilon} \geq 0$ such that $\sum_{\epsilon} \lambda_{\epsilon} = 1$ and $\vec{p} = \sum_{\epsilon} \lambda_{\epsilon} u^{\epsilon}$. We define $\Omega = \{0, 1\}^n$ and $\sigma(\Omega)$ the power set of Ω . For $X \subset \Omega$ we define then $P(X) = \sum_{\epsilon \in X} \lambda_{\epsilon}$. Then we choose $A_i = \{\epsilon, \epsilon_i = 1\}$ which gives that $P(A_i) = \sum_{\epsilon} \lambda_{\epsilon} \epsilon_i = \sum_{\epsilon} \lambda_{\epsilon} u_i^{\epsilon} = p_i$ and $P(A_i \cap A_j) = \sum_{\epsilon} \lambda_{\epsilon} \epsilon_i \epsilon_j = \sum_{\epsilon} \lambda_{\epsilon} u_{ij}^{\epsilon} = p_{ij}$. This shows that we have a classical set of weights. \square

5 The correlation polytopes for pairs of concepts and their conjunctions

In the case of two concepts A_1, A_2 and their conjunction ‘ A_1 and A_2 ’ the set of indices is $S = \{(1, 2)\}$ and the correlation polytope $c(2, S)$ is contained in the $2 + |S| = 3$ dimensional euclidean space, i.e. $R(2, \{1, 2\}) = \mathbb{R}^3$. Further we have four vectors $\epsilon \in \{0, 1\}^n$, namely $(0, 0), (0, 1), (1, 0)$ and $(1, 1)$, and hence the four vectors $\vec{u}^{\epsilon} \in \mathbb{R}^3$ which are the following

$$\vec{u}^{(0,0)} = (0, 0, 0) \quad \vec{u}^{(1,0)} = (1, 0, 0) \quad \vec{u}^{(0,1)} = (0, 1, 0) \quad \vec{u}^{(1,1)} = (1, 1, 1) \quad (6)$$

This means that the correlation polytope $c(2, \{1, 2\})$ is the convex region spanned by the convex combinations of the vectors $(0, 0, 0), (1, 0, 0), (0, 1, 0)$ and $(1, 1, 1)$, and the correlation vector is given by $\vec{p} = (\mu(A_1), \mu(A_2), \mu(A_1 \text{ and } A_2))$. It is well-known that every polytope admits two dual descriptions: one in terms of convex combinations of its vertices, and one in terms of the inequalities that define its boundaries [43]. For the polytope $c(2, \{1, 2\})$ the inequalities defining its boundaries are $0 \leq \mu(A_1 \text{ and } A_2); \mu(A_1 \text{ and } A_2) \leq \mu(A_1); \mu(A_1 \text{ and } A_2) \leq \mu(A_2)$ and $\mu(A_1) + \mu(A_2) - \mu(A_1 \text{ and } A_2) \leq 1$.

In Figures 1, 2 and 3 we have represented this correlation polytope $c(2, \{1, 2\})$ and all the correlation vectors \vec{p} for the different items (we have presented the vectors as points not to overload the figure). If the point of the correlation vector corresponding to the data of a specific item lies inside the polytope spanned by $(0, 0, 0), (1, 0, 0), (0, 1, 0)$ and $(1, 1, 1)$, it is a classical item, for which the membership weights can be represented within a normed measure space. If the point does not lie inside the polytope, the corresponding item is non-classical, indicating that perhaps a quantum representation, for example the one presented in [18], can be elaborated for its weights.

Since the polytope is also given by the inequalities defining its boundaries, points lying inside and outside the polytope can be characterized by their coordinates satisfying or violating these inequalities. The inequalities that define the boundaries of polytope $c(2, \{1, 2\})$ are a lower dimensional variant of the well-known Bell inequalities studied in the foundations of quantum mechanics. This means that the violation of these inequalities, such as it happens by the data corresponding to items for which the points lie outside the polytope, has from a probabilistic perspective an analogous meaning as the

Table 1: The list of pairs of concepts and their conjunction used in [29]

A_1	A_2	A_1 and A_2
<i>Furniture</i>	<i>Household Appliances</i>	<i>Furniture and Household Appliances</i>
<i>Food</i>	<i>Plant</i>	<i>Food and Plant</i>
<i>Weapon</i>	<i>Tool</i>	<i>Weapon and Tool</i>
<i>Building</i>	<i>Dwelling</i>	<i>Building and Dwelling</i>
<i>Machine</i>	<i>Vehicle</i>	<i>Machine and Vehicle</i>
<i>Bird</i>	<i>Pet</i>	<i>Bird and Pet</i>

violation of Bell inequalities. Hence these violations may indicate the presence of quantum structures in the domain where the data is collected, which makes it plausible that a quantum model, such as for example the one proposed in [18], can be used to model the data.




Figure 1: The polytopes for the concepts *Furniture* and *Household Appliances* and the concepts *Building* and *Dwelling*. The classical item are *Castle*, *Cave*, *Phone Box*, *Synagogue*, *Log Cabin* and *House*. The other items are non classical.

6 Conclusion

If two concepts are combined to form a conjunction we can measure the membership weights of items with respect to each of these concepts and also with respect to their conjunction. If the ‘conjunction of concepts’ would behave like a classical logical conjunction of propositions does, we would expect that the membership weight of an item with respect to the conjunction is never bigger than the membership weight of this item with respect to one of the concepts. Experiments show that this is not the case, and the counter intuitive effect is referred to as the guppy effect [27] and also called overextension [29]. It has been shown that a quantum description can model this overextension while classical measure theoretic structures cannot [18]. In the present paper we elaborate a simple

Table 2: Three of the pairs of concepts and items of experiment 4 in [29]. The non classical items are labeled by q and the classical items by c .

$\mu(A_1)$ $\mu(A_2)$ $\mu(A_1 \text{ and } A_2)$					$\mu(A_1)$ $\mu(A_2)$ $\mu(A_1 \text{ and } A_2)$				
$A_1 = \text{Furniture}, A_2 = \text{Household Appliances}$					$A_1 = \text{Building}, A_2 = \text{Dwelling}$				
<i>Filing Cabinet</i>	q	0.9744	0.3077	0.5263	<i>Castle</i>	c	1	1	1
<i>Clothes Washer</i>	q	0.15	1	0.725	<i>Cave</i>	c	0.2821	0.95	0.2821
<i>Vacuum Cleaner</i>	q	0.075	1	0.3846	<i>Phone box</i>	c	0.2308	0.0526	0.02778
<i>Hifi</i>	q	0.5789	0.7895	0.7895	<i>Apartment Block</i>	q	0.9231	0.8718	0.9231
<i>Heated Water Bed</i>	q	1	0.4872	0.775	<i>Library</i>	q	0.95	0.175	0.3077
<i>Sewing Chest</i>	q	0.8718	0.5	0.55	<i>Trailer</i>	q	0.35	1	0.6154
<i>Floor Mat</i>	q	0.5641	0.15	0.2051	<i>Jeep</i>	q	0	0.05	0.05
<i>Coffee Table</i>	q	1	0.15	0.3846	<i>Palena</i>	q	0.975	1	1
<i>Piano</i>	q	0.95	0.1282	0.3333	<i>Igloo</i>	q	0.875	1	0.9
<i>Rug</i>	q	0.5897	0.05128	0.1842	<i>Synagogue</i>	c	0.925	0.4872	0.4474
<i>Painting</i>	q	0.6154	0.0513	0.1053	<i>Tent</i>	q	0.5	0.9	0.55
<i>Chair</i>	q	0.975	0.175	0.3590	<i>Bown</i>	q	0.9487	0.8205	0.8974
<i>Fridge</i>	q	0.4103	1	0.775	<i>Theatre</i>	q	0.95	0.1282	0.2821
<i>Desk Lamp</i>	q	0.725	0.825	0.825	<i>LogCabin</i>	c	1	1	1
<i>Cooking Stove</i>	q	0.3333	1	0.825	<i>House</i>	c	1	1	1
<i>TV</i>	q	0.7	0.9	0.925	<i>Tree House</i>	q	0.7692	0.8462	0.85
$A = \text{Food}, B = \text{Plant}$					$A = \text{Machine}, B = \text{Vehicle}$				
<i>Garlic</i>	q	0.9487	0.7105	0.8514	<i>Dog Sled</i>	q	0.1795	0.925	0.275
<i>Toadstool</i>	q	0.1429	0.6061	0.2727	<i>Dishwasher</i>	q	1	0.025	0
<i>Steak</i>	c	1	0	0	<i>Backpack</i>	c	0	0	0
<i>Peppercorn</i>	q	0.875	0.6207	0.7586	<i>Bicycle</i>	q	0.85	0.975	0.95
<i>Potato</i>	q	1	0.7436	0.9	<i>Sailboat</i>	c	0.5641	0.8	0.4211
<i>Raisin</i>	q	1	0.3846	0.775	<i>Roadroller</i>	q	0.9375	0.9063	0.9091
<i>Mint</i>	q	0.8718	0.8056	0.8974	<i>Raft</i>	c	0.2051	0.725	0.2
<i>Sunflower</i>	q	0.7692	1	0.775	<i>Elevator</i>	q	0.9744	0.6	0.7949
<i>Seaweed</i>	q	0.825	0.9744	0.8684	<i>Course liner</i>	q	0.875	0.875	0.95
<i>Sponge</i>	q	0.0263	0.3421	0.0882	<i>Automobile</i>	c	1	1	1
<i>Bread</i>	q	1	0.0769	0.2051	<i>Horsecart</i>	q	0.3846	0.95	0.2895
<i>Cabbage</i>	q	1	0.9	1	<i>Skateboard</i>	q	0.2821	0.8421	0.3421
<i>Eucalyptus</i>	q	0.1622	0.8974	0.3243	<i>Bus</i>	c	1	1	1
<i>Poppy</i>	q	0.3784	0.8947	0.5405	<i>Bulldozer</i>	q	1	0.925	0.95
<i>Mushroom</i>	q	1	0.6667	0.9	<i>Lawn-mower</i>	q	0.975	0.1053	0.2632
<i>Lettuce</i>	q	1	0.925	1	<i>Ski Lift</i>	q	1	0.5897	0.875
$A = \text{Weapon}, B = \text{Tool}$					$A = \text{Bird}, B = \text{Pet}$				
<i>Ruler</i>	q	0.05	0.9	0.1538	<i>Dog</i>	c	0	1	0
<i>Toothbrush</i>	c	0	0.55	0	<i>Cuckoo</i>	q	1	0.575	0.8421
<i>Chisel</i>	q	0.4	0.975	0.6410	<i>Parakeet</i>	c	1	1	1
<i>Axe</i>	q	0.875	1	0.975	<i>Cat</i>	c	0	1	0
<i>Screwdriver</i>	q	0.3	1	0.625	<i>Lark</i>	q	1	0.275	0.4872
<i>Arrow</i>	q	1	0.225	0.575	<i>Heron</i>	q	0.9412	0.1515	0.2581
<i>Knife</i>	c	1	0.975	0.975	<i>Peacock</i>	q	1	0.4	0.5789
<i>Rifle</i>	q	1	0.35	0.5	<i>Cow</i>	q	0	0.425	0.025
<i>Whip</i>	q	0.875	0.2632	0.625	<i>Toucan</i>	q	1	0.6154	0.8026
<i>Hammer</i>	q	0.575	1	0.8	<i>Parrot</i>	c	1	1	1
<i>Scissors</i>	q	0.6053	0.9744	0.7692	<i>Mynah Bird</i>	q	1	0.8710	0.8438
<i>Spoon</i>	q	0	0.752	0.075	<i>Raven</i>	q	1	0.2368	0.4
<i>Spear</i>	q	1	0.275	0.7179	<i>Elephant</i>	c	0	0.25	0
<i>Chain-saw</i>	q	0.55	1	0.75	<i>Goldfish</i>	c	0	1	0
<i>Club</i>	q	1	0.3590	0.775	<i>Homing Pigeon</i>	q	1	0.775	0.8974
<i>Razor</i>	q	0.625	0.775	0.825	<i>Canary</i>	c	1	1	1




Figure 2: The polytopes for the concepts *Food* and *Plant* and the concepts *Machine* and *Vehicle*. The classical item are *Steak*, *Backpack*, *Automobile*, *Bus*, *Sailboat* and *Raft*. The other items are non classical.




Figure 3: The polytopes for the concepts *Weapon* and *Tool* and the concepts *Bird* and *Pet*. The classical item are *Knife*, *Toothbrush*, *Elephant*, *Dog*, *Cat*, *Goldfish*, *Parakeet*, *Parrot* and *Canary*. The other items are non classical.

geometric method to identify the membership weights of items with respect to the conjunction of concepts that cannot be modeled within a classical measure theoretic structure. We do this for the general situation of a set of n concepts and a set of conjunctions between these concepts. The method consists of determining a convex polytope, and making correspond each of the items with a correlation vector in the real vector space where also the polytope is defined. We prove that if the correlation vector is contained in the polytope the considered set of membership weights can be modeled within a classical measure space, while if the correlation vector is not contained in the polytope it cannot. We apply this geometrical characterization method to the set of data collected in [29] and see that most of the tested items have membership weights for which the correlation vector falls outside of the polytope, and hence these membership weights cannot be modeled within a classical measure space (see Figures 1, 2 and 3). We explain in which way this geometrical identification method gives rise to a demarcation equivalent to the violation of the well known Bell inequalities in physics. The violation of Bell inequalities in physics proves that no classical (in the sense physicists and mathematicians in the field of probability use the word) model can explain or describe the data obtained experiments in physics that test Bell inequalities. The majority of physicists agree nowadays that the violation of Bell inequalities represent experimental proof of the need of a fundamental change in the classical paradigm to describe the process under consideration. Hence, Hampton's membership weight data giving rise to a situation equivalent to the violation of Bell inequalities, constitutes a strong argument in favor of the fact that genuine quantum structure would be at work within the mechanism giving rise to these data, hence within human cognition. If so, then these experiments constitute a pioneering example of experimentally tested quantum structure in cognition performed by a psychologist in tempore non suspecto. It also would mean that only non classical description, for example one based on quantum mechanics eventually as the one elaborated in [18], is able to model the mechanism giving rise to these experimental data.

7 Acknowledgments

This work was supported by grant G.040508N to Diederik Aerts from the Fund for Scientific Research-Flanders, Belgium (F.W.O. Vlaanderen), and a grant to Liane Gabora from Foundation for the Future.

References

- [1] Schaden, M.: Quantum finance: A quantum approach to stock price fluctuations. *Physica A*, 316, 511–538 (2002).
- [2] Baaquie, B. E.: *Quantum Finance: Path Integrals and Hamiltonians for Options and Interest Rates*. Cambridge UK: Cambridge University Press (2004).
- [3] Bagarello, F.: An operational approach to stock markets. *Journal of Physics A* 39, 6823–6840 (2006).
- [4] Bordley, R. F.: Quantum mechanical and human violations of compound probability principles: Toward a generalized Heisenberg uncertainty principle. *Operations Research* 46, 923–926 (1998).
- [5] Bordley, R. F., & Kadane, J. B.: Experiment dependent priors in psychology and physics. *Theory and Decision* 47, 213–227 (1999).

- [6] Aerts, D., & Aerts, S.: Applications of quantum statistics in psychological studies of decision processes. *Foundations of Science*, 1, 85–97 (1994). Reprinted in B. Van Fraassen, (Eds.), *Topics in the Foundation of Statistics* (pp. 111-122). Dordrecht: Kluwer Academic (1997).
- [7] Grossberg, S.: The complementary brain: Unifying brain dynamics and modularity. *Trends in Cognitive Science* 4, 233–246 (2000).
- [8] Gabora, L., & Aerts, D.: Contextualizing concepts. In *Proceedings of the 15th International FLAIRS Conference. Special track: Categorization and Concept Representation: Models and Implications*, Pensacola Florida, May 14-17, American Association for Artificial Intelligence (pp. 148–152) (2002).
- [9] Gabora, L., & Aerts, D.: Contextualizing concepts using a mathematical generalization of the quantum formalism. *Journal of Experimental and Theoretical Artificial Intelligence* 14, 327–358 (2002).
- [10] Aerts, D., & Gabora, L.: A theory of concepts and their combinations I: The structure of the sets of contexts and properties. *Kybernetes* 34, 167–191 (2005).
- [11] Aerts, D., & Gabora, L.: A theory of concepts and their combinations II: A Hilbert space representation. *Kybernetes*, 34, 192–221 (2005).
- [12] Busemeyer, J. R., Wang, Z., & Townsend, J. T.: Quantum dynamics of human decision making. *Journal of Mathematical Psychology* 50, 220–241 (2006).
- [13] Busemeyer, J. R., Matthew, M., Wang, Z.: A quantum information processing theory explanation of disjunction effects. *Proceedings of the Cognitive Science Society* (2006).
- [14] Aerts, D.: Quantum interference and superposition in cognition: Development of a theory for the disjunction of concepts. Archive address and link: <http://arxiv.org/abs/0705.0975> (2007).
- [15] Aerts, D.: General quantum modeling of combining concepts: A quantum field model in Fock space. Archive address and link: <http://arxiv.org/abs/0705.1740> (2007).
- [16] Franco, R.: The conjunction fallacy and interference effects. Archive address and link: <http://arxiv.org/abs/0708.3948> (2007).
- [17] Franco, R.: Quantum mechanics and rational ignorance. Archive address and link: <http://arxiv.org/abs/physics/0702163> (2007).
- [18] Aerts, D.: Quantum interference and superposition in cognition: Development of a theory for the disjunction of concepts. Submitted to *Journal of Mathematical Psychology*, archive reference and link: <http://uk.arxiv.org/abs/0705.0975> (2008).
- [19] Eisert, J., Wilkens, M., & Lewenstein, M.: Quantum games and quantum strategies. *Physical Review Letters* 83, 3077–3080 (1999).
- [20] Piotrowski, E. W., & Sladkowski, J.: An invitation to quantum game theory. *International Journal of Theoretical Physics* 42, 1089 (2003).
- [21] Aerts, D., & Czachor, M.: Quantum aspects of semantic analysis and symbolic artificial intelligence. *Journal of Physics A, Mathematical and Theoretical* 37, L123-L132 (2004).

- [22] Widdows, D.: Orthogonal negation in vector spaces for modelling word-meanings and document retrieval. In *Proceedings of the 41st Annual Meeting of the Association for Computational Linguistics* (pp. 136-143). Sapporo, Japan, July 7-12 (2003).
- [23] Widdows, D., & Peters, S.: Word vectors and quantum logic: Experiments with negation and disjunction. In *Mathematics of Language 8* (pp. 141-154). Indiana: Bloomington (2003).
- [24] Van Rijsbergen, K.: *The Geometry of Information Retrieval*. Cambridge UK: Cambridge University Press (2004).
- [25] Aerts, D., Czachor, M., & D’Hooghe, B.: Towards a quantum evolutionary scheme: violating bell’s inequalities in language. In N. Gontier, J. P. Van Bendegem, & D. Aerts (Eds.), *Evolutionary Epistemology, Language and Culture - A Non Adaptationist Systems Theoretical Approach*. Dordrecht: Springer (2006).
- [26] Widdows, D.: *Geometry and Meaning*. CSLI Publications: University of Chicago Press (2006).
- [27] Osherson, D. N., Smith, E. E.: On the adequacy of prototype theory as a theory of concepts. *Cognition* 9, 35–58 (1981).
- [28] Hampton, J. A.: Inheritance of attributes in natural concept conjunctions. *Memory & Cognition* 15, 55–71 (1987).
- [29] Hampton, J. A.: Overextension of conjunctive concepts: Evidence for a unitary model for concept typicality and class inclusion. *Journal of Experimental Psychology: Learning, Memory, and Cognition* 14, 12–32 (1988).
- [30] Hampton, J. A.: Disjunction of natural concepts. *Memory & Cognition* 16, 579–591 (1988).
- [31] Hampton, J. A.: The combination of prototype concepts. In P. Schwanenflugel (Ed.), *The Psychology of Word Meanings*. Hillsdale, NJ: Erlbaum (1991).
- [32] Hampton, J. A.: Conceptual combination: Conjunction and negation of natural concepts. *Memory & Cognition* 25, 888–909 (1997).
- [33] Hampton, J. A.: Conceptual combination. In K. Lamberts & D. Shanks (Eds.), *Knowledge, Concepts, and Categories* (pp. 133-159). Hove: Psychology Press (1997).
- [34] Osherson, D. N., Smith, E. E.: Gradedness and conceptual combination. *Cognition* 12, 299–318 (1982).
- [35] Rips, L. J.: The current status of research on concept combination. *Mind & Language* 10, 72–104 (1995).
- [36] Smith, E. E., & Osherson, D. N.: Conceptual combination with prototype concepts. *Cognitive Science* 8, 357–361 (1988).
- [37] Springer, K., & Murphy, G. L.: Feature availability in conceptual combination. *Psychological Science* 3, 111–117 (1992).
- [38] Storms, G., De Boeck, P., Van Mechelen, I., & Geeraerts, D. (1993). Dominance and non-commutativity effects in concept conjunctions: Extensional or intensional basis? *Memory & Cognition*, 21, 752-762.

- [39] Storms, G., De Boeck, P., Van Mechelen, I. & Ruts, W. (1996). The dominance effect in concept conjunctions: Generality and interaction aspects. *Journal of Experimental Psychology: Learning, Memory & Cognition*, 22, 1-15.
- [40] Storms, G., De Boeck, P., Van Mechelen, I., Ruts, W.: Not guppies, nor goldfish, but tumble dryers, Noriega, Jesse Jackson, panties, car crashes, bird books, and Stevie Wonder. *Memory & Cognition* 26, 143–145 (1998).
- [41] Storms, G., de Boeck, P., Hampton, J.A., & van Mechelen, I. (1999). Predicting conjunction typicalities by component typicalities. *Psychonomic Bulletin and Review*, 6, 677-684.
- [42] Pitowsky, I.: Quantum Probability, Quantum Logic. Lecture Notes in Physics 321, Heidelberg: Springer (1989).
- [43] Aerts, S., Aerts, D. When is a set of data best described by a quantum (like) theory. Proceedings of the Quantum Interaction Symposium, University of Oxford, March 2008.