

Fractional recurrence in quantum walk

C. M. Chandrashekar

*Institute for Quantum Computing, University of Waterloo, Ontario N2L 3G1, Canada and
Perimeter Institute for Theoretical Physics, Waterloo, ON, N2L 2Y5, Canada*

Quantum recurrence theorem holds for quantum systems with discrete energy eigenvalues and fails to hold in general for systems with continuous energy. We show that during quantum walk process dominated by interference of amplitude corresponding to different paths fail to satisfy the complete quantum recurrence theorem. Due to the revival of the fractional wave packet, a fractional recurrence characterized using quantum Pólya number can be seen.

I. INTRODUCTION

In the dynamics of physical systems, from free particles to stellar dynamics, the recurrence phenomenon have significantly contributed to the better understanding of the systems dynamics [1]. For a classical conservative system, whether discrete or continuous in time, the Poincaré recurrence theorem states that any phase-space configuration (q, p) of a system enclosed in a finite volume will be repeated as accurately as one wishes after a finite interval of time (with no restriction on the interval)[2]. A similar recurrence theorem is shown to hold in quantum theory as well [3]. In a system with a discrete energy eigenvalue spectrum $\{E_n\}$; if $\Psi(t_0)$ is its state vector at the time t_0 and ϵ is any positive number, at least one time T will exist such that the norm of the vector $\Psi(T) - \Psi(t_0)$,

$$|\Psi(T) - \Psi(t_0)| < \epsilon. \quad (1)$$

Recurrence in quantum systems has also been extensively studied [4].

Quantum walks (QWs) are the quantum analog of the dynamics of classical random walks (CRWs) [5, 6, 7]. The QW involves superposition of states and moves simultaneously exploring multiple possible paths with the amplitudes corresponding to different paths interfering with one another. This makes the variance of the QW on a line to grow quadratically with the number of steps compared to the linear growth for the CRW. Due to the speedup, the QW has emerged as a powerful approach to quantum algorithm design [8, 9, 10, 11]. The dynamics of QWs seems to serve as a potential tool to understand and explain various phenomenon in physical systems beyond quantum computation. It has been used to demonstrate the coherent quantum control over atoms and realize quantum phase transition from Mott insulator to superfluid state and vice versa [12], to explain the phenomenon such as, the breakdown of an electric-field driven system [13] and the direct experimental evidence for wavelike energy transfer within photosynthetic systems[14, 15].

Many properties of the dynamics of the CRW are very well understood and several analogous properties have been studied in QWs. One such property is the recurrence probability of the QW [16]. Recurrence in quantum walk differ significantly from the recurrence in classical random walk. For CRW we can consider the recurrence probability $p_0(t)$, i.e., the probability of the periodicity of the dynamics that the particle returns to the origin during the time evolution (t steps). It is characterized by the Pólya number

$$P_{crw} \equiv 1 - \frac{1}{\sum_{t=0}^{\infty} p_0(t)}. \quad (2)$$

If the Pólya number equals one, the CRW is recurrent, otherwise the walk is transient, i.e., with nonzero probability the particle never returns to the origin. For a CRW to be transient the series $\sum_{t=0}^{\infty} p_0(t)$ must converge [17]. Pólya proved that the one- and two- dimensional CRW are recurrent [18], and for each higher dimension a unique Pólya number is associated while the CRWs are transient.

In standard quantum mechanics, initially localized wave packet in state $|\Psi(t_0)\rangle$ which can spread significantly in a closed system can also reform later in the form of a quantum revival in which the spreading reverses itself and the wave packet relocalizes [4]. The relocalized wave packet can again spread and the periodicity in the dynamics can be seen validating the quantum recurrence theorem. The time evolution of a state $|\Psi(t)\rangle$, or a wave function $\Psi(x, t)$ is given by a deterministic unitary transformation associated with the Hamiltonian. During quantum evolution we deal with the amplitudes and the probability density $p(x, t) = |\Psi(x, t)|^2$ at position x after time t appears only when we collapse the wave packet to perform measurement.

Unlike standard wave packet spreading[30], the QW spreads the wave packet in multiple possible paths with the amplitudes corresponding to different paths interfering. In this paper we show that due to the entanglement between the particle and the position space and interference effect during the evolution of the QW the wave packet delocalizes over the position space as a small copies of the initial wave packet. These delocalized copies of fractional wave packet

fails to satisfy the complete quantum recurrence theorem during the QW evolution. However, due to the revival of the fractional wave packets, a fractional recurrence can be seen in QW. The probabilistic characterization of the QW using quantum Pólya number defined in Ref.[19] shows fractional recurrence nature of QW. We also show the exceptional cases of QW that can be constructed by suppressing or minimizing the interference effect and get close to the complete recurrence.

In Sec. II we briefly describe the standard variants of the discrete-time QW (DTQW) and the continuous-time QW (CTQW). In Sec. III we revisit the quantum recurrence theorem. In Sec. IV we discuss the quantum recurrence theorem for QW and show the fractional recurrence nature of the QW on a line and an n -cycle before concluding in Sec. V.

II. QUANTUM WALK

The study of QWs has been largely divided into two standard variants, the discrete-time QW (DTQW)[7, 20] and the continuous-time QW (CTQW) [21]. Considering the limiting value of the DTQW the connection between the two variants was established [22] and recently the present author has presented a generic model of the QW which can be conveniently used to retrieve both the standard DTQW and CTQW variants [23].

The DTQW is defined on the Hilbert space $\mathcal{H} = \mathcal{H}_c \otimes \mathcal{H}_p$ where, \mathcal{H}_c is the *coin* Hilbert space and the \mathcal{H}_p is the *position* Hilbert space. For a QW in one dimension, the \mathcal{H}_c is spanned by the basis state of the particle $|0\rangle$ and $|1\rangle$ and the \mathcal{H}_p is spanned by the basis state of the position $|x\rangle$, $x \in \mathbb{Z}$. To implement the DTQW, a three parameter quantum coin toss operation $C_{\xi, \theta, \zeta}$ of the form

$$C_{\xi, \theta, \zeta} \equiv \begin{pmatrix} e^{i\xi} \cos(\theta) & e^{i\zeta} \sin(\theta) \\ e^{-i\zeta} \sin(\theta) & -e^{-i\xi} \cos(\theta) \end{pmatrix} \quad (3)$$

is applied on the particle at origin in state

$$|\Psi_0\rangle = (\cos(\delta)|0\rangle + e^{i\eta} \sin(\delta)|1\rangle) \otimes |\psi_0\rangle. \quad (4)$$

The quantum coin operation on the particle is followed by the conditional unitary shift operation S which delocalizes the wave packet over the position $(x-1)$ and $(x+1)$,

$$S = \exp(-i\sigma_z \otimes Pl), \quad (5)$$

P being the momentum operator and σ_z , the operator corresponding to step of length l depending on the state of the particle respectively. The eigenstates of σ_z are denoted by $|0\rangle$ and $|1\rangle$. Therefore, S can also be written as,

$$S = |0\rangle\langle 0| \otimes \sum_{x \in \mathbb{Z}} |\psi_{x-1}\rangle\langle \psi_x| + |1\rangle\langle 1| \otimes \sum_{x \in \mathbb{Z}} |\psi_{x+1}\rangle\langle \psi_x|. \quad (6)$$

The process of

$$W_{\xi, \theta, \zeta} = S(C_{\xi, \theta, \zeta} \otimes \mathbb{1}) \quad (7)$$

is iterated without resorting to the intermediate measurement to realize large number of steps of the QW. δ and η in Eq. (4) can be varied to get different initial state of the particle. The three variable parameters of the quantum coin, ξ , θ and ζ in Eq. (3) can be varied to change the probability amplitude distribution in the position space. By varying the parameter θ the variance can be increased or decreased as a functional form, $\sigma^2 \approx (1 - \sin(\theta))t^2$, where t is the number of steps (iteration). For a particle with symmetric superposition as initial state the parameter ξ and ζ introduces asymmetry in the probability distribution and their effect on the variance is very small [24].

In the CTQW [21], the walk is defined on the *position* Hilbert space \mathcal{H}_p spanned by the basis state $|\psi_x\rangle$, $x \in \mathbb{Z}$. To implement the CTQW the Hamiltonian H is defined such that

$$H|\psi_x\rangle = -\gamma|\psi_{x-1}\rangle + 2\gamma|\psi_x\rangle - \gamma|\psi_{x+1}\rangle \quad (8)$$

and is made to evolve with time τ by applying the transformation

$$U(\tau) = \exp(iH\tau). \quad (9)$$

The Hamiltonian H of the process acts as the generator matrix which will transform the probability amplitude at the rate of γ to the neighboring sites. γ is a fixed, time-independent, constant. In this paper we will only consider the study using the DTQW evolution.

III. QUANTUM RECURRENCE THEOREM

Quantum recurrence theorem in the dynamics of the closed system states that there exist a time T when

$$|\Psi(T) - \Psi(t_0)| < \epsilon, \quad (10)$$

where $\Psi(T) = |\Psi_T\rangle$ is the state of the system after time T , $\Psi(t_0) = |\Psi_0\rangle$ is the initial state of the system and ϵ is any positive number ≤ 2 (both $\Psi(T)$ and $\Psi(t_0)$ are normalized functions)[3].

The recurrence of complete state of the system or the exact revival happens when all the expectation values of observables A of the two states $|\Psi_T\rangle$ and $|\Psi_0\rangle$ are equal to one another, that is,

$$\langle \Psi_T | A | \Psi_T \rangle = \langle \Psi_0 | A | \Psi_0 \rangle. \quad (11)$$

In classical dynamics, the characterization of the recurrence nature can be conveniently done using probabilistic measures. Measurements on quantum system leaves the state in one of its basis states with certain probability. Therefore, recurrence in quantum systems can be analyzed using the analysis of two cases of the comparative evolution of the two identically prepared quantum system with the initial states $|\Psi_{0,0}\rangle$ at position x_0 and time t_0 .

Case 1: Consider two identically prepared particle wave packet which revive completely in position space at time T . One of the two particle wave packet at position x_0 and time t_0 is first evolved to spread in position space and then reverse the spreading till it relocalizes completely at position x_0 at time T . The measurement performed on this particle will collapse the wave packet at the relocalized position with an expectation value

$$\langle \Psi_{0,T} | X | \Psi_{0,T} \rangle = \langle \Psi_{0,0} | X | \Psi_{0,0} \rangle = 1, \quad (12)$$

where X is the position operator. After the measurement, the system is further evolved for an additional unit time and the corresponding state can be given by $|\Psi_{x,T_{M+1}}\rangle$, the subscript M stands for the measurement being performed at time T . The second particle wave packet at position x_0 and time t_0 is evolved up to time $(T+1)$ directly without any measurement being performed at time T and the state can be written as $|\Psi_{x,T+1}\rangle$. Since both the wave packet completely relocalize at position x_0 after the evolution for the time period T , irrespective of the measurement being performed, the expectation value for both the particle after time $(T+1)$ would be identical,

$$\langle \Psi_{x,T_{M+1}} | X | \Psi_{x,T_{M+1}} \rangle = \langle \Psi_{x,T+1} | X | \Psi_{x,T+1} \rangle \quad (13)$$

with x spanning over all the position space.

Case 2: Consider two identically prepared particle wave packet which does not relocalize completely at position x_0 at time T , i.e., revive fractionally or does not revive at all. The measurement will collapse the wave packet and return the expectation value

$$\langle \Psi_{0,T} | X | \Psi_{0,T} \rangle = \delta \quad ; \quad 0 < \delta < 1. \quad (14)$$

In this case the two identically prepared particle evolved to time $(T+1)$, one with a measurement being performed at time T and an other without any measurement being performed, will not return the same expectation value, that is

$$\langle \Psi_{x,T_{M+1}} | X | \Psi_{x,T_{M+1}} \rangle \neq \langle \Psi_{x,T+1} | X | \Psi_{x,T+1} \rangle \quad (15)$$

with x spanning over all position space. A non-zero values in both the expectation values in the inequality can act as a signature of the fractional recurrence of the quantum state at time T and a zero expectation value on the left hand side shows the transient dynamics.

From the above analysis we can conclude that, if the system is completely recurrent irrespective of the measurement at time T , the two states $|\Psi_{x,T_{M+1}}\rangle$ and $|\Psi_{x,T+1}\rangle$ will be equal to one another.

IV. FRACTIONAL RECURRENCE OF QUANTUM WALK

A. On a line

The state of the particle wave packet after implementing the DTQW of t steps on a line with unit time required to implement each step can be written as

$$\sum_x |\Psi_{x,t}\rangle = W_{\xi,\theta,\zeta}^t |\Psi_{0,0}\rangle = \sum_x a_x |\Phi_{x,t}\rangle, \quad (16)$$




FIG. 1: The plot of $[1 - p_0(t)]$, where $p_0(t)$ is the probability of particle at the origin with t being the number of steps in the DTQW evolution. The plot for QW on a line using different coin operation parameter $\theta = 15^\circ, 45^\circ$, and 75° is shown. With increase in θ , the quantum Pólya number which characterizes the fractional recurrence nature of the QW also increases. The plot with $\theta = 0^\circ$ is for a walk on an n -cycle with a completely suppressed interference effect, here $n = 51$. For a walk to be completely recurrent their should exist a $t = T$ where $[1 - p_0(T)] = 0$, any non-zero value can be used to characterize the fractional recurrence nature of the QW and an absolute 1 for all t would show that the QW is completely transient.

$|\Phi_{x,t}\rangle$ is the state of the delocalized wave packet at each position x in \mathcal{H}_p of any dimension and $|\Psi_{0,0}\rangle$ is the state of the wave packet before implementing the QW. In the QW process which involves a deterministic unitary evolution, the particle wave packet spreads over the position space forming a mini wave packets. During this delocalization process the mini wave packets interfere and entangle the position and coin Hilbert space, \mathcal{H}_p and \mathcal{H}_c . The interference and the entanglement between the \mathcal{H}_p and \mathcal{H}_c during the standard QW evolution does not permit complete relocalization of the wave packet at initial position after any given number of steps t . Therefore the argument leading to Eq.(14) and Eq.(15) holds to show that the complete recurrence of the quantum state does not occur during the evolution of the QW process on a line. That is,

In QW evolution on a line and an n -cycle dominated by interference of quantum amplitude[31], there exist no time T where the quantum state of the system revive completely and repeat the delocalization and complete revival at regular interval of time.

The above statement can also be quantified in the following way: After implementing quantum walk, the wave function describing the particle at position x and time t can be written as a two component vector of amplitudes of particle being at position x at time t with left and right propagating component

$$|\Psi_{x,t}\rangle = \begin{pmatrix} \Psi^L(x,t) \\ \Psi^R(x,t) \end{pmatrix}. \quad (17)$$

Lets analyze the dynamics of DTQW on wave packet $\Psi(x,t)$ driven by single parameter quantum coin

$$C = \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix} \quad (18)$$

and shift operator S ($W = SC$). In terms of left and right propagating component it is given by

$$\begin{aligned} \Psi^R(x,t+1) &= \cos \theta \Psi^R(x-1,t) + \sin \theta \Psi^L(x-1,t) \\ \Psi^L(x,t+1) &= \sin \theta \Psi^R(x+1,t) - \cos \theta \Psi^L(x+1,t) \end{aligned} \quad (19)$$

Then the probability of being at position x and t is

$$P(x,t) = |\Psi^L(x,t)|^2 + |\Psi^R(x,t)|^2. \quad (20)$$

and sum of probability over the entire position space is

$$\sum_x P(x, t) = 1. \quad (21)$$

After time t with unit time required to implement each step of the quantum walk on a line will be spread between $x = -t$ to $x = +t$ and the wave function over the position space is

$$\sum_{x=-t}^t \Psi(x, t) = \sum_{x=-t}^t [\Psi^R(x, t) + \Psi^L(x, t)] \quad (22)$$

It should be noted that for even number of steps the amplitude at odd labeled positions is 0 and for odd number of steps amplitude at even labeled position is 0.

For CRW, each step of walk is associated with the *randomness* and the *probability* of the entire particle therefore, however small the probability is at the origin, it is attributed to the recurrence of the entire particle. Whereas, for the DTQW evolution shown in Eq. (19) we find that the coin degree of freedom is carried over during the dynamics of the walk making it reversible. The *randomness* and the *probability* comes into consideration only when the wave packet is collapsed to discard the signature of the coin degree of freedom. Therefore, for a DTQW to be completely recurrent, the condition

$$P(0, t) = |\Psi^L(0, t)|^2 + |\Psi^R(0, t)|^2 = 1. \quad (23)$$

has to be satisfied for some time t . For any $P(0, t) < 1$ shows the fractional recurrence nature of the QW. From Eqs. (19) and (20) we can conclude that the Eq. (23) is satisfied only when $\theta = \pi/2$ and t is even, that is when there is no interference of the quantum amplitudes. For $\theta = 0$ the two left and right component move in opposite direction without returning and for any $0 < \theta < \pi/2$ we get $P(0, t) < 1$ showing the fractional recurrence of the QW.

Alternatively, using quantum Fourier analysis to study the evolution of the DTQW on a line, it is shown that the amplitude at the origin decreases by $O(1/\sqrt{t})$. To a very good approximation, independent of position x it has also been shown that the amplitude decrease by $O(1/\sqrt{t})$ (x being the points between the two dominating peaks in the distribution)[16]. Therefore,

$$\langle \Psi_{x,t} | \Psi_{x,t} \rangle = O\left(\frac{1}{t}\right) < 1. \quad (24)$$

and there exists no time $T = t$ where the walk is completely recurrent, whereas QW on a line shows a fractional recurrence nature.

A probability based characterization of the recurrence nature of the QW, quantum Pólya number was defined for an ensemble of identically prepared QW systems by the expression

$$P_{qw} = 1 - \prod_{t=1}^{\infty} [1 - p_0(t)] \quad (25)$$

where $p_0(t)$ is the post measurement recurrence probability of the particle. Each identically prepared particle is subjected to different number of steps of QW from 1 to ∞ and the probability of the particle at the origin is measured and discarded [19]. The probability that the particle is found at the origin in a single series of such measurement records is the quantum Pólya number. The quantum Pólya number was calculated for various coined QWs and it is shown that in the higher dimension it depends both on the initial state and the parameters in the coin operator whereas, for CRW the Pólya number is uniquely determined by its dimensionality [25].

To show the QW to be completely recurrent adopting a probability based characterization given by Eq. (25) needs to have at least one of the many particle, each evolved to different steps t from 1 to ∞ to return $p_0(t) = 1$. If $0 < p_0(t) < 1$ for any t , then only with certain probability $p_0(t)$ the wave packet collapses at x_0 , that is, prior to measurement the particle existed in superposition of position space. If $0 < p_0(t) < 1$ for all the particles evolved to different steps from 1 to ∞ then the fractional recurrence of the QW is characterized by the P_{qw} .

In Fig. (1) the plot of $[1 - p_0(t)]$ is shown for a DTQW on a line where the different coin operation parameter $\theta = 15^\circ, 45^\circ, 75^\circ$. With increase in θ the quantum Pólya number which can also be called as fractional recurrence number P_{qw} also increases.




FIG. 2: Mixing time M of the probability distribution of a quantum walker on a n -cycle for different value of θ using coin operation $U_{0,\theta,0}$, where n , the number of position, is 101. Mixing is faster for lower value of θ . The distribution is for 200 cycles.

B. On an n -cycle

The n -cycle is the simplest finite Cayley graph with n vertices, number of position in the \mathcal{H}_p . This example has most of the features of the walks on the general closed graphs. The unitary shift operation S , Eq. (6) for a QW on an n -cycle is defined by

$$S^c = |0\rangle\langle 0| \otimes \sum_{x=0}^{n-1} |\psi_{x-1 \bmod n}\rangle\langle \psi_x| + |1\rangle\langle 1| \otimes \sum_{x=0}^{n-1} |\psi_{x+1 \bmod n}\rangle\langle \psi_x| \quad (26)$$

and the quantum state after t steps of QW is written as

$$|\Psi_{x,t}\rangle = W_{\xi,\theta,\zeta}^t |\Psi_{0,0}\rangle = \sum_{x=0}^{n-1} a_x |\Phi_{x,t}\rangle. \quad (27)$$

The CRW approaches a stationary probability distribution independent of its initial state on a finite graph. The time required to approach a stationary distribution is known as mixing time [26]. As we have discussed earlier in case of CRW walk on a line, a non-zero probability is sufficient to show the recurrence of the entire particle subjected to CRW and stationary distribution returns a non-zero probability.

Unitary QW, does not converge to any stationary distribution on an n -cycle. But by defining a time-averaged distribution,

$$\overline{P(x,T)} = \frac{1}{T} \sum_{t=0}^{T-1} P(x,t), \quad (28)$$

obtained by uniformly picking a random time t between 0 and $(T-1)$, and evolving for t time steps and measuring to see which vertex it is at, a convergence in the probability distribution can be seen even in quantum case. It has been shown that the QW on an n -cycle mixes in time $M = O(n \log n)$, almost quadratically faster than the classical case which is $O(n^2)$ [26]. The mixing time can be optimized on an n -cycle by choosing a lower value of θ in $W_{0,\theta,0}$ [24]. Figure (2) is the time averaged probability distribution of a quantum walk on an n -cycle graph after $n \log n$ time where n is 101. It can be seen that the variation of the probability distribution over the position space is least for $\theta = 15^\circ$ compared to $\theta = 45^\circ$ and $\theta = 75^\circ$.

Uniform distribution in quantum case does not reveal the complete recurrence nature of the particle like it does in classical case. In Fig. (3) we show numerically that the probability of finding the particle at the initial position x_0 after different number of steps of QW on 15- and 24- cycle. The distribution is obtained without time-averaging for up to steps as large as 5000. At no time t =number of steps, a unit probability value is returned showing that wave packet evolved using QW fail to revive complete at the initial position.




FIG. 3: Probability at the initial position after different number of steps of quantum walk on 15– and 24– cycle. This clearly shows that even for steps as large as 5000, there is no signature of complete recurrence. The distribution is obtained using Hadamard operation, $C_{0,\pi/4,0}$ as quantum coin operation.




FIG. 4: Probability at the initial position on 8–cycle. (a) The distribution is obtained using Hadamard coin operation $C_{0,\pi/4,0}$, due to return of amplitudes to initial position (constructive interference at the origin) before interference of amplitude dominates uniformly over the entire vertices, recurrence is seen. (b) The use of quantum coin, $C_{0,\pi/6,0}$ during the evolution does not lead to complete constructive interference at origin and hence complete recurrence is not seen.

The failure of the wave packet to completely revive at initial position and recur can be attributed to the interference effect caused by the mixing of the left and right propagating components of the amplitude. By suppressing the interference effect during the evolution in a closed path one can get closer to the complete relocalization, revival at initial position x_0 . For example, the wave packet can be completely relocalized at x_0 on an n –cycle after $T = n$ and make the QW recurrent by choosing an extreme value of coin parameters $(\xi, \theta, \zeta) = (0^\circ, 0^\circ, 0^\circ)$ in Eq.(3) during the QW evolution. However $\theta = \delta$, δ being very small and close to 0° , where the interference effect is minimized, will also return a near complete recurrence on an n –cycle. In Fig. (1) the plot of $[1 - p_0(t)]$ is also shown for a particular case where the QW shows the complete recurrence nature, a walk on an n –cycle with $\theta = 0^\circ$ and $n = 50$. The interference effect is completely suppressed and the quantum recurs after every 50 steps.

From the numerically data for upto 5000 steps of quantum walk we note that for small n –, especially when n is even, the left and right propagating amplitude return back completely to the initial position (origin) before the mixing and repeated interference of the amplitudes takes over at non-initial position in the evolution process. Therefore, for a QW on an n –cycle with n being even upto 8 a complete revival and recurrence of wave packet at initial position $j = 0$ is seen, Figure 4 [27]. For QW on particle initially in symmetric superposition state

$$|\Psi_0\rangle = \frac{1}{\sqrt{2}}(|0\rangle + i|1\rangle) \otimes |\psi_0\rangle \quad (29)$$




FIG. 5: Probability at the initial position on 10-cycle. The distribution is obtained using Hadamard operation $C_{0,\pi/4,0}$. A small deviation from complete recurrence can be seen. (a) and (b) are probability at initial position for quantum walk up to 250 steps and 5000 steps respectively and (c) is the close up of the probability and we note that the probability is not exact 1 at any time within 5000 steps.

with coin operation C , Eq. (18) and even $n > 8$, due to larger position Hilbert space the interference effect at the non-initial position dominates reducing the recurrence nature of the dynamics. In Fig. 5 for QW on 10-cycle, small deviation from complete recurrence is shown.

For example, we will consider a small odd number, $n = 5$, if the position are marked as $x = 0, 1, 2, 3, 4$, during the third step of the QW, the left propagating amplitude move from position 2 to 3 and the right propagating amplitude move from 3 to 2. That is, after the second and third step of quantum walk using shift operator of the form S^c , Eq. (26) and Hadamard operation $H = C_{0,\pi/2,0}$ as coin operation on a particle initially in superposition state Eq. (29) takes the form,

$$S^c H S^c |\Psi_0\rangle = \frac{1}{2} \{ |0\rangle \otimes |\psi_3\rangle + (|1\rangle + i|0\rangle) \otimes |\psi_0\rangle - i|1\rangle \otimes |\psi_2\rangle \} \quad (30)$$

$$S^c H S^c H S^c |\Psi_0\rangle = \frac{1}{2\sqrt{2}} \{ |0\rangle \otimes |\psi_2\rangle + (|0\rangle + |1\rangle + i|0\rangle) \otimes |\psi_4\rangle - (|1\rangle - i|0\rangle + i|1\rangle) \otimes |\psi_1\rangle + i|1\rangle \otimes |\psi_3\rangle \}. \quad (31)$$

The left and right propagating amplitude crossover without suppressing the mixing, therefore the constructive interference effect continues to exist at position other than origin during the evolution. In Fig. (6) the probability of finding the particle at the initial position is shown. Due to the small size of the Hilbert space, the probability is seen to be close to unity but the closer look reveals that its only a fraction revival.




FIG. 6: Probability at the initial position after different number of steps of QW on 5-cycle, (a) is the complete plot whereas (b) is a close up of probability values between 0.9 and 1. At no step in the plot, probability is a unit value. This numerically shows that the QW on an n -cycle does not recur for $n=5$.

We also note that localization effect found in 2D [28] or found in QWs using multi quantum coins to diminish the interference effect [29] can result in increasing the fractional recurrence number P_{qw} on a line and higher dimension. If the particle wave packet is evolved in a *position* Hilbert space \mathcal{H}_P with the edges that permits the wave packet to escape, the fractional recurrence nature of the QW does not allow the QW to be completely transient. Therefore, the fractional transient nature of the QW is seen to complement the fractional recurrence nature.

V. CONCLUSION

In summary we show that, as long as the wave packet spread in position space interfering, forming mini wave packets during the evolution of the QW process, it fails to satisfy the complete recurrence theorem. We show this by analyzing the dynamics of quantum walk process on a line and on an n -cycle. We have shown that due to the revival of the fractional wave packet, a fractional recurrence can be seen during the QW process which can be characterized using the quantum Pólya number or fractional recurrence number.

Acknowledgement:

I thank Raymond Laflamme for his encouragement and support. I also acknowledge Mike and Ophelia Lezaridis for the financial support at IQC.

-
- [1] S. Chandrasekhar, Rev. Mod. Phys. **15**, 1 (1943).
 - [2] Luis Barreira, *Poincaré recurrence: old and new*, IVth International Congress on Mathematical Physics, World Scientific, pp. 415-422 (2006) ; U. Krengel, *Ergodic Theorems*, Walter de Gruyter, Berlin, New York, 1985.
 - [3] P. Bocchieri and A. Loinger, Phys. Rev. **107**, 337 (1957) ; L.S. Schulman, Phys. Rev. A **18** 2379 (1978).
 - [4] R.W. Robinett, Phys. Rep. **392**, 1 (2004).
 - [5] G.V. Riazanov, Sov. Phys. JETP **6** 1107 (1958).
 - [6] R. P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
 - [7] Y. Aharonov, L. Davidovich and N. Zagury, Phys. Rev. A **48**, 1687, (1993).
 - [8] A.M. Childs, R. Cleve, E. Deotto, E. Farhi, S. Gutmann and D.A. Spielman, in *Proceedings of the 35th ACM Symposium on Theory of Computing* (ACM Press, New York, 2003), p.59.
 - [9] N. Shenvi, J. Kempe and K. Birgitta Whaley, Phys. Rev. A **67**, 052307, (2003).
 - [10] A.M. Childs, J. Goldstone, Phys. Rev. A **70**, 022314, (2004).
 - [11] A. Ambainis, J. Kempe, and A. Rivosh, *Proceedings of ACM-SIAM Symp. on Discrete Algorithms (SODA)*, (AMC Press, New York, 2005), pp.1099-1108.
 - [12] C.M. Chandrashekar and R. Laflamme, Phys. Rev. A **78** 022314 (2008).

- [13] Takashi Oka, Norio Konno, Ryotaro Arita, and Hideo Aoki, *Phys. Rev. Lett.* **94** 100602 (2005).
- [14] Gregory S. Engel *et. al.*, *Nature*, **446**, 782-786 (2007).
- [15] M. Mohseni, P. Rebentrost, S. Lloyd, A. Aspuru-Guzik, *J. Chem. Phys.* **129**, 174106 (2008).
- [16] A. Ambainis, E. Bach, A. Nayak, A. Vishwanath and J. Watrous, *Proceeding of the 33rd ACM Symposium on Theory of Computing* (ACM Press, New York, 2001), p.60.
- [17] P. Révész. *Random walk in Random and non-random Environments* (World Scientific, Singapore, 1990).
- [18] G. Pólya, *Math. Ann.* **84** 149 (1921).
- [19] M. Stefanak, I. Jex, and T. Kiss, *Phys. Rev. Lett.* **100** 020501 (2008).
- [20] David A. Meyer, *J. Stat. Phys.* **85**, 551 (1996).
- [21] E. Farhi and S. Gutmann, *Phys. Rev. A* **58**, 915 (1998).
- [22] Fredrick Strauch, *Phys. Rev. A* **74**, 030310 (2006).
- [23] C.M. Chandrashekar, *Phys. Rev. A* **78**, 052309 (2008).
- [24] C.M. Chandrashekar, R. Srikanth, and R. Laflamme, *Phys. Rev. A* **77** 032326 (2008).
- [25] M. Stefanak, I. Jex, and T. Kiss, *Phys. Rev. A* **78**, 032306 (2008).
- [26] D. Aharonov, A. Ambainis, J. Kempe, and U. Vazirani, *Proceeding of the 33rd ACM Symposium on Theory of Computing* (ACM Press, New York, 2001) p.50.
- [27] Ben Tregenna, Will Flanagan, Rik Maile, and Viv Kendon, *New J. Phys.* **5**, 83 (2003).
- [28] N. Inui, Y. Konishi, and N. Konno, *Phys. Rev. A* **69** 052323 (2004).
- [29] Norio Inui, Norio Konno, and Etsuo Segawa, *Phys. Rev. E* **72** 056112 (2005).
- [30] During standard wave packet spreading, a wave packet which is initially Gaussian, spreads retaining the Gaussian shape causing increase in the full-width at half maxima (FWHM).
- [31] By choosing extreme value of θ (0 or $\pi/2$) in the quantum coin operation, quantum walk on a line can be evolved without constructive or distractive interference taking place.