VERY TWISTED STABLE MAPS

QILE CHEN, STEFFEN MARCUS, AND HENNING ÚLFARSSON

ABSTRACT. Let X be a smooth projective Deligne-Mumford stack over an algebraically closed field k of characteristic 0. In this paper we construct the moduli stack $\mathcal{K}_{q,n}(X,\beta)$ of very twisted stable maps, extending the notion of twisted stable maps from [\[AV02\]](#page-16-0) to allow for generic stabilizers on the source curves. We also consider the Gromov-Witten theory given by this construction.

CONTENTS

1. INTRODUCTION

Throughout, we let X be a smooth, projective Deligne-Mumford stack (DM-stack from now on) over an algebraically closed field k of characteristic 0.

The Gromov-Witten theory of orbifolds was first introduced in the symplectic setting in [\[CR05\]](#page-16-2). This was followed by an adaptation to the algebraic setting in [\[AGV02\]](#page-16-3) and [\[AGV08\]](#page-16-4), where the Gromov-Witten theory of DM-stacks was developed, and heavy use is made of the moduli stack of twisted stable maps into X, denoted $\mathcal{K}_{q,n}(X,\beta)$. This stack was constructed in [\[AV02\]](#page-16-0) and is the necessary analogue of Kontsevich's moduli stack of stable maps for smooth projective varieties when replacing the variety with a DM-stack. The main purpose of this note is to provide a further extension of these spaces by allowing generic stabilizers on the source curves of the twisted stable maps.

Following [\[AGV08\]](#page-16-4), we have a diagram:

$$
\Sigma_i^{\mathcal{C}} \subset \mathcal{C} \xrightarrow{f} X
$$

$$
\downarrow \qquad \qquad \downarrow
$$

$$
\mathcal{K}_{g,n}(X,\beta)
$$

where $\Sigma_i^{\mathcal{C}} \subset \mathcal{C} \longrightarrow X$ is the universal *n*-pointed twisted stable map. This gives rise to evaluation maps $e_i: \mathcal{K}_{g,n}(X,\beta) \longrightarrow \mathcal{I}_{\mu}(X)$ mapping into the rigidified cyclotomic inertia

Date: October 29, 2018.

stack $\overline{\mathcal{I}}_{\mu}(X)$ of X. If $\gamma_1,\ldots,\gamma_n\in A^*(\overline{\mathcal{I}}_{\mu}(X))_{\mathbb{Q}}$ then the Gromov-Witten numbers are defined to be

$$
\langle \gamma_1, \ldots, \gamma_n \rangle_{g,\beta}^X = \int\limits_{\left[\mathcal{K}_{g,n}(X,\beta) \right]^{vir}} \prod_i e_i^* \gamma_i
$$

where $[\mathcal{K}_{g,n}(X,\beta)]^{\text{vir}}$ is the virtual fundamental class of $\mathcal{K}_{g,n}(X,\beta)$ as in [\[BF97\]](#page-16-5).

In the case when X is a 3-dimensional Calabi-Yau *variety*, we also have Donaldson-Thomas theory (originating from [\[Tho00\]](#page-16-6),[\[DT98\]](#page-16-7)) which in contrast to Gromov-Witten theory gives invariants by counting sheaves. We get the following diagram:

$$
\mathcal{Y} \subset X \times \mathcal{H}\mathrm{ilb}_{\chi,\beta}(X)
$$

$$
\downarrow
$$

$$
\mathcal{H}\mathrm{ilb}_{\chi,\beta}(X)
$$

where the virtual dimension of $\mathcal{H}\text{ilb}_{\chi,\beta}(X)$ is zero as in [\[BF97\]](#page-16-5). The conjectural Donaldson-Thomas / Gromov-Witten correspondence of [\[MNOP06\]](#page-16-8) predicts a correspondence (in the case $n=0$:

(Z [Kg,0(X,β)]vir 1 = GW(g, β)) (Z [Hilbχ,β(X)]vir 1 = DT(χ, β)) .

This correspondence is manifested in a subtle relationship between generating functions of the invariants. One wishes to discover a similar correspondence when X is a 3-dimensional Calabi-Yau *orbifold*. The Hilbert scheme of a stack was constructed by Olsson and Starr [\[OS03\]](#page-16-9), but notice that in general it contains components corresponding to substacks with nontrivial generic stabilizers. In our definition of $\mathcal{K}_{g,n}(X,\beta)$ the twisted curves used as the sources of our maps have stacky structure only at the nodes and marked points. To allow for the above correspondence, one needs to extend the notion of the space of twisted stable maps as defined in [\[AV02\]](#page-16-0) and [\[AGV08\]](#page-16-4) so that our twisted curves have generic stabilizers. We approach this problem in three steps:

(1) Construct the stack \mathcal{G}_X of étale gerbes in X as a rigidification of the stack \mathcal{S}_X of subgroups of the inertia stack $\mathcal{I}(X)$. We exhibit \mathcal{S}_X as the universal gerbe sitting over \mathcal{G}_X , giving a diagram:

$$
\mathcal{S}_X \xrightarrow{\phi} X
$$

$$
\downarrow^{\alpha}
$$

$$
\mathcal{G}_X.
$$

This is done in section [2.](#page-2-0)

(2) Define the moduli stack of very twisted stable maps $\widetilde{\mathcal{K}}_{g,n}(X,\beta)$ by setting

$$
\widetilde{\mathcal{K}}_{g,n}(X,\beta):=\coprod_{\beta_{\mathcal{G}}}\mathcal{K}_{g,n}(\mathcal{G}_X,\beta_{\mathcal{G}})
$$

where the disjoint union is taken over all curve classes $\beta_{\mathcal{G}} \in H^*(\mathcal{G}_X)$ such that $\phi_* \alpha^* \beta_{\mathcal{G}} = \beta$. By pulling back, we see that each $\mathcal{K}_{g,n}(\mathcal{G}_X, \beta_{\mathcal{G}})$ has two different universal objects sitting above it, one giving twisted stable maps into \mathcal{G}_X and the other

giving "very twisted" stable maps into X . Through our disjoint union, we get the two corresponding universal objects sitting above $\widetilde{\mathcal{K}}_{g,n}(X,\beta)$. This is done in section [3.](#page-11-0)

(3) Relate the Gromov-Witten theory of X given by the two different universal objects sitting above $\widetilde{\mathcal{K}}_{g,n}(X,\beta) = \coprod_{\beta_{\mathcal{G}}}\mathcal{K}_{g,n}(\mathcal{G}_X,\beta_{\mathcal{G}})$ and show that they give the same invariants. This is done in section [4.](#page-13-0)

1.1. Acknowledgements. We would like to thank Jim Bryan and Rahul Pandharipande for suggesting the problem, and Angelo Vistoli for his great help in understanding the constructions of section [2.](#page-2-0) We would also like to thank our advisor, Dan Abramovich, for his continuous support and encouragement.

Finally, Ezra Getzler has suggested earlier that very twisted curves should be of independent interest; this note may give further motivation for following this suggestion.

2. CONSTRUCTING THE STACKS \mathcal{S}_X and \mathcal{G}_X

In this section we construct DM-stacks \mathcal{S}_X and \mathcal{G}_X and show that \mathcal{G}_X is presented as the rigidification of \mathcal{S}_X along a group scheme. Our construction follows closely the construction of $\mathcal{I}_{\mu}(X)$ and its rigidification, from [\[AGV08\]](#page-16-4).

2.1. The stack of subgroups of the inertia stack. Recall that X is a smooth, projective DM-stack over an algebraically closed field k of characteristic 0. We begin by defining the stack \mathcal{S}_X of subgroups of the inertia stack $\mathcal{I}(X)$.

Definition 2.1.1. We define a category S_X , fibered over the category of schemes, as follows:

- (a) An object of $\mathcal{S}_X(T)$ consists of a pair $(\xi, \alpha : G \hookrightarrow \mathcal{A}ut_T(\xi))$, where $\xi \in X(T)$, the arrow α is an injective morphism of group schemes, and G is finite and étale over T.
- (b) An arrow sitting over $T \longrightarrow T'$ from $(\xi, \alpha) \in \mathcal{S}_X(T)$ to $(\xi', \alpha') \in \mathcal{S}_X(T')$ is a morphism $F: \xi \longrightarrow \xi'$ making the following diagram commutative:

where $Aut_T(\xi) \longrightarrow Aut_{T'}(\xi')$ is the morphism induced by F and the diagram is cartesian.

This definition follows Definition 3.1.1 in [\[AGV08\]](#page-16-4) except we are allowing our subgroup to vary.

There are several things that need mention or proof.

Remark. Notice that there is an obvious morphism (of fibered categories) $S_X \longrightarrow X$ which sends (ξ, α) to ξ .

Proposition 2.1.2. \mathcal{S}_X *is fibered in groupoids over the category of schemes.*

Proof. By Proposition 3.22 in Part 1 of [\[FGI](#page-16-10)+05] S_X is fibered in groupoids if and only if

- every arrow is cartesian; and
- given ξ' over T' and $f: T \to T'$ there exists an arrow $\phi: \xi \to \xi'$ over f.

The first condition is automatically satisfied since every arrow in S_X is an arrow in X which is fibered in groupoids. For the second we set $\xi = f^*\xi'$ and $G = f^*G'$, so α is automatically defined. We now clearly have an arrow $(\xi, \alpha) \to (\xi', \alpha')$ making the necessary diagram \Box commute.

Proposition 2.1.3. *The category* S_X *is a DM-stack, and the functor* $S_X \longrightarrow X$ *is representable and finite.*

Proof. We prove that the functor $S_X \longrightarrow X$ is representable and finite, which implies that \mathcal{S}_X is a DM-stack. The inertia stack $\mathcal{I}(X) \to X$ is finite, unramified and representable, since X is assumed to be a separated DM-stack. We can therefore let M be the maximal degree of a fiber of $\mathcal{I}(X)$. We begin by showing that the relative Hilbert scheme $\mathcal{H}ilb(\mathcal{I}(X)/X)$ is unramified and finite:

- The fiber over a geometric point of X is unramified since a subscheme of a finite étale scheme has only trivial infinitesimal deformations.
- The relevant Hilbert polynomials here are just integers since the schemes in question are all finite. These integers are bounded above by M , so there are only finitely many of them. This together with the fact that each component is proper implies finiteness.

The relative Hilbert scheme is representable by definition. The last part we need is to notice that S_X is an open and a closed subscheme in $\mathcal{H}ilb(\mathcal{I}(X)/X)$ since the condition on a finite subset to be a group is an open and closed condition.

Proposition 2.1.4. (1) *When X is smooth,* S_X *is smooth as well.* (2) When X is proper, S_X is proper as well.

Proof. We start with (1). Let x be a geometric point of X and G_x its stabilizer group. We know from [\[AOV08\]](#page-16-11) that we can view X in a local chart around x as $[U/G_x]$. S_X has a local chart $[U^H/N(H)]$, where $H \subset G_x$ is a subgroup and we quotient out by the normalizer subgroup $N(H)$. To prove that this chart is smooth note that $T_{U^H} = (T_U)^H$ and that $\dim(T_U)^H = \dim(T_U^*)^H$ and T_V^* is generated by

$$
\overline{y_1},\ldots,\overline{y_d},\overline{y_{d+1}},\ldots,\overline{y_n},
$$

where $\overline{y_1}, \ldots, \overline{y_d}$ are the invariant generators. Since H is reductive we have a section of the projection map $m \to m/m^2$ which allows us to lift all the generators to

$$
y_1,\ldots,y_d,y_{d+1},\ldots,y_n,
$$

in such a manner that y_1, \ldots, y_d are still the invariant generators and y_{d+1}, \ldots, y_n span a finite-dimensional representation. Let J be the ideal generated by y_{d+1}, \ldots, y_n . Then the quotient by that ideal corresponds to an invariant subscheme, which is smooth since formally it is generated by y_1, \ldots, y_d . This proves that the dimension of U^H is at least, and thus equal, to dim T_{U^H} .

(2) follows from the proof of Proposition [2.1.3,](#page-3-0) since S_X is a closed subscheme of $\mathcal{H}ilb_X(\mathcal{I}(X)),$ which is proper over X .

2.2. Alternative description of S_X . It will be useful to provide another, less obvious, description of S_X . We start with a definition.

Definition 2.2.1. An *étale gerbe* over a scheme S is an DM-stack A over S such that

- (1) there exists an étale covering $S_i \longrightarrow S$ such that each $\mathcal{A}(S_i)$ is not empty.
- (2) given two objects a and b of $\mathcal{A}(T)$, where T is an S-scheme, there exists a covering $T_i \longrightarrow T$ such that the pullbacks a_{T_i} and b_{T_i} are isomorphic in $F(T_i)$.

More generally, a stack F over a stack X is an étale gerbe if for any morphism $V \longrightarrow \mathcal{X}$ with V a scheme, the pull-back $\mathcal{F}_V \longrightarrow V$ along $V \longrightarrow \mathcal{X}$ is an étale gerbe.

Definition 2.2.2. We define a 2-category S'_X , fibered over the category of schemes, as follows:

(a) An object of $S_X'(T)$ consist of representable morphisms $\phi : A \longrightarrow X$, where A is an étale gerbe over T, with a section $\tau : T \longrightarrow A$.

(b) A 1-arrow $(F, \rho) : (\mathcal{A}, \phi) \longrightarrow (\mathcal{A}', \phi')$ consists of a morphism $F : \mathcal{A} \longrightarrow \mathcal{A}'$ over some $f: T \longrightarrow T'$ making a cartesian square, and a natural transformation $\rho: \phi \Rightarrow \phi' \circ F$ making the following diagram commutative:

(c) A 2-arrow $(F, \rho) \longrightarrow (F_1, \rho_1)$ is an natural transformation $\sigma : F \Rightarrow F_1$ giving an equivalence, and compatible with ρ and ρ_1 in the sense that the following diagram is commutative:

- *Remark.* (1) By Lemma 3.21 in [\[LMB00\]](#page-16-12), the section τ gives a neutral section, hence the gerbe A is isomorphic to $\mathcal{B}_T G_\tau$ where $G_\tau = \mathcal{A}ut_{T,\mathcal{A}}(\tau,\tau)$ is étale over T. The section τ corresponds to the trivial G_{τ} -torsor $G_{\tau} \longrightarrow T$.
	- (2) Since T gives a moduli space of A, the arrow $A \rightarrow T$ is proper. This implies that the group scheme G_{τ} is finite over T.

Lemma 2.2.3. The 2-category S'_X is equivalent to a category.

Proof. Lemma 3.3.3 in [\[AGV08\]](#page-16-4) shows that the isomorphism group of a 1-arrow in S'_X is trivial. Since all 2-arrows are isomorphisms, the result follows. **Definition 2.2.4.** By abuse of notation, we denote by \mathcal{S}'_X the 1-category associated to the above 2-category. Our arrows become 2-isomorphism classes of 1-arrows.

Definition 2.2.5. We define a morphism of fibered categories $\mathcal{S}'_X \longrightarrow \mathcal{S}_X$ as follows:

(a) Given an object

$$
\begin{array}{c}\nA \xrightarrow{\phi} X \\
\downarrow \\
T\n\end{array}
$$

with a section $\tau : T \longrightarrow \mathcal{A}$, we obtain a pair (ξ, α) as follows: ξ is obtained by composing ϕ with τ . Note that τ gives an element $\zeta_{\tau,T} \in \mathcal{A}(T)$. Then α is the associated map of automorphisms

$$
G = \mathcal{A}ut_T(\zeta_{\tau,T}) \longrightarrow \mathcal{A}ut_T(\xi),
$$

which is injective since ϕ is representable.

(b) Given an arrow ρ as above, we obtain an arrow $F : \phi(\zeta_{\tau,T}) \longrightarrow \phi'(\zeta_{\tau',T'})$ by completing the following diagram:

The proof of the following proposition is almost the same as Proposition 3.2.3 in [\[AGV08\]](#page-16-4). For completeness we rewrite it in our case.

Proposition 2.2.6. The morphism $S'_X \longrightarrow S_X$ is an equivalence of fibered categories.

Proof. By Proposition 3.36 in Part 1 of [\[FGI](#page-16-10)⁺05], it is enough to show that the induced functor on the fiber $S_X'(T) \longrightarrow S_X(T)$ is an equivalence for any given scheme T.

(1) The functor is faithful.

Assume we are given two elements in $\mathcal{S}'_X(T)$ and a 2-arrow $\rho : \phi \Rightarrow \phi' \circ F$ making the following diagram commutative:

We need to show that for any T-scheme U and any $G_{\tau} \times_T U$ -torsor $P \longrightarrow U$, the arrow $\rho(P \to U) : \phi(P \to U) \to \phi' \circ F(P \to U)$ is uniquely determined by $\rho(G_{\tau} \longrightarrow T) : \phi(G_{\tau} \longrightarrow T) \longrightarrow \phi' \circ F(G_{\tau} \longrightarrow T)$. Note that $G_{\tau} \longrightarrow T$ is the trivial torsor given by the section τ .

Let $\{U_i \longrightarrow U\}$ be an étale covering such that the pull-backs $P_i \longrightarrow U_i$ are trivial (i.e., $P_i \simeq (G_\tau \times_T U) \times_U U_i \simeq G_\tau \times_T U_i$). Since X is a DM-stack, the arrow $\phi(P \longrightarrow U) \longrightarrow \phi' \circ F(P \longrightarrow U)$ is determined by $\phi(P_i \longrightarrow U_i) \longrightarrow \phi' \circ F(P_i \longrightarrow U_i)$. Hence, we can assume that $P \longrightarrow U$ is trivial. But then the cartesian arrow $P \longrightarrow G_{\tau}$ and the induced diagram

$$
\phi(P \longrightarrow U) \longrightarrow \phi' \circ F(P \longrightarrow U)
$$

\n
$$
\downarrow \qquad \qquad \downarrow
$$

\n
$$
\phi(G_{\tau} \longrightarrow T) \longrightarrow \phi' \circ F(G_{\tau} \longrightarrow T)
$$

proves what we want.

(2) The functor is fully faithful.

Assume we are given an arrow $\beta : \phi(G_{\tau} \to T) \to \phi' \circ F(G_{\tau} \to T)$ in $X(T)$ which is compatible with the action of G_{τ} and $G_{\tau'}$. First, consider the trivial torsor $P \simeq G_\tau \times_T U \longrightarrow U$, which induces a cartesian arrow $P \longrightarrow G_\tau$. By the definition of a cartesian arrow, there is a unique arrow $\rho(P \longrightarrow U)$ that we can insert in the diagram

$$
\phi(P \longrightarrow U) - \xrightarrow{\rho(P \longrightarrow U)} \phi' \circ F(P \longrightarrow U)
$$

$$
\downarrow \qquad \qquad \downarrow
$$

$$
\phi(G_{\tau} \longrightarrow T) \xrightarrow{\beta} \phi' \circ F(G_{\tau} \longrightarrow T)
$$

making it commutative. This arrow $\rho(P \longrightarrow U)$ is independent of the chosen trivialization, since β compatible with the group action.

Now, if the G_{τ} -torsor $P \longrightarrow U$ is not necessarily trivial, choose a covering $U_i \longrightarrow U$ such that the pull-backs $P_i \longrightarrow U_i$ are trivial. We have arrows $\phi(P_i \longrightarrow U_i) \longrightarrow$ $\phi' \circ F(P_i \longrightarrow U_i)$ in $X(U_i)$, and their pullbacks to $U_i \times_U U_j$ coincide; hence they glue together to given an arrow $\rho(P \to U) : \phi(P \to U) \to \phi' \circ F(P \to U)$. It is not hard to see that $\rho(P \longrightarrow U)$ does not depend on the choice of covering, and defines a 2-arrow $\phi \longrightarrow \phi' \circ F$ whose image in $\mathcal{S}_X(T)$ coincides with β .

(3) The functor is essentially surjective.

Note that since G_{τ} is étale over T, and $\mathcal{A} \simeq \mathcal{B}_T G_{\tau}$, then $T \longrightarrow \mathcal{A}$ gives an atlas of A. Given an object $(\xi, \alpha : G \hookrightarrow Aut_{T,X}(\xi))$ of $\mathcal{S}_X(T)$, we need to construct $\phi : \mathcal{B}_T G \longrightarrow X$, whose image in $\mathcal{S}_X(T)$ is isomorphic to (ξ, α) .

Let $P \longrightarrow U$ be a $G \times_T U$ -torsor, where U is a T-scheme. The morphism $G \times_T U$ $P \longrightarrow P \times_{U} P$ given by $(g, p) \mapsto (gp, p)$, is an isomorphism. The pull-back of ξ to $P \times_U P \simeq G \times_T P$ via the first and the second projection coincide with the pull-back $\xi_{G\times_T P}$, and the isomorphism is given by the projection $G\times_T P\longrightarrow G$. These give descent data for ξ_P along the étale covering $P \longrightarrow U$. The descent data are effective and define an object η of $X(U)$, so we assigned to every object of $\mathcal{B}_T G(U)$ an object of $X(U)$. This extends to a morphism of fibered categories $\phi : \mathcal{B}_T G \longrightarrow X$.

Now let $G \longrightarrow T$ be the trivial G-torsor over T. We claim that $\eta := \phi(G \longrightarrow T)$ of $X(T)$ is isomorphic to ξ . In fact, the object with descent data defining ξ is ξ_G , with the descent data given by the identity on $\xi_{G\times G}$. Those descend to $\eta \simeq \xi$ in $X(T)$. Hence the image of ϕ in $\mathcal{S}_X(T)$ is isomorphic to (ξ, G) , as we wanted.

 \Box

2.3. The stack of gerbes in X. In the following we introduce a stack \mathcal{G}_X which is closely related to S_X and will play an important role in the construction of the moduli stack of very twisted stable maps.

Definition 2.3.1. Define as before a 2-category \mathcal{G}_X with a functor to the category of schemes as follows:

- (a) An object over a scheme T is a pair (\mathcal{A}, ϕ) where A is an étale gerbe over T and $\mathcal{A} \longrightarrow X$ is a representable morphism.
- (b) A 1-arrow $(F, \rho) : (\mathcal{A}, \phi) \longrightarrow (\mathcal{A}', \phi')$ consists of a morphism $F : \mathcal{A} \longrightarrow \mathcal{A}'$ over some $f: T \longrightarrow T'$ making a cartesian square, and a natural transformation $\rho: \phi \Rightarrow \phi' \circ F$ making the following diagram commutative:

(c) A 2-arrow $(F, \rho) \longrightarrow (F_1, \rho_1)$ is a natural transformation $\sigma : F \Rightarrow F_1$ giving an equivalence, and compatible with ρ and ρ_1 in the sense that the following diagram is commutative:

Definition 2.3.2. Exactly as above, the 2-category \mathcal{G}_X is equivalent to a 1-category. By abuse of notation, we denote by \mathcal{G}_X its associated 1-category.

It will follow from our results below on rigidification that \mathcal{G}_X is a DM-stack.

Remark. There is a tautological functor of categories $S_X \longrightarrow \mathcal{G}_X$, sending the pair (ϕ, τ) in $\mathcal{S}_X(T)$ to the representable morphism $\phi : \mathcal{A} \longrightarrow X$ in $\mathcal{G}_X(T)$.

2.4. Rigidification. Rigidification of algebraic stacks was first defined and used in [\[ACV03\]](#page-16-13), [\[AOV08\]](#page-16-11) and [\[Rom05\]](#page-16-14). It was foreseen in [\[Art74\]](#page-16-15).

In this section, we discuss the rigidification of DM-stacks, and prove a proposition that the morphism $S_X \longrightarrow \mathcal{G}_X$ defined in the above remark gives a rigidification of S_X along an appropriate relative group scheme, which all our constructions rely on.

We begin by restating the theorem in [\[AOV08\]](#page-16-11) on the existence of the rigidification of an algebraic stack. We stick to the case of DM-stacks since this is all we will need for our purposes. It follows from Theorem 5.1.5 in [\[ACV03\]](#page-16-13) that the rigidification of a DM-stack is a DM-stack, and the quotient map is étale.

Let S be a scheme, $X \longrightarrow S$ a DM-stack of finite type over (Sch/S) . Suppose $G \subseteq \mathcal{I}(X)$, where $\mathcal{I}(X)$ is the inertia stack of X, and G is a subgroup stack of $\mathcal{I}(X)$ of finite type and \forall tale over X.

Theorem 2.4.1 ([\[AOV08\]](#page-16-11) Theorem A.1). *There exists a finite type DM-stack* $X/\!\!/ G$ *over* S *with a morphism* $\rho: X \longrightarrow X \mathbf{f}$ *G satisfying the following properties:*

- (1) X *is an étale gerbe over* $X/\!\!/ G$.
- (2) *For each object* ξ *of* X(T)*, the morphism of group schemes*

$$
\rho: \mathcal{A}ut_T(\xi) \longrightarrow \mathcal{A}ut_T(\rho(\xi))
$$

is surjective with kernel G_{ξ} .

Furthermore, if G *is finite over* X *then* ρ *is proper; while since* G *is étale,* ρ *is also étale. These properties characterize* $X \mathcal{J} G$ *uniquely up to equivalence.*

*Remark.*From ([\[AOV08\]](#page-16-11) Theorem A.1) we already know that $X/\!\!/ G$ is an algebraic stack. To show that it is also a DM-stack, we only need an étale atlas. But since the arrow $\rho: X \longrightarrow X/\!\!/ \!\!/ G$ is *étale* and surjective, the atlas of $X/\!\!/ \!\!/ G$ is obtained from that of X.

We add to this a proposition about pulling back rigidifications along representable morphisms.

Proposition 2.4.2. *Let* X *and* G *be defined as above, and* Y *a DM-stack of finite type over* (Sch/S) *. Given any representable morphism* $Y \longrightarrow X \mathbf{f}$ *G, the pullback*

$$
\tilde{\rho}: Y \times_{X \not\parallel G} X \longrightarrow Y
$$

of ρ *satisfies the properties in the above theorem, hence is a rigidification of* $Y \times_{X/\mathbb{C}} X$ *along the pull-back* $G_Y := (Y \times_{X \mid G} X) \times_X G$.

Proof. First, any $A \longrightarrow T$ which is a pullback of $Y \times_{X/\!\!/ G} X \longrightarrow Y$ along $T \longrightarrow Y$ can be viewed as a pullback of $X \longrightarrow X/\!\!/ G$ along the composition $T \longrightarrow Y \longrightarrow X/\!\!/ G$. Hence, by Definition [2.2.1,](#page-4-0) $\tilde{\rho}: Y \times_{X/\!\!/ G} X \longrightarrow Y$ is an étale gerbe.

Second, given any $T \longrightarrow Y \times_{X/G} X$, where T is any scheme, consider the following cartesian diagram:

where $\xi = \psi \circ (\zeta, \xi, \mu) \in X(T)$, $\zeta = \tilde{\rho} \circ (\zeta, \xi, \mu) \in Y(T)$ and $\mu : \phi \circ \zeta \simeq \rho \circ \xi$ is an isomorphism in $X/\!\!/ \!\!/ G(T)$. Note that $G_T \hookrightarrow \mathcal{A}ut_T(\xi)$ gives the action of G_T on ξ , and lies in the kernel of $\rho: \mathcal{A}ut_T(\xi) \longrightarrow \mathcal{A}ut_T(\rho(\xi))$. There is an natural action of G_T on (ζ, ξ, μ) , which acts on ξ and fixes ζ and μ . Hence there is a injection $G_T \hookrightarrow \mathcal{A}ut_T((\zeta,\xi,\mu))$. Since G_T is étale and of finite type over T, this implies that G_Y is a subgroup stack of the inertia stack $I_{Y\times_{X/\!\!/ G}X}$, and étale and finite type over $Y \times_{X/\!\!/ G} X$.

Finally, consider the following diagram:

$$
id \longrightarrow G_T \longrightarrow Aut_T(\xi) \longrightarrow Aut_T(\rho(\xi)) \longrightarrow id
$$

\n
$$
\simeq \int_{G_T}^{\uparrow} f \int_{G_T}^{\uparrow} g \int_{\mathcal{A}}^{g} g
$$

\n
$$
G_T \longrightarrow Aut_T((\zeta, \xi, \mu)) \longrightarrow Aut_T(\tilde{\rho}(\zeta, \xi, \mu)).
$$

Here, f and q are injective since the corresponding arrow is representable and the first row is exact. An easy diagram chase shows that the second row is also exact, which implies that G_T is the kernel of $\mathcal{A}ut_T((\zeta,\xi,\mu)) \longrightarrow \mathcal{A}ut_T(\tilde{\rho}(\zeta,\xi,\mu)).$

By the uniqueness of the theorem above, $\tilde{\rho}: Y \times_{X/\!\!/ G} X \longrightarrow Y$ is the rigidification along $G_Y \longrightarrow Y.$

Now let's consider the stack S_X . For each object (ξ, G) of $S_X(T)$ over a scheme T, we associate a subgroup scheme $G \subset Aut_T((\xi, G))$. Note that G is finite and étale over T. Take $(\xi, G) \longrightarrow (\xi', G)$ in \mathcal{S}_X over $T \longrightarrow T'$. By the definition of \mathcal{S}_X , G is the pullback of G' along $T \longrightarrow T'$. By Appendix A of [\[AOV08\]](#page-16-11), there is a unique étale finite subgroup stack, denoted $\Gamma_X \subset I_{\mathcal{S}_X}$, where $I_{\mathcal{S}_X}$ is the inertia stack of \mathcal{S}_X , such that for any object (ξ, G) of $\mathcal{S}_X(T)$ the pullback of Γ_X to T coincides with G.

The following is the main result of this section.

Proposition 2.4.3. *There is an equivalence* $\mathcal{S}_X/\!\!\Gamma_X \longrightarrow \mathcal{G}_X$ *of fibered categories, hence we can view* \mathcal{G}_X *as a rigidification of* \mathcal{S}_X *along* $\Gamma_X \longrightarrow \mathcal{S}_X$ *.*

Proof. Given any scheme T and an arrow $T \longrightarrow S_X/\!\!/ \Gamma_X$, consider the following cartesian diagram:

where A is the gerbe obtained by pulling back along $T \longrightarrow S_X/\!\!/ \Gamma_X$. Also, note that the arrows $A \longrightarrow S_X$ and $S_X \longrightarrow X$ are representable, so their composition $A \longrightarrow S_X \longrightarrow X$ is also representable. Since Γ_X is finite étale over \mathcal{S}_X , by the above theorem the arrow $\mathcal{S}_X \longrightarrow \mathcal{S}_X/\!\!/ \Gamma_X$ is étale and proper, hence A is an étale proper gerbe over T with a representable arrow $A \longrightarrow X$. This construction extends to an arrow $S_X/\!\!\!\!\int \Gamma_X \longrightarrow \mathcal{G}_X$ of fibered categories.

To show that the arrow $\mathcal{S}_X/\!\!\!\Gamma_X \longrightarrow \mathcal{G}_X$ constructed above is an equivalence, we construct its inverse. Given any object in $\mathcal{G}_X(T)$

consider a T-scheme U, and an arrow $\zeta: U \longrightarrow \mathcal{A}$. Denoting $\phi(\zeta)$ by ξ for simplicity, since ϕ is representable we have an injection $\alpha : \mathcal{A}ut_{U,\mathcal{A}}(\zeta) \hookrightarrow \mathcal{A}ut_{U,X}(\zeta)$. Hence we have a pair (ξ, α) which is an object in $\mathcal{S}_X(U)$. Note that this construction gives an arrow of stacks $\psi: \mathcal{A} \longrightarrow \mathcal{S}_X$. From this construction, it is easy to see that the composition $\mathcal{A} \longrightarrow \mathcal{S}_X \longrightarrow X$ is the representable arrow ϕ . By Lemma 1.3 in [\[Vis89\]](#page-16-16), ψ is also representable.

Next we will construct an arrow $T \longrightarrow S_X/\!\!\!\Gamma_X$ and show that $\mathcal{A} \longrightarrow T$ is the pullback of $\mathcal{S}_X \longrightarrow \mathcal{S}_X /\!\!/ \Gamma_X$ along this arrow, proving our result.

According to the definition of a gerbe, we can choose a covering $\{T_i\}$ of T such that there exists $\zeta_i \in \mathcal{A}(T_i)$ non-zero and isomorphisms $f_{ij}: \zeta_i|_{T_{ij}} \simeq \zeta_j|_{T_{ij}}$, where $T_{ij} \simeq T_i \times_T T_j$ and $f_{ji} = f_{ij}^{-1}$. Note that $f_{ki} \circ f_{jk} \circ f_{ij} = g_i \in Aut_{T_i,A}(\zeta_i)|_{T_{ij}}$. Since g_i might not be the identity, $\{f_{ij}\}\right\}$ might not satisfy the cocycle condition, becoming a obstruction to glue the ζ_i . Let us denote $\rho \circ \psi \circ \zeta_i$ by ξ_i , and $\rho \circ \psi \circ f_{ij}$ by f'_{ij} . Note that $f'_{ij} : \xi_i |_{T_{ij}} \simeq \xi_j |_{T_{ij}}$ gives an isomorphism. Now we want to glue $\{\xi_i\}$, and for this we only need to show that $f'_{ki} \circ f'_{jk} \circ f'_{ij} = \rho \circ \psi \circ g_i = id$. But we have the following exact sequence:

$$
1 \longrightarrow \mathcal{A}ut_{T_i,\mathcal{A}}(\zeta_i)|_{T_{ij}} \stackrel{\psi}{\longrightarrow} \mathcal{A}ut_{T_i,\mathcal{S}_X}(\psi \circ \zeta_i)|_{Tij} \stackrel{\rho}{\longrightarrow} \mathcal{A}ut_{T_i,\mathcal{S}_X/\hspace{-0.1cm}/\hspace{-0.1cm}T_X}(\xi_i)|_{T_{ij}} \longrightarrow 1
$$

which shows that $\rho \circ \psi \circ g_i = id \in Aut_{T_i, \mathcal{S}_X/\!\!/ \Gamma_X}(\xi_i)|_{T_{ij}}$, hence we can glue $\{\xi_i\}$ and get an arrow $\sigma: T \longrightarrow \mathcal{S}_X/\!\!/ \Gamma_X$. The difference of two choices of $\{f_{ij}\}\$ is an element in $\mathcal{A}ut_{T_i,\mathcal{A}}(\zeta_i)|_{T_i}$, and becomes the identity when composed with $\rho \circ \psi$. Hence the arrow $\sigma : T \longrightarrow S_X/\!\!\!\!\!\int \Gamma_X$ is unique.

The last step is to show that the following diagram is cartesian:

This is equivalent to showing that the following is cartesian:

where $A_i \longrightarrow T_i$ is obtained by pulling back $A \longrightarrow T$ along the covering $T_i \longrightarrow T$. We also note that π_i has a section $\zeta'_i: T_i \longrightarrow \mathcal{A}_i$, which is obtained from the following diagram:

Now, assume we are given a T_i -scheme U and the diagram

where $\xi = (\xi_U, G_U) \in \mathcal{S}_X(U)$.

We need to show that $h = \zeta_i' \circ f$ is the unique arrow making the diagram commutative. Note that $\pi_i \circ h = \pi_i \circ \zeta_i \circ f = id_{T_i} \circ f = f$. So we only need to show that $\xi = \psi \circ h$. But by our construction, $\sigma \circ f = \rho \circ \psi \circ \zeta_i' \circ f = \rho \circ \psi \circ h$, so $\rho \circ \psi \circ h = \rho \circ \xi$. Since our construction is by gluing étale locally, we can restrict to a small étale neighborhood, and assume that ρ is an isomorphism, hence $\psi \circ h = \xi$. Since both π_i and ζ'_i are étale and surjective, h is unique by our construction.

- *Remark.* (1) In the proof of this proposition, we actually proved that every pair (A, ψ) over a scheme T, where $\mathcal A$ is an étale gerbe over T and $\psi : \mathcal A \longrightarrow \mathcal S_X$ is a representable arrow, corresponds to an arrow $T \longrightarrow \mathcal{S}_X/\!\!/ \Gamma_X$. Also, \mathcal{S}_X is the universal gerbe over $\mathcal{S}_X/\!\!/ \Gamma_X$.
	- (2) We can identify \mathcal{G}_X with $\mathcal{S}_X/\!\!/ \Gamma_X$. By Theorem [2.4.1,](#page-7-0) \mathcal{G}_X is a DM-stack.

3. Very twisted curves and their maps

Recall from [\[AGV08\]](#page-16-4) that a twisted stable map over a scheme T is given by a family of twisted curves $C \longrightarrow T$ and n gerbes $\Sigma_i \subset C$, with a representable morphism $f: C \longrightarrow X$. $\mathcal{K}_{q,n}(X,\beta)$ is the moduli stack parameterizing maps of this type with the image of f lying in the curve class $\beta \in H^*(X)$. Our goal is to extend the notion of stable maps into X by allowing the source twisted curve $\mathcal C$ to have generic stabilizers. This leads us to the following definitions:

Definition 3.0.4. A *very twisted curve* over a scheme T is an étale gerbe $\widetilde{\mathcal{C}} \longrightarrow \mathcal{C}$ over a twisted curve $\Sigma_i \subset \mathcal{C} \longrightarrow T$. We define a *very twisted stable map* to be a diagram of the form:

where $\tilde{C} \longrightarrow C \longrightarrow T$ is a very twisted curve, the markings $\tilde{\Sigma}_i \subset \tilde{C}$ are given by taking the preimage of Σ_i , and such that the diagram admits finitely many automorphisms (our stability condition). Stability is equivalent to the corresponding map $\mathcal{C} \longrightarrow \mathcal{G}_X$ being a twisted stable map, and \tilde{f} factoring through the projection $\tilde{C} \longrightarrow S_X$.

Remark. Notice that in our definitions we are using a single gerbe structure over the whole curve. We are not considering curves with different generic stabilizers over different components.

Given a curve class $\beta \in H^*(X)$, we want to define the moduli stack $\mathcal{K}_{g,n}(X,\beta)$ parametrizing very twisted stable maps with image class β . A map $\mathcal{C} \to \mathcal{G}_X$ from a twisted curve \mathcal{C} to our stack \mathcal{G}_X corresponds to a diagram of the form:

Where $\widetilde{\mathcal{C}} \to \mathcal{C}$ is an étale gerbe. As stated above, étale gerbes over twisted curves are our very twisted curves. Let $\{\beta_j\} \subset H^*(\mathcal{G}_{\mathcal{X}})$ be the collection of curve classes satisfying $\phi_* \alpha^* \beta_j = \beta$. We are led to consider, for each j, the diagram

Here \mathcal{C}_j is the universal twisted curve sitting above $\mathcal{K}_{g,n}(\mathcal{G}_X,\beta_j)$ and \mathcal{C}_j is the universal very twisted curve given by pulling back. The compositions:

give us a parametrization of very twisted stable maps

whose images lie in β and whose corresponding map $C \longrightarrow \mathcal{G}_X$ has image lying in β_j . We get our desired moduli space by taking a disjoint union over all j :

Definition 3.0.5. With X as above, let

$$
\widetilde{\mathcal{K}}_{g,n}(X,\beta) := \coprod_j \mathcal{K}_{g,n}(\mathcal{G}_X,\beta_j).
$$

This moduli space has all the nice properties we want given for free by the construction of $\mathcal{K}_{g,n}(X,\beta)$. Sitting above it are two universal objects, one corresponding to very twisted stable maps into X by taking \tilde{C} over each $\mathcal{K}_{g,n}(\mathcal{G}_X,\beta_{\mathcal{G}})$, and the other corresponding to twisted stable maps into \mathcal{G}_X by taking \mathcal{C} . We use the following notation to distinguish them:

$$
\begin{array}{ccc}\n\coprod_{j} C_{j} & \xrightarrow{f} & \mathcal{G}_{X} & \text{and} & \coprod_{j} \widetilde{C}_{j} & \xrightarrow{\tilde{f}} & X \\
\downarrow^{\pi} & & & \downarrow^{\tilde{\pi}} \\
\coprod_{j} \mathcal{K}_{g,n}(\mathcal{G}_{X}, \beta_{j}) & & & \widetilde{\mathcal{K}}_{g,n}(X, \beta).\n\end{array}
$$

4. Equality of the Gromov-Witten Theories

As discussed in section [3,](#page-11-0) we have moduli spaces with two different universal curves, each part of the diagram

In this section, we show that the construction of the virtual fundamental classes and the evaluation maps give the same answer for $\coprod_j \mathcal{K}_{g,n}(\mathcal{G}_X, \beta_j)$ and $\widetilde{\mathcal{K}}_{g,n}(X, \beta)$. The virtual fundamental classes are constructed by looking at the relative obstruction theories $(R\pi_* f^*T_{\mathcal{G}_X})^{\vee}$ and $(R\tilde{\pi}_*\tilde{f}^*T_X)^\vee$ over $\mathfrak{M}_{g,n}^{tw}$ as in [\[Beh97\]](#page-16-17), [\[BF97\]](#page-16-5) and [\[AGV08\]](#page-16-4). We simply need to show that $R\pi_* f^*T_{\mathcal{G}_X} \cong R\tilde{\pi}_* \tilde{f}^*T_X$. Since push-forward is functorial, it will be enough to show that $f^*T_{\mathcal{G}_X} \cong \delta_*\tilde{f}^*\tilde{T}_X$, for each j, in the diagram:

To do this, we prove an analogue to the tangent bundle lemma (Lemma 3.6.1) of [\[AGV08\]](#page-16-4). We follow their proof closely.

Lemma 4.0.6. Let S be a scheme, and $f : S \longrightarrow \mathcal{G}_X$ a morphism with its associated diagram

Then there is a canonical isomorphism $\delta_*(\widetilde{f}^*(T_X)) \cong f^*(T_{\mathcal{G}_X})$.

Proof. We showed above that by rigidification we have the universal gerbe $S_X \longrightarrow \mathcal{G}_X$. Consider the following diagram of smooth stacks:

Given $f: S \longrightarrow \mathcal{G}_X$ as in the statement of the lemma, we have the diagram

The morphism ω is flat and, by Lemma 2.3.4 in [\[AV02\]](#page-16-0), ω_* is exact on coherent sheaves, so for any locally free sheaf F on S_X we have $f^*\omega_*\mathcal{F} = \pi_*g^*\mathcal{F}$. All that is needed is to check that $\omega_* F^* T_X \cong T_{\mathcal{G}_X}$.

By applying ω_* to the natural morphism $T_{\mathcal{S}_X} \longrightarrow F^*T_X$, we get $\omega_*T_{\mathcal{S}_X} \longrightarrow \omega_*F^*T_X$. But ω : $S_X \longrightarrow \mathcal{G}_X$ is an étale gerbe, so $\omega_* T_{\mathcal{S}_X} \cong T_{\mathcal{G}_X}$. Through this isomorphism, we get a morphism $T_{\mathcal{G}_X} \longrightarrow \omega_* F^* T_X$. The claim is that this is an isomorphism. We show this by looking at the geometric points.

The fiber of a geometric point y of \mathcal{G}_X through ω can be identified with BH for some subgroup $H \subset G_x$. Here x is a geometric point of X with stabilizer group G_x . This lifts y to \mathcal{S}_X , mapping y to x. Let T be the pullback of T_X to our lift of y. There is a natural action of H on T, and the fiber of $\omega_* F^* T_X$ at y is given by the space of invariants T^H . We must show that we get an isomorphism of this fiber with the fiber of T_{G_X} at y.

We can view X in a local chart around x as $[U/G_x]$. Then \mathcal{S}_X has a local chart $[U^H/N(H)]$, where we quotient out by the normalizer subgroup $N(H)$. But $T_{U^H} = T_U^H$ and $S_X \longrightarrow \mathcal{G}_X$ is an étale gerbe, so we obtain our desired isomorphism $T_{G_X,y} \cong T$ H .

 $\textbf{Corollary 4.0.7.}$ The virtual fundamental classes $\left[\coprod_j \mathcal{K}_{g,n}(\mathcal{G}_X,\beta_j)\right]^{vir}$ and $\left[\widetilde{\mathcal{K}}_{g,n}(X,\beta)\right]^{vir}$ *are equal. In particular, for* $n = 0$ *we have equality of the Gromov-Witten invariants:*

$$
\int 1 = \int 1.
$$

$$
\left[\prod_j K_{g,n}(g_X,\beta_j)\right]^{vir} \left[\widetilde{K}_{g,n}(X,\beta)\right]^{vir}
$$

Proof. Immediate from the lemma. □

In general, our Gromov-Witten invariants will be defined by taking $\gamma_i \in A^*(\overline{\mathcal{I}}_\mu(\mathcal{G}_X))_{\mathbb{Q}},$ pulling back along the evaluation maps $e_i: \coprod_j \mathcal{K}_{g,n}(\mathcal{G}_X, \beta_j) \longrightarrow \overline{\mathcal{I}}_\mu(\mathcal{G}_X)$, and taking a product:

$$
\langle \gamma_1, \ldots, \gamma_n \rangle_{g, \beta_j}^X = \left(\prod_{i=1}^n e_i^* \gamma_i \right) \cap \left[\coprod_j \mathcal{K}_{g, n}(\mathcal{G}_X, \beta_j) \right]^{vir}
$$

.

Recall from [\[AGV08\]](#page-16-4) that $\overline{\mathcal{I}}_{\mu}(\mathcal{G}_X)$ is given by the disjoint union

$$
\overline{\mathcal{I}}_{\mu}(\mathcal{G}_X) := \coprod_r \overline{\mathcal{I}}_{\mu_r}(\mathcal{G}_X).
$$

Each component $\mathcal{I}_{\mu_r}(\mathcal{G}_X)$ can be reinterpreted as parametrizing diagrams of the form

where $\Sigma \longrightarrow T$ is a gerbe over a scheme T banded by μ_r and ϕ is a representable morphism. This allows us to extent our evaluation map to a map $\tilde{e}_i : \mathcal{K}_{g,n}(X,\beta) \longrightarrow \mathcal{I}_{\mu}(\mathcal{G}_X)$. Given a very twisted stable map $f: \mathcal{C} \longrightarrow X$ over T, its image $\widetilde{e}_i(f) \in \mathcal{I}_{\mu}(\mathcal{G}_X)(T)$ is the object associated with the diagram:

One can see that this agrees with the image $e_i(f) \in \overline{\mathcal{I}}_{\mu}(\mathcal{G}_X)(T)$ of the twisted stable map $f: \mathcal{C} \longrightarrow \overline{\mathcal{I}}_{\mu}(\mathcal{G}_X)$ given by the diagram:

$$
\bigcup_{\substack{\sum_{i} \downarrow \\ \downarrow \\ T}} \mathcal{G}_X
$$

Our desired equality of Gromov-Witten invariants follows.

Remark. Although the naive GW-theory described above is the natural first approach to take, there is evidence from the DT-side of the conjectural correspondence for orbifolds that some adjustment to the theory is needed. In particular, the moduli spaces for orbifold Calabi-Yau 3-folds in the DT-theory have virtual dimension 0, and the correspondence would hope for the same to be true on the GW-side. An example sugested by Jim Bryan is the following. Let E be the total space of $\mathcal{O}_{\mathbb{P}^1}(-1) \bigoplus \mathcal{O}_{\mathbb{P}^1}(-1)$ over \mathbb{P}^1 . The quotient $X = [E/(\mathbb{Z}/2\mathbb{Z})]$, where the action along each fiber is component-wise, is an orbi-Calabi-Yau 3-fold with the zerosection giving an embedded $\mathbb{P}^1 = [\mathbb{P}^1/(\mathbb{Z}/2\mathbb{Z})]$. In this case, $\mathcal{S}_X = X \sqcup \mathbb{P}^1$ and $\mathcal{G}_X = X \sqcup \mathbb{P}^1$. Stable maps into \mathcal{G}_X (in particular, into \mathbb{P}^1) will not have virtual dimension zero. So even in this straightforward case we have a virtual fundamental class $\left[\widetilde{\mathcal{K}}_{g,n}(X,\beta)\right]^{vir}$ with nonzero virtual dimension. A nicer theory might adjust the relative obstruction theory used to construct the virtual fundamental class in order to fix the problem with the virtual dimension. This is something we hope to look into.

REFERENCES

- [ACV03] D. Abramovich, A. Corti, and A. Vistoli, Twisted bundles and admissible covers, Comm. Algebra 31 (2003), no. 8, 3547–3618, Special issue in honor of Steven L. Kleiman. MR MR2007376 (2005b:14049)
- [AGV02] D. Abramovich, T. Graber, and A. Vistoli, Algebraic orbifold quantum products, Orbifolds in mathematics and physics (Madison, WI, 2001), Contemp. Math., vol. 310, Amer. Math. Soc., Providence, RI, 2002, pp. 1–24. MR MR1950940 (2004c:14104)
- [AGV08] $\qquad \qquad \qquad \ldots$, Gromov-Witten theory of Deligne-Mumford stacks, American Journal of Mathematics 130 (2008), no. 5, 1337–1398.
- [AOV08] D. Abramovich, M. Olsson, and A. Vistoli, *Tame stacks in positive characteristic*, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 4, 1057–1091. MR MR2427954
- [Art74] M. Artin, Versal deformations and algebraic stacks, Invent. Math. 27 (1974), 165–189. MR MR0399094 (53 #2945)
- [AV02] D. Abramovich and A. Vistoli, *Compactifying the space of stable maps*, J. Amer. Math. Soc. 15 (2002), no. 1, 27–75 (electronic). MR MR1862797 (2002i:14030)
- [Beh97] K. Behrend, Gromov-Witten invariants in algebraic geometry, Inventiones Mathematicae 127 (1997), 601–617.
- [BF97] K. Behrend and B. Fantechi, The intrinsic normal cone, Invent. Math. 128 (1997), no. 1, 45–88. MR MR1437495 (98e:14022)
- [CR05] W. Chen and Y. Ruan, *Orbifold Gromov-Witten theory*, Orbifolds in mathematics and physics (Madison, WI, 2001) J.53 (2005), no. 1, 25–85.
- [DT98] S. K. Donaldson and R. P. Thomas, Gauge theory in higher dimensions, The geometric universe (Oxford, 1996), no. 1, Oxford Univ. Press, Oxford, 1998, pp. 31–47. MR MR1634503 (2000a:57085)
- [FGI+05] B. Fantechi, L. Göttsche, L. Illusie, S. L. Kleiman, N. Nitsure, and A. Vistoli, Fundamental algebraic geometry, Mathematical Surveys and Monographs, vol. 123, American Mathematical Society, Providence, RI, 2005, Grothendieck's FGA explained. MR MR2222646 (2007f:14001)
- [LMB00] G. Laumon and L. Moret-Bailly, Champs algebrique, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 39, Springer-Verlag, 2000.
- [MNOP06] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande, Gromov-Witten theory and *Donaldson-Thomas theory.* I, Compos. Math. 142 (2006), no. 5, 1263–1285. MR MR2264664 (2007i:14061)
- [OS03] M. Olsson and J. Starr, Quot functors for Deligne-Mumford stacks, Comm. Algebra 31 (2003), no. 8, 4069–4096, Special issue in honor of Steven L. Kleiman. MR MR2007396 (2004i:14002)
- [Rom05] M. Romagny, Group actions on stacks and applications, Michigan Math J.53 (2005), no. 1, 209–236.
- [Tho00] R. P. Thomas, A holomorphic Casson invariant for Calabi-Yau 3-folds, and bundles on K3 fibrations, J. Differential Geom. 54 (2000), no. 2, 367–438. MR MR1818182 (2002b:14049)
- [Vis89] A. Vistoli, Intersection theory on algebraic stacks and on their moduli spaces, Invent. Math. 97 (1989), no. 3, 613–670. MR MR1005008 (90k:14004)

Qile Chen, Department of Mathematics, Box 1917, Brown University, Providence, RI, 02912, U.S.A

 $\it E\mbox{-}mail\;address\mbox{:}\;q\mbox{.chen@math.brown.edu}$

Steffen Marcus, Department of Mathematics, Box 1917, Brown University, Providence, RI, 02912, U.S.A

E-mail address: ssmarcus@math.brown.edu

HENNING ÚLFARSSON, SCHOOL OF COMPUTER SCIENCE, REYKJAVIK UNIVERSITY, REYKJAVIK, ICE-LAND

E-mail address: henningu@ru.is