

ABOUT THE NON-INTEGER PROPERTY OF HYPERHARMONIC NUMBERS

ISTVÁN MEZŐ

ABSTRACT. It was proven in 1915 by Leopold Theisinger that the H_n harmonic numbers are never integers. In 1996 Conway and Guy have defined the concept of hyperharmonic numbers. The question naturally arises: are there any integer hyperharmonic numbers? The author gives a partial answer to this question and conjectures that the answer is “no”.

The n -th harmonic number is the n -th partial sum of the harmonic series:

$$H_n = \sum_{k=1}^n \frac{1}{k}.$$

Conway and Guy in [4] defined the harmonic numbers of higher orders, also known as the hyperharmonic numbers: $H_n^{(1)} := H_n$, and for all $r > 1$ let

$$H_n^{(r)} = \sum_{k=1}^n H_k^{(r-1)}$$

be the n -th harmonic number of order r . These numbers can be expressed by binomial coefficients and ordinary harmonic numbers:

$$H_n^{(r)} = \binom{n+r-1}{r-1} (H_{n+r-1} - H_{r-1}).$$

The prominent role of these numbers has been realized recently in combinatorics. The $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]_r$ r -Stirling number is the number of the permutations of the set $\{1, \dots, n\}$ having k disjoint, non-empty cycles, in which the elements 1 through r are restricted to appear in different cycles.

In [1] one can find the following interesting equality:

$$H_n^{(r)} = \frac{\left[\begin{smallmatrix} n+r \\ r+1 \end{smallmatrix} \right]_r}{n!}.$$

Let us turn our attention to the main question of this paper. It is known that any number of consecutive terms not necessarily beginning with 1 will never sum to an integer (see [7]). As a corollary, we get that the H_n harmonic numbers are never integers ($n > 1$). Theisinger

2000 *Mathematics Subject Classification.* 11B83.

Key words and phrases. harmonic numbers, hyperharmonic numbers, 2-adic norm.

proved this latter result directly in 1915 [6]. The question appears obviously: are there any integer hyperharmonic numbers?

Theisinger's main tool was the 2-adic norm. We give a short summary of his method. Every rational number $x \neq 0$ can be represented by $x = \frac{p^\alpha r}{s}$, where p is a fixed prime number, r and s are relative prime integers to p . α is a unique integer. We can define the p -adic norm of x by

$$|x|_p = p^{-\alpha}, \text{ and let } |0|_p = 0.$$

This norm fulfills the properties of the usual norms, namely

$$\begin{aligned} |x|_p = 0 &\iff x = 0, \\ |xy|_p &= |x|_p |y|_p \quad (x, y \in \mathbb{Q}), \\ |x + y|_p &\leq |x|_p + |y|_p \quad (x, y \in \mathbb{Q}). \end{aligned}$$

Furthermore, the so-called strong triangle inequality also holds:

$$|x + y|_p \leq \max\{|x|_p, |y|_p\} (\leq |x|_p + |y|_p).$$

We shall use the following property of integer numbers:

$$x \in \mathbb{Z} \implies x = p^\alpha r \implies |x|_p = \frac{1}{p^\alpha} \leq 1,$$

where r and the prime p are relative prime integers. This means that if the p -norm of a rational x is greater than 1 then x is necessarily non-integer.

Let us introduce the order of a natural number n : if $2^m \leq n < 2^{m+1}$, then $\text{Ord}_2(n) := m$. It is obvious that $\text{Ord}_2(n) = \lfloor \log_2(n) \rfloor$.

Theorem 1.

$$|H_n|_2 = 2^{\text{Ord}_2(n)} \quad (n \in \mathbb{N}),$$

that is – by our observation above – H_n is never integer.

Proof. First, let n be even. Since $|x|_2 = |-x|_2$ for all $x \in \mathbb{Q}$, by the strong triangle inequality we get

$$\begin{aligned} \max\{|H_n|_2, |1|_2\} &= \max\left\{|H_n|_2, |1|_2, \left|\frac{1}{3}\right|_2, \left|\frac{1}{5}\right|_2, \dots, \left|\frac{1}{n-1}\right|_2\right\} \geq \\ &\geq \left|H_n - 1 - \frac{1}{3} - \frac{1}{5} - \dots - \frac{1}{n-1}\right|_2 = \\ &= \left|\frac{1}{2} + \frac{1}{4} + \dots + \frac{1}{n-2} + \frac{1}{n}\right|_2 = \\ &= \left|\frac{1}{2}\right|_2 \left|1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n/2}\right|_2 = 2 |H_{n/2}|_2. \end{aligned}$$

If n is odd, the situation is the same:

$$\max\{|H_n|_2, |1|_2\} \geq 2 |H_{(n-1)/2}|_2.$$

The reader may verify it.

So we get that the 2-adic norm of the harmonic numbers is monotone increasing. Since $|H_2|_2 = \left|\frac{3}{2}\right|_2 = 2$, the 2-adic norm of all the harmonic numbers are greater than 1. As a corollary, this means that the harmonic numbers are not integers because of the property of the 2-adic norm mentioned above.

We can continue the calculations on $H_{n/2}$ (or on $H_{(n-1)/2}$) instead of H_n . For instance let us consider that $n/2$ is even. Then the method described above gives that

$$|H_{n/2}|_2 \geq \left|\frac{1}{2}\right|_2 |H_{n/4}|_2 = 2|H_{n/4}|_2.$$

This and the previous estimation implies that

$$|H_n|_2 \geq \left|\frac{1}{2}\right|_2 |H_{n/2}|_2 \geq \left|\frac{1}{2}\right|_2 \left|\frac{1}{2}\right|_2 |H_{n/4}|_2 = 4|H_{n/4}|_2.$$

And so on. If $n/2$ is odd then we choose $(n/2 - 1)/2$ instead of $n/4$. We can perform these steps exactly $\text{Ord}_2(n)$ times.

After all, we shall have the following:

$$|H_n|_2 \geq \left|\frac{1}{2^{\text{Ord}_2(n)}}\right|_2 |H_1|_2 = 2^{\text{Ord}_2(n)}.$$

On the other hand,

$$|H_n|_2 \leq \max \left\{ |1|_2, \left|\frac{1}{2}\right|_2, \dots, \left|\frac{1}{n}\right|_2 \right\} = \left|\frac{1}{2^{\text{Ord}_2(n)}}\right|_2 = 2^{\text{Ord}_2(n)},$$

because the greatest 2-power occurring between 1 and n is $\text{Ord}_2(n)$.

The inequalities detailed above give the statement. \square

A different approach can be found in [5], [2] and in their references. The next proof comes from these sources.

Proof. Let us fix the order of n , i.e.: $\text{Ord}_2(n) := m$. This implies that the denominator of $\frac{2^{m-1}}{n}$ is odd, unless $n = 2^m$. We get that the number

$$2^{m-1}H_n - \frac{1}{2}$$

can be represented by the sum of rationals with odd denominators. For example,

$$2^{m-1}H_n - \frac{1}{2} = \frac{a_1}{b_1} + \dots + \frac{a_s}{b_s} = \frac{c}{\text{lcm}(b_1, \dots, b_s)},$$

where b_i is odd for all $i = 1, \dots, s$. It means that $b := \text{lcm}(b_1, \dots, b_s)$ is odd. The last formula gives the result

$$H_n = \frac{\frac{c}{b} + \frac{1}{2}}{2^{m-1}} = \frac{2c + b}{2^m b}.$$

\square

Let us turn our attention to hyperharmonic numbers. We need a lemma which can be found in [3]:

Lemma 2.

$$|n!|_p = p^{(A_p(n)-n)/(p-1)},$$

where $A_p(n)$ is the sum of the digits of the p -adic expansion of n .

Example 3. Let $p = 2$ and $n = 11$. Then $n = 1011_2$, that is, $A_2(n) = 3$.

$$|n!|_2 = |39916800|_2 = |256 \cdot 155925|_2 = |256|_2 |155925|_2 = |2^8|_2 \cdot 1 = 2^{-8}.$$

We can apply the lemma: $A_2(n) - n = 3 - 11 = -8$, whence $|n!|_2 = 2^{-8}$.

Theorem 4. If $\text{Ord}_2(n + r - 1) > \text{Ord}_2(r - 1)$ then

$$|H_n^{(r)}|_2 = 2^{A_2(n+r-1) - A_2(n) - A_2(r-1) + \text{Ord}_2(n+r-1)},$$

else

$$|H_n^{(r)}|_2 = 2^{A_2(n+r-1) - A_2(n) - A_2(r-1) + \max\{\lfloor \frac{1}{r} \rfloor_2, \lfloor \frac{1}{r+1} \rfloor_2, \dots, \lfloor \frac{1}{n+r-1} \rfloor_2\}}.$$

Proof.

$$\begin{aligned} |H_n^{(r)}|_2 &= \left| \binom{n+r-1}{r-1} (H_{n+r-1} - H_{r-1}) \right|_2 = \\ &= \left| \binom{n+r-1}{r-1} \right|_2 \left| \frac{a}{2^{\text{Ord}_2(n+r-1)}b} - \frac{c}{2^{\text{Ord}_2(r-1)}d} \right|_2 = \\ &= \left| \binom{n+r-1}{r-1} \right|_2 \left| \frac{2^{\text{Ord}_2(r-1)}ad - 2^{\text{Ord}_2(n+r-1)}bc}{2^{\text{Ord}_2(n+r-1) + \text{Ord}_2(r-1)}bd} \right|_2. \end{aligned}$$

Because of the condition $\text{Ord}_2(n + r - 1) > \text{Ord}_2(r - 1)$ we get

$$|H_n^{(r)}|_2 = \left| \binom{n+r-1}{r-1} \right|_2 \left| \frac{ad - 2^{\text{Ord}_2(n+r-1) - \text{Ord}_2(r-1)}bc}{2^{\text{Ord}_2(n+r-1)}bd} \right|_2.$$

Since the nominator is odd, we get the following:

$$\left| \frac{ad - 2^{\text{Ord}_2(n+r-1) - \text{Ord}_2(r-1)}bc}{2^{\text{Ord}_2(n+r-1)}bd} \right|_2 = 2^{\text{Ord}_2(n+r-1)}.$$

To compute the 2-adic norm of the binomial coefficient, we use the previous lemma.

$$\begin{aligned} \left| \binom{n+r-1}{r-1} \right|_2 &= \left| \frac{(n+r-1)!}{(r-1)!n!} \right|_2 = \\ &= \frac{2^{A_2(n+r-1) - n - r + 1}}{2^{A_2(r-1) - r + 1} 2^{A_2(n) - n}} = 2^{A_2(n+r-1) - A_2(n) - A_2(r-1)}. \end{aligned}$$

This, and the previous equality give the result with respect to the condition $\text{Ord}_2(n + r - 1) > \text{Ord}_2(r - 1)$.

Let us fix an arbitrary n for which $\text{Ord}_2(n + r - 1) = \text{Ord}_2(r - 1)$.

$$H_{n+r-1} - H_{r-1} = \frac{1}{r} + \frac{1}{r+1} + \dots + \frac{1}{n+r-1}.$$

Let us subtract all of the fractions with odd denominators. Then we can take $\frac{1}{2}$ out of the remainder and continue the recursive method described in the first proof of Theorem 1. We can make such subtraction steps

$$\max \left\{ \left| \frac{1}{r} \right|_2, \left| \frac{1}{r+1} \right|_2, \dots, \left| \frac{1}{n+r-1} \right|_2 \right\}$$

times. The result:

$$|H_{n+r-1} - H_{r-1}|_2 \geq \max \left\{ \left| \frac{1}{r} \right|_2, \left| \frac{1}{r+1} \right|_2, \dots, \left| \frac{1}{n+r-1} \right|_2 \right\}.$$

On the other hand, by the strong triangle inequality

$$|H_{n+r-1} - H_{r-1}|_2 \leq \max \left\{ \left| \frac{1}{r} \right|_2, \left| \frac{1}{r+1} \right|_2, \dots, \left| \frac{1}{n+r-1} \right|_2 \right\}.$$

□

Corollary 5. *The sum of the harmonic numbers cannot be integer:*

$$H_1 + H_2 + \dots + H_n \notin \mathbb{N} \quad (n > 1).$$

Or, which is the same,

$$H_n^{(2)} \notin \mathbb{N} \quad (n > 1).$$

Proof. $H_1 + H_2 + \dots + H_n = H_n^{(2)}$. The condition with respect to the order of n and r holds because $\text{Ord}_2(n+2-1) > \text{Ord}_2(2-1) = 0$ for all $n \geq 1$. Furthermore,

$$|H_n^{(2)}|_2 = 2^{A_2(n+1) - A_2(n) - A_2(1) + \text{Ord}_2(n+1)}.$$

Let $m = \text{Ord}_2(n+1)$. Our goal is to minimize the power of 2. $\text{Ord}_2(n+1) = m$ implies that $n+1 < 2^{m+1}$, therefore $1 \leq A_2(n+1) \leq m+1$ and $1 \leq A_2(n) \leq m$. The minimum in the power is taken when $A_2(n) = m$ and $A_2(n+1) = 1$. It is possible if and only if $n = 2^m - 1$. In this case

$$A_2(n+1) - A_2(n) - A_2(1) + \text{Ord}_2(n+1) = 1 - m - 1 + m = 0.$$

We get that if $n \neq 2^m - 1$ for some m , then $|H_n^{(2)}|_2 > 1$, that is, $H_n^{(2)} \notin \mathbb{N}$. On the other hand, let us assume that n has the form $2^m - 1$. This implies that

$$\begin{aligned} H_n^{(2)} &= \binom{n+2-1}{2-1} (H_{n+2-1} - H_{2-1}) = \\ &= (n+1)(H_{n+1} - 1) = 2^m \left(\frac{a}{2^{\text{Ord}_2(n+1)} b} - 1 \right) = \frac{a}{b} - 2^m \notin \mathbb{N}. \end{aligned}$$

□

One can easily prove the following, using the method in the previous proof.

Corollary 6. $H_n^{(3)} \notin \mathbb{N}$ for all $n > 1$.

As we can see, the method to prove the non-integer property of harmonic numbers does not work for hyperharmonic numbers, because there are n and r integers for which $|H_n^{(r)}|_2 = 1$. In spite of this fact, we believe that Theisinger's theorem holds for all hyperharmonic numbers, too.

Conjecture 7. *None of the hyperharmonic numbers can be integers ($r, n \geq 2$).*

Example 8. We demonstrate that the theorem described above simplifies the calculation of the 2-norm of hyperharmonic numbers.

For instance,

$$\begin{aligned} H_{18}^{(8)} &= \binom{18+8-1}{8-1} (H_{18+8-1} - H_{8-1}) = \\ &= 480700 \left(\frac{34052522467}{8923714800} - \frac{363}{140} \right) = \frac{10914604807}{18564}. \end{aligned}$$

Since $|18564|_2 = 2^{-2}$, we get that $|H_{18}^{(8)}|_2 = 2^2 = 4$.

On the other hand, $A_2(18+8-1) = A_2(16+8+1) = 3$, $A_2(8-1) = A_2(4+2+1) = 3$, $A_2(18) = A_2(16+2) = 2$ and $\text{Ord}_2(18+8-1) = \text{Ord}_2(16+9) = 4$. By theorem 5,

$$|H_{18}^{(8)}|_2 = 2^{3-3-2+4} = 2^2 = 4.$$

Finally, we pose an interesting question:

Problem 9. For which $n_1 \neq n_2$ and $r_1 \neq r_2$ does the equality

$$H_{n_1}^{(r_1)} = H_{n_2}^{(r_2)}$$

stand?

REFERENCES

- [1] A. T. Benjamin, D. Gaebler and R. Gaebler, A combinatorial approach to hyperharmonic numbers, *INTEGERS: The Electronic Journal of Combinatorial Number Theory*, Vol 3 (2003), 1-9, #A15.
- [2] B. H. Brown and B. Rosenbaum, E46, *Amer. Math. Monthly* 41 (1934), 48-49.
- [3] E. B. Burger and T. Struppeck, Does $\sum_{n=0}^{\infty} \frac{1}{n!}$ Really Converge? Infinite Series and p -adic Analysis, *Amer. Math. Monthly* 103 (1996), 565-577.
- [4] J. H. Conway and R. K. Guy, *The book of numbers*, New York, Springer-Verlag, p. 258-259, 1996.
- [5] R. L. Graham et al. *Concrete Mathematics*, Addison Wesley, 1993.
- [6] L. Theisinger, Bemerkung über die harmonische Reihe, *Monatshefte für Mathematik und Physik* 26 (1915), 132-134.
- [7] E. W. Weisstein et al. "Harmonic Number." From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/HarmonicNumber.html>

DEPARTMENT OF ALGEBRA AND NUMBER THEORY, INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN, HUNGARY

E-mail address: imezo@math.klte.hu

URL: <http://www.math.klte.hu/algebra/mezo.htm>