

Representations and characterizations of polynomial functions on chains

Miguel Couceiro

Mathematics Research Unit, University of Luxembourg
162A, avenue de la Faiencerie, L-1511 Luxembourg, Luxembourg
miguel.couceiro[at]uni.lu

Jean-Luc Marichal

Mathematics Research Unit, University of Luxembourg
162A, avenue de la Faiencerie, L-1511 Luxembourg, Luxembourg
jean-luc.marichal[at]uni.lu

November 3, 2008

Abstract

We are interested in representations and characterizations of lattice polynomial functions $f: L^n \rightarrow L$, where L is a given bounded distributive lattice. In a companion paper [4], we investigated certain representations and provided various characterizations of these functions both as solutions of certain functional equations and in terms of necessary and sufficient conditions. In the present paper, we investigate these representations and characterizations in the special case when L is a chain, i.e., a totally ordered lattice. More precisely, we discuss representations of lattice polynomial functions given in terms of standard simplices and we present new axiomatizations of these functions by relaxing some of the conditions given in [4] and by considering further conditions, namely comonotonic minitivity and maxitivity.

Keywords: Lattice polynomial function, discrete Sugeno integral, term function, normal form, standard simplex, homogeneity, strong idempotency, median decomposability, comonotonicity.

1 Introduction

In [4], the class of (*lattice*) *polynomial functions*, i.e., functions representable by combinations of variables and constants using the lattice operations \wedge and \vee , was considered and characterized both as solutions of certain functional equations and in terms of necessary and sufficient conditions rooted in aggregation theory.

Formally, let L be a bounded distributive lattice with operations \wedge and \vee , and with least and greatest elements 0 and 1, respectively. An *n-ary polynomial function* on L is any function $f: L^n \rightarrow L$ which can be obtained by finitely many applications of the following rules:

- (i) For each $i \in [n] = \{1, \dots, n\}$ and each $c \in L$, the projection $\mathbf{x} \mapsto x_i$ and the constant function $\mathbf{x} \mapsto c$ are polynomial functions from L^n to L .
- (ii) If f and g are polynomial functions from L^n to L , then $f \vee g$ and $f \wedge g$ are polynomial functions from L^n to L .

Polynomial functions are also called lattice functions (Goodstein [9]), algebraic functions (Burris and Sankappanavar [3]) or weighted lattice polynomial functions (Marichal [13]). Polynomial functions obtained from projections by finitely many applications of (ii) are referred to as *lattice term functions*.

The recent interest by aggregation theorists in this class of polynomial functions is partially motivated by its connection to noteworthy aggregation functions such as the (*discrete*) *Sugeno integral*, which was introduced by Sugeno [19, 20] and widely investigated in aggregation theory, due to the many applications in fuzzy set theory, data fusion, decision making, image analysis, etc. As shown by Marichal [13], the discrete Sugeno integrals are nothing other than those polynomial functions $f : L^n \rightarrow L$ which are idempotent, that is, satisfying $f(x, \dots, x) = x$. For general background on aggregation theory, see [1, 11] and for a recent reference, see [10].

In this paper, we refine our previous results in the particular case when L is a chain, by relaxing our conditions and proposing weak analogues of those properties used in [4], and then providing characterizations of polynomial functions, accordingly. Moreover, and motivated by the axiomatizations of the discrete Sugeno integrals established by de Campos and Bolaños [5] (in the case when $L = [0, 1]$ is the unit real interval), we present further and alternative characterizations of polynomial functions given in terms of comonotonic minitivity and maxitivity. As particular cases, we consider the subclasses of discrete Sugeno integrals as well as that of term functions.

The current paper is organized as follows. We start in §2 by introducing the basic notions needed in this paper and presenting the characterizations of lattice polynomial functions on arbitrary (possibly infinite) bounded distributive lattices, established in [4]. Those characterizations are reassembled in Theorem 1. We discuss representations of polynomial functions in normal form (such as the classical disjunctive and conjunctive normal forms) and introduce variant representations in the case of chains and given in terms of standard simplices in §3. In §4, we provide characterizations of polynomial functions on chains given in terms of weak analogues of the properties used in Theorem 1 as well as in terms of comonotonic minitivity and maxitivity. The subclasses of discrete Sugeno integrals and of term functions are then axiomatized in §5 using the results obtained in the previous section.

2 Basic notions and terminology

Throughout this paper, let L be a bounded distributive lattice with operations \wedge and \vee , and with least and greatest elements 0 and 1, respectively. For $a, b \in L$, $a \leq b$ simply means that $a \wedge b = a$ or, equivalently, $a \vee b = b$. A *chain* is simply a lattice such that for every $a, b \in L$ we have $a \leq b$ or $b \leq a$. A subset S of a lattice L is said to be *convex* if for every $a, b \in S$ and every $c \in L$ such that $a \leq c \leq b$, we have $c \in S$. For any subset $S \subseteq L$, we denote by \overline{S} the convex hull of S , that is, the smallest convex subset of L containing S . For every $a, b \in S$ such that $a \leq b$, we denote by $[a, b]$ the *interval*

$$[a, b] = \{c \in L : a \leq c \leq b\}.$$

For any integer $n \geq 1$, we set $[n] = \{1, \dots, n\}$.

For an arbitrary nonempty set A and a lattice L , the set L^A of all functions from A to L constitutes a lattice under the operations

$$(f \wedge g)(x) = f(x) \wedge g(x) \quad \text{and} \quad (f \vee g)(x) = f(x) \vee g(x),$$

for every $f, g \in L^A$. In particular, any lattice L induces a lattice structure on the Cartesian product L^n , $n \geq 1$, by defining \wedge and \vee componentwise, i.e.,

$$\begin{aligned} (a_1, \dots, a_n) \wedge (b_1, \dots, b_n) &= (a_1 \wedge b_1, \dots, a_n \wedge b_n), \\ (a_1, \dots, a_n) \vee (b_1, \dots, b_n) &= (a_1 \vee b_1, \dots, a_n \vee b_n). \end{aligned}$$

We denote the elements of L by lower case letters a, b, c, \dots , and the elements of L^n , $n > 1$, by bold face letters $\mathbf{a}, \mathbf{b}, \mathbf{c}, \dots$. We also use $\mathbf{0}$ and $\mathbf{1}$ to denote the least element and greatest element, respectively, of L^n . For $c \in L$ and $\mathbf{x} = (x_1, \dots, x_n) \in L^n$, set

$$\mathbf{x} \wedge c = (x_1 \wedge c, \dots, x_n \wedge c) \quad \text{and} \quad \mathbf{x} \vee c = (x_1 \vee c, \dots, x_n \vee c).$$

The *range* of a function $f: L^n \rightarrow L$ is defined by $\mathcal{R}_f = \{f(\mathbf{x}) : \mathbf{x} \in L^n\}$. A function $f: L^n \rightarrow L$ is said to be *nondecreasing (in each variable)* if, for every $\mathbf{a}, \mathbf{b} \in L^n$ such that $\mathbf{a} \leq \mathbf{b}$, we have $f(\mathbf{a}) \leq f(\mathbf{b})$. Note that if f is nondecreasing, then $\overline{\mathcal{R}}_f = [f(\mathbf{0}), f(\mathbf{1})]$.

Let S be a nonempty subset of L . A function $f: L^n \rightarrow L$ is said to be

- *S-idempotent* if for every $c \in S$, we have $f(c, \dots, c) = c$.
- *S-min homogenous* if for every $\mathbf{x} \in L^n$ and every $c \in S$, we have

$$f(\mathbf{x} \wedge c) = f(\mathbf{x}) \wedge c. \tag{1}$$

- *S-max homogenous* if for every $\mathbf{x} \in L^n$ and every $c \in S$, we have

$$f(\mathbf{x} \vee c) = f(\mathbf{x}) \vee c. \tag{2}$$

- *horizontally S-minitive* if for every $\mathbf{x} \in L^n$ and every $c \in S$, we have

$$f(\mathbf{x}) = f(\mathbf{x} \vee c) \wedge f([\mathbf{x}]^c), \tag{3}$$

where $[\mathbf{x}]^c$ is the n -tuple whose i th component is 1, if $x_i \geq c$, and x_i , otherwise.

- *horizontally S-maxitive* if for every $\mathbf{x} \in L^n$ and every $c \in S$, we have

$$f(\mathbf{x}) = f(\mathbf{x} \wedge c) \vee f([\mathbf{x}]_c), \tag{4}$$

where $[\mathbf{x}]_c$ is the n -tuple whose i th component is 0, if $x_i \leq c$, and x_i , otherwise.

- *median decomposable* if, for every $\mathbf{x} \in L^n$ and every $k \in [n]$, we have

$$f(\mathbf{x}) = \text{median} \left(f(\mathbf{x}_k^0), x_k, f(\mathbf{x}_k^1) \right), \tag{5}$$

where $\mathbf{x}_k^c = (x_1, \dots, x_{k-1}, c, x_{k+1}, \dots, x_n)$ for any $c \in L$.

- *strongly idempotent* if, for every $\mathbf{x} \in L^n$ and every $k \in [n]$, we have

$$f(x_1, \dots, x_{k-1}, f(\mathbf{x}), x_{k+1}, \dots, x_n) = f(\mathbf{x}).$$

Remark 1. In the case when $S = L$ is the real interval $[0, 1]$, the concepts of S -min and S -max homogeneity were introduced by Fodor and Roubens [8] to specify certain classes of aggregation functions, and the concept of horizontal S -maxitivity was introduced by Benvenuti et al. [2] as a general property of the Sugeno integral. The concept of median decomposition was introduced by Marichal [13] and that of strong idempotency by Couceiro and Marichal [4] as properties of polynomial functions.

We say that a function $f: L^n \rightarrow L$ has a *componentwise convex range* if either $n = 1$ and f has a convex range, or $n > 1$ and for every $\mathbf{a} = (a_1, \dots, a_n) \in L^n$ and every $k \in [n]$, the unary function $f_{\mathbf{a}}^k: L \rightarrow L$, given by

$$f_{\mathbf{a}}^k(x) = f(a_1, \dots, a_{k-1}, x, a_{k+1}, \dots, a_n).$$

has a convex range.

The following theorem reassembles the various characterizations of polynomial functions, defined on an arbitrary (possibly infinite) bounded distributive lattice L , established in [4] and given in terms of the properties introduced above.

Theorem 1. *Let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

- (i) f is a polynomial function.
- (ii) f is median decomposable.
- (iii) f is nondecreasing, strongly idempotent, has a convex range and a componentwise convex range.
- (iv) f is nondecreasing, $\overline{\mathcal{R}}_f$ -min homogeneous, and $\overline{\mathcal{R}}_f$ -max homogeneous.
- (v) f is nondecreasing, $\overline{\mathcal{R}}_f$ -min homogeneous, and horizontally $\overline{\mathcal{R}}_f$ -maxitive.
- (vi) f is nondecreasing, horizontally $\overline{\mathcal{R}}_f$ -minitive, and $\overline{\mathcal{R}}_f$ -max homogeneous.
- (vii) f is nondecreasing, $\overline{\mathcal{R}}_f$ -idempotent, horizontally $\overline{\mathcal{R}}_f$ -minitive, and horizontally $\overline{\mathcal{R}}_f$ -maxitive.

3 Representations of polynomial functions

Polynomial functions are known to be exactly those functions which can be represented by formulas in disjunctive and conjunctive normal forms. This fact was first observed by Goodstein [9] who, in fact, showed that each polynomial function $f: L^n \rightarrow L$ is uniquely determined by its restriction to $\{0, 1\}^n$. For a recent reference, see Rudeanu [18].

In this section we recall and refine some known results concerning normal forms of polynomial functions and, in the special case when L is a chain, we provide variant representations given in terms of standard simplices of L^n .

The following three results are due to Goodstein [9].

Corollary 2. *Every polynomial function is completely determined by its restriction to $\{0, 1\}^n$.*

Corollary 3. *A function $g: \{0, 1\}^n \rightarrow L$ can be extended to a polynomial function $f: L^n \rightarrow L$ if and only if it is nondecreasing. In this case, the extension is unique.*

Proposition 4. *Let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

(i) *f is a polynomial function.*

(ii) *There exists $\alpha: 2^{[n]} \rightarrow L$ such that*

$$f(\mathbf{x}) = \bigvee_{I \subseteq [n]} \left(\alpha(I) \wedge \bigwedge_{i \in I} x_i \right).$$

(iii) *There exists $\beta: 2^{[n]} \rightarrow L$ such that*

$$f(\mathbf{x}) = \bigwedge_{I \subseteq [n]} \left(\beta(I) \vee \bigvee_{i \in I} x_i \right).$$

We shall refer to the expressions given in (ii) and (iii) of Proposition 4 as the *disjunctive normal form* (DNF) representation and the *conjunctive normal form* (CNF) representation, respectively, of the polynomial function f .

Remark 2. By requiring α and β to be nonconstant functions from $2^{[n]}$ to $\{0, 1\}$ and satisfying $\alpha(\emptyset) = 0$ and $\beta(\emptyset) = 1$, respectively, we obtain the analogue of Proposition 4 for term functions.

For each polynomial function $f: L^n \rightarrow L$, set

$$\begin{aligned} \text{DNF}(f) &= \left\{ \alpha \in L^{2^{[n]}} : f(\mathbf{x}) = \bigvee_{I \subseteq [n]} \left(\alpha(I) \wedge \bigwedge_{i \in I} x_i \right) \right\}, \\ \text{CNF}(f) &= \left\{ \beta \in L^{2^{[n]}} : f(\mathbf{x}) = \bigwedge_{I \subseteq [n]} \left(\beta(I) \vee \bigvee_{i \in I} x_i \right) \right\}. \end{aligned}$$

A complete description of the sets $\text{DNF}(f)$ and $\text{CNF}(f)$ can be found in Couceiro and Marichal [4, §3.1]. As we are concerned by the case when L is a chain, we recall the description only in this special case; see Marichal [13, §3].

For each $I \subseteq [n]$, let \mathbf{e}_I be the element of L^n whose i th component is 1, if $i \in I$, and 0, otherwise. Let $\alpha_f: 2^{[n]} \rightarrow L$ be the function given by $\alpha_f(I) = f(\mathbf{e}_I)$ and consider the function $\alpha_f^*: 2^{[n]} \rightarrow L$ defined by

$$\alpha_f^*(I) = \begin{cases} \alpha_f(I), & \text{if } \bigvee_{J \subsetneq I} \alpha_f(J) < \alpha_f(I), \\ 0, & \text{otherwise.} \end{cases}$$

Dually, Let $\beta_f: 2^{[n]} \rightarrow L$ be the function given by $\beta_f(I) = f(\mathbf{e}_{[n] \setminus I})$ and consider the function $\beta_f^*: 2^{[n]} \rightarrow L$ defined by

$$\beta_f^*(I) = \begin{cases} \beta_f(I), & \text{if } \bigwedge_{J \subsetneq I} \beta_f(J) > \beta_f(I), \\ 1, & \text{otherwise.} \end{cases}$$

Proposition 5. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a polynomial function. Then*

- (i) $DNF(f) = [\alpha_f^*, \alpha_f]$ and f has a unique DNF representation if and only if $\bigvee_{J \subsetneq I} \alpha_f(J) < \alpha_f(I)$ for every $I \subseteq [n]$,
- (ii) $CNF(f) = [\beta_f, \beta_f^*]$ and f has a unique CNF representation if and only if $\bigwedge_{J \subsetneq I} \beta_f(J) > \beta_f(I)$ for every $I \subseteq [n]$.

In particular, α_f and β_f are the unique isotone and antitone, respectively, maps in $DNF(f)$ and $CNF(f)$, respectively.

In the case of chains, the DNF and CNF representations of polynomial functions $f: L^n \rightarrow L$ can be refined and given in terms of standard simplices of L^n (see Proposition 7 below). To provide these variants, we first need the following lemma due to Dubois and Prade [7]. For the sake of self-containment, we provide a simpler proof. Recall that

$$\text{median}(x_1, \dots, x_{2n+1}) = \bigvee_{\substack{I \subseteq [2n+1] \\ |I|=n+1}} \bigwedge_{i \in I} x_i.$$

Lemma 6. *Let $a_1, \dots, a_{n+1}, b_1, \dots, b_{n+1} \in L$ such that $a_1 \leq \dots \leq a_{n+1}$ and $b_1 \geq \dots \geq b_{n+1}$. If $a_{n+1} \geq b_{n+1}$, then*

$$\bigvee_{i \in [n+1]} (a_i \wedge b_i) = \text{median}(a_1, \dots, a_n, b_1, \dots, b_{n+1}).$$

Proof. Clearly, we have $\bigvee_{i \in [n+1]} (a_i \wedge b_i) \leq \text{median}(a_1, \dots, a_n, b_1, \dots, b_{n+1})$. Hence, to complete the proof it is enough to show that for every $I \subseteq [n]$ and every $J \subseteq [n+1]$, with $|I| + |J| = n+1$, there is $k \in [n+1]$ such that

$$\left(\bigwedge_{i \in I} a_i \right) \wedge \left(\bigwedge_{j \in J} b_j \right) \leq a_k \wedge b_k. \quad (6)$$

Let $i' = \bigwedge_{i \in I} i$ and $j' = \bigvee_{j \in J} j$. Since $|I| + |J| = n+1$, we have that $[i', n] \cap [1, j'] \neq \emptyset$. Clearly, any $k \in [i', n] \cap [1, j']$ fulfills (6). \square

Let σ be a permutation on $[n]$. The *standard simplex* of L^n associated with σ is the subset $L_\sigma^n \subseteq L^n$ defined by

$$L_\sigma^n = \{\mathbf{x} = (x_1, \dots, x_n) \in L^n : x_{\sigma(1)} \leq x_{\sigma(2)} \leq \dots \leq x_{\sigma(n)}\}.$$

For each $i \in [n]$, define $S_\sigma^\uparrow(i) = \{\sigma(i), \dots, \sigma(n)\}$ and $S_\sigma^\downarrow(i) = \{\sigma(1), \dots, \sigma(i)\}$. As a matter of convenience, set $S_\sigma^\uparrow(n+1) = S_\sigma^\downarrow(0) = \emptyset$.

Proposition 7. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

- (i) f is a polynomial function.
- (ii) For any permutation σ on $[n]$ and every $\mathbf{x} \in L_\sigma^n$, we have

$$\begin{aligned} f(\mathbf{x}) &= \bigvee_{i \in [n+1]} \left(\alpha_f(S_\sigma^\uparrow(i)) \wedge x_{\sigma(i)} \right) = \bigwedge_{i \in [n+1]} \left(\alpha_f(S_\sigma^\uparrow(i)) \vee x_{\sigma(i-1)} \right) \\ &= \text{median} \left(x_1, \dots, x_n, \alpha_f(S_\sigma^\uparrow(1)), \dots, \alpha_f(S_\sigma^\uparrow(n+1)) \right), \end{aligned} \quad (7)$$

where $x_{\sigma(0)} = 0$ and $x_{\sigma(n+1)} = 1$.

(iii) For any permutation σ on $[n]$ and every $\mathbf{x} \in L_\sigma^n$, we have

$$\begin{aligned} f(\mathbf{x}) &= \bigvee_{i \in [n+1]} \left(\beta_f(S_\sigma^\downarrow(i-1)) \wedge x_{\sigma(i)} \right) = \bigwedge_{i \in [n+1]} \left(\beta_f(S_\sigma^\downarrow(i-1)) \wedge x_{\sigma(i-1)} \right) \\ &= \text{median} \left(x_1, \dots, x_n, \beta_f(S_\sigma^\downarrow(0)), \dots, \beta_f(S_\sigma^\downarrow(n)) \right), \end{aligned}$$

where $x_{\sigma(0)} = 0$ and $x_{\sigma(n+1)} = 1$.

Proof. The equivalence (ii) \Leftrightarrow (iii) follows immediately from the facts $\beta_f(I) = \alpha_f([n] \setminus I)$, for every $I \subseteq [n]$, and $S_\sigma^\downarrow(i-1) = [n] \setminus S_\sigma^\uparrow(i)$.

To show that (i) \Rightarrow (ii), suppose that $f: L^n \rightarrow L$ is a polynomial function, that is,

$$f(\mathbf{x}) = \bigvee_{I \subseteq [n]} \left(\alpha_f(I) \wedge \bigwedge_{i \in I} x_i \right). \quad (8)$$

Clearly, for each permutation σ on $[n]$ and each $\mathbf{x} \in L_\sigma^n$, we have that (8) becomes

$$f(\mathbf{x}) = \alpha_f(\emptyset) \vee \bigvee_{\substack{i \in [n] \\ \sigma(i) \in I}} \bigvee_{I \subseteq S_\sigma^\uparrow(i)} \left(\alpha_f(I) \wedge x_{\sigma(i)} \right) = \alpha_f(\emptyset) \vee \bigvee_{i \in [n]} \left(\alpha_f(S_\sigma^\uparrow(i)) \wedge x_{\sigma(i)} \right).$$

This, together with Lemma 6, shows that (i) \Rightarrow (ii).

To show that (ii) \Rightarrow (i) also holds, let $f': L^n \rightarrow L$ be the polynomial function given by

$$f'(\mathbf{x}) = \bigvee_{I \subseteq [n]} \left(\alpha_f(I) \wedge \bigwedge_{i \in I} x_i \right).$$

By Corollary 3, f' is the unique extension of α_f to a polynomial function. Let $\mathbf{x} \in L^n$ and let σ be permutation on $[n]$ such that $\mathbf{x} \in L_\sigma^n$. As in the proof of (i) \Rightarrow (ii), we have

$$f'(\mathbf{x}) = \bigvee_{I \subseteq [n]} \left(\alpha_f(I) \wedge \bigwedge_{i \in I} x_i \right) = \bigvee_{i \in [n+1]} \left(\alpha_f(S_\sigma^\uparrow(i)) \wedge x_{\sigma(i)} \right) = f(\mathbf{x}). \quad (9)$$

Since (9) holds for any $\mathbf{x} \in L^n$, it follows that $f = f'$ and thus f is a polynomial function. \square

Remark 3. The equivalence between (i) and (ii) of Proposition 7 was already observed by Marichal [13, §5]. Prior to this, Propositions 5 and 7 were already established in Marichal [15] for idempotent polynomial functions (discrete Sugeno integrals) in the case when L is the unit real interval $[0, 1]$; see also Marichal [14, §4.3].

4 Characterizations of polynomial functions

In this section, we propose weak analogues of the properties used in Theorem 1 and provide characterizations of polynomial functions on chains, accordingly. Moreover, we introduce further properties, namely, comonotonic minitivity and maxitivity, which we then use to provide further characterizations of polynomial functions.

For integers $0 \leq p \leq q \leq n$, define

$$L_n^{(p,q)} = \{ \mathbf{x} \in L^n : |\{x_1, \dots, x_n\} \cap \{0, 1\}| \geq p \text{ and } |\{x_1, \dots, x_n\}| \leq q \}.$$

For instance, $L_n^{(0,2)}$ is the set of Boolean vectors of L^n that are two-sided trimmed by constant vectors, that is

$$L_n^{(0,2)} = \bigcup_{\substack{\mathbf{e} \in \{0,1\}^n \\ c, d \in L}} \{ \text{median}(c, \mathbf{e}, d) \}.$$

4.1 Weak homogeneity

Let S be a nonempty subset of L . We say that a function $f: L^n \rightarrow L$ is *weakly S -min homogeneous* (resp. *weakly S -max homogeneous*) if (1) (resp. (2)) holds for every $\mathbf{x} \in L_n^{(0,2)}$ and every $c \in S$.

For every integer $m \geq 1$, every $\mathbf{x} \in L^m$, and every $f: L^n \rightarrow L$, we define $\langle \mathbf{x} \rangle_f \in L^m$ as the m -tuple

$$\langle \mathbf{x} \rangle_f = \text{median}(f(\mathbf{0}), \mathbf{x}, f(\mathbf{1})),$$

where the right-hand side median is taken componentwise. As observed in [4], for every nonempty subset $S \subseteq L$, we have that f is S -min homogeneous and S -max homogeneous if and only if it satisfies

$$f(\text{median}(r, \mathbf{x}, s)) = \text{median}(r, f(\mathbf{x}), s) \quad (10)$$

for every $\mathbf{x} \in L^n$ and every $r, s \in S$. In particular, if $f(\mathbf{0}), f(\mathbf{1}) \in S$ then, for any $\mathbf{x} \in L^n$ such that $f(\mathbf{0}) \leq f(\mathbf{x}) \leq f(\mathbf{1})$, we have $f(\mathbf{x}) = f(\langle \mathbf{x} \rangle_f)$.

It was also shown in [4] that, for every nonempty subset $S \subseteq L$, if f is S -min homogeneous and S -max homogeneous, then it is S -idempotent. The following lemma shows that the weak analogue also holds.

Lemma 8. *Let S be a nonempty subset of L . If $f: L^n \rightarrow L$ is weakly S -min homogeneous and weakly S -max homogeneous, then it is S -idempotent. Moreover, if $f(\mathbf{0}), f(\mathbf{1}) \in S$ then, for any $\mathbf{x} \in L_n^{(0,2)}$ such that $f(\mathbf{0}) \leq f(\mathbf{x}) \leq f(\mathbf{1})$, we have $f(\mathbf{x}) = f(\langle \mathbf{x} \rangle_f)$.*

Proof. If $f: L^n \rightarrow L$ is weakly S -min homogeneous and weakly S -max homogeneous then, for any $c \in S$, we have $f(\mathbf{1} \wedge c) \wedge c = f(\mathbf{1} \wedge c) = f(\mathbf{1} \wedge c) \vee c$, and thus f is S -idempotent. The second statement follows from formula (10) when restricted to vectors $\mathbf{x} \in L_n^{(0,2)}$. \square

As we are going to see, of particular interest is when $S = \overline{\mathcal{R}}_f$, for which we have the following result (see [4]).

Proposition 9. *For any function $f: L^n \rightarrow L$ the following hold:*

- (i) *If f is $\overline{\mathcal{R}}_f$ -idempotent, then f has a convex range (i.e., $\overline{\mathcal{R}}_f = \mathcal{R}_f$).*
- (ii) *If f is a polynomial function, then it is $\overline{\mathcal{R}}_f$ -min homogeneous, $\overline{\mathcal{R}}_f$ -max homogeneous, $\overline{\mathcal{R}}_f$ -idempotent, and has a convex range.*
- (iii) *The function f is $\overline{\mathcal{R}}_f$ -min homogeneous (resp. $\overline{\mathcal{R}}_f$ -max homogeneous) if and only if it is \mathcal{R}_f -min homogeneous (resp. \mathcal{R}_f -max homogeneous) and has a convex range. In this case, $\mathcal{R}_f = \overline{\mathcal{R}}_f = [f(\mathbf{0}), f(\mathbf{1})]$.*

Let $f: L^n \rightarrow L$ be a nondecreasing function so that $\overline{\mathcal{R}}_f = [f(\mathbf{0}), f(\mathbf{1})]$. Now, if f is weakly $\overline{\mathcal{R}}_f$ -min homogeneous, then for any $c \in \overline{\mathcal{R}}_f$, we have $f(\mathbf{1} \wedge c) = f(\mathbf{1}) \wedge c = c$, and thus f is $\overline{\mathcal{R}}_f$ -idempotent. Dually, if f is weakly $\overline{\mathcal{R}}_f$ -max homogeneous, then it is also $\overline{\mathcal{R}}_f$ -idempotent. Hence we have the following result.

Lemma 10. *Let $f: L^n \rightarrow L$ be nondecreasing. If f is weakly $\overline{\mathcal{R}}_f$ -min homogeneous or weakly $\overline{\mathcal{R}}_f$ -max homogeneous, then it is $\overline{\mathcal{R}}_f$ -idempotent.*

We now provide our first characterization of polynomial functions which shows that, in the case of chains, the conditions in (iv) of Theorem 1 can be replaced with their weak analogues.

Theorem 11. *Let L be a bounded chain. A function $f: L^n \rightarrow L$ is a polynomial function if and only if it is nondecreasing, weakly $\overline{\mathcal{R}}_f$ -min homogenous, and weakly $\overline{\mathcal{R}}_f$ -max homogenous.*

Proof. From Proposition 9 (ii) it follows that each condition is necessary. To show that they are also sufficient, let $\mathbf{x} \in L^n$. By nondecreasing monotonicity, Lemma 8, and weak $\overline{\mathcal{R}}_f$ -min homogeneity, for every $I \subseteq [n]$ we have

$$\begin{aligned} f(\mathbf{x}) &\geq f\left(\mathbf{e}_I \wedge \left(\bigwedge_{i \in I} x_i\right)\right) = f\left(\left\langle \mathbf{e}_I \wedge \left(\bigwedge_{i \in I} x_i\right) \right\rangle_f\right) = f\left(\left\langle \mathbf{e}_I \right\rangle_f \wedge \left\langle \bigwedge_{i \in I} x_i \right\rangle_f\right) \\ &= f(\mathbf{e}_I) \wedge \left\langle \bigwedge_{i \in I} x_i \right\rangle_f \end{aligned}$$

and thus $f(\mathbf{x}) \geq \bigvee_{I \subseteq [n]} (f(\mathbf{e}_I) \wedge \langle \bigwedge_{i \in I} x_i \rangle_f)$. To complete the proof, it is enough to establish the converse inequality. Let $I^* \subseteq [n]$ be such that $f(\mathbf{e}_{I^*}) \wedge \langle \bigwedge_{i \in I^*} x_i \rangle_f$ is maximum. Define

$$J = \left\{ j \in [n] : x_j \leq f(\mathbf{e}_{I^*}) \wedge \left\langle \bigwedge_{i \in I^*} x_i \right\rangle_f \right\}.$$

We claim that $J \neq \emptyset$. For the sake of contradiction, suppose that $x_j > f(\mathbf{e}_{I^*}) \wedge \langle \bigwedge_{i \in I^*} x_i \rangle_f$ for every $j \in [n]$. Then, by nondecreasing monotonicity, we have $f(\mathbf{e}_{[n]}) \geq f(\mathbf{e}_{I^*})$, and since $f(\mathbf{e}_{[n]}) = f(\mathbf{1}) \geq \langle \bigwedge_{i \in [n]} x_i \rangle_f$,

$$f(\mathbf{e}_{[n]}) \wedge \left\langle \bigwedge_{i \in [n]} x_i \right\rangle_f > f(\mathbf{e}_{I^*}) \wedge \left\langle \bigwedge_{i \in I^*} x_i \right\rangle_f,$$

which contradicts the definition of I^* . Thus $J \neq \emptyset$.

Now, by nondecreasing monotonicity and weak $\overline{\mathcal{R}}_f$ -max homogeneity, we have

$$f(\mathbf{x}) \leq f\left(\left(f(\mathbf{e}_{I^*}) \wedge \left\langle \bigwedge_{i \in I^*} x_i \right\rangle_f\right) \vee \mathbf{e}_{[n] \setminus J}\right) = \left(f(\mathbf{e}_{I^*}) \wedge \left\langle \bigwedge_{i \in I^*} x_i \right\rangle_f\right) \vee f(\mathbf{e}_{[n] \setminus J}).$$

We claim that $f(\mathbf{e}_{[n] \setminus J}) \leq f(\mathbf{e}_{I^*}) \wedge \langle \bigwedge_{i \in I^*} x_i \rangle_f$. Indeed, otherwise by definition of J we would have

$$f(\mathbf{e}_{[n] \setminus J}) \wedge \left\langle \bigwedge_{i \in [n] \setminus J} x_i \right\rangle_f > f(\mathbf{e}_{I^*}) \wedge \left\langle \bigwedge_{i \in I^*} x_i \right\rangle_f,$$

again contradicting the definition of I^* .

Finally,

$$f(\mathbf{x}) \leq f(\mathbf{e}_{I^*}) \wedge \left\langle \bigwedge_{i \in I^*} x_i \right\rangle_f = \bigvee_{I \subseteq [n]} \left(f(\mathbf{e}_I) \wedge \left\langle \bigwedge_{i \in I} x_i \right\rangle_f\right). \quad \square$$

Remark 4. (i) Note that Theorem 11 does not generally hold in the case of bounded distributive lattices. To see this, let $L = \{0, a, b, 1\}$ where $a \wedge b = 0$ and $a \vee b = 1$, and consider the binary function $f: L^2 \rightarrow L$ defined by

$$f(x_1, x_2) = \begin{cases} 1, & \text{if } x_1 = 1 \text{ or } x_2 = 1, \\ 1, & \text{if } x_1 = x_2 = b, \\ a, & \text{if } (x_1 = a \text{ and } x_2 \neq 1) \text{ or } (x_2 = a \text{ and } x_1 \neq 1), \\ 0, & \text{otherwise.} \end{cases}$$

It is easy to verify that f is nondecreasing and both weakly $\overline{\mathcal{R}}_f$ -min homogenous and weakly $\overline{\mathcal{R}}_f$ -max homogenous, but it is not $\overline{\mathcal{R}}_f$ -max homogenous. For instance, $f(a \vee a, b \vee a) = f(a, 1) = 1 \neq a = f(a, b) \vee a$. Thus, f is not a polynomial function by Theorem 1.

- (ii) The proof technique of Proposition 11 was already used in Marichal [15] to prove a similar result for idempotent polynomial functions (discrete Sugeno integrals) in the case when L is the unit real interval $[0, 1]$.

4.2 Weak horizontal minitivity and maxitivity

Let S be a nonempty subset of L . We say that a function $f: L^n \rightarrow L$ is *weakly horizontally S -minitive* (resp. *weakly horizontally S -maxitive*) if (3) (resp. (4)) holds for every $\mathbf{x} \in L_n^{(0,2)}$ and every $c \in S$.

It was shown in [4] that if a function $f: L^n \rightarrow L$ is nondecreasing, S -idempotent, and horizontally S -minitive (resp. horizontally S -maxitive) then it is S -min homogenous (resp. S -max homogenous). The proof of this result can be easily adapted to derive the following weak analogue.

Lemma 12. *Let S be a nonempty subset of L . If $f: L^n \rightarrow L$ is nondecreasing, S -idempotent, and weakly horizontally S -minitive (resp. weakly horizontally S -maxitive) then it is weakly S -min homogenous (resp. weakly S -max homogenous).*

Proof. Let $f: L^n \rightarrow L$ be nondecreasing, S -idempotent, and weakly horizontally S -minitive. Then, for any $\mathbf{x} \in L_n^{(0,2)}$ and any $c \in S$,

$$\begin{aligned} f(\mathbf{x}) \wedge c &= f(\mathbf{x}) \wedge f(c, \dots, c) \geq f(\mathbf{x} \wedge c) = f((\mathbf{x} \wedge c) \vee c) \wedge f([\mathbf{x} \wedge c]^c) \\ &= f(c, \dots, c) \wedge f([\mathbf{x}]^c) \geq f(c, \dots, c) \wedge f(\mathbf{x}) = f(\mathbf{x}) \wedge c. \end{aligned}$$

Hence f is weakly S -min homogenous. The other statement can be proved similarly. \square

In [4] it was also shown that if $f: L^n \rightarrow L$ is nondecreasing and $\overline{\mathcal{R}}_f$ -min homogenous (resp. $\overline{\mathcal{R}}_f$ -max homogenous), then f is $\overline{\mathcal{R}}_f$ -max homogenous (resp. $\overline{\mathcal{R}}_f$ -min homogenous) if and only if it is horizontally $\overline{\mathcal{R}}_f$ -maxitive (resp. horizontally $\overline{\mathcal{R}}_f$ -minitive). Once again, the proof of this result can be adjusted to obtain the following weak analogue.

Lemma 13. *Let $f: L^n \rightarrow L$ be nondecreasing and weakly $\overline{\mathcal{R}}_f$ -min homogenous (resp. weakly $\overline{\mathcal{R}}_f$ -max homogenous). Then f is weakly $\overline{\mathcal{R}}_f$ -max homogenous (resp. weakly $\overline{\mathcal{R}}_f$ -min homogenous) if and only if it is weakly horizontally $\overline{\mathcal{R}}_f$ -maxitive (resp. weakly horizontally $\overline{\mathcal{R}}_f$ -minitive).*

Proof. Let $f: L^n \rightarrow L$ be nondecreasing and weakly $\overline{\mathcal{R}}_f$ -min homogenous. By Lemma 10, f is $\overline{\mathcal{R}}_f$ -idempotent.

Assume first that f is also weakly $\overline{\mathcal{R}}_f$ -max homogenous. For any $\mathbf{x} \in L_n^{(0,2)}$ and any $c \in \overline{\mathcal{R}}_f$, we have $[\mathbf{x}]_c \in L_n^{(0,2)}$ and hence

$$\begin{aligned} f(\mathbf{x} \wedge c) \vee f([\mathbf{x}]_c) &= (f(\mathbf{x}) \wedge c) \vee f([\mathbf{x}]_c) = (f(\mathbf{x}) \vee f([\mathbf{x}]_c)) \wedge (c \vee f([\mathbf{x}]_c)) \\ &= f(\mathbf{x}) \wedge f(c \vee [\mathbf{x}]_c) \\ &= f(\mathbf{x}). \end{aligned}$$

Therefore, f is weakly horizontally $\overline{\mathcal{R}}_f$ -maxitive.

Now assume that f is weakly horizontally $\overline{\mathcal{R}}_f$ -maxitive and let us prove that it is weakly $\overline{\mathcal{R}}_f$ -max homogenous. For any $\mathbf{x} \in L_n^{(0,2)}$ and any $c \in \overline{\mathcal{R}}_f$, we have

$$f(\mathbf{x} \vee c) = f((\mathbf{x} \vee c) \wedge c) \vee f([\mathbf{x} \vee c]_c) = f(c, \dots, c) \vee f([\mathbf{x}]_c) = c \vee f([\mathbf{x}]_c).$$

Therefore, we have

$$f(\mathbf{x} \vee c) = f(\mathbf{x} \wedge c) \vee f(\mathbf{x} \vee c) = f(\mathbf{x} \wedge c) \vee f([\mathbf{x}]_c) \vee c = f(\mathbf{x}) \vee c$$

and hence f is weakly $\overline{\mathcal{R}}_f$ -max homogenous. The other claim can be verified dually. \square

The following result reassembles Theorem 11, Lemmas 12 and 13, and provides characterizations of the n -ary polynomial functions on a chain L , given in terms of weak homogeneity and weak horizontal minitivity and maxitivity.

Theorem 14. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

- (i) f is a polynomial function.
- (ii) f is nondecreasing, weakly $\overline{\mathcal{R}}_f$ -min homogeneous, and weakly $\overline{\mathcal{R}}_f$ -max homogeneous.
- (iii) f is nondecreasing, weakly $\overline{\mathcal{R}}_f$ -min homogeneous, and weakly horizontally $\overline{\mathcal{R}}_f$ -maxitive.
- (iv) f is nondecreasing, weakly horizontally $\overline{\mathcal{R}}_f$ -minitive, and weakly $\overline{\mathcal{R}}_f$ -max homogeneous.
- (v) f is nondecreasing, $\overline{\mathcal{R}}_f$ -idempotent, weakly horizontally $\overline{\mathcal{R}}_f$ -minitive, and weakly horizontally $\overline{\mathcal{R}}_f$ -maxitive.

4.3 Weak median decomposability

As we saw in Theorem 1, in the case of distributive lattices L , the n -ary polynomial functions on L are exactly those which satisfy the median decomposition formula (5). As we are going to see, in the case of chains, this condition can be relaxed by restricting the satisfaction of (5) by a function $f: L^n \rightarrow L$ to the vectors of $L_n^{(0,2)} \cup L_n^{(1,3)}$. In the latter case, we say that $f: L^n \rightarrow L$ is *weakly median decomposable*.

Lemma 15. *Assume L is a chain and let $f: L^n \rightarrow L$ be nondecreasing. If f is weakly median decomposable, then it is $\overline{\mathcal{R}}_f$ -idempotent.*

Proof. Let $c \in \overline{\mathcal{R}}_f$ and suppose $f(c, \dots, c) > c$. Then, by weak median decomposability, we have

$$f(c, \dots, c) = \text{median}(f(0, c, \dots, c), c, f(1, c, \dots, c)) = f(0, c, \dots, c).$$

By applying the same argument, we obtain $f(c, \dots, c) = f(0, 0, c, \dots, c)$, and finally

$$c < f(c, \dots, c) = f(\mathbf{0}) \leq c,$$

that is a contradiction. The case $f(c, \dots, c) < c$ can be dealt with similarly. \square

Proposition 16. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a nondecreasing function. The following conditions are equivalent:*

- (i) f is weakly median decomposable.
- (ii) f weakly $\overline{\mathcal{R}}_f$ -min homogeneous and weakly $\overline{\mathcal{R}}_f$ -max homogeneous.

Proof. We first prove that (ii) \Rightarrow (i). By Theorem 11, f is a polynomial function, and thus, by Theorem 1, it is median decomposable. In particular, it is weakly median decomposable.

Now we prove (i) \Rightarrow (ii). We only need to prove that f is weak $\overline{\mathcal{R}}_f$ -min homogeneous. The other property can be proved dually. Let $\mathbf{x} = \text{median}(c, \mathbf{e}, d) \in L_n^{(0,2)}$, where $c, d \in L$ and $c \leq d$, and let $r \in \overline{\mathcal{R}}_f$. By Lemma 15, we have $f(r, \dots, r) = r$.

- If $r \geq d$ then $f(\mathbf{x} \wedge r) = f(\mathbf{x}) = f(\mathbf{x}) \wedge f(r, \dots, r) = f(\mathbf{x}) \wedge r$.
- If $r \leq c$ then $f(\mathbf{x} \wedge r) = f(r, \dots, r) = f(\mathbf{x}) \wedge f(r, \dots, r) = f(\mathbf{x}) \wedge r$.
- Assume $r \in]c, d[$ and let $K = \{k \in [n] : x_k = d\}$. By weak median decomposability, for any $k \in K$, we have

$$\begin{aligned} f(\mathbf{x} \wedge r) &= f(\text{median}(c, \mathbf{e}, r)) \\ &= f(\text{median}(c, \mathbf{e}, r)_k^1) \wedge (r \vee f(\text{median}(c, \mathbf{e}, r)_k^0)) \\ &= f(\text{median}(c, \mathbf{e}, r)_k^1) \wedge r. \end{aligned}$$

By repeating this process, we finally obtain $f(\mathbf{x} \wedge r) = f(\text{median}(c, \mathbf{e}, 1)) \wedge r$. Since f is nondecreasing, we finally obtain

$$\begin{aligned} f(\mathbf{x} \wedge r) &= f(\text{median}(c, \mathbf{e}, r)) \wedge r \leq f(\text{median}(c, \mathbf{e}, d)) \wedge r \\ &\leq f(\text{median}(c, \mathbf{e}, 1)) \wedge r = f(\mathbf{x} \wedge r) \end{aligned}$$

that is $f(\mathbf{x} \wedge r) = f(\mathbf{x}) \wedge r$. □

Remark 5. Using the binary function f given in Remark 4, we can see that Proposition 16 does not hold in the general case of bounded distributive lattices. Indeed, as observed, f is nondecreasing and both weakly $\overline{\mathcal{R}}_f$ -min homogenous and weakly $\overline{\mathcal{R}}_f$ -max homogenous, but $f(b, b) = 1 \neq b = \text{median}(f(0, b), b, f(1, b))$ which shows that f is not weakly median decomposable.

From Proposition 16 and Theorem 11, we obtain the following description of polynomial functions given in terms of weak median decomposability.

Theorem 17. *Let L be a bounded chain. A nondecreasing function $f: L^n \rightarrow L$ is a polynomial function if and only if it is weakly median decomposable.*

Remark 6. Note that Theorem 17 does not hold if weak median decomposability would have been defined in terms of vectors in $L_n^{(0,2)}$ only. To see this, let $L = \{0, c, 1\}$ and consider the following nondecreasing function $f: L^3 \rightarrow L$, defined by

$$f(x_1, x_2, x_3) = \begin{cases} 1, & \text{if } \text{median}(x_1, x_2, x_3) = 1, \\ c, & \text{if } \text{median}(x_1, x_2, x_3) = x_1 \wedge x_2 \wedge x_3 = c, \\ 0, & \text{otherwise.} \end{cases}$$

It is easy to this that f is median decomposable for vectors in $L_n^{(0,2)}$, but it is not a polynomial function, e.g., we have $f(0, c, c) = 0$ but $f(0, 1, 1) \wedge c = c$.

4.4 Strong idempotency and componentwise range convexity

By Theorem 1, a nondecreasing function $f: L^n \rightarrow L$ is a polynomial function if and only if it is strongly idempotent, has a convex range, and a componentwise convex range. Our next result shows that the condition requiring a convex range becomes redundant in the case when L is a chain, since it becomes a consequence of componentwise range convexity.

Lemma 18. *Let L be a bounded chain. If a nondecreasing function $f: L^n \rightarrow L$ has a componentwise convex range, then it has a convex range.*

Proof. Since L is a chain and f has a componentwise convex range, we have

$$\overline{\mathcal{R}}_f = [f(\mathbf{0}), f(\mathbf{1})] \subseteq \bigcup_{i=1}^n [f(\mathbf{e}_{\{1, \dots, i-1\}}), f(\mathbf{e}_{\{1, \dots, i\}})] \subseteq \mathcal{R}_f \subseteq \overline{\mathcal{R}}_f.$$

Therefore $\overline{\mathcal{R}}_f = \mathcal{R}_f$ and f has a convex range. \square

Using Lemma 18, we obtain the following characterization of polynomial functions which weakens condition (iii) of Theorem 1 when L is a chain.

Theorem 19. *Let L be a bounded chain. A function $f: L^n \rightarrow L$ is a polynomial function if and only if it is nondecreasing, strongly idempotent, and has a componentwise convex range.*

Remark 7. None of the conditions provided in Theorem 19 can be dropped off. For instance, let L be the real interval $[0, 1]$. Clearly, the unary function $f(x) = x^2$ is nondecreasing and has a componentwise convex range, but it is not strongly idempotent. On the other hand, the function $f: L^2 \rightarrow L$ defined by

$$f(x_1, x_2) = \begin{cases} 1, & \text{if } x_1 = x_2 = 1, \\ 0, & \text{otherwise,} \end{cases}$$

is nondecreasing and strongly idempotent but it does not have a componentwise convex range, e.g., both f_1^1 and f_1^2 do not have convex ranges.

In the special case of real interval lattices, i.e., where $L = [a, b]$ for reals $a \leq b$, the property of having a convex range, as well as the property of having a componentwise convex range, are consequences of continuity. More precisely, for nondecreasing functions $f: [a, b]^n \rightarrow \mathbb{R}$, being continuous reduces to being continuous in each variable, and this latter property is equivalent to having a componentwise convex range. In fact, since polynomial functions are continuous, the condition of having a componentwise convex range can be replaced in Theorem 19 by continuity in each variable. Also, by Proposition 9 (iii), we can add continuity and replace $\overline{\mathcal{R}}_f$ by \mathcal{R}_f in Theorems 11 and 14.

Corollary 20. *Assume that L is a bounded real interval $[a, b]$. A function $f: L^n \rightarrow L$ is a polynomial function if and only if it is nondecreasing, strongly idempotent, and continuous (in each variable).*

4.5 Comonotonic maxitivity and minitivity

Let L be a bounded chain. Two vectors $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{x}' = (x'_1, \dots, x'_n)$ in L^n are said to be *comonotonic* if there exists a permutation σ on $[n]$ such that

$$x_{\sigma(1)} \leq x_{\sigma(2)} \leq \dots \leq x_{\sigma(n)} \quad \text{and} \quad x'_{\sigma(1)} \leq x'_{\sigma(2)} \leq \dots \leq x'_{\sigma(n)}.$$

Equivalently, \mathbf{x} and \mathbf{x}' are comonotonic if $(x_i - x_j)(x'_i - x'_j) \geq 0$ for every $i, j \in [n]$. Thus if $x_i < x_j$ for some $i, j \in [n]$ then $x'_i \leq x'_j$.

A function $f: L^n \rightarrow L$ is said to be

- *comonotonic minitive* if, for any two comonotonic vectors $\mathbf{x}, \mathbf{x}' \in L^n$, we have

$$f(\mathbf{x} \wedge \mathbf{x}') = f(\mathbf{x}) \wedge f(\mathbf{x}').$$

- *comonotonic maxitive* if, for any two comonotonic vectors $\mathbf{x}, \mathbf{x}' \in L^n$, we have

$$f(\mathbf{x} \vee \mathbf{x}') = f(\mathbf{x}) \vee f(\mathbf{x}').$$

Note that for any $\mathbf{x} \in L^n$ and any $c \in L$, we have that \mathbf{x} and (c, \dots, c) are comonotonic and that $\mathbf{x} \vee c$ and $[\mathbf{x}]^c$ are comonotonic. These facts lead to the following result.

Lemma 21. *Let L be a bounded chain and let S be a nonempty subset of L . If a function $f: L^n \rightarrow L$ is comonotonic minitive (resp. comonotonic maxitive), then it is horizontally S -minitive (resp. horizontally S -maxitive). Moreover, if f is S -idempotent, then it is S -min homogeneous (resp. S -max homogeneous).*

Let σ be a permutation on $[n]$. Clearly, every comonotonic minitive (or comonotonic maxitive) function $f: L^n \rightarrow L$ is nondecreasing on the standard simplex

$$L_\sigma^n = \{\mathbf{x} = (x_1, \dots, x_n) \in L^n : x_{\sigma(1)} \leq x_{\sigma(2)} \leq \dots \leq x_{\sigma(n)}\}.$$

The following lemma shows that this fact can be extended to the whole domain L^n .

Lemma 22. *Let L be a bounded chain. If $f: L^n \rightarrow L$ is comonotonic minitive or comonotonic maxitive, then it is nondecreasing. Furthermore, every nondecreasing unary function is comonotonic minitive and comonotonic maxitive.*

Proof. To see that the last claim holds just note that, on any chain L , we necessarily have $x \leq y$ or $x \geq y$ for every $x, y \in L$. For instance, if $x \leq y$ then we have $f(x \wedge y) = f(x) = f(x) \wedge f(y)$ and $f(x \vee y) = f(y) = f(x) \vee f(y)$.

We now prove the first claim for comonotonic minitive functions. The case of comonotonic maxitive functions is shown similarly. Let $f: L^n \rightarrow L$ be a comonotonic minitive function and consider $\mathbf{x}, \mathbf{x}' \in L^n$ such that $\mathbf{x} \leq \mathbf{x}'$ and $\mathbf{x} \neq \mathbf{x}'$. We show that $f(\mathbf{x}) \leq f(\mathbf{x}')$. For each $i \in [n]$, we denote by \mathbf{y}^i the vector in L^n whose j th component is x'_j if $j \leq i$, and x_j otherwise. As a matter of convenience, let $\mathbf{y}^0 = \mathbf{x}$. Clearly, we have

$$\mathbf{x} = \mathbf{y}^0 \leq \mathbf{y}^1 \leq \dots \leq \mathbf{y}^{n-1} \leq \mathbf{y}^n = \mathbf{x}'.$$

Let $k \in [n]$. If \mathbf{y}^{k-1} and \mathbf{y}^k are comonotonic, then

$$f(\mathbf{y}^{k-1}) = f(\mathbf{y}^{k-1} \wedge \mathbf{y}^k) = f(\mathbf{y}^{k-1}) \wedge f(\mathbf{y}^k) \leq f(\mathbf{y}^k).$$

Otherwise, either there is $j > k$ such that $x_k < x_j < x'_k$ or there is $j < k$ such that $x_k < x'_j < x'_k$. Let \mathbf{y}' be the vector obtained from \mathbf{y}^k by replacing the k th component with x_j if $j > k$ and with x'_j if $j < k$. Then both $\{\mathbf{y}^{k-1}, \mathbf{y}'\}$ and $\{\mathbf{y}', \mathbf{y}^k\}$ constitute pairs of comonotonic vectors and, since $\mathbf{y}^{k-1} < \mathbf{y}' < \mathbf{y}^k$, it follows that $f(\mathbf{y}^{k-1}) \leq f(\mathbf{y}') \leq f(\mathbf{y}^k)$. Since the same argument holds for any $k \in [n]$, we have that $f(\mathbf{x}) \leq f(\mathbf{x}')$. \square

We now have the following characterization of polynomial functions.

Theorem 23. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

- (i) f is a polynomial function.
- (ii) f is weakly $\overline{\mathcal{R}}_f$ -min homogeneous and comonotonic maxitive.
- (iii) f is comonotonic minitive and weakly $\overline{\mathcal{R}}_f$ -max homogeneous.
- (iv) f is $\overline{\mathcal{R}}_f$ -idempotent, weakly horizontally $\overline{\mathcal{R}}_f$ -minitive, and comonotonic maxitive.
- (v) f is $\overline{\mathcal{R}}_f$ -idempotent, comonotonic minitive, and weakly horizontally $\overline{\mathcal{R}}_f$ -maxitive.
- (vi) f is $\overline{\mathcal{R}}_f$ -idempotent, comonotonic minitive, and comonotonic maxitive.

Proof. Using distributivity and the first equality in (7) of Proposition 7, it can be easily verified that every polynomial function is comonotonic maxitive. By the dual argument, it follows that every polynomial function is also comonotonic minitive. Thus we have (i) \Rightarrow (vi). The implications (vi) \Rightarrow (v) and (vi) \Rightarrow (iv) immediately follow from Lemma 21. Then, the implications (iv) \Rightarrow (ii) and (v) \Rightarrow (iii) immediately follow from Lemmas 12 and 22. Finally, the implications (ii) \Rightarrow (i) and (iii) \Rightarrow (i) follow from Lemmas 10, 21, 22, and Theorem 11. \square

Remark 8. (i) The condition requiring $\overline{\mathcal{R}}_f$ -idempotency is necessary in conditions (iv)–(vi) of Theorem 23. For instance, let L be the unit interval $[0, 1]$. Clearly, the unary function $f(x) = x^2$ is nondecreasing and thus comonotonic minitive and comonotonic maxitive. By Lemma 21, it is also horizontally $\overline{\mathcal{R}}_f$ -minitive and horizontally $\overline{\mathcal{R}}_f$ -maxitive. However, it is not a polynomial function.

- (ii) The concept of comonotonic vectors appeared as early as 1952 in Hardy et al. [12]. Comonotonic minitivity and maxitivity were introduced in the context of Sugeno integrals in de Campos et al. [6]. An interpretation of these properties was given by Ralescu and Ralescu [17] in the framework of aggregation of fuzzy subsets.

5 Some special classes of polynomial functions

In this final section, we consider two noteworthy subclasses of polynomial functions, namely, those of discrete Sugeno integrals and of term functions, and provide their characterizations. Further subclasses, such as those of symmetric polynomial functions and weighted minimum and maximum functions, were investigated and characterized in [4].

5.1 Discrete Sugeno integrals

A function $f: L^n \rightarrow L$ is said to be *idempotent* if it is L -idempotent.

Fact 24. *A polynomial function is $\{0, 1\}$ -idempotent if and only if it is idempotent.*

In Marichal [13, §4], $\{0, 1\}$ -idempotent polynomial functions are referred to as *discrete Sugeno integrals*. They coincide exactly with those functions $\mathcal{S}_\mu: L^n \rightarrow L$ for which there is a fuzzy measure μ such that

$$\mathcal{S}_\mu(\mathbf{x}) = \bigvee_{I \subseteq [n]} \left(\mu(I) \wedge \bigwedge_{i \in I} x_i \right).$$

(A fuzzy measure μ is simply a set function $\mu: 2^{[n]} \rightarrow L$ satisfying $\mu(I) \leq \mu(I')$ whenever $I \subseteq I'$, and $\mu(\emptyset) = 0$ and $\mu([n]) = 1$.)

For idempotent polynomial functions, Proposition 7 reduces to the following statement.

Proposition 25. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

(i) *f is a discrete Sugeno integral.*

(ii) *For any permutation σ on $[n]$ and every $\mathbf{x} \in L_\sigma^n$, we have*

$$\begin{aligned} f(\mathbf{x}) &= \bigvee_{i \in [n]} \left(\alpha_f(S_\sigma^\uparrow(i)) \wedge x_{\sigma(i)} \right) = \bigwedge_{i \in [n]} \left(\alpha_f(S_\sigma^\uparrow(i+1)) \vee x_{\sigma(i)} \right) \\ &= \text{median} \left(x_1, \dots, x_n, \alpha_f(S_\sigma^\uparrow(2)), \dots, \alpha_f(S_\sigma^\uparrow(n)) \right). \end{aligned}$$

(iii) *For any permutation σ on $[n]$ and every $\mathbf{x} \in L_\sigma^n$, we have*

$$\begin{aligned} f(\mathbf{x}) &= \bigvee_{i \in [n]} \left(\beta_f(S_\sigma^\downarrow(i-1)) \wedge x_{\sigma(i)} \right) = \bigwedge_{i \in [n]} \left(\beta_f(S_\sigma^\downarrow(i)) \wedge x_{\sigma(i)} \right) \\ &= \text{median} \left(x_1, \dots, x_n, \beta_f(S_\sigma^\downarrow(1)), \dots, \beta_f(S_\sigma^\downarrow(n-1)) \right). \end{aligned}$$

The following proposition shows how polynomial functions relate to Sugeno integrals; see Marichal [13, Proposition 12].

Proposition 26. *For any polynomial function $f: L^n \rightarrow L$ there is a fuzzy measure $\mu: 2^{[n]} \rightarrow L$ such that $f(\mathbf{x}) = \langle \mathcal{S}_\mu(\mathbf{x}) \rangle_f$.*

We say that a function $f: L^n \rightarrow L$ is *Boolean min homogeneous* (resp. *Boolean max homogeneous*) if (1) (resp. (2)) holds for every $\mathbf{x} \in \{0, 1\}^n$ and every $c \in L$. Note that every weakly L -min homogeneous (resp. weakly L -max homogeneous) function is Boolean min homogeneous (resp. Boolean max homogeneous).

Lemma 27. *If a function $f: L^n \rightarrow L$ is Boolean min homogeneous and Boolean max homogeneous, then it is idempotent.*

Proof. For any $c \in L$, we have $c \leq f(\mathbf{0}) \vee c = f(\mathbf{0} \vee c) = f(\mathbf{1} \wedge c) = f(\mathbf{1}) \wedge c \leq c$. \square

The following result provides a variant of Theorem 11.

Theorem 28. *Let L be a bounded chain. A function $f: L^n \rightarrow L$ is a discrete Sugeno integral if and only if it is nondecreasing, Boolean min homogenous, and Boolean max homogenous.*

Proof. Since any discrete Sugeno integral is comonotonic minitive and idempotent, we have $f(\mathbf{e} \wedge c) = f(\mathbf{e}) \wedge f(c, \dots, c) = f(\mathbf{e}) \wedge c$ for every $\mathbf{e} \in \{0, 1\}^n$ and every $c \in L$. Thus, any discrete Sugeno integral is Boolean min homogeneous. We can prove similarly that it is also Boolean max homogeneous.

To show that the conditions are sufficient, let $\mathbf{x} \in L^n$. By nondecreasing monotonicity and binary min homogeneity, for every $I \subseteq [n]$ we have

$$f(\mathbf{x}) \geq f\left(\mathbf{e}_I \wedge \left(\bigwedge_{i \in I} x_i\right)\right) = f(\mathbf{e}_I) \wedge \left(\bigwedge_{i \in I} x_i\right)$$

and thus $f(\mathbf{x}) \geq \bigvee_{I \subseteq [n]} (f(\mathbf{e}_I) \wedge (\bigwedge_{i \in I} x_i))$. The converse inequality can be obtained by following exactly the same steps as in the proof of Theorem 11. \square

Remark 9. (i) Theorem 28 as well as the characterization of the discrete Sugeno integrals obtained by combining $\{0, 1\}$ -idempotency with (vi) in Theorem 23 were presented in the case of real variables by Marichal [14, §4.3]; see also Marichal [15].

(ii) Even though Theorem 28 can be derived from condition (ii) of Theorem 14 by simply modifying the two homogeneity properties, to proceed similarly with conditions (iii) and (iv), it is necessary to add the conditions of $\{1\}$ -idempotency and $\{0\}$ -idempotency, respectively (and similarly with conditions (ii) and (iii) of Theorem 23). To see this, let L be a bounded chain with at least three elements and consider the unary functions $f(x) = x \wedge d$ and $g(x) = x \vee d$, where $d \in L \setminus \{0, 1\}$. Clearly, f is L -min homogeneous and horizontally L -maxitive and g is L -max homogeneous and horizontally L -minitive. However, neither f nor g is a discrete Sugeno integral. To see that these additions are sufficient, just note that L -min homogeneity (resp. L -max homogeneity) implies $\{0\}$ -idempotency (resp. $\{1\}$ -idempotency).

(iii) Marichal [14, §2.2.3] showed that, when L is a chain, a nondecreasing and idempotent function $f: L^n \rightarrow L$ is Boolean min homogeneous (resp. Boolean max homogeneous) if and only if we have $f(\mathbf{e} \wedge c) \in \{f(\mathbf{e}), c\}$ (resp. $f(\mathbf{e} \vee c) \in \{f(\mathbf{e}), c\}$) for every $\mathbf{e} \in \{0, 1\}^n$ and every $c \in L$.

5.2 Lattice term functions

A function $f: L^n \rightarrow L$ is said to be

- *conservative* if, for every $\mathbf{x} \in L^n$, we have

$$f(\mathbf{x}) \in \{x_1, \dots, x_n\}. \tag{11}$$

- *weakly conservative* if (11) holds for every $\mathbf{x} \in \{0, 1\}^n$.

Remark 10. Conservative functions (also called *quasi-projections*) were defined in the binary case by Pouzet et al. [16].

Proposition 29. *Let $f: L^n \rightarrow L$ be a function. The following conditions are equivalent:*

(i) f is conservative.

(ii) For every nonempty $S \subseteq L$, we have $f(S^n) \subseteq S$.

(iii) For every nonempty $S \subseteq L$ and every $\mathbf{x} \in L^n$, if $f(\mathbf{x}) \in S$ then there exists $i \in [n]$ such that $x_i \in S$.

Proof. Assume that f is conservative. For every nonempty $S \subseteq L$ and every vector $\mathbf{x} \in S^n$, we have $f(\mathbf{x}) \in S$, which proves that (i) \Rightarrow (ii). Now, assume that (ii) holds and suppose that there exist $S \subsetneq L$ and $\mathbf{x} \in (L \setminus S)^n$ such that $f(\mathbf{x}) \in S$. By (ii), we have $f(\mathbf{x}) \in L \setminus S$, a contradiction. Thus (ii) \Rightarrow (iii). Finally, assume that (iii) holds and that f is not conservative, that is, there exists $\mathbf{x} \in L^n$ such that $f(\mathbf{x}) \notin \{x_1, \dots, x_n\}$. Then, choosing $S = L \setminus \{x_1, \dots, x_n\}$ contradicts (iii). This proves (iii) \Rightarrow (i). \square

Clearly, every conservative function is idempotent. Similarly, every weakly conservative function is $\{0, 1\}$ -idempotent. As observed in [4], the term functions are exactly the weakly conservative discrete Sugeno integrals. Similarly, we can readily see that, when L is a chain, the term functions $f: L^n \rightarrow L$ are exactly the conservative discrete Sugeno integrals.

Theorem 30. *Let L be a bounded chain and let $f: L^n \rightarrow L$ be a discrete Sugeno integral. The following conditions are equivalent:*

(i) f is a term function.

(ii) f is conservative.

(iii) f is weakly conservative.

Remark 11. Not all nondecreasing and conservative functions are term functions. For instance, if L is the real unit interval $[0, 1]$, the binary function $f: L^2 \rightarrow L$, given by $f(x_1, x_2) = x_1 \vee x_2$ on $[0.5, 1]^2$ and by $f(x_1, x_2) = x_1 \wedge x_2$ everywhere else, is nondecreasing and conservative, but it is not a term function.

References

- [1] G. Beliakov, A. Pradera, and T. Calvo. *Aggregation Functions: A Guide for Practitioners*. Studies in Fuziness and Soft Computing. Springer, Berlin, 2007.
- [2] P. Benvenuti, R. Mesiar, and D. Vivona. Monotone set functions-based integrals. In *Handbook of measure theory, Vol. I, II*, pages 1329–1379. North-Holland, Amsterdam, 2002.
- [3] S. Burris and H. P. Sankappanavar. *A course in universal algebra*, volume 78 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1981.
- [4] M. Couceiro and J.-L. Marichal. Representations and characterizations of lattice polynomial functions. <http://arxiv.org/abs/0808.2619>
- [5] L. M. de Campos and M. J. Bolaños. Characterization and comparison of Sugeno and Choquet integrals. *Fuzzy Sets and Systems*, 52(1):61–67, 1992.

- [6] L. M. de Campos, M. T. Lamata, and S. Moral. A unified approach to define fuzzy integrals. *Fuzzy Sets and Systems*, 39(1):75–90, 1991.
- [7] D. Dubois and H. Prade. Weighted minimum and maximum operations in fuzzy set theory. *Inform. Sci.*, 39(2):205–210, 1986.
- [8] J. C. Fodor and M. Roubens. Characterization of weighted maximum and some related operations. *Inform. Sci.*, 84(3-4):173–180, 1995.
- [9] R. L. Goodstein. The solution of equations in a lattice. *Proc. Roy. Soc. Edinburgh Sect. A*, 67:231–242, 1965/1967.
- [10] M. Grabisch, J.-L. Marichal, R. Mesiar, and E. Pap. *Aggregation functions*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, UK, to appear in 2009.
- [11] M. Grabisch, T. Murofushi, and M. Sugeno, editors. *Fuzzy measures and integrals*, volume 40 of *Studies in Fuzziness and Soft Computing*. Physica-Verlag, Heidelberg, 2000.
- [12] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge, at the University Press, 1952. 2d ed.
- [13] J.-L. Marichal. Weighted lattice polynomials. *Discrete Mathematics*. to appear. <http://arxiv.org/abs/0706.0570>
- [14] J.-L. Marichal. *Aggregation operators for multicriteria decision aid*. PhD thesis, Institute of Mathematics, University of Liège, Liège, Belgium, December 1998.
- [15] J.-L. Marichal. On Sugeno integral as an aggregation function. *Fuzzy Sets and Systems*, 114(3):347–365, 2000.
- [16] M. Pouzet, I. G. Rosenberg, and M. G. Stone. A projection property. *Algebra Universalis*, 36(2):159–184, 1996.
- [17] A. L. Ralescu and D. A. Ralescu. Extensions of fuzzy aggregation. *Fuzzy Sets and Systems*, 86(3):321–330, 1997.
- [18] S. Rudeanu. *Lattice functions and equations*. Springer Series in Discrete Mathematics and Theoretical Computer Science. Springer-Verlag London Ltd., London, 2001.
- [19] M. Sugeno. *Theory of fuzzy integrals and its applications*. PhD thesis, Tokyo Institute of Technology, Tokyo, 1974.
- [20] M. Sugeno. Fuzzy measures and fuzzy integrals—a survey. In M. M. Gupta, G. N. Saridis, and B. R. Gaines, editors, *Fuzzy automata and decision processes*, pages 89–102. North-Holland, New York, 1977.