

Schwinger Pair Production at Finite Temperature in QED

Sang Pyo Kim*

*Department of Physics, Kunsan National University, Kunsan 573-701, Korea and
Asia Pacific Center for Theoretical Physics, Pohang 790-784, Korea*

Hyun Kyu Lee[†] and Yongsung Yoon[‡]

Department of Physics, Hanyang University, Seoul 133-791, Korea

(Dated: November 24, 2018)

We use the evolution operator method to find the Schwinger pair-production rate at finite temperature in scalar and spinor QED by counting the vacuum production, the induced production and the stimulated annihilation from the initial ensemble. It is shown that the pair-production rate for each state is factorized into the mean number at zero temperature and the initial thermal distribution for bosons and fermions.

PACS numbers: 12.20.-m, 13.40.-f, 11.10.Wx, 11.15.Tk

I. INTRODUCTION

Vacuum polarization and pair production has been an issue of continuous concern since the early works by Sauter, Heisenberg and Euler, and Weisskopf [1], and then by Schwinger [2] (for a review and references, see Ref. [3]). The task of directly computing, without relying on the electromagnetic duality, the effective action in electric field backgrounds, however, has been a challenging problem due to the vacuum instability. Dunne and Hall used the resolvent method to directly find the effective action in time-dependent electric fields [4]. In the previous paper, employing the evolution operator method, we found the exact one-loop effective actions of scalar and spinor QED at zero temperature in a constant or a pulsed electric field of Sauter-type, which satisfy the exact relation $2\text{Im}\mathcal{L}_{\text{eff}} = \pm \sum_n \ln(1 \pm \overline{\mathcal{N}}_n)$ (with + for scalar and – for spinor) between the imaginary part of the effective Lagrangian density \mathcal{L}_{eff} and the mean number of created pairs $\overline{\mathcal{N}}_n$ at state n [5].

To calculate the effective action and thereby Schwinger pair production at finite temperature is another challenging problem in QED. In a constant pure magnetic field the QED effective action was studied at finite temperature [6] and at finite temperature and density [7]. However, the presence of an additional electric field raised Schwinger pair production at debate depending on the formalism employed. The effective action in both a constant magnetic and electric field in the proper time formalism has an imaginary part that shows thermal enhancement of pair production at high temperature [8]. On the other hand, in the real-time formalism no imaginary part was found in the QED effective action in the presence of both magnetic and electric field [9]. In different formalism, thermal effects have been found on the pair-production rate [10, 11]. The recent calculation of effective action in the imaginary-time formalism shows an imaginary part only at two-loop but not at one-loop in both a constant electric and magnetic field [12].

In this paper, using the evolution operator method, we find the pair-production rate at finite temperature in time-dependent electric fields both in scalar and spinor QED. The evolution operator, unitarily transforming the particle and antiparticle operators from the ingoing vacuum to the outgoing vacuum, carries all the information of quantum evolution. In fact, the evolution operator is completely determined by the Bogoliubov coefficients. The advantage of the evolution operator is the readiness to calculate the probability for transitions among multi-particles states. This allows us to compute the mean number of created pairs at finite temperature in scalar and spinor QED by counting the pairs from the vacuum and the induced production and the stimulated annihilation in a thermal ensemble of bosons and fermions. We find that the mean number of created pairs is factorized into the mean number of created pairs at zero temperature and the initial thermal distribution for bosons and fermions.

The organization of this paper is as follows. In Sec. II, rewriting the Bogoliubov transformation as a unitary transformation by the evolution operator, we find the mean number of created pairs at zero temperature in scalar and spinor QED. In Sec. III, we calculate the mean number of created pairs at finite temperature.

*Electronic address: sangkim@kunsan.ac.kr

†Electronic address: hyunkyuu@hanyang.ac.kr

‡Electronic address: cem@hanyang.ac.kr

II. EVOLUTION OPERATOR AND PAIR PRODUCTION AT $T = 0$

We first consider scalar QED for spinless charged bosons under an external electric field with the gauge field A_μ . The electric field is assumed to be acting on for a finite period of time so that the ingoing and the outgoing vacua are well-defined. Thus, at $t_{\text{in}} = -\infty$, before the external electric field being turned on, the scalar field is free and the Hamiltonian takes the usual form

$$H_{\text{in}}^{\text{sc}} = \int \frac{d^3\mathbf{k}}{(2\pi)^3} \omega_{\mathbf{k},\text{in}} N_{\mathbf{k},\text{in}} = \int \frac{d^3\mathbf{k}}{(2\pi)^3} \omega_{\mathbf{k},\text{in}} (a_{\mathbf{k},\text{in}}^\dagger a_{\mathbf{k},\text{in}} + b_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}}), \quad (1)$$

where $\omega_{\mathbf{k},\text{in}}$ is the initial frequency at momentum \mathbf{k} . Here, the gauge is chosen $A_\mu = 0$, so that the ingoing vacuum $|0; t_{\text{in}}\rangle$ is the Minkowski vacuum $|0\rangle_{\text{M}}$, annihilated by $a_{\mathbf{k}}(t_{\text{in}})$ and $b_{\mathbf{k}}(t_{\text{in}})$ for each momentum \mathbf{k} . Similarly, the outgoing vacuum at $t_{\text{out}} = \infty$ is defined by $a_{\mathbf{k}}(t_{\text{out}})$ and $b_{\mathbf{k}}(t_{\text{out}})$. These operators are related through the Bogoliubov transformations [13]

$$a_{\mathbf{k},\text{out}} = \mu_{\mathbf{k}} a_{\mathbf{k},\text{in}} + \nu_{\mathbf{k}}^* b_{\mathbf{k},\text{in}}^\dagger, \quad b_{\mathbf{k},\text{out}} = \mu_{\mathbf{k}} b_{\mathbf{k},\text{in}} + \nu_{\mathbf{k}}^* a_{\mathbf{k},\text{in}}^\dagger, \quad (2)$$

where $|\mu_{\mathbf{k}}|^2 - |\nu_{\mathbf{k}}|^2 = 1$.

To express the outgoing vacuum as multi-particle states of the ingoing vacuum, we rewrite the Bogoliubov transformations (2) as a unitary transformation [5]

$$a_{\mathbf{k},\text{out}}(A) = U_{\mathbf{k}}(A) a_{\mathbf{k},\text{in}}(0) U_{\mathbf{k}}^\dagger(A), \quad b_{\mathbf{k},\text{out}}(A) = U_{\mathbf{k}}(A) b_{\mathbf{k},\text{in}}(0) U_{\mathbf{k}}^\dagger(A). \quad (3)$$

Here, the evolution operator $U_{\mathbf{k}}$ is factorized into the overall phase factor and the two-mode squeeze operator as [14, 15]

$$U_{\mathbf{k}}(A) = e^{i\theta_{\mathbf{k}}(a_{\mathbf{k},\text{in}}^\dagger a_{\mathbf{k},\text{in}} + b_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}} + 1)} e^{\xi_{\mathbf{k}} a_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}}^\dagger} e^{\frac{\gamma_{\mathbf{k}}}{2}(a_{\mathbf{k},\text{in}}^\dagger a_{\mathbf{k},\text{in}} + b_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}} + 1)} e^{-\xi_{\mathbf{k}}^* a_{\mathbf{k},\text{in}} b_{\mathbf{k},\text{in}}}, \quad (4)$$

where

$$e^{2i\theta_{\mathbf{k}}} = \frac{\mu_{\mathbf{k}}^*}{\mu_{\mathbf{k}}}, \quad \xi_{\mathbf{k}} = \frac{\nu_{\mathbf{k}}^*}{\mu_{\mathbf{k}}}, \quad \gamma_{\mathbf{k}} = -2 \ln(\mu_{\mathbf{k}}). \quad (5)$$

From the charge neutrality of the vacuum, equal numbers of particles and antiparticles are produced at zero temperature and they carry the opposite momenta due to the momentum conservation. The multi-particle state of n -pairs can be concisely denoted as $|n_{\mathbf{k}}, t\rangle = |n_{\mathbf{k}}; \bar{n}_{\mathbf{k}}; t\rangle$. The probability for n -pairs with momentum \mathbf{k} to be created from the vacuum is

$$P_n(\mathbf{k}) = |\langle n_{\mathbf{k}}, \text{out} | 0, \text{in} \rangle|^2 = |\langle n_{\mathbf{k}}, \text{in} | U_{\mathbf{k}}^\dagger | 0, \text{in} \rangle|^2 = e^{\gamma_{\mathbf{k}}} |\xi_{\mathbf{k}}|^{2n}. \quad (6)$$

Note that $P_0 = e^{\gamma_{\mathbf{k}}}$ and $P_1 = e^{\gamma_{\mathbf{k}}} |\xi_{\mathbf{k}}|^2$ so that $P_n = P_0 (P_1/P_0)^n$ and $\sum_{n=0}^{\infty} P_n = 1$ for each \mathbf{k} . Thus, at zero temperature, the mean number of pairs created from the vacuum for each momentum per unit volume is

$$\bar{N}_{\mathbf{k}}^{\text{sc}}(T=0) = \sum_{n=0}^{\infty} n P_n(\mathbf{k}) = |\nu_{\mathbf{k}}|^2. \quad (7)$$

Next, in spinor QED, before the interaction of an external electric field, the spinor field is free without the gauge potential ($A_\mu = 0$), and has the Hamiltonian given by

$$H_{\text{in}}^{\text{sp}} = \sum_{\sigma} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \omega_{n,\text{in}} N_{n,\text{in}} = \sum_{\sigma} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \omega_{n,\text{in}} (b_{n,\text{in}}^\dagger b_{n,\text{in}} + d_{n,\text{in}}^\dagger d_{n,\text{in}}), \quad (8)$$

where $b_{n,\text{in}}$ and $d_{n,\text{in}}$ are particle and antiparticle operators in the ingoing vacuum. After the interaction of the electric field, the ingoing vacuum evolves to the outgoing vacuum, whose particle and antiparticle operator are $b_{n,\text{out}}$ and $d_{n,\text{out}}$. The Bogoliubov transformations between the ingoing and the outgoing particle and antiparticle operators, b_n, d_n , are similarly given by

$$b_{n,\text{out}} = \mu_n b_{n,\text{in}} + i\nu_n^* d_{n,\text{in}}^\dagger, \quad d_{n,\text{out}} = \mu_n d_{n,\text{in}} - i\nu_n^* b_{n,\text{in}}^\dagger, \quad (9)$$

where $|\mu_n|^2 + |\nu_n|^2 = 1$ and $n = (\mathbf{k}, \sigma)$ with $\sigma = \pm 1/2$. The Bogoliubov transformation can be also written as a unitary transformation [16]

$$b_{n,\text{out}} = U_n b_{n,\text{in}} U_n^\dagger, \quad d_{n,\text{out}} = U_n d_{n,\text{in}} U_n^\dagger, \quad (10)$$

where

$$U_n = e^{\xi_n b_{n,\text{in}}^\dagger d_{n,\text{in}}^\dagger} e^{(\frac{\gamma_n}{2} + i\theta_n)(b_{n,\text{in}}^\dagger b_{n,\text{in}} + d_{n,\text{in}}^\dagger d_{n,\text{in}} - 1)} e^{e^{2i\theta_n} \xi_n^* b_{n,\text{in}} d_{n,\text{in}}}. \quad (11)$$

Here, the three parameters ξ_n , γ_n and θ_n are determined by the Bogoliubov coefficients as

$$\xi_n = -i \frac{\nu_n^*}{\mu_n}, \quad \gamma_n = -2 \ln(|\mu_n|), \quad e^{2i\theta_n} = \frac{\mu_n^*}{\mu_n}. \quad (12)$$

Note that the pair production on spinor QED is restricted to only one pair of particle and antiparticle for a given quantum number n due to the Pauli exclusion principle. Thus, the mean number of pairs created from the vacuum for each state n at zero temperature is calculated as

$$\bar{\mathcal{N}}_n^{\text{SP}}(T=0) = |\langle 1_n, \text{out} | 0, \text{in} \rangle|^2 = |\nu_n|^2. \quad (13)$$

III. PAIR PRODUCTION AT $T \neq 0$

We now calculate the mean number of pairs at finite temperature from the probability for each transition, as in the case of zero temperature. As there is no mode-mixing, we separately calculate the mean number of created pairs for each mode. For an initial thermal ensemble at $\beta = 1/kT$, which might not be charge neutral, the mean number of produced pairs consists of the vacuum pair production, the induced pair production, and the stimulated pair annihilation as shown in Fig. 1:

$$\begin{aligned} \bar{\mathcal{N}}_n(T) = \frac{1}{Z_n} & \left[\sum_{n_n > 0}^{\infty} e^{-\beta E_{0n,0n}} n_n P_{0n,0n \rightarrow n_n, n_n} - \sum_{p_n > n_n \geq 0, q_n > m_n \geq 0}^{\infty} e^{-\beta E_{p_n, q_n}} (p_n - n_n) P_{p_n, q_n \rightarrow n_n, m_n} \right. \\ & \left. + \sum_{p_n > n_n \geq 0, q_n > m_n \geq 0, n_n + m_n \neq 0}^{\infty} e^{-\beta E_{n_n, m_n}} (p_n - n_n) P_{n_n, m_n \rightarrow p_n, q_n} \right], \quad (14) \end{aligned}$$

where Z_n is the partition function for the initial ensemble and $P_{n_n, m_n \rightarrow p_n, q_n} = |\langle p_n, q_n, \text{in} | U_n | n_n, m_n, \text{in} \rangle|^2$ is the transition probability from $|n_n, m_n, \text{in}\rangle$ to $|p_n, q_n, \text{in}\rangle$. Using $|\langle p_n, q_n, \text{in} | U_n | n_n, m_n, \text{in} \rangle|^2 = |\langle p_n, q_n, \text{in} | U_n^\dagger | n_n, m_n, \text{in} \rangle|^2$, which implies $P_{p_n, q_n \rightarrow n_n, m_n} = P_{n_n, m_n \rightarrow p_n, q_n}$, the mean number of pairs created at finite temperature, Eq. (14), can be written as

$$\begin{aligned} \bar{\mathcal{N}}_n(T) &= \frac{1}{2Z_n} \sum_{p_n > n_n, q_n > m_n}^{\infty} \left[\langle n_n, m_n, \text{in} | e^{-\beta H_{n,\text{in}}} U_n | p_n, q_n, \text{in} \rangle \langle p_n, q_n, \text{in} | (N_{n,\text{in}} U_n^\dagger - U_n^\dagger N_{n,\text{in}}) | n_n, m_n, \text{in} \rangle \right. \\ & \quad \left. - \langle n_n, m_n, \text{in} | U_n^\dagger e^{-\beta H_{n,\text{in}}} | p_n, q_n, \text{in} \rangle \langle p_n, q_n, \text{in} | (N_{n,\text{in}} U_n - U_n N_{n,\text{in}}) | n_n, m_n, \text{in} \rangle \right] \\ &= \frac{1}{2Z_n} \text{Tr}(U_{n,\text{in}} N_{n,\text{in}} U_{n,\text{in}}^\dagger - N_{n,\text{in}}) e^{-\beta H_{n,\text{in}}}, \quad (15) \end{aligned}$$

where $N_{n,\text{in}} = a_{\mathbf{k},\text{in}}^\dagger a_{\mathbf{k},\text{in}} + b_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}}$ for scalar particles and $N_{n,\text{in}} = b_{n,\text{in}}^\dagger b_{n,\text{in}} + d_{n,\text{in}}^\dagger d_{n,\text{in}}$ for spinor particles and $H_{n,\text{in}} = \omega_{n,\text{in}} N_{n,\text{in}}$ for both.

From the Bogoliubov transformations (3) and (10), we have $N_{n,\text{out}} = U_n N_{n,\text{in}} U_n^\dagger$. Thus, the mean number of pairs created at finite temperature can be written concisely as

$$\bar{\mathcal{N}}_n(T) = \frac{1}{2Z_n} \text{Tr}(N_{n,\text{out}} - N_{n,\text{in}}) e^{-\beta H_{n,\text{in}}}. \quad (16)$$

First, for scalar particles, from Eq. (2),

$$N_{\mathbf{k},\text{out}} - N_{\mathbf{k},\text{in}} = 2|\nu_{\mathbf{k}}|^2 (a_{\mathbf{k},\text{in}}^\dagger a_{\mathbf{k},\text{in}} + b_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}} + 1) + 2\mu_{\mathbf{k}} \nu_{\mathbf{k}} a_{\mathbf{k},\text{in}} b_{\mathbf{k},\text{in}} + 2\mu_{\mathbf{k}}^* \nu_{\mathbf{k}}^* a_{\mathbf{k},\text{in}}^\dagger b_{\mathbf{k},\text{in}}^\dagger, \quad (17)$$




FIG. 1: (a) the vacuum production, (b) the induced production, and (c) the stimulated annihilation

and $Z_{\mathbf{k}} = e^{\beta\omega_{\mathbf{k},\text{in}}}(2 \sinh \beta\omega_{\mathbf{k},\text{in}}/2)^{-2}$, the mean number of scalar pairs created in momentum \mathbf{k} at finite temperature is given by

$$\bar{N}_{\mathbf{k}}^{\text{sc}}(T) = \frac{|\nu_{\mathbf{k}}|^2}{Z_{\mathbf{k}}} \sum_{n,m=0}^{\infty} \langle n, m | (n+m+1) e^{-(n+m)\beta\omega_{\mathbf{k},\text{in}}} | n, m \rangle = |\nu_{\mathbf{k}}|^2 \coth \frac{\beta\omega_{\mathbf{k},\text{in}}}{2}. \quad (18)$$

Second, for spinor particles, from Eq. (9),

$$N_{n,\text{out}} - N_{n,\text{in}} = -2|\nu_n|^2 (b_{n,\text{in}}^\dagger b_{n,\text{in}} + d_{n,\text{in}}^\dagger d_{n,\text{in}} - 1) + 2i\mu_n \nu_n b_{n,\text{in}} d_{n,\text{in}} + 2i\mu_n^* \nu_n^* b_{n,\text{in}}^\dagger d_{n,\text{in}}^\dagger, \quad (19)$$

and $Z_n = (1 + e^{-\beta\omega_{n,\text{in}}})^2$, we find the mean number of spinor pairs created in state n at finite temperature

$$\bar{N}_n^{\text{sp}}(T) = -\frac{|\nu_n|^2}{Z_n} \sum_{n,m=0}^1 \langle n, m | (n+m-1) e^{-(n+m)\beta\omega_{n,\text{in}}} | n, m \rangle = |\nu_n|^2 \tanh \frac{\beta\omega_{n,\text{in}}}{2}. \quad (20)$$

Note that contrary to the bosonic pair production (18), the fermionic pair production (20) decreases as temperature increases as shown in Fig.2, which is expected from the Pauli exclusion principle.




FIG. 2: The mean number of produced pairs for scalar(a) and spinor(b): $\bar{N}_n(T)/\bar{N}_n(0)$ versus temperature $kT/\omega_{n,\text{in}}$

IV. CONCLUSION

In this paper, using the evolution operator method, we found that the mean number of created pairs in state n at finite temperature is given by

$$\bar{N}_n(T) = \frac{1}{2Z_n} \text{Tr}(N_{n,\text{out}} - N_{n,\text{in}}) e^{-\beta H_{n,\text{in}}}. \quad (21)$$

For scalar and spinor QED in external electric fields, the total mean number density of created pairs at finite temperature is given by

$$\bar{\mathcal{N}}^{\text{sc}}(T) = \int \frac{d^3\mathbf{k}}{(2\pi)^3} \bar{\mathcal{N}}_{\mathbf{k}}^{\text{sc}}(0) \coth \frac{\beta\omega_{\mathbf{k},\text{in}}}{2} \quad (\text{for scalar}), \quad (22)$$

$$\bar{\mathcal{N}}^{\text{sp}}(T) = \sum_{\sigma} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \bar{\mathcal{N}}_n^{\text{sp}}(0) \tanh \frac{\beta\omega_{n,\text{in}}}{2} \quad (\text{for spinor}), \quad (23)$$

where $\bar{\mathcal{N}}_{\mathbf{k}}^{\text{sc}}(0)$ and $\bar{\mathcal{N}}_n^{\text{sp}}(0)$ are the mean number of created pairs at zero temperature for scalar and spinor QED, respectively.

Acknowledgments

S. P. K. would like to appreciate the hospitality of Hanyang University, and H. K. L. and Y. Y. would like to appreciate the hospitality of Kunsan National University. The work of S. P. K. was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) (KRF-2007-C00167) and the work of H. K. L. was supported by the Korea Science and Engineering Foundation (KOSEF) grant funded by the Korea government (MOST) (No. R01-2006-000-10651-0).

-
- [1] F. Sauter, Z. Phys. **69**, 742 (1931); W. Heisenberg and H. Euler, Z. Physik **98**, 714 (1936); V. Weisskopf, K. Dan. Vidensk. Selsk. Mat. Fys. Medd. **XIV**, No. 6 (1936).
 - [2] J. Schwinger, Phys. Rev. **82**, 664 (1951).
 - [3] G. V. Dunne, "Heisenberg-Euler Effective Lagrangians: Basics and Extensions," *From Fields to Strings: Circumnavigating Theoretical Physics*, edited by M. Shifman, A. Vainshtein, and J. Wheeler, (World Scientific, Singapore, 2005), Vol. I, pp. 445-522, hep-th/0406216.
 - [4] G. V. Dunne and T. Hall, Phys. Rev. D **58**, 105022 (1998).
 - [5] S. P. Kim, H. K. Lee, and Y. Yoon, "Effective Action of QED in Electric Field Backgrounds," arXiv:0807.2696.
 - [6] W. Dittrich, Phys. Rev. D **19**, 2385 (1979).
 - [7] P. Elmfors, D. Persson, and B.-S. Skagerstam, Phys. Rev. Lett. **71**, 480 (1993); P. Elmfors, P. Liljenberg, D. Persson, and B.-S. Skagerstam, Phys. Rev. D **51**, 5885 (1995).
 - [8] M. Loewe and J. C. Rojas, Phys. Rev. D **46**, 2689 (1992).
 - [9] P. Elmfors and B.-S. Skagerstam, Phys. Lett. B **348**, 141 (1995).
 - [10] A. K. Ganguly, P. K. Kaw, and J. C. Parikh, Phys. Rev. C **51**, 2091 (1995).
 - [11] J. Hallin and P. Liljenberg, Phys. Rev. D **52**, 1150 (1995).
 - [12] H. Gies, Phys. Rev. D **60**, 105002 (1999); Phys. Rev. D **61**, 085021 (2000).
 - [13] S. P. Kim and H. K. Lee, Phys. Rev. D **76**, 125002 (2007).
 - [14] C. M. Caves and B. L. Schumaker, Phys. Rev. A **31**, 3068 (1985).
 - [15] B. L. Schumaker and C. M. Caves, Phys. Rev. A **31**, 3093 (1985).
 - [16] H.-Y. Fan and C. Li, J. Opt. B: Quantum Semiclass. Opt **6**, S502 (2004).