

Dipole emission and coherent transport in random media II. Density of states vs. index of refraction

M. Donaire^{1,*}

¹*Departamento de Física de la Materia Condensada,
Universidad Autónoma de Madrid, E-28049 Madrid, Spain.*

This is the second of a series of papers devoted to develop a microscopical approach to the dipole emission process and its relation to coherent transport in random media. In this Letter, we deduce a relation between the transverse decay rate of an emitter in a virtual cavity and the complex refraction index of the host medium. We argue on the possibility of a criterion for inhibition/enhancement of spontaneous emission in function of the transition frequency and the correlation length of the host scatterers. In addition, we study the radiative/non-radiative nature of the net power emission through a microscopical analysis of the scattering events involved. This study reveals essential discrepancies with previous interpretations.

PACS numbers: 42.25.Dd,05.60.Cd,42.25.Bs,42.25.Hz,03.75.Nt

In a previous work [1] we computed the average local density of states (LDOS) within an isotropic and homogeneous medium. It is the aim of this Letter to use those equations to deduce a relation between LDOS and coherent transport parameters. Recent experiments [2] have put forward evidences of such a relation. In [1], several scenarios were addressed for an emitter seated in either a virtual or a real cavity. In this Letter we focus on a virtual cavity scenario. We will consider the host medium made of a set of spatially disconnected point scatterers of density ρ and the emitter as a fluorescence particle seated on top of a host scatterer. We will refer to that scatterer as *host particle*. The emitter transition frequency and the resonance frequency of the host particle are assumed far apart from each other. Dipole induction is completely cast in the host particle polarizability. A bulk propagator \bar{G} and electrical susceptibility $\bar{\chi}$ can be unambiguously defined, where overlines denote rank-2 tensors. The decay rate Γ reads [1]

$$\Gamma_{\omega}^{\alpha,\mu} = \frac{-2\omega^2}{3\hbar\epsilon_0 c^2} \frac{|\mu|^2}{|1 + \frac{1}{3}k_0^2\alpha_0[2\gamma_{\perp} + \gamma_{\parallel}]|^2} \left[\Im\{2\gamma_{\perp} + \gamma_{\parallel}\} - \frac{k_0^2}{3} \Im\{\alpha_0\} |2\gamma_{\perp} + \gamma_{\parallel}|^2 \right], \quad (1)$$

where the factors $\gamma_{\perp,\parallel}$ are given by

$$2\gamma_{\perp} = \int \frac{d^3k}{(2\pi)^3} \left[\frac{2\chi_{\perp}(k)/(\rho\alpha)}{k_0^2[1 + \chi_{\perp}(k)] - k^2} \right] - 2\Re\{\gamma_{\perp}^{(0)}\}, \quad (2)$$

$$\gamma_{\parallel} = \int \frac{d^3k}{(2\pi)^3} \left[\frac{1}{\rho\alpha} \frac{\chi_{\parallel}(k)}{k_0^2[1 + \chi_{\parallel}(k)]} - \frac{1}{k_0^2} \right]. \quad (3)$$

In the above formulae, $\alpha_0 \equiv 4\pi a^3 \frac{\epsilon_e - 1}{\epsilon_e + 2}$ is the electrostatic polarizability of the host scatterers, being $\epsilon_e(\omega)$ the single-particle dielectric contrast; $\alpha = \frac{\alpha_0}{1 - \frac{ik_0^3}{6\pi}\alpha_0}$ is the in-vacuum renormalized single-particle polarizability; μ is the spontaneous transition amplitude of the emitter and ω the transition frequency, with $k_0 = \omega/c$.

The term $-2\Re\{\gamma_{\perp}^{(0)}\}$ accounts for the singularity of the radiative part of $\bar{G}^{(0)}(r)$ at $r = 0$ in vacuum (denoted by the script (0)). The denominator in Eq.(1) amounts to self-polarization cycles over the host particle. It gives rise to an effective renormalization of $|\mu|^2$, $|\tilde{\mu}|^2 \equiv \frac{|\mu|^2}{|1 + \frac{1}{3}k_0^2\alpha_0[2\gamma_{\perp} + \gamma_{\parallel}]|^2}$, which depends weakly on γ factors. In a first approach, we can distinguish three contributions to $\Gamma_{\omega}^{\alpha,\mu}$ —we drop the scripts on Γ hereafter, namely

$$\Gamma_{abs.}^0 = \frac{2\omega^4}{9\hbar\epsilon_0 c^4} |\tilde{\mu}|^2 |2\gamma_{\perp} + \gamma_{\parallel}|^2 \Im\{\alpha_0\}, \quad (4)$$

$$2\Gamma_{\perp} = \frac{-4\omega^2}{3\hbar\epsilon_0 c^2} |\tilde{\mu}|^2 \Im\left\{ \int \frac{d^3k}{(2\pi)^3} \frac{\frac{\chi_{\perp}(k)}{\rho\alpha}}{k_0^2[1 + \chi_{\perp}] - k^2} \right\}, \quad (5)$$

$$\Gamma_{\parallel} = \frac{-2\omega^2}{3\hbar\epsilon_0 c^2} |\tilde{\mu}|^2 \Im\left\{ \int \frac{d^3k}{(2\pi)^3} \frac{\frac{\chi_{\parallel}(k)}{\rho\alpha}}{k_0^2[1 + \chi_{\parallel}]} \right\}. \quad (6)$$

Eq.(4) is the decay rate due to absorption within the host particle. The other two equations correspond to transverse and longitudinal decay respectively. Propagating modes are determined by the singularities of the integrands in Eqs.(5,6) in the complex plane, that is, poles and branch cuts. Poles are modes k_{\perp}, k_{\parallel} satisfying the dispersion relations in the effective medium,

$$k^2 - k_0^2[1 + \chi_{\perp}(k)]|_{k=k_{\perp}} = 0, \quad 1 + \chi_{\parallel}(k)|_{k=k_{\parallel}} = 0. \quad (7)$$

The refraction index, \bar{n} , and the extinction mean free path, l_{ext} , parametrize propagation over long distances. That is, they correspond to the longest wave length solutions for k_{\perp} . We will disregard the possibility of propagating longitudinal modes as these do not affect directly the relation between Γ and the index of refraction. As a matter of fact, propagation of longitudinal modes requires some fine tuning on the arrangement of scatterers [3]. Because of isotropy, $\bar{\chi}$ is a function of k^2 . It is only for $k_0\xi \ll 1$ that this dependence is weak, ξ being the typical correlation length between host scatterers. For

a virtual cavity, the cavity radius R can be identified with ξ as well. It is for $k_0\xi \ll 1$ that $\bar{\chi}$ admits an expansion in powers of k^2/k_0^2 and it suffices to truncate the series at order k^2/k_0^2 . In doing so, Eq.(5) presents a unique pole k_\perp and the whole transverse propagator is characterized solely by \bar{n} and l_{ext} , $\bar{n} = \Re\{\sqrt{1 + \chi_\perp(k_\perp)}\}$, $l_{ext}^{-1} = 2k_0\Im\{\sqrt{1 + \chi_\perp(k_\perp)}\}$. On the other hand, we can distinguish two components in Γ_\perp , namely,

$$2\Gamma_\perp^{NP} = \frac{-4\omega^2|\tilde{\mu}|^2}{3\hbar\epsilon_0c^2} \int \frac{d^3k}{(2\pi)^3} \Im\left\{\frac{\chi_\perp}{\rho\alpha}\right\} \Re\left\{\frac{1}{k_0^2[1 + \chi_\perp] - k^2}\right\}, \quad (8)$$

$$2\Gamma_\perp^P = \frac{-4\omega^2|\tilde{\mu}|^2}{3\hbar\epsilon_0c^2} \int \frac{d^3k}{(2\pi)^3} \Re\left\{\frac{\chi_\perp}{\rho\alpha}\right\} \Im\left\{\frac{1}{k_0^2[1 + \chi_\perp] - k^2}\right\}, \quad (9)$$

where the scripts P and NP stand for *propagating* and *non-propagating* respectively. In Eq.(9) the 'normalized' susceptibility function $\Re\left\{\frac{\chi_\perp(k)}{\rho\alpha}\right\}$ plays a role analogous to that of the field-strength renormalization factor Z in a quantum field theory (QFT). In QFT it is the analytical structure of $G(k)$ together with Z in the complex plane that determines the mass spectrum of quantum particles propagating in space-time [4]. Correspondingly, it is the analytical structure of $G_\perp(k)$ together with $\Re\left\{\frac{\chi_\perp(k)}{\rho\alpha}\right\}$ that determines the spectrum of propagating coherent photons in a random medium. In view of the above classification, we will refer to Γ_{abs}^0 , Γ_\parallel and $2\Gamma_\perp^{NP}$ all together as *non-propagating decay rate*, $\Gamma^{NP} = \Gamma_{abs}^0 + \Gamma_\parallel + 2\Gamma_\perp^{NP}$; while $2\Gamma_\perp^P$ is the only propagating component, $\Gamma^P = 2\Gamma_\perp^P$ –compare with other interpretations [15]. It is however not clear, except for Γ_{abs}^0 , which contribution of Γ^{NP} is associated to absorption by host scatterers and which part corresponds to incoherent radiation. That needs a deeper microscopical analysis. We proceed to compute Γ_\perp^P in function of the extinction coefficient, $\kappa = l_{ext}^{-1}/(2k_0)$, and \bar{n} . On the one hand, we have that $\Re\{\chi_\perp(k_\perp)\} = \bar{n}^2 - 1 - \kappa^2$ and $\Im\{\chi_\perp(k_\perp)\} = 2\bar{n}\kappa$. On the other hand, Eq.(9) yields $\frac{\omega^3|\tilde{\mu}|^2}{3\pi\hbar\epsilon_0c^3} \Re\left\{\frac{\chi_\perp(k)}{\rho\alpha}\right\} \bar{n}$. Putting everything together we end up with

$$\begin{aligned} 2\Gamma_\perp^P &= \frac{2\omega^3|\tilde{\mu}|^2}{3\pi\hbar\epsilon_0c^3} \bar{n} \Re\{\chi_\perp/\rho\alpha\} \\ &= \frac{2\omega^3|\tilde{\mu}|^2}{3\pi\hbar\epsilon_0c^3\rho|\alpha|^2} \bar{n} \left[(\bar{n}^2 - 1 - \kappa^2) \Re\{\alpha\} \right. \\ &\quad \left. + 2\bar{n}\kappa\Im\{\alpha\} \right]. \end{aligned} \quad (10)$$

Taking the approximation $\epsilon \approx \bar{n}^2$ and assuming that Maxwell-Garnett relation (MG) holds, $\epsilon = 1 + \frac{\rho\alpha}{1 - \frac{1}{3}\rho\alpha}$, Eq.(10) goes as $\sim \frac{\epsilon+2}{3}\sqrt{\epsilon}$. The missing local factor with respect to the usual virtual cavity formula [5, 11, 17] implies that only part of the decay rate there is transverse and radiative. We will go over this point later on. Pro-

ceeding in a similar manner, $2\Gamma_\perp^{NP}$ reads

$$\begin{aligned} 2\Gamma_\perp^{NP} &= \frac{-2\omega^3|\tilde{\mu}|^2}{3\pi\hbar\epsilon_0c^3} \left[\kappa\Im\{\chi_\perp/\rho\alpha\} - \frac{\bar{n}\kappa}{2k_0\xi} \right] \\ &= \frac{2\omega^3|\tilde{\mu}|^2}{3\pi\hbar\epsilon_0\rho|\alpha|^2c^3} \kappa \left[(\bar{n}^2 - 1 - \kappa^2) \Im\{\alpha\} \right. \\ &\quad \left. - 2\bar{n}\kappa\Re\{\alpha\} \right] + \frac{\omega^3|\tilde{\mu}|^2}{3\pi\hbar\epsilon_0c^3} \frac{\bar{n}\kappa}{k_0\xi}, \end{aligned} \quad (11)$$

where the ξ dependent terms are approximate as explained below. Further on, invoking causality, $\bar{n}(\omega)$ and




FIG. 1: (a) Feynman's rules –see also Fig.2(a). (b) Diagrammatic representation of the total decay rate in a dilute system, Eqs.(15,23).

$\kappa(\omega)$ are related through integral equations in ω -modes. That is, by Kramers-Kronig relationships (KK). Therefore, one can in principle control the variations in $2\Gamma_\perp$ as a function of ω and so optimize its value once the variations of either $\bar{n}(\omega)$ or $l_{ext}(\omega)$ are known [8]. On the other hand, in order to get full control over the total spontaneous emission we have to find similar expressions for Γ_\parallel . In the limit $k_0\xi \ll 1$ we can approximate $\chi_\parallel \approx \chi_\perp(k_\perp)$. MG belongs indeed to this case. In such a situation it is possible to give a formula for total $\Gamma(\omega)$ in closed form once $\alpha(\omega)$ is known. As the medium surrounding the emitter is continuous beyond some distance $R = \xi$ in application of $k_0\xi \ll 1$, Eq.(6) and the leading order ξ -dependent term of Eq.(8) approximate the formulae for a real cavity in [1] with $\chi(\omega) = \frac{\rho\alpha(\omega)}{1 - \frac{1}{3}\rho\alpha(\omega)}$, yielding the second term in Eq.(11) and (see also [16])

$$\begin{aligned} \Gamma_\parallel &\approx \frac{2\omega^3|\tilde{\mu}|^2}{6\pi\hbar\epsilon_0c^3} \left\{ \frac{2\bar{n}\kappa}{3} \left[\frac{1}{(k_0\xi)^3} + \frac{1}{k_0\xi} \right] \left[1 + \frac{2}{(\bar{n}^2 + \kappa^2)^2} \right] \right. \\ &\quad \left. - \frac{\bar{n}\kappa}{k_0\xi} + \frac{5}{9} \left[1 + \frac{\kappa^6 - \bar{n}^6 + (\kappa^2 - \bar{n}^2)(4 - \kappa^2\bar{n}^2)}{5(\kappa^2 + \bar{n}^2)^2} \right] \right\} \quad (12) \end{aligned}$$

The total decay rate is thus a function of ω and ξ . Optimization of Γ with respect to ω, ξ leads this way to inhibition/enhancement of spontaneous emission.

On the other hand, it is expected that not only the functional form of $\epsilon_e(\omega)$ in host scatterers, but also the spatial correlations among these, cooperate towards inhibition/enhancement. For $\xi \sim k_0^{-1}$ the above treatment fails as, in general, additional poles in Eq.(5) enter the problem and \bar{n} , l_{ext} do not suffice to compute Γ . Full knowledge of $\bar{\chi}(k)$ is needed. Nevertheless, by invoking

energy conservation, Ward's identities for vector waves [9] formulated in a suitable manner may yield integral expressions – sum rules over k -modes – relating real and imaginary parts of $\chi_{\parallel}(k)$ and $\chi_{\perp}(k)$.

Next, we consider a simple case to study microscopically the nature of each of the terms contributing to Γ in Eqs.(4-6). The setup consists of an emitter on top of a host particle in a dilute medium made of spherical inclusions of polarizability α . A similar model was studied in [10, 11]. In this simple model, the only relevant correlation is that given by the two-point function $h(r) = -\Theta(r - \xi)$, ξ being the radius of the exclusion volume surrounding each scatterer. Therefore, the electrical susceptibility reads $\chi_{\parallel, \perp} \simeq \rho\alpha + \chi_{\parallel, \perp}^{(2)}$, where

$$\begin{aligned}\chi_{\perp}^{(2)}(k) &= \frac{-(\alpha\rho)^2 k_0^2}{2} \int d^3r e^{i\vec{k}\cdot\vec{r}} h(r) \text{Tr}\{\bar{G}^{(0)}(r)[\bar{I} - \hat{k} \otimes \hat{k}]\} \\ &= -\frac{(\alpha\rho)^2 k_0^2}{2} \int \frac{d^3k'}{(2\pi)^3} h(|\vec{k}' - \vec{k}|) \left[G_{\perp}^{(0)}(k') \right. \\ &\quad \left. + G_{\perp}^{(0)}(k') \cos^2 \theta + G_{\parallel}^{(0)}(k') \sin^2 \theta \right],\end{aligned}\quad (13)$$

$$\begin{aligned}\chi_{\parallel}^{(2)}(k) &= -(\alpha\rho)^2 k_0^2 \int d^3r e^{i\vec{k}\cdot\vec{r}} h(r) \text{Tr}\{\bar{G}^{(0)}(r)[\hat{k} \otimes \hat{k}]\} \\ &= -(\alpha\rho)^2 k_0^2 \int \frac{d^3k'}{(2\pi)^3} h(|\vec{k}' - \vec{k}|) \\ &\quad \times \left[G_{\parallel}^{(0)}(k') \cos^2 \theta + G_{\perp}^{(0)}(k') \sin^2 \theta \right],\end{aligned}\quad (14)$$

with $\cos \theta \equiv \hat{k} \cdot \hat{k}'$. In the above equations, $G_{\perp, \parallel}^{(0)}$ are the transverse and longitudinal components of the propagator of the electric field in free space, $\bar{G}_{\perp}^{(0)}(k) = \frac{\Delta(\hat{k})}{k_0^2 - k^2}$, $\bar{G}_{\parallel}^{(0)}(k) = \frac{\hat{k} \otimes \hat{k}}{k_0^2}$; \hat{k} being a unitary vector along the propagation direction and $\Delta(\hat{k}) \equiv I - \hat{k} \otimes \hat{k}$ being the projective tensor orthogonal to the propagation direction. In spatial space, $\bar{G}^{(0)}(r) = \frac{e^{ik_0 r}}{-4\pi r} \bar{I} + \left[\frac{1}{k_0} \bar{\nabla} \otimes \bar{\nabla} \right] \frac{e^{ik_0 r}}{-4\pi r}$. At leading order in $\rho\alpha$, the problem becomes one of single scattering. Therefore, the only correlation that matters is that of the emitter with the host scatterers, $h(r)$.

Pictorially, upon self-polarization cycles, Γ is given by the first three diagrams of Fig.1(b). It reads,

$$\begin{aligned}\Gamma &= \Gamma_0 - \frac{2\omega^2 |\mu|^2}{3\hbar\epsilon_0 c^2} \Im \left\{ \text{Tr} \left\{ \int d^3r_1 \bar{G}^{(0)}(\vec{r}_1) (-k_0^2 \rho\alpha) \right. \right. \\ &\quad \left. \left. \times \bar{G}^{(0)}(\vec{r}_1) [1 - \Theta(r_1 - \xi)] \right\} \right\},\end{aligned}\quad (15)$$

where $\Gamma_0 \equiv \frac{\omega^3 |\mu|^2}{3\pi \hbar \epsilon_0 c^3}$ is the total decay rate in vacuum disregarding self-polarization effects. In the limit $k_0 \xi \ll 1$, we expand Γ in powers of $k_0 \xi$,

$$\Gamma \simeq \Gamma_0 \left\{ 1 + \Re\{\alpha\rho\} \left[\frac{7}{6} - \frac{11}{15} (k_0 \xi)^2 \right] \right. \quad (16)$$

$$\left. + \Im\{\alpha\rho\} \left[\frac{1}{(k_0 \xi)^3} + \frac{1}{k_0 \xi} \right] \right\}. \quad (17)$$

The first two terms on the right of Eq.(16) equal the usual Lorentz-Lorenz (LL) and Onsager-Böttcher (OB) formulae in absence of self-polarization [5, 7, 13], $\Gamma_{LL} = \Gamma_0 \left(\frac{\epsilon+2}{3} \right)^2 \sqrt{\epsilon}$, $\Gamma_{OB} = \Gamma_0 \left(\frac{3\epsilon}{2\epsilon+1} \right)^2 \sqrt{\epsilon} \simeq \Gamma_0 (1 + \frac{7}{6} \Re\{\alpha\rho\})$, with $\epsilon = 1 + \Re\{\alpha\rho\}$. A detailed analysis shows that a contribution $\frac{\Gamma_0}{3} \Re\{\alpha\rho\}$ comes from Γ^{\parallel} while a contribution $\frac{5\Gamma_0}{6} \Re\{\alpha\rho\}$ comes from $2\Gamma^{\perp}$. We show this as follows. The vertex provided by Eqs.(13,14), which gives rise to coupling of longitudinal and transverse modes – see Fig.2(b), yields a term common to $2\Gamma^{\perp}$ and Γ_{\parallel} ,

$$\begin{aligned}\Gamma_0 \Re\{\alpha\rho\} k_0 \Im \left\{ \int \frac{d^3k'}{(2\pi)^3} \frac{d^3k}{(2\pi)^3} h(|\vec{k}' - \vec{k}|) \right. \\ \left. \times G_{\perp}^{(0)}(k') G_{\parallel}^{(0)}(k) \sin^2 \theta \right\}.\end{aligned}\quad (18)$$

In the limit $k_0 \xi \ll 1$, the above expression reduces to $\frac{\Gamma_0}{3} \Re\{\alpha\rho\}$. It equals the contribution of each local field factor, $\frac{\epsilon+2}{3}$ in Γ_{LL} and $\frac{3\epsilon}{2\epsilon+1}$ in Γ_{OB} . On the other hand, $2\Gamma^{\perp}$ includes the term

$$\Gamma_0 \Re\{\alpha\rho\} k_0 \Im \left\{ \int \frac{d^3k}{(2\pi)^3} 2[G_{\perp}^{(0)}(k)]^2 \right\}, \quad (19)$$

which amounts to $\frac{\Gamma_0}{2} \Re\{\alpha\rho\}$. Its corresponding counterpart in Γ_{LL} and Γ_{OB} is in the bulk factor $\sqrt{\epsilon}$. In some works only the bulk factor is associated to propagating decay rate [8, 15] or even to the total transverse decay rate [14]. Our microscopic approach reveals clear discrepancies with those interpretations.

Moreover, one of the common terms given in Eq.(18), $\frac{\Gamma_0}{3} \Re\{\alpha\rho\}$, corresponds to Γ_{\parallel} and is associated to $\Im\{\chi_{\parallel}/(\rho\alpha)\}$ in Eq.(6). It is therefore part of the incoherent emission which is scattered away as bare longitudinal modes couple to bare transverse-radiation modes at the cavity surface. The existence of the common term in Eq.(18) is a consequence of reciprocity. However, while it gives rise to propagating emission when the coupling reads $G_{\parallel} - G_{\perp}$, it gives rise to extinction as read in opposite direction as in Fig.2(b). This longitudinal term corresponds to the remaining local field factor in Γ_{LL} and Γ_{OB} at leading order. Some works [6, 11, 17] make the identification $\Gamma_{LL, OB} = 2\Gamma_{LL, OB}^{\perp}$, which is erroneous according to our approach.

In the presence of absorption, the electrostatic polarizability is complex and additional non-propagating (NP) terms in both $2\Gamma^{\perp}$ and Γ^{\parallel} contribute to the total decay rate. These are,

$$\begin{aligned}2\Gamma_{\perp}^{NP} &= \Gamma_0 \Im\{\alpha\rho\} k_0 \int \frac{d^3k'}{(2\pi)^3} \frac{d^3k}{(2\pi)^3} h(|\vec{k}' - \vec{k}|) G_{\perp}^{(0)}(k') \\ &\quad \times (1 + \cos^2 \theta) G_{\perp}^{(0)}(k),\end{aligned}\quad (20)$$

$$\begin{aligned}\Gamma_{\parallel} &= \Gamma_0 \Im\{\alpha\rho\} k_0 \int \frac{d^3k'}{(2\pi)^3} \frac{d^3k}{(2\pi)^3} G_{\parallel}^{(0)}(k) [h(|\vec{k}' - \vec{k}|) \\ &\quad \times G_{\parallel}^{(0)}(k') \cos^2 \theta + \delta^{(3)}(\vec{k} - \vec{k}') G_{\parallel}^{(0)}(k')].\end{aligned}\quad (21)$$

At leading order in $k_0\xi$, the dominant term is that of Eq.(21) due to absorption in the Coulomb-Coulomb coupling, $\Gamma^{NP} \approx \Gamma_{\parallel}^{abs.} \simeq \Gamma_0 \Im\{\alpha_0\rho\} \frac{1}{(k_0\xi)^3}$ [10, 11, 12]. The remaining term $\Gamma_0 \frac{\Im\{\alpha_0\rho\}}{k_0\xi}$ of Eq.(17) contains equal contributions from $2\Gamma_{\perp}^{NP}$ and Γ_{\parallel} in agreement with Eqs.(11,12).

In absence of absorption, the terms in Eqs.(20,21) still contribute as considering radiative corrections in α . These corrections affect both the host medium and the host particle. In the former case, radiative corrections show up as a consequence of the coupling between transverse and longitudinal modes in single scattering events [1]. This is due to the 'internal' correlation function $h_a(r) = \Theta(r-a)$ –see Fig.2(d). Fig.2(d) illustrates how, for an incoming and an outgoing photon being longitudinal, the coupling to a transverse internal mode gives rise to radiation. Hence, the effective vertex in Fig.2(c) with α including radiative corrections. This kind of terms amount to scattered incoherent radiation. Eq.(20) is in fact part of Eq.(8) and so of Eq.(11).

Finally, we consider the back-reaction of the host medium on the host particle, with electrostatic polarizability α_e –eventually equal to α_0 . To this aim and at first order, we have to compute the self-polarization diagrams, i.e. 4th to 6th diagrams in Fig.1(b). For the sake of simplicity, we disregard absorption and radiative corrections in α and keep leading order terms in α_0 –denoted by the script (1). We can approximate

$$\gamma_{\parallel}^{(1)} \simeq \frac{-k_0}{2\pi} \rho \alpha_0 \left[\frac{i}{3} + \frac{1}{(k_0\xi)^3} \right], \quad 2\gamma_{\perp}^{(1)} \simeq \frac{-k_0}{2\pi} \left[i + \rho \alpha_0 \frac{5i}{6} \right], \quad (22)$$

so that, in absence of self-polarization, we obtain just $\Gamma_{LL}^{(1)} = \Gamma_{OB}^{(1)} = \Gamma_0(1 + \frac{7}{6}\Re\{\alpha_0\rho\})$ as given above. Plugging Eq.(22) into Eq.(1), we get

$$\Gamma^{(1)} \simeq \Gamma_{LL}^{(1)} \left[1 + \frac{4}{9} \frac{\alpha_e}{V_{\xi}} \rho \alpha_0 \right], \quad (23)$$

where V_{ξ} is the volume of the cavity. Making the identification $\epsilon = 1 + \alpha_0\rho$, one can verify the agreement of the above expression with the OB formula at leading order in α_0, α_e [12, 13], $\Gamma_{OB} = \Gamma_0 \left(\frac{3\epsilon}{2\epsilon+1 - \frac{2}{3} \frac{\alpha_e}{V_{\xi}} (\epsilon-1)} \right)^2 \sqrt{\epsilon}$.

In summary, we have derived equations which relate the spontaneous decay rate of an emitter within a virtual cavity, both propagating –Eq.(10)– and non-propagating –Eqs.(11,12), to the mean free path l_{ext} and the refraction index \bar{n} of the host medium. The last equation being strictly valid only in the range of validity of MG. Because \bar{n} and l_{ext} depend ultimately on ω and ξ , it is argued that variations on Γ towards inhibition/enhancement can be achieved by applying KK relations and Ward's identities. Our derivation is based on the approach of [1]. That is shown to include both LL and OB at leading order in ρ . Microscopical analysis of the scattering events involved in the formulae of [1] reveals that local field

factors present longitudinal decay rate contributions. A non-propagating decay rate has been identified. It includes both absorption and incoherent scattered radiation. The latter involves coupling of bare transverse to bare longitudinal modes which takes place at different levels. Those are, at the emitter cavity, at single scatterers, at the correlation boundary between host scatterers and at the boundary of the sample. The latter was not considered in this paper.

We thank S.Albaladejo, R.Carminati and J.J.Saenz for fruitful discussions and suggestions. This work has been supported by the Spanish integrated project Consolider-NanoLight CSD2007-00046 and the EU project NanoMagMa EU FP7-NMP-2007-SMALL-1.




FIG. 2: (a) Feynman's rules. (b) Common vertex derived from Eqs.(13,14) which couples longitudinal to transverse modes, with $|\hat{k} \times \hat{k}'| = \sin \theta$. (c) Longitudinal-longitudinal vertex from Eq.(14), with $|\hat{k} \cdot \hat{k}'| = \cos \theta$. (d) Diagrammatic representation of the origin of the coupling of single scattering radiative corrections to longitudinal external modes. Integration of internal radiative corrections yields the vertex in Fig.(e).

* Electronic address: manuel.donaire@uam.es

- [1] M. Donaire, e-print arXiv:0811.0323.
- [2] C. Toninelli *et al.*, Phys. Rev. Lett. **101**, 123901 (2008); P.D.Garcia *et al.*, e-print arXiv:0811.4494.
- [3] D.S. Citrin, Nano Lett. **4**, 1561 (2004).
- [4] M. Peskin and D.V. Schroeder *Introduction to Quantum Field Theory* Westview Press (1995).
- [5] H.A. Lorentz, Wiedem. Ann. **9**, 641 (1880); L. Lorenz, Wiedem. Ann. **11**, 70 (1881).
- [6] S.M. Barnett, B. Huttner and R. Loudon, Phys. Rev. Lett. **68**, 3698 (1992).
- [7] L. Onsager, J. Am. Chem. Soc. **58**, 1486 (1936).
- [8] S.M. Barnett and R. Loudon, Phys. Rev. Lett. **77**, 2444 (1996); R. Carminati and J.J. Saenz, to appear in Phys.Rev. Lett. (2009).
- [9] Y.N. Barabanenkov and V.D. Ozrin Phys. Lett. A **206**, 116 (1995).
- [10] L.S. Froufe-Perez, R. Carminati and J.J. Saenz, Phys. Rev. A **76**, 013835 (2007).
- [11] S.M. Barnett, B. Huttner, R. Loudon and R. Matloob, J. Phys. B **29**, 3763 (1996).
- [12] F.Hynne and R.K.Bullough J. Phys. A **5**, 1272 (1972); F.Hynne and R.K.Bullough, Phil. Trans. R. Soc. Lond. A **321**, 305 (1987).

- [13] P. de Vries and A.Legendijk, Phys. Rev. Lett. **81**, 1381 (1998); R. Glauber and M. Lewenstein, Phys.Rev. A **43**, 467 (1991).
- [14] C.A. Guérin, B. Gralak and A. Tip, Phys. Rev. E **75**, 056601 (2007).
- [15] A.Legendijk and B. van Tiggelen, Phys.Rep. **270**, 143 (1996); P. de Vries, D.V. van Coevorden and A. Legendijk, Rev. Mod. Phys. **70**, 447 (1998).
- [16] M. Donaire, e-print arXiv:0902.1783.
- [17] S. Scheel *et al.*, Phys. Rev. A **60**, 1590 (1999).