Simplifying the algebra of first class constraints, examples on SO(3) and SO(4)

M. Dehghani*

A. Shirzad^{\dagger}

Department of Physics, Isfahan University of Technology P.O.Box 84156-83111, Isfahan, IRAN, School of Physics, Institute for Research in Fundamental Sciences (IPM) P.O.Box 19395-5531, Tehran, IRAN.

(Dated: October 31, 2018)

We discuss the problem of non abelian constrained systems and the origin of appearance of non abelian algebras. We show that it is possible, in principle, to change a non abelian system to an abelian one, at least locally. Our method is based on solutions of the differential equations due to the algebra of first class constraints. We suggest that multiplicative constraints should be regularized at each step of calculation. Some examples on SO(3) and SO(4) algebras are discussed. We show that they can be replaced by abelian constraints.

Keywords: First class system, Constraint algebra, Abelianization.

PACS numbers:

I. INTRODUCTION

The general formalism of constrained systems seems sometimes very complicated. The reason is the complicated and extensive algebra of constraints. Consider a first class system given by constraints ϕ_a and the canonical Hamiltonian H_c . The most general form of the algebra of Poisson brackets is [1]:

$$\{\phi_a, \phi_b\} = C^c_{ab}\phi_c$$

$$\{\phi_a, H_c\} = V^b_a\phi_b$$

(I.1)

The coefficients C_{ab}^c and V_a^b are called structure functions. In general these coefficients may be functions of phase space variables (q, p). If one wishes to keep the track of levels of consistency, additional labels showing the level should be assigned to the constraints and the algebra would be much more complicated. This complication causes so many difficulties in proving general statements in the context of constrained systems and makes this field of study difficult to follow. However, in the real models of gauge systems we never encounter a problem in which the complete set of structure functions C_{ab}^c and V_a^b are present or depend on (q, p). The simplest possibility is $C_{ab}^c = 0$ and $V_a^b =$ cons.. Such systems are called abelian first class constrained systems. As we will see in the next section this is the case for quadratic Lagrangians (with respect to velocities and coordinates) which include most physical models. A non abelian system requires a Lagrangian or Hamiltonian with higher powers, or more complicated functional dependence on the corresponding variables.

On the other hand, it is well known [2] that the choice of constraints in not unique. In other words, different sets of constraints may describe the same constraint surface. We say that the two sets of constraints $\phi_a(q, p)$, $a = 1, \dots m$ and $\phi_{a'}(q, p), a' = 1, \dots m'$ are equivalent if $\phi_a(q, p) = 0 \Leftrightarrow \phi_{a'}(q, p) = 0$. Then it is obvious that in general, one can write:

$$\phi_a(q,p) = \sum_{a'=1}^{m'} M_{aa'} \phi_{a'}(q,p) \qquad a = 1, \cdots m.$$
 (I.2)

A system is called reducible if it can be converted to one with less number of constraints. Transforming from the set $\phi_a(q, p)$ to the set $\phi_{a'}(q, p)$ is called a "redefinition" of constraints.

We wish to use the redefinition process to make the algebra of the constraints as simple as possible. The best situation is one with abelian algebra. Then the system is said to be abelianized. We can also make use of the "canonical transformations" (CT's) whenever necessary. The first tool, i.e. redefinition of the constraints, changes the variables describing the constraint surface only, while the second one, i.e. CT, change the coordinates of the whole phase space canonically. It should be emphasized that the Poisson brackets are invariant under CT, while the redefinition procedure may lead to a different algebra of Poisson brackets.

Converting a system of first class constraints to an abelian system, if it is possible, has so many advantages. In fact in any analysis based on the constraint structure of a gauge system one encounters the complicated algebra of Poisson brackets of first class constraints and one can not show the essential features of the problem in clear formulas. For example a closed formula for the generating function of gauge transformations in terms of the constraints, is not given so far. There have been proposed only some rules and instructions in this regard for the

^{*}mdehghani@ph.iut.ac.ir

[†]shirzad@ipm.ir

There is a similar situation for the BRST charge which is needed to quantize a gauge system. In fact, this charge can be written as an infinite expansion in terms of recursive Poisson brackets of structure functions which in general make an open algebra which does not terminate. For an abelian system, on the other hand, besides clarity and simplicity in understanding the structure of physical as well as gauged degrees of freedom, one is able to write down the generator of gauge transformations in a closed form [4] and the expansion of the BRST charge terminates after the first term [5].

As we will see through the paper, non abelian constraints may stem from the choice of coordinates for description of constraint surface. In this view, the abelianization procedure is a redefinition and purgation process for the constraints which make it possible to describe the constraint surface with a set of suitable (i.e. commuting) coordinates.

Our aim in this paper is to show that some first class constrained systems can in principle be abelianized, at least locally. In fact, one can say that the origin of non abelianity stem from bad choice of the canonical coordinates of the whole phase space as well as the variables describing the constraint surface. Therefore, the abelianization means that one tries to find the suitable coordinates in which the algebra of constraints is abelian.

Local abelianization has been shown, in principle, to be possible [1, 6] through solving constraint equations, $\phi_a = 0$ to find a number of coordinates ξ_a in terms of other variables $\tilde{\xi}$ as:

$$\xi_{a'} = f_{a'}(\xi) \tag{I.3}$$

Then the new constraints $\psi_{a'} = \xi_{a'} - f_{a'}(\xi)$ have Poisson brackets which are independent of $\xi_{a'}$. Hence, the only way that the Poisson brackets may vanish on the constraint surface $\psi_{a'} = 0$ is that they vanish strongly. In [7] it is argued that if one maps each constraint to the surface of other constraints they would be abelianized. However, it is asserted in [8] that for gauge systems such as SO(3) the maximality conditions is violated, hence the sufficient condition for SO(3) to be abelianizable is not satisfied. Another method is proposed for abelianization [6] which is based on finding a complicated solution for the matrix M in (I.2) such that the new set of constraints are abelian.

In this paper we try to study the problem of abelianization in a systematic way from the point of view of differential equations coming from the algebra of constraints. Our method is based on finding suitable coordinates to describe the constraint surface. We do this by solving differential equations due to Poisson brackets of constraints with one momentum constraint. We will find that in this way one would naturally lead to simple coordinates of the constraint surface. This method will be discussed in details in section III. Before that we will discuss in the next section some general aspects of abelian and non abelian nature of first class systems. We will also discuss the problem of regularity of the constrains. We suggest that the multiplicative constraints should be regularized before abelianization. Section IV denotes detailed calculations concerning abelianization of a first class system with SO(3) algebra. This important example shows the general features of the abelianization procedure. In section V we give our conclusions.

II. HOW NON ABELIAN CONSTRAINTS MAY HAPPEN?

Let us first consider a simple example to see in what sense a non abelian system of first class constraints may emerge. Suppose, in a system with q_1 and q_2 as coordinates, we are given two first class constraints

$$\phi_1 = p_1 e^{\alpha q_2} \quad \phi_2 = p_2 e^{-\beta q_1} \tag{II.4}$$

where α and β are constants. Clearly we have

$$\{\phi_1, \phi_2\} = (\alpha e^{-\beta q_1})\phi_1 + (\beta e^{\alpha q_2})\phi_2 \qquad (\text{II.5})$$

which exhibits the non abelian feature of the system. It is, however, obvious that the constraints surface $\phi_1 = 0$ and $\phi_2 = 0$ is equivalent to the surface described by $p_1 =$ 0 and p_2 , since the exponentials does not vanish in the finite range of their arguments. Clearly, the constraints p_1 and p_2 are abelian. In general the situation is not so obvious, and the algebraic structure of the Poisson brackets may be so complicated that one is not able to recognize the best and simplest phase space coordinates describing the constraints surface.

It is also possible to inspect some features of non abelian systems by power counting. Suppose $\phi_i(z^m)$ is any constraint which can be written as a polynomial of order m with respect to phase space coordinates $z_{\mu}, \mu = 1, \dots 2N$. For a first class system the algebra of Poisson brackets reads

$$\{\phi_i(z^m), \phi_j(z^n)\} = a_{ij}^k(z^r)\phi_k(z^s).$$
 (II.6)

Since Poisson bracket requires two times of differentiation, we should have

$$m + n - 2 = r + s.$$
 (II.7)

If the constraints are linear with respect to z_{μ} , i.e. m = n = 1, then a first class system can be achieved just for $a_{ij}^k = 0$. In other words linear first class constraints are essentially abelian. For quadratic Lagrangians, which is the case for most physical systems, the primary constraints which emerge due to singularity of Hessian (the matrix of second derivatives with respect to velocities), are necessarily linear. Since the Hamiltonian is also quadratic, the secondary constraints at any level would be linear, too. So, for the wide class of gauge systems with quadratic Lagrangians the system is abelian by itself. The above analysis shows that the problem of abelianization maybe converted to the problem of linearization. In other words if we are able to abelianize a first class system this means that there can be found suitable coordinates in which the constraint surface is described by constant values of some phase space coordinates. Specially we can choose a basis in which the constraints are some momenta.

For example a constrained system given by $\phi_1 = xp_x + yp_y$, $\phi_2 = \frac{1}{2}p_x^2$ and $\phi_3 = \frac{1}{2}p_y^2$ obeys the non abelian algebra $\{\phi_1, \phi_2\} = \phi_2$ and $\{\phi_1, \phi_3\} = \phi_3$. This system reduces to the constraints p_x and p_y , which are linear as well as abelian.

Coming back to the Eq's. (II.6) and (II.7), assume a system in which the constraints are quadratic homogenous functions of z_{μ} which leads to r = 0. In other words, for quadratic constraints we may have a non abelian closed algebra of Poisson brackets only with constant structure functions a_{ij}^k . For example in a system with x, y and z as coordinates, the quadratic constraints $\phi_1 = yp_z - zp_y, \ \phi_2 = zp_x - xp_z \ \text{and} \ \phi_3 = xp_y - yp_x$ exhibit the SO(3) algebra. Such systems require cubic terms in Lagrangian and/or Hamiltonian, assuming the primary constraints are linear. Yang-Mils models fall in this category. More complicated examples in which constraints of different powers constitute a closed algebra may be imagined. But such strange systems are not met in concrete physical models, and it seems that following sophisticated discussions in this direction does not give us more insight about gauge theories.

One important point should be added here. It is well known that [1], multiplicative expressions of constraints (first or second class) are first class, i.e. their Poisson brackets with all constraints are at least linear with respect to the constraints and vanish weakly. It is possible to construct, for instance, a set of quadratic expressions out of a smaller set of constraints such that they make a closed non abelian Lie algebra of Poisson brackets. Assume, for example, four second class constraints x, p_x and y, p_y . One can write ten quadratic monomials such as xp_x,xy, p_xp_y , etc. Clearly these constraints are first class and show up a non abelian closed Lie algebra. We consider a subset of them as follows:

They obey the following closed algebra:

$$\begin{cases} , \} & \phi_1 & \phi_2 & \phi_3 & \phi_4 & \phi_5 & \phi_6 \\ \phi_1 & 0 & 2\phi_1 & 0 & 4\phi_2 & 2\phi_3 & 0 \\ \phi_2 & -2\phi_1 & 0 & -\phi_3 & 2\phi_4 & \phi_5 & 0 \\ \phi_3 & 0 & \phi_3 & 0 & 2\phi_5 & \phi_6 & 0 \\ \phi_4 & -4\phi_2 & -2\phi_4 & -2\phi_5 & 0 & 0 & 0 \\ \phi_5 & -2\phi_3 & -\phi_5 & -\phi_6 & 0 & 0 & 0 \\ \phi_6 & 0 & 0 & 0 & 0 & 0 \\ \end{cases}$$
(II.9)

It is clear that the constraints surface described by $\phi_1 \cdots \phi_6$ in (II.8) is the same as given by $\psi_1 \equiv x \approx 0$,

 $\psi_2 \equiv p_x \approx 0$ and $\psi_3 \equiv p_y \approx 0$ which is a mixed system composed of a pair of second class constraints (x,p_x) together with the first class constraint ψ_3 . Such systems are recognized in the literature [1, 9] as irregular constraints, which are defined as systems of constraints whose gradients vanish on the constraint surface. More precisely their Jacobian $\frac{\partial \phi_a}{\partial \xi_j}$ (where ξ_j are the phase space coordinates) is not full rank on the constraint surface.

If the non abelian algebra of the first class constraints has originated from the irregular nature of the constraints, then the most direct way toward an abelian equivalent system is regularizing the system, i.e. replacing the multiplicative constraints by the equivalent linear ones. For this reason we should classify multiplicative constraints into three different categories:

- 1. Nonlinear constraints of the form $[f(q, p)]^k \approx 0$. Such a constraint should be linearized, i.e. replaced by linear expression $f(q, p) \approx 0$. Our evidence for this replacement is that a constraint of the form $[f(q, p)]^k \approx 0$ has no meaning other than $f(q, p) \approx$ 0. This suggestion does not mean any change in the Hamiltonian or Lagrangian (as in ref. [9]).
- 2. Bifurcating systems of the form $f(q, p)g(q, p) \approx 0$. In this case the constraint surface obviously "bifurcates" into two different branches $f(q, p) \approx 0$ and $g(q, p) \approx 0$. These systems should be considered as the union of different branching $f(q, p) \approx 0$ and $g(q, p) \approx 0$. Each branch should be treated individually. For example, the system given by $\phi_1 = x^2, \phi_2 = xp_x$ and $\phi_3 = xy$ may be regularized to $\psi_1 = x$ (first class), or $\psi_1 = x, \psi_2 = p_x$ (second class) or $\psi'_1 = y, \psi_2 = p_x$ (first class). In each branch the dynamics of the system is different from the others.
- 3. Reducible systems of the form $f(q, p)G(q, p) \approx 0$ where G(q, p) has no finite root. One can replace such a constraint with the simpler one $f(q, p) \approx 0$. This reduction is in fact, the essence of the abelianization procedure explained in the next section.

As the consequence of three kinds of simplifications given above the number of constraints, as well as their algebra, may change. In most cases the regularization and reduction procedures simplifies the algebra of constraints as the above examples show. However, the more important point is clarifying the first or second class nature of the system. For example the system given in (II.8) is not in reality a first class system. Besides simplifying the algebra, the important point is that the system given by (II.8) is partly second class (in the x, p_x plane). Conclusively, given a set of irregular first class constraints satisfying a non abelian algebra, there is no guarantee that the system remains first class after regularization.

One problem that makes the above analysis more difficult is the possibility of combining the multiplicative constraints in the form of more complicated functions. Another problem is that the simple multiplicative forms may be hidden by changing the variables describing the phase space or the constraint surface. Such difficulties may be resolved during the process of abelianization in which one tries to find the most suitable coordinates for describing the constraint surface.

As a prescription toward an abelian algebra we propose to regularize a system of multiplicative constraints in advance. Then the remaining algebra of first class constraints, if it is still non abelian, can be abelianized following the method of the next section. Therefore, in the following we assume that the system is already regularized. We restrict ourselves to a pure and truly first class system and try to abelianize the algebra of Poisson brackets. By truly first class system we mean a system that remains first class in every description of the constraint surface.

III. ABELIANIZATION

Suppose we are given a set of first class constraints $\phi_i(q, p), i = 1, \cdots m$, satisfying the algebra

$$\{\phi_i, \phi_j\} = \alpha_{ijk}\phi_k \tag{III.10}$$

where in general α_{ijk} depends on (q, p). We wish to find an equivalent set of constraints $\phi_r(q, p)$ where

$$\begin{aligned} \phi_i(q,p) &= 0 \Leftrightarrow \phi_r(q,p) = 0 \\ \{ \widetilde{\phi_r}(q,p), \widetilde{\phi_s}(q,p) \} &= 0, \end{aligned}$$
(III.11)

assuming that the system remains first class under the reduction $\phi_i \to \widetilde{\phi_r}$. If we are succeeded in this regard, then it is in principle possible to find a suitable canonical transformation which transforms all the $\widetilde{\phi_r}$ to a set of momenta, i.e.

$$\widetilde{\phi_r} \to P_r.$$
 (III.12)

In the first step we can choose the first constraint momentum P_1 the same as $\phi_1(q, p)$. It is justified that in principle there exists a CT which does this task, although it may require heavy algebraic manipulations. Therefore, suppose under the desired CT we have

$$P_1 \equiv \phi_1(q, p). \tag{III.13}$$

Suppose Q_1 is the coordinate conjugate to P_1 and the other coordinates change to $\tilde{q}_2, \tilde{p}_2, \cdots$ under the above CT. In practice Q_1 should be determined by solving the differential equation due to $\{Q, P\} = 1$ and \tilde{q}, \tilde{p} are derived after determining the corresponding generating function of CT, as we will show in the example of the next section. In the new coordinates all the remaining constraints may change, so as to say

We first show that P_1 should be absent in the remaining constraints $\phi_i(Q_1, P_1, \tilde{q}, \tilde{p})$. The reason is that in general we can expand ϕ_i such that:

$$\phi_i(Q_1, P_1, \widetilde{q}, \widetilde{p}) = \phi_i(Q_1, 0, \widetilde{q}, \widetilde{p}) + P_1 f_i(Q_1, P_1, \widetilde{q}, \widetilde{p})$$
(III.15)

Excluding irregular cases where the gradient of constraints go to infinity on the constraint surfaces; the Taylor expansion always makes sense. On the constraint surface we have $P_1 = 0$. So we have the equivalency:

Now we are at the point to consider the powerful requirement that the constraints are first class. For example the Poisson brackets $\{\phi_i(Q_1, \tilde{q}, \tilde{p}), P_1\}$ vanishes on the constraint surface described yet in Eq. (III.16):

$$\frac{\partial \phi_i(Q_1, \tilde{q}, \tilde{p})}{\partial Q_1} = \sum_{j=2}^m C_{ij}(Q_1, \tilde{q}, \tilde{p})\phi_j(Q_1, \tilde{q}, \tilde{p}) \quad i = 2, \cdots m.$$
(III.17)

Since P_1 is absent from LHS of the above equation, no term proportional to P_1 has been written in the RHS. This system of coupled ordinary linear first order differential equations has an analytic solution when C_{ij} 's are independent of Q_1 which reads in the matrix notation as

$$\phi(Q_1, \widetilde{q}, \widetilde{p}) = e^{CQ_1} \phi(0, \widetilde{q}, \widetilde{p})$$
(III.18)

where C is a $(m-1) \times (m-1)$ matrix with elements $C_{ij}(\tilde{q}, \tilde{p})$. This is the case for models in which the structure functions are constant. Since the exponential part in (III.18) can not vanish for any finite value of Q_1 , the necessary and sufficient condition for vanishing of $\phi_i(Q_1, \tilde{q}, \tilde{p})$ is vanishing of $\phi_i(0, \tilde{q}, \tilde{p})$ which we can rename them as $\eta_i(\tilde{q}, \tilde{p})$. So, the reduction shown in Eq. (III.16) goes on one step further to

The above case with constant C_{ij} gives the general feature of the problem. The result is that given P_1 as a constraint, the remaining constraints after redefinition should be independent of Q_1 . In other words the Q_1 dependent part of them has no root and can be omitted.

This feature can also be shown for the general case through the following argument. We come back to the system of linear ODE's given by Eq. (III.17). From the theory of differential equations [10] it is well-known that the general solutions should contain m-1 constants, which can be chosen as the initial values $\phi_i(0, \tilde{q}, \tilde{p})$. In fact, it can be shown that the general solution $\phi_i(Q_1, \tilde{q}, \tilde{p})$ is linear with respect to the initial values $\phi_i(0, \tilde{q}, \tilde{p})$. Just to remind the reader, the proof can be achieved by dividing the interval $(0, Q_1)$ into an infinite number of segments δQ_1 ; then the linear differential equations (III.17) show that after *n* steps we have

$$\phi_i(n\delta Q_1, \widetilde{q}, \widetilde{p}) = (1 + C((n-1)\delta Q_1, \widetilde{q}, \widetilde{p}))_{ij}(1 + C((n-2)\delta Q_1, \widetilde{q}, \widetilde{p}))_{jk} \cdots_n (1 + C(0, \widetilde{q}, \widetilde{p}))_{nl}\phi_l(0, \widetilde{q}, \widetilde{p}).$$
(III.20)

Although the product of parentheses above can not be exhibited by a simple exponential (as in Eq.(III.18)), the linearity of the final values of $\phi_i(Q_1, \tilde{q}, \tilde{p})$ with respect to initial values $\phi_i(0, \tilde{q}, \tilde{p})$ is established. In this way one can propose the general form of the solution of Eqs. (III.17) as

$$\phi_i(Q_1, \tilde{q}, \tilde{p}) = \sum_j \phi_i^{(j)}(Q_1, \tilde{q}, \tilde{p})\eta_j(\tilde{q}, \tilde{p})$$
(III.21)

where

$$\eta_j(\widetilde{q}, \widetilde{p}) \equiv \phi_j(0, \widetilde{q}, \widetilde{p}). \tag{III.22}$$

The functions $\phi_i^{(j)}(Q_1, \tilde{q}, \tilde{p})$ are special solutions of Eqs. (III.17) with the initial conditions

$$\phi_i^{(j)}(0,\widetilde{q},\widetilde{p}) = \delta_i^j \quad i, j = 2, \cdots m - 1.$$
(III.23)

The general solution (III.21) can be viewed as an expansion in terms of special solution $\phi_i^{(j)}(Q_1, \tilde{q}, \tilde{p})$ of the ODE's (III.17). From this point of view the functions $\eta_j(\tilde{q}, \tilde{p})$ can be interpreted as the constant (with respect to Q_1) coefficients of expansion. It should be noticed that we did not solve practically the differential equations (III.17) which are resulted from the algebra of the constraints. The point is that the constraints ϕ_2, \dots, ϕ_m in terms of coordinates $(Q_1, P_1, \tilde{q}, \tilde{p})$ automatically should appear in the form of solutions of Eqs. (III.17). This point will be seen clearly in the example given in the next section.

Now we claim that the constraint surface given by $\phi_i(Q_1, \tilde{q}, \tilde{p})$ is the same as one given by $\eta_i(\tilde{q}, \tilde{p})$. It is obvious from (III.21) that $\eta_i(\tilde{q},\tilde{p}) = 0$ give rise to $\phi_i(Q_1, \tilde{q}, \tilde{p}) = 0$. What about the inverse deduction? Our assertion here is that if in some direction the constraint surface can not be described unless some definite function of Q_1 say $f(Q_1, \tilde{q}, \tilde{p})$ vanishes, then the equation $f(Q_1, \tilde{q}, \tilde{p}) = 0$ can in principle be solved to give Q_1 as $Q_1 = g(\tilde{q}, \tilde{p})$. Then P_1 and $Q_1 - g(\tilde{q}, \tilde{p})$ constitute a second class constrained system, which we have excluded it from our consideration. Hence, the necessary and sufficient condition for $\phi_i(Q_1, \tilde{q}, \tilde{p}) = 0$ is $\eta_i(\tilde{q}, \tilde{p}) = 0$. So we come to a noticeable result that the redefinition procedure has brought us to the set of equivalent constraints $P_1, \eta_2(\widetilde{q}, \widetilde{p}), \cdots, \eta_m(\widetilde{q}, \widetilde{p})$, with the property that P_1 commutes with all other constraints. By this procedure we have decoupled the constraint $\phi_1 = P_1$ from others. Decoupling is done by purging other constraints from canonical conjugate pair (Q_1, P_1) .

Now we can restrict our attention to constraints η_2, \dots, η_m which are defined in a smaller phase space (\tilde{q}, \tilde{p}) where the canonical pair (Q_1, P_1) are no longer present. Any canonical transformation in the (\tilde{q}, \tilde{p}) subspace does not affect the subspace (Q_1, P_1) . Therefore, one can in principle repeat the same procedure once more and in this time assumes that $\eta_2(\tilde{q}, \tilde{p})$ is the momentum P_2 in some suitable coordinates. In this way after several stages all the constraints would be reduced to a set of momenta.

Note should be added that the number of constraints may be changed, in fact reduced, at any stage of the above process of abelianization. The reason is that, for example in the first stage, linear independence of the constraints $\phi_i(Q_1, \tilde{q}, \tilde{p})$, does not necessarily require that $\phi_i(0, \tilde{q}, \tilde{p})$ are linearly independent. Hence, from Eq. (III.22) the number of independent $\eta_i(\tilde{q}, \tilde{p})$ may be less than $\phi_i(Q_1, \tilde{q}, \tilde{p})$.

As a concrete example consider two first class non abelian constraints ϕ_1 and ϕ_2 with the algebra

$$\{\phi_1, \phi_2\} = \alpha \phi_1 + \beta \phi_2. \tag{III.24}$$

By a canonical transformation we map the constraint ϕ_1 to momentum P_1 . After projection ϕ_2 on the surface $P_1 = 0$, the algebra (III.24) turns to:

$$\{P_1, \phi_2\} = \beta \phi_2(Q_1, \widetilde{q}, \widetilde{p}) \tag{III.25}$$

The constraint ϕ_2 can be found from the differential equation

$$\frac{\partial \phi_2}{\partial Q_1} = -\beta \phi_2 \tag{III.26}$$

as

$$\phi_2(Q_1, \tilde{q}, \tilde{p}) = \eta(\tilde{q}, \tilde{p}) \exp\left(-\int_0^{Q_1} \beta dQ_1'\right) \qquad \text{(III.27)}$$

where $\eta(\tilde{q}, \tilde{p}) = \phi_2(0, \tilde{q}, \tilde{p})$. This is a realization of the solutions given in Eq. (III.21) for the general case. Since the exponential function does not vanish for finite values of its argument, vanishing of ϕ_2 could be only due to $\eta(\tilde{q}, \tilde{p})$. The constraints ϕ_1 and ϕ_2 are equivalent to P_1 and $\eta(\tilde{q}, \tilde{p})$, where

$$\{P_1, \eta\} = 0$$
 (III.28)

Then, we can make a CT to canonical variables in which $\eta(\tilde{q}, \tilde{p})$ is P_2 . One may wonder if $\beta(Q_1, q, p)$ is such that $\int \beta dQ_1 = \ln f(Q_1, q, p)$ and the equation $f(Q_1, q, p) = 0$ has some finite roots, then one may no longer exclude vanishing of the exponential part in (III.27). If this is the case, the equivalent constraints are P_1 and $f(Q_1, q, p)$ where $f(Q_1, q, p)$ can be solved for Q_1 . This leads to a second class system which has been excluded before.

IV. EXAMPLES ON SO(3) AND SO(4) ALGEBRAS

In this section we apply our method to some examples. A famous non abelian algebra is the angular momentum algebra in three dimensional configuration space. Let us at first stage show how a SO(3) algebra of constraints may emerge. We may assume that for a rotational invariant Hamiltonian, L_x , L_y and L_z are given as primary constraints. Consistency condition of primary constraints then gives no further secondary constraints and the set of constraints terminate here.

It is also possible to consider more realistic examples in which the angular momentum algebra emerge in a natural way. For example the Lagrangian

$$L = \frac{1}{2}\dot{\mathbf{X}}^2 - V(\mathbf{X}^2) - \xi.\mathbf{L} \qquad (\text{IV.29})$$

where $\mathbf{X} \equiv (x, y, z)$ and $\xi = (\xi_x, \xi_y, \xi_z)$ constitute a six dimensional configuration space, in which $L_i = \epsilon_{ijk} x_j \dot{x_k}$. In phase space π_x , π_y and π_z , the momenta conjugate to ξ_x , ξ_y and ξ_z , are primary constraints and the total Hamiltonian reads

$$H_T = \frac{1}{2}\mathbf{P}^2 + V(\mathbf{X}^2) + \xi.\mathbf{L} + \lambda.\pi \qquad (\text{IV.30})$$

where $\mathbf{P} \equiv (P_x, P_y, P_z)$ represents the momenta conjugate to $\mathbf{X}, \lambda \equiv (\lambda_x, \lambda_y, \lambda_z)$ shows Lagrange multipliers and $L_i = \epsilon_{ijk} x_j p_k$. The consistency conditions of primary constraints π_i give secondary constraints L_i and no further constraint emerges from the consistency of L_i . First level constraints are abelian while the second level constraints L_i obey the non abelian SO(3) algebra with constant structure functions ϵ_{ijk} , i.e.

$$\{L_i, L_j\} = \epsilon_{ijk} L_k \qquad i, j = 1, 2, 3.$$
 (IV.31)

It is also possible to get the SO(3) algebra from the Lagrangian

$$L = \frac{1}{2}\dot{\mathbf{X}}^2 - V(\mathbf{X}^2) - e^w L_x - L_y$$
 (IV.32)

where w is a variable. Consistency of p_w gives L_x , L_y and L_z respectively as the second, third and forth level constraints.

The above expressions are related to a one particle system with rotational invariance. Similar treatments may be considered for multi-particle systems. Now let us examine our method of abelianization to different examples of this character.

A. One particle SO(3) model

Now let us go through the abelianization procedure of the SO(3) algebra of constraints. As stated in the previous section we should first find a CT that transforms for example $L_1 = xp_y - yp_x$ to a momentum P_1 . The conjugate coordinate Q_1 should be determined such that

$$\{Q_1, P_1\} = 1. \tag{IV.33}$$

A possible solution for Q_1 is

$$Q_1 = \tan^{-1}(\frac{y}{x}). \tag{IV.34}$$

As is apparent, P_1 and Q_1 are functions of subspace $(x, y; p_x, p_y)$. Hence, we can exclude the subspace (z, p_z) . Reminding the standard method [11] for extracting a CT from a generating function, the following generator can be used

$$F(x, y, P_1, P_2) = P_1 \tan^{-1}(\frac{y}{x}) + P_2 f(x, y).$$
 (IV.35)

Transformation relations then gives Q_1 as in (IV.34) and $Q_2 = f(x, y)$. Imposing the task $xp_y - yp_x = P_1$ on the relations $p_x = \frac{\partial F}{\partial x}$ and $p_y = \frac{\partial F}{\partial y}$ also gives $x\frac{\partial f}{\partial x} - y\frac{\partial f}{\partial y} = 0$. In this way the canonical pair (Q_2, P_2) can be given as

$$Q_{2} = \frac{1}{2} \ln(x^{2} + y^{2})$$

$$P_{2} = xp_{x} + yp_{y}.$$
(IV.36)

Renaming the variables $(Q_1, P_1; Q_2, P_2)$ as $(\varphi, p_{\varphi}; \psi, p_{\psi})$ the old variables can be written in terms of the new ones as

$$z = Z \qquad p_z = P_Z x = e^{\psi} \cos \varphi \quad p_x = e^{-\psi} (p_{\psi} \cos \varphi - p_{\varphi} \sin \varphi)$$
(IV.37)
$$y = e^{\psi} \sin \varphi \quad p_y = e^{-\psi} (p_{\varphi} \cos \varphi + p_{\psi} \sin \varphi).$$

In terms of the new variables the constraints (L_1, L_2, L_3) are:

$$L_{1} = p_{\varphi}$$

$$L_{2} = \eta_{2}(z, p_{z}, \psi, p_{\psi}) \cos \varphi - \eta_{3}(z, p_{z}, \psi, p_{\psi}) \sin \varphi$$

$$L_{3} = \eta_{2}(z, p_{z}, \psi, p_{\psi}) \sin \varphi + \eta_{3}(z, p_{z}, \psi, p_{\psi}) \cos \varphi$$
(IV.38)

where

$$\eta_2(z, p_z; \psi, p_\psi, p_\varphi) \equiv -zp_\varphi e^{-\psi},$$

$$\eta_3(z, p_z; \psi, p_\psi) \equiv e^{-\psi} zp_\psi - e^\psi p_z.$$
(IV.39)

From the angular momentum algebra we have

$$\frac{\partial L_2}{\partial \varphi} = -L_3, \qquad \frac{\partial L_3}{\partial \varphi} = L_2.$$
 (IV.40)

These are the same differential equations as (III.17). Eqs. (IV.38) are in fact the solutions of Eqs. (IV.40) with respect to the variable $Q_1 = \varphi$. As is seen, Eqs. (IV.38) are in the form given in Eqs. (III.21). The solution $L_2 \propto -\sin\varphi$ and $L_3 \propto \cos\varphi$ is the one with initial condition $L_2(\varphi = 0) = 0$ and $L_3(\varphi = 0) = 1$, as stated in Eq. (III.23), and the solution $L_2 \propto \cos\varphi$ and $L_3 \propto \sin\varphi$ satisfy $L_2(\varphi = 0) = 1$ and $L_3(\varphi = 0) = 0$. Since $\eta_2 \approx 0$ on the surface $p_{\varphi} = 0$ we see that the set of constraints (L_1, L_3) finally reduces to p_{φ} and η_3 which commute with each other.

Important notice should be added that our transformation here is not acceptable globally. In fact, at x = y = 0the transformation is singular. Therefore as indicated in some references [12] the abelianization process of SO(3)algebra can be done just locally. We remind the reader that far from the origin the constraint surface given by functions L_1 , L_2 and L_3 is the same as given by two of them. In fact, since $\mathbf{x}.\mathbf{L} = \mathbf{p}.\mathbf{L} = 0$ the constraints L_1 , L_2 and L_3 are reducible, provided that $\mathbf{x} \neq 0$ and/or $\mathbf{p} \neq 0$. It seems that this subtle point is the essence that the reference [8] has not given a clear statement that the SO(3) gauge system is abelianizable or not. However, an expanded version of SO(3) gauge system is shown to be abelianizable in [13].

B. Two particle SO(3) model

We can extend the above model to a system of two particles with coordinates (x, y, z) and (a, b, c). The angular momentum components of this system read:

$$L_1 = xp_y - yp_x + ap_b - bp_a$$

$$L_2 = yp_z - zp_y + bp_c - cp_b$$

$$L_3 = zp_x - xp_z + cp_a - ap_c$$

(IV.41)

with the same algebra as before. At the first stage we consider L_1 as (twice) the momentum p_{φ} in a new coordinate system. Using the results of one particle system it is easily seen that $\varphi = \varphi_1 + \varphi_2$ is the coordinate conjugate to p_{φ} where

$$\varphi_1 = \arctan(\frac{y}{x}) \quad \varphi_2 = \arctan(\frac{b}{a})$$
 (IV.42)

In this way we should consider

$$\begin{array}{l}
\varphi = \varphi_1 + \varphi_2 & \chi = \varphi_1 - \varphi_2 \\
p_{\varphi} = \frac{1}{2}(p_{\varphi_1} + p_{\varphi_2}) & p_{\chi} = \frac{1}{2}(p_{\varphi_1} - p_{\varphi_2})
\end{array} \tag{IV.43}$$

as new coordinates, where

$$p_{\varphi_1} = xp_y - yp_x \qquad p_{\varphi_2} = ap_b - bp_a \qquad (IV.44)$$

The canonical transformation which gives the above new coordinates act in the four dimensional space of (x, y; a.b) and the corresponding momenta. This transformation should be accompanied by the following new variables

$$\psi_1 = \frac{1}{2} \ln(x^2 + y^2) \quad \psi_2 = \frac{1}{2} \ln(a^2 + b^2) p_{\psi_1} = xp_x + yp_y \qquad p_{\psi_2} = ap_a + bp_b$$
 (IV.45)

Inverting equations (IV.42), (IV.43), (IV.44) and (IV.45) gives old variables in terms of new ones. Inserting them into (IV.41) gives L_2 and L_3 projected on the subspace $p_{\varphi} = 0$ as:

$$L_{2} = -p_{z}r_{1}\sin\frac{\varphi+\chi}{2} + \frac{z}{r_{1}}(p_{\chi_{1}}\cos\frac{\varphi+\chi}{2} + p_{r_{1}}\sin\frac{\varphi+\chi}{2}) -\frac{1}{r_{2}}(cp_{\chi}\cos\frac{\varphi-\chi}{2} - (r_{2}^{2}p_{c} - cp_{r_{2}})\sin\frac{\varphi-\chi}{2}) L_{3} = p_{c}r_{2}\cos\frac{\varphi-\chi}{2} - \frac{c}{r_{2}}(p_{\chi}\sin\frac{\varphi-\chi}{2} + p_{r_{2}}\cos\frac{\varphi-\chi}{2}) -\frac{1}{r_{1}}(zp_{\chi}\sin\frac{\varphi+\chi}{2} + (r_{1}^{2}p_{z} - zp_{r_{1}})\cos\frac{\varphi+\chi}{2}).$$
(IV.46)

These can be rewritten as

$$L_{2} = -M_{2}\sin\frac{\varphi}{2} + M_{3}\cos\frac{\varphi}{2} L_{3} = M_{3}\sin\frac{\varphi}{2} + M_{2}\cos\frac{\varphi}{2}$$
(IV.47)

where

$$M_{2} = (r_{2}p_{c} + r_{1}p_{z} - \frac{zp_{r_{1}}}{r_{1}} - \frac{cp_{r_{2}}}{r_{2}})\cos\frac{\chi}{2} + (\frac{z}{r_{1}} + \frac{c}{r_{2}})p_{\chi}\sin\frac{\chi}{2} M_{3} = (r_{2}p_{c} - r_{1}p_{z} + \frac{zp_{r_{1}}}{r_{1}} - \frac{cp_{r_{2}}}{r_{2}})\sin\frac{\chi}{2} + (\frac{z}{r_{1}} - \frac{c}{r_{2}})p_{\chi}\cos\frac{\chi}{2}$$
(IV.48)

As in (IV.40) we see that $L_2^{(2)} = \cos \frac{\varphi}{2}, L_3^{(2)} = \sin \frac{\varphi}{2}$ and $L_2^{(3)} = -\sin \frac{\varphi}{2}, L_3^{(3)} = \cos \frac{\varphi}{2}$ are special solutions of angular momentum algebra with initial condition (III.23) provided that $L_1 = 2p_{\varphi}$. As discussed in the previous section the constraints L_2 and L_3 reduce to M_2 and M_3 in order that the system remain first class. Hence, the constraint p_{φ} commutes with M_2 and M_3 and decouples from the algebra of first class constraints.

Now it seems that we should repeat our procedure to abelianize the remaining two constraints. However, a little care shows that we can combine M_2 and M_3 to get

$$S_{2} = (r_{2}p_{c} - r_{1}p_{z} + \frac{zp_{r_{1}}}{r_{1}} - \frac{cp_{r_{2}}}{r_{2}})M_{2} - (\frac{z}{r_{1}} + \frac{c}{r_{2}})p_{\chi}M_{3}$$

$$S_{3} = (r_{2}p_{c} + r_{1}p_{z} - \frac{zp_{r_{1}}}{r_{1}} - \frac{cp_{r_{2}}}{r_{2}})M_{3} - (\frac{z}{r_{1}} - \frac{c}{r_{2}})p_{\chi}M_{2}$$

(IV.49)

so that

$$S_2 = F \cos \frac{\chi}{2} \qquad S_3 = F \sin \frac{\chi}{2} \qquad (\text{IV.50})$$

where

$$F = (r_2^2 p_c^2 - r_1^2 p_z^2) + (\frac{c^2 (p_\chi^2 + p_{r_2}^2)}{r_2^2} - \frac{z^2 (p_\chi^2 + p_{r_1}^2)}{r_1^2}) -2(cp_c p_{r_2} - zp_z p_{r_1})$$
(IV.51)

If one demands either $\sin \frac{\chi}{2} = 0$ or $\cos \frac{\chi}{2} = 0$ to satisfy one of the constraints, then the condition F = 0 should also be imposed to satisfy the other constraint, which leads to a seconds class system. Therefore the only way is to consider just F = 0 as the only remaining constraint. In this way the system is reduced to two constraints p_{φ} and F, which are apparently abelian.

C. One particle SO(4) model

As a third example consider a constrained system with SO(4) gauge symmetry with

$$L_{ij} = x_i p_j - x_j p_i, \quad i, j = 1, 2, 3, 4$$
 (IV.52)

The algebra of Poisson brackets of constraints read:

$$\{L_{ij}, L_{kl}\} = \delta_{ik}L_{jl} - \delta_{il}L_{jk} + \delta_{jl}L_{ik} - \delta_{jk}L_{il} \quad (IV.53)$$

Clearly constraints with no common indexes commute. For example $\{L_{12}, L_{34}\} = 0$. Assume simultaneous change of variables in (1-2) as well as (3-4) surfaces similar to SO(3) case, as follows:

$$\phi = \arctan(\frac{x_2}{x_1}) \qquad p_{\phi} = x_1 p_2 - x_2 p_1
\sigma = \frac{1}{2} \ln(x_1^2 + x_2^2) \qquad p_{\sigma} = x_1 p_1 + x_2 p_2
\theta = \arctan(\frac{x_4}{x_3}) \qquad p_{\theta} = x_3 p_4 - x_4 p_3
\rho = \frac{1}{2} \ln(x_3^2 + x_4^2) \qquad p_{\rho} = x_3 p_3 + x_4 p_4$$
(IV.54)

Inverting the above relations and rewriting the constraints in terms of the new variables, we get:

$$L_{12} = p_{\varphi} \qquad L_{23} = F \cos \theta \sin \varphi$$

$$L_{13} = F \cos \theta \cos \varphi \qquad L_{24} = F \sin \theta \sin \varphi \qquad (IV.55)$$

$$L_{14} = F \sin \theta \cos \varphi \qquad L_{34} = p_{\theta}$$

where we have reduced the expressions on the surface $p_{\varphi} = p_{\theta} = 0$ and

$$F = e^{\sigma - \rho} p_{\sigma} - e^{\rho - \sigma} p_{\rho}.$$
 (IV.56)

A little care shows that the sin's and cos's in (IV.54) provide special solutions of the differential equations due to the algebra of constraints (in terms of new variables). In this way the non-abelian system of 6 constraints L_{ij} reduced to a system of 3 abelian constraints $p_{\varphi}, p_{\theta}, F$.

V. CONCLUDING REMARKS

Our main objective in this paper is that the algebra of Poisson brackets of first class systems may be simplified considerably. In fact, a complicated algebra may be escaped during the procedure of determining the constraints from the very beginning. This possibility has not been discussed here. The other possibility is simplifying the given constraints (regardless how they are produced) by using legal algebraic methods. The best thing that one may desire is an abelian system in which all of the constraints commute. For this reason one should try to find the most suitable coordinates describing either the whole phase space or the constraint surface.

One may use canonical transformations to change the coordinates of phase space. This process does not change the algebra of constraints. However, it may be used, for example, to clarify the multiplicative nature of the constraints. On the other hand, the constraint surface may be demonstrated by vanishing different sets of functions, which may have different algebras of Poisson brackets. This possibility, i.e. "redefinition" of constraint surface, motivates our desire of abelianization of the constraint systems.

In this paper we first observed that ordinary quadratic Lagrangians lead to abelian constraints which are linear with respect to phase space coordinates. Non abelian algebras require at least cubic or more complicated functions of coordinates and velocities in the Lagrangian.

We also showed with some examples that non abelian systems may appear because of unsuitable choice of constraints. In other words, one may solve the constraint equations to find the most simple way to describe the constraint surface. This procedure is in fact redefinition of the constraint surface. The main reason behind these calculations is to find the most suitable coordinates in which the constraint surface is described by vanishing or constancy of some phase space coordinates.

Although we were not able to prove a wide-standing theorem, we observed that sometimes the non abelian constraints (in suitable coordinates) contain factors, such as exponentials, that have no root in the finite region of the range of variables. Therefore, one may redefine the constraints by omitting such factors. This procedure is formulated in a systematic way by solving the differential equations due to the algebra of first class constraints. This is in fact the essence of our method of abelianization of the constraints.

The method may involve heavy algebraic manipulations, even in the seemingly simple step of canonical transformations. However, one can use it in any concrete example of constrained systems, despite the fact that for more complicated systems, like any other mathematical method, lengthy calculations are needed to get the final result. In other words it seems that all of the constrained systems may in principal be abelianized and no description can be found about a system which is "intrinsically non-abelianizable".

We showed that our method works well for a oneparticle as well as two-particle system with SO(3) symmetry. The problem is also solved for SO(4) symmetry and one may find it straightforward to follow the same procedure for SO(5) and more generally for SO(N). Other simple Lie algebras may also be considered in the same way.

There exist, however, two important points which should be considered carefully, i.e. "regularity" and "locality". The first point is the case when we encounter multiplicative constraints. We suggest that multiplicative constraints should be regularized at each step of calculation. This means that they should be replaced by simple roots of the corresponding expressions.

One important feature in this regard is that during simplifying the multiplicative constraints, the first class nature of the system may be altered. In general, there is no guarantee that the system remains first class after such simplification of the constraints. This point makes us to take the necessity of regularization of the constraints more serious, since otherwise it is possible to have a first class algebra of constraints out of a number of second class ones.

The next important point is that most of the above mathematical manipulations (i.e. regularization, canonical transformation and redefinition of constraints) which we use to find a simple description of constraint surface are valid locally, and may fail for some singular points or finite regions of the constraint surface. For example, in the case of SO(3) algebra we showed that the system can be abelianized everywhere except the origin of the phase space. In this way we can divide first class systems into "globally abelianizable" and "locally abelianizable".

We mention briefly that both the above problems are not difficulties with respect to our method of abelianization. They may be considered as intrinsic features of the constrained system itself, as it is described from the very beginning. For example, the problem of locality arises whenever the algebra of constraints is not valid globally throughout the whole phase space. Therefore, it is obvious that any method may encounter difficulties in the regions where the algebra is different.

The question may arise, however, that is there any advantage in employing the abelianized system instead of the original one? For instance what is preference of using the abelian constraints p_{φ} and η in the case of

SO(3), instead L_1 , L_2 and L_3 ? The answer depends on different applications of the first class constraints in physical problems such as gauge symmetry, quantization procedure, counting the physical degrees of freedom and so on. We postpone such analysis to future works. We reserve this possibility that maybe in some cases it is better to keep a non abelian and reducible algebra of

- M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princton Univ. Press, 1992.
- [2] J. M. Pons, J. Antonio Garcia, Int. J. Mod. Phys. A15 (2000) 4681-4721, [arXiv: hep-th/9908151].
- [3] M. Henneaux, C. Teitelboim, J. D. Vergara, Nucl. Phys. B387(1992) 391-418, [arXiv: hep-th/9205092].
- [4] A. Shirzad and M. Shabani Moghadam, J. Phys. A32 (1999) 8185-8195.
- [5] M. Henneaux, Phys. Rep. 126 (1985) 1.
- [6] S. A. Gogilidze, A. M. Khvedelidze and V. N. Pervushin, J. Math. Phys. **37**(1996) 1760-1771, [arXiv: hep-th/9504153].
- [7] F. Loran, Phys. Lett. B547 (2002) 63-68, [arXiv: hep-th/0209180].

constraints instead of change it into a non reducible and abelian one. The reason may be the better possibility of tracking the physical symmetries such as rotation.

Acknowledgements: The authors would like to thanks Institute for Research in Fundamental Sciences (IPM) for financial support.

- [8] F. Loran, Commun. Math. Phys. 254 (2005) 167-178, [arXiv: hep-th/0303014].
- [9] O. Miskovic, J. Zanelli, J. Math. Phys. 44 (2003) 3876-3887, [arXiv: hep-th/0302033].
- [10] R. Courant and D. Hilbert, Methods of Mathematical Physics, Interscience Poblishers John Wiley and Sons, 1989.
- [11] H. Goldstein, C. Poole and J. Safko, *Classical Mechanics* (3rd ed), Addison wesley, 2002.
- [12] J. Gomis, M. Henneaux, J. M. Pons, Class. Quantum. Grav. 7 (1990) 1089-1096.
- [13] F. Loran, Int. J. Mod. Phys. A23 (2008) 4051-4062, [arXiv:0808.3654].