

# Uniqueness of positive solutions to semilinear elliptic equations with double power nonlinearities

Shinji Kawano

## Abstract

We consider the problem

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \mathbb{R}^n, \\ \lim_{|x| \rightarrow \infty} u(x) = 0, \end{cases} \quad (1)$$

where

$$f(u) = -\omega u + u^p - u^q, \quad \omega > 0, \quad q > p > 1.$$

It is known that a positive solution to (1) exists if and only if  $F(u) := \int_0^u f(s)ds > 0$  for some  $u > 0$ . Moreover, Ouyang and Shi in 1998 found that the solution is unique if  $f$  satisfies furthermore the condition that  $\tilde{f}(u) := (uf'(u))'f(u) - uf'(u)^2 < 0$  for any  $u > 0$ . In the present paper we remark that this additional condition is unnecessary.

## 1 Introduction

We shall consider a boundary value problem

$$\begin{cases} u_{rr} + \frac{n-1}{r}u_r + f(u) = 0 & \text{for } r > 0, \\ u_r(0) = 0, \\ \lim_{r \rightarrow \infty} u(r) = 0, \end{cases} \quad (2)$$

where  $n \in \mathbb{N}$  and

$$f(u) = -\omega u + u^p - u^q, \quad \omega > 0, \quad q > p > 1.$$

The above problem arises in the study of

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \mathbb{R}^n, \\ \lim_{|x| \rightarrow \infty} u(x) = 0. \end{cases} \quad (3)$$

Indeed, the classical work of Gidas, Ni and Nirenberg [4, 5] tells us that any positive solution to (3) is radially symmetric. On the other hand, for a solution  $u(r)$  of (2),  $v(x) := u(|x|)$  is a solution to (3).

The condition to assure the existence of positive solutions to (3) (and so (2)) was given by Berestycki and Lions [1] and Berestycki, Lions and Peletier [2]:

**Proposition 1.** *A positive solution to (2) exists if and only if*

$$F(u) := \int_0^u f(s)ds > 0, \quad \text{for some } u > 0. \quad (4)$$

Uniqueness of positive solutions to (2) had long remained unknown. Finally in 1998 Ouyang and Shi [8] proved uniqueness for (2) with  $f$  satisfying the additional condition (See also Kwong and Zhang [7]):

**Proposition 2.** *If  $f$  satisfies furthermore the following condition, then the positive solution is unique;*

$$\tilde{f}(u) := (uf'(u))'f(u) - uf'(u)^2 < 0, \quad \text{for any } u > 0. \quad (5)$$

Following is the main result of the present paper:

**Theorem 1.** *If the nonlinearity  $f$  satisfies the existence condition (4), then the uniqueness condition (5) is automatically fulfilled.*

## 2 Proof of Theorem 1.

**Lemma 1.** *The existence condition (4) is equivalent to*

$$\omega < \omega_{p,q},$$

where

$$\omega_{p,q} = \frac{2(q-p)}{(p+1)(q-1)} \left[ \frac{(p-1)(q+1)}{(p+1)(q-1)} \right]^{\frac{p-1}{q-p}}.$$

(See Ouyang and Shi [8] and the appendix of Fukuizumi [3].)

**Lemma 2.** *The uniqueness condition (5) is equivalent to*

$$\omega < \eta_{p,q},$$

where

$$\eta_{p,q} = \frac{q-p}{q-1} \left[ \frac{p-1}{q-1} \right]^{\frac{p-1}{q-p}}.$$

The proofs of these Lemmas are nothing but straightforward calculation and shall be omitted.

*Proof of Theorem 1.* It is apparent that

$$0 < \omega_{p,q} < \eta_{p,q},$$

which asserts the theorem. □

## References

- [1] H. Berestycki and P. L. Lions, Nonlinear scalar field equation, I., *Arch. Rat. Math. Anal.* **82** (1983), 313-345.
- [2] H. Berestycki, P. L. Lions and L. A. Peletier, An ODE approach to the existence of positive solutions for semilinear problems in  $\mathbb{R}^N$ , *Indiana University Math. J.* **30** (1981), 141-157.
- [3] R. Fukuizumi, Stability and instability of standing waves for nonlinear Schrödinger equations, *Tohoku Mathematical Publications*, **No.25**, 2003.
- [4] B. Gidas, W. M. Ni and L. Nirenberg, Symmetry and related properties via the maximal principle, *Comm. Math. Phys.* **68** (1979), 209-243.
- [5] B. Gidas, W. M. Ni and L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbb{R}^n$ , Mathematical analysis and applications, Part A, *Adv. in Math. Suppl. stud.* **7a**, Academic press, 1981, 369-402.
- [6] S. Kawano, A remark on the uniqueness of positive solutions to semilinear elliptic equations with double power nonlinearities, preprint.
- [7] M. K. Kwong and L. Zhang, Uniqueness of positive solutions of  $\Delta u + f(u) = 0$  in an annulus, *Differential Integral Equations* **4** (1991), 583-599.
- [8] T. Ouyang and J. Shi, Exact multiplicity of positive solutions for a class of semilinear problems, *J. Differential Equations* **146** (1998), 121-156.
- [9] J. Wei and M. Winter, On a cubic-quintic Ginzburg-Landau equation with global coupling, *Proc. Amer. Math. Soc.* **133** (2005), 1787-1796.