

Existence and uniqueness conditions of positive solutions to semilinear elliptic equations with double power nonlinearities

Shinji Kawano

Abstract

We consider the problem

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \mathbb{R}^n, \\ \lim_{|x| \rightarrow \infty} u(x) = 0, \end{cases} \quad (1)$$

where

$$f(u) = -\omega u + u^p - u^q, \quad \omega > 0, \quad q > p > 1.$$

It is known that a positive solution to (1) exists if and only if $F(u) := \int_0^u f(s)ds > 0$ for some $u > 0$. Moreover, Ouyang and Shi in 1998 found that the solution is unique if f satisfies furthermore the condition that $\tilde{f}(u) := (uf'(u))'f(u) - uf'(u)^2 < 0$ for any $u > 0$. In the present paper we find the equivalent conditions of these. As a corollary we have a simpler proof of the result presented in the former paper that the existence condition asserts the uniqueness condition.

1 Introduction

We shall consider a boundary value problem

$$\begin{cases} u_{rr} + \frac{n-1}{r}u_r + f(u) = 0 & \text{for } r > 0, \\ u_r(0) = 0, \\ \lim_{r \rightarrow \infty} u(r) = 0, \end{cases} \quad (2)$$

where $n \in \mathbb{N}$ and

$$f(u) = -\omega u + u^p - u^q, \quad \omega > 0, \quad q > p > 1.$$

The above problem arises in the study of

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \mathbb{R}^n, \\ \lim_{|x| \rightarrow \infty} u(x) = 0. \end{cases} \quad (3)$$

Indeed, the classical work of Gidas, Ni and Nirenberg [4, 5] tells us that any positive solution to (3) is radially symmetric. On the other hand, for a solution $u(r)$ of (2), $v(x) := u(|x|)$ is a solution to (3).

The condition to assure the existence of positive solutions to (3) (and so (2)) was given by Berestycki and Lions [1] and Berestycki, Lions and Peletier [2]:

Proposition 1. *A positive solution to (2) exists if and only if*

$$F(u) := \int_0^u f(s)ds > 0, \quad \text{for some } u > 0. \quad (4)$$

Uniqueness of positive solutions to (2) had long remained unknown. Finally in 1998 Ouyang and Shi [9] proved uniqueness for (2) with f satisfying the additional condition (See also Kwong and Zhang [8]):

Proposition 2. *If f satisfies furthermore the following condition, then the positive solution is unique;*

$$\tilde{f}(u) := (uf'(u))'f(u) - uf'(u)^2 < 0, \quad \text{for any } u > 0. \quad (5)$$

In the present paper we find the equivalent conditions of these. Following is the main result of the present paper:

Theorem 1. *The existence condition (4) is equivalent to the following condition;*

$$\tilde{F}(u) = (uf(u))'F(u) - uf(u)^2 < 0, \quad \text{for any } u > 0. \quad (6)$$

The uniqueness condition (5) is equivalent to the following condition;

$$f(u) > 0, \quad \text{for some } u > 0. \quad (7)$$

As a corollary of the Theorem 1 above we have a simpler proof of the result presented in the former paper that the existence condition asserts the uniqueness condition [7]:

Corollary 1. *If the nonlinearity f satisfies the existence condition (4), then the uniqueness condition (5) is automatically fulfilled.*

This paper is organized as follows. In section 2 we present the proof of the theorem 1. In section 3 we present a simpler proof of the corollary.

2 Proof of Theorem 1.

The following two lemmas asserts the Theorem 1.

Lemma 1. *Both the existence condition (4) and the condition (6) are equivalent to*

$$\omega < \omega_{p,q},$$

where

$$\omega_{p,q} = \frac{2(q-p)}{(p+1)(q-1)} \left[\frac{(p-1)(q+1)}{(p+1)(q-1)} \right]^{\frac{p-1}{q-p}}.$$

(See Ouyang and Shi [9] and the appendix of Fukuizumi [3].)

Lemma 2. *Both the uniqueness condition (5) and the condition (7) are equivalent to*

$$\omega < \eta_{p,q},$$

where

$$\eta_{p,q} = \frac{q-p}{q-1} \left[\frac{p-1}{q-1} \right]^{\frac{p-1}{q-p}}.$$

The proofs of these Lemmas are nothing but straightforward calculation and shall be omitted.

3 Proof of Corollary 1.

Proof of Corollary 1. From Theorem 1 it is enough to show that if F has positive parts (the existence condition (4)) then f has positive parts (the condition (7)). The contraposition is clear by the monotonicity of the integral. \square

References

- [1] H. Berestycki and P. L. Lions, Nonlinear scalar field equation, I., *Arch. Rat. Math. Anal.* **82** (1983), 313-345.
- [2] H. Berestycki, P. L. Lions and L. A. Peletier, An ODE approach to the existence of positive solutions for semilinear problems in \mathbb{R}^N , *Indiana University Math. J.* **30** (1981), 141-157.
- [3] R. Fukuizumi, Stability and instability of standing waves for nonlinear Schrödinger equations, *Tohoku Mathematical Publications*, **No.25**, 2003.
- [4] B. Gidas, W. M. Ni and L. Nirenberg, Symmetry and related properties via the maximal principle, *Comm. Math. Phys.* **68** (1979), 209-243.
- [5] B. Gidas, W. M. Ni and L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n , *Mathematical analysis and applications*, Part A, *Adv. in Math. Suppl. stud.* **7a**, Academic press, 1981, 369-402.
- [6] S. Kawano, A remark on the uniqueness of positive solutions to semilinear elliptic equations with double power nonlinearities, preprint.
- [7] S. Kawano, Uniqueness of positive solutions to semilinear elliptic equations with double power nonlinearities, preprint.
- [8] M. K. Kwong and L. Zhang, Uniqueness of positive solutions of $\Delta u + f(u) = 0$ in an annulus, *Differential Integral Equations* **4** (1991), 583-599.
- [9] T. Ouyang and J. Shi, Exact multiplicity of positive solutions for a class of semilinear problems, *J. Differential Equations* **146** (1998), 121-156.
- [10] J. Wei and M. Winter, On a cubic-quintic Ginzburg-Landau equation with global coupling, *Proc. Amer. Math. Soc.* **133** (2005), 1787-1796.