

# SOME EXAMPLES OF LIFTING PROBLEMS FROM QUOTIENT ALGEBRAS

HYUN HO LEE

ABSTRACT. We consider three lifting questions: Given a  $C^*$ -algebra  $I$ , if there is a unital  $C^*$ -algebra  $A$  contains  $I$  as an ideal, is every unitary from  $A/I$  lifted to a unitary in  $A$ ? is every unitary from  $A/I$  lifted to an extremal partial isometry? is every extremal partial isometry from  $A/I$  lifted to an extremal partial isometry? We show several constructions of  $I$  which serve as working examples or counter-examples for above questions.

## 1. INTRODUCTION

Let  $\mathcal{E}(A)$ , or just  $\mathcal{E}$ , denote the set of extreme points of the unit ball of a unital  $C^*$ -algebra  $A$ . Recall that elements in  $\mathcal{E}$  are characterized as the partial isometries  $u$  in  $A$  satisfying  $(1 - uu^*)A(1 - u^*u) = 0$  by R. V. Kadison [10]. We call them extremal partial isometries and call the projections  $1 - uu^*$ ,  $1 - u^*u$  defect projections. In [4], Brown and Pedersen defined the notion of extremal richness for  $C^*$ -algebra  $A$  which means quasi-invertible elements are dense in  $A$  as an analogue of stable rank one for possibly infinite  $C^*$ -algebras. (We say  $T$  in  $A$  is quasi-invertible if  $T$  has closed range and the kernel projections of  $T^*$  and  $T$  are centrally orthogonal in  $A$ , or if  $T$  is in  $A^{-1}\mathcal{E}A^{-1}$ . For more equivalent definitions, see Theorem 1.1 in [4].) We denote by  $A_q^{-1}$  the set of quasi-invertible elements. As a result, stable rank one  $C^*$ -algebras are characterized within the class of extremally rich  $C^*$ -algebras by the property that all extreme points of the unit ball are unitaries or  $A_q^{-1} = A^{-1}$  where  $A^{-1}$  is the set of invertible elements of  $A$ .

Suppose we have an extension of  $C^*$ -algebras:

$$0 \longrightarrow I \longrightarrow A \longrightarrow B \longrightarrow 0$$

---

*Date:* Sep 15, 2008.

*2000 Mathematics Subject Classification.* Primary:46L05.

*Key words and phrases.* Extremally rich  $C^*$ -algebras, Good index theory, Stable rank one .

It is well known that if we have an extension of an extremally rich  $C^*$ -algebra  $I$  by an extremally rich  $C^*$ -algebra  $B$ , we cannot deduce  $A$  is extremally rich even in the finite case. The obstacle, as in the analogous problem with stable rank one, can be expressed as a lifting problem but with special properties. In fact, Brown and Pedersen proved the following theorem [3, Theorem 6.1].

**Theorem 1.1.** *If  $A$  is an extension of  $I$  by  $B$  and both  $I$  and  $B$  are extremally rich, then  $A$  is extremally rich if and only if extremal partial isometries in  $B$  are liftable to extremal partial isometries in  $A$  and  $PAQ$  is extremally rich whenever  $P$  is projection of the form  $1 - uu^*$  where  $u \in \mathcal{E}(A)$  and  $Q$  projection of the form  $1 - v^*v$  where  $v \in \mathcal{E}(\tilde{I})$ .*

However, there are some special ideals  $I$  for which the hypotheses in Theorem 1.1 can be simplified:

**Corollary 1.2.** *Let  $I$  be a  $C^*$ -algebra of stable rank one. Then  $A$  is extremally rich if and only if  $B$  is extremally rich and extremal partial isometries in  $B$  lift.*

This corollary implies the following: Whenever  $I$  embedded as an ideal in a unital extremally rich  $C^*$ -algebra and  $u$  is an extremal partial isometry in a extremally rich  $C^*$ -algebra  $A/I$ , there is an extremal partial isometry in  $A$  which lifts  $u$ .

Motivated by above example, but not necessarily related to extension, we can ask whether we can find examples of  $I$  such that whenever there exists  $A$  containing  $I$  as a closed ideal, certain lifting questions below from any quotient  $C^*$ -algebra  $B = A/I$  to  $A$  are affirmative:

- (1) Is every unitary in  $B$  lifted to a unitary in  $A$ ?
- (2) Is every unitary in  $B$  lifted to an extremal partial isometry in  $A$ ?
- (3) Is every extremal partial isometry in  $B$  lifted to an extremal partial isometry in  $A$ ?

We cannot expect a positive answer in great generality due to the nature of the problem; In fact, it is not very difficult to find counter examples for these questions. Interesting direction, however, is to find an affirmative answer. For example, it is well-known that if  $I$  has an approximate identity consisting of projections, then every unitary in  $B$  lifts to a partial isometry in  $A$ .

**Remark 1.3.** (i) When  $I$  and  $B$  are of stable rank one, an affirmative answer to (1) is equivalent to the stable rank one property for  $A$ .

- (ii) When  $I$  and  $B$  are of stable rank one, an affirmative answer to (2) is equivalent to extremal richness for  $A$ .
- (iii) When  $I$  has stable rank one and  $B$  is extremally rich, an affirmative answer to (3) is equivalent to extremal richness for  $A$ .

## 2. EXAMPLES

Throughout this article  $H$  will denote an infinite dimensional Hilbert space and  $B(H)$  the set of bounded operators on  $H$  and  $K(H)$  (shortly  $K$ ) the set of compact operators on  $H$ . Given a  $C^*$ -algebra  $I$ ,  $M(I)$  will denote the multiplier algebra of  $I$  and  $C(I)$  will denote the corona algebra of  $I$ , that is,  $M(I)/I$ . In view of Remark 1.3-(i), it is still interesting if we restrict  $I$  to be the class of stable rank one  $C^*$ -algebras. But, even in this case, question (1) is not true for every stable rank one  $C^*$ -algebras. In fact, there is a well-known counter example: the Toeplitz extension

$$0 \longrightarrow K \longrightarrow \mathcal{T} \longrightarrow C(S^1) \longrightarrow 0$$

where  $\mathcal{T}$  is the  $C^*$ -algebra generated by the unilateral shift  $S$  on  $H$  is such an example by the remark 1.3-(i) and the fact  $\text{tsr}(\mathcal{T}) \neq 1$ . (See [14, 4.13].) However, there is a  $C^*$ -algebra of stable rank one which serve as an answer for question (1).

**Definition 2.1.** ([5, p.2]) We say a (non-unital)  $C^*$ -algebra  $I$  has good index theory if whenever  $I$  is embedded as an ideal in a unital  $C^*$ -algebra  $A$  and  $u$  is a unitary in  $A/I$  such that  $\partial_1([u]) = 0$  where  $\partial_1 : K_1(A/I) \rightarrow K_0(I)$ , there is a unitary in  $A$  which lifts  $u$ .

It was noted by Brown and Pedersen that G. Nagy proved that any stable rank one  $C^*$ -algebra has good index theory [5]. Thus if we can show the existence of  $I$  with  $\text{tsr}(I) = 1$  such that  $\partial_1 : K_1(B) \rightarrow K_0(I)$  is trivial, question (1) holds for  $I$ .

**Example 2.2.** Let

$$I = \{(f_n) \mid f_n \in C(\mathbb{T}, M_n(\mathbb{C})) \text{ and } f_n \rightarrow f \text{ in } C(\mathbb{T}, \mathcal{K})\}$$

Here  $f_n \rightarrow f$  in  $C(\mathbb{T}, \mathcal{K})$  means  $\sup_{t \in \mathbb{T}} \|f_n(t) - f(t)\|$  goes to 0 as  $n$  goes to infinity. Note that  $I$  has stable rank one. Let  $C(\mathbb{T}, B(H)_{*-S})$  be the set of functions  $m : \mathbb{T} \rightarrow B(H)$  which are continuous with respect to the  $*$ -strong operator topology on  $B(H)$ , and which satisfy  $\|m\|_\infty := \sup_{t \in \mathbb{T}} \|m(t)\| < \infty$ .

It is not hard to show that

$$M(I) = \{(F_n) \mid F_n \in C(\mathbb{T}, M_n(\mathbb{C})) \text{ and } F_n \rightarrow_s F \text{ in } C(\mathbb{T}, B(H)_{*-S})\}$$

Here  $F_n \rightarrow_s F$  means  $F_n f_n \rightarrow Ff$  and  $F_n^* f_n \rightarrow F^* f$  in  $I$  for each  $(f_n) \in I$ . Next we want to show  $K_0(\iota) : K_0(I) \rightarrow K_0(M(I))$  is injective where  $\iota : I \rightarrow M(I)$ ; since the sequence  $\alpha = (z_n)$  in  $K_0(I)$  is eventually constant, we may assume  $I$  has only finite dimensional irreducible representations. Let  $\pi$  be such an irreducible representation and suppose  $K_0(\iota)(\alpha) = 0$ . Then  $K_0(\bar{\pi})(K_0(\iota)(\alpha)) = 0$  where  $\bar{\pi} : M(I) \rightarrow B(H)$  is a unique map such that  $\bar{\pi} \circ \iota = \pi$ . Note that we have  $\pi(I) = \bar{\pi}(I)$  since  $\pi$  is finite dimensional representation. Consequently, we have  $K_0(\pi)(\alpha) = 0$  in  $K_0(\pi(I))$ . By the Remark 5.12 in [3],  $\alpha = 0$ .

From the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I & \xrightarrow{j} & A & \longrightarrow & B & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & I & \xrightarrow{\iota} & M(I) & \longrightarrow & C(I) & \longrightarrow & 0 \end{array}$$

we have

$$\begin{array}{ccc} K_0(I) & \xrightarrow{K_0(j)} & K_0(A) \\ \parallel & & \downarrow \\ K_0(I) & \xrightarrow{K_0(\iota)} & K_0(M(I)) \end{array}$$

Thus it follows that  $K_0(j)$  is also injective. Thus from the exactness of six term exact sequence of K-theory we know the map  $\partial_1 : K_1(B) \rightarrow K_0(I)$  is trivial.

Next we show there are counter-examples to question (2).

**Example 2.3.** Let

$$I = \{(a_n) \in \prod M_{2n}(\mathbb{C}) \mid a_n \rightarrow \begin{pmatrix} D & 0 \\ 0 & E \end{pmatrix} \text{ in } M_2(K)\}$$

Then

$$M(I) = \{(T_n) \in \prod M_{2n}(\mathbb{C}) \mid T_n \rightarrow \begin{pmatrix} M & 0 \\ 0 & N \end{pmatrix} \text{ in } M_2(B(H)_{*-s})\}$$

Now let's fix the basis  $\mathfrak{B}$  of Hilbert space  $H$  as follows;

$$\{\dots, w_n, \dots, w_2, w_1, v_1, v_2, \dots, v_n, \dots\}$$

Let  $e_{xy}$  be a rank one projection such that  $e_{xy}(z) := \langle y, z \rangle x$ , and take  $T_n = \sum_{i=1}^{n-1} e_{w_i w_{i+1}} + \sum_{j=1}^{n-1} e_{v_{j+1} v_j} + e_{w_n v_n}$ . In fact,  $T_n$  is of the

following form with respect to the basis  $\mathfrak{B}$ :

$$\begin{pmatrix} 0 & 0 & \dots & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & 0 & \dots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & \dots & \dots & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & 0 & 0 & \ddots & 0 & 0 \\ 0 & \dots & \dots & 0 & 0 & \dots & 1 & 0 \end{pmatrix}$$

Then  $1 - T_n^*T_n = e_{w_1 w_1}$ ,  $1 - T_n T_n^* = e_{v_1 v_1}$ . Therefore  $\pi((T_n))$  is an unitary element of  $C(I)$  where  $\pi$  is the natural quotient map. But it cannot be liftable to an extremal partial isometry: Assume there is an element  $(a_n) \in I$  such that  $T_n + a_n$  is extremal partial isometry in  $M(I)$ . Since  $M_{2n}(\mathbb{C})$  has stable rank one,  $T_n + a_n$  must be a unitary, and it happens only when  $a_n = e_{v_1 w_1}$ . Then  $T_n + a_n$  cannot converge to the operator of the form  $\begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$ . It is a contradiction.

More sophisticated example can be found as follows.

**Example 2.4.** Let  $D$  be the stabilization of cone or suspension. Assume there exist a projection  $\mathbf{p}$  in the corona algebra of  $D$  such that which does not lift but its  $K_0$ -class does lift.(For the construction of such a projection, see the example 5.13 in [11].) If we let  $a$  be the self adjoint element which lifts  $\mathbf{p}$  in  $M(D)$ , we take  $I$  to be the  $C^*$ -algebra generated by  $a$  so that the quotient  $I/D$  is isomorphic to  $\mathbb{C}$ . Then the Busby invariant is determined by sending 1 to  $\mathbf{p}$ , and we have the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & D & \xrightarrow{j} & I & \longrightarrow & \mathbb{C} & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow \mathbf{p} & & \\ 0 & \longrightarrow & D & \longrightarrow & M(D) & \longrightarrow & C(D) & \longrightarrow & 0 \end{array}$$

By the long exact sequence, we have

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_0(I) & \longrightarrow & K_0(\mathbb{C}) & \xrightarrow{\partial_0} & K_1(D) \\ & & \downarrow & & \downarrow \mathbf{p} & & \parallel \\ 0 & \longrightarrow & K_0(M(D)) & \longrightarrow & K_0(C(D)) & \xrightarrow{\partial_0} & K_1(D) \end{array}$$

Since  $\partial_0([\mathbf{p}]_0) = 0$ ,  $\partial_0 : K_0(\mathbb{C}) \rightarrow K_1(D)$  becomes trivial. Thus  $K_0(I) \cong K_0(\mathbb{C})$ . In particular,  $K_0(I)$  is non-trivial. Consequently, we found a (stably) projectionless stable rank one  $C^*$ -algebra such that its

$K_0$ -group is non-trivial. Now stabilize this algebra and call it  $I$  again. We consider an extension of  $I$  by  $C(\mathbb{T})$  corresponding to a unitary  $u$  in the coronal algebra with non-trivial class  $K_1$ -class.  $u$  can't be lifted to a unitary (If so,  $[u]_1 = 0$  which is a contradiction), and it can't be lifted to a partial isometry either because there are no non-zero projections available to defect projections of the partial isometry.

On the other hand, we have a certain class of  $C^*$ -algebras (e.g. elementary  $C^*$ -algebras) such that question (2) is true.

**Proposition 2.5.** *Let  $I$  be the  $C^*$ -algebra such that  $M(I)$  is extremally rich. Suppose a unital  $C^*$ -algebra  $A$  contains  $I$  as an ideal. Then any unitary  $u$  in  $A/I$  can be liftable to an extremal partial isometry.*

*Proof.* From the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I & \longrightarrow & A & \longrightarrow & A/I \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \tau \\ 0 & \longrightarrow & I & \longrightarrow & M(I) & \longrightarrow & C(I) \longrightarrow 0 \end{array}$$

we see  $\tau(u)$  is also a unitary in  $C(I)$ . Since  $M(I)$  is an extremally rich  $C^*$ -algebra, we can find an extremal partial isometry  $w$  which is an inverse image of  $\tau(u)$  under the map  $\pi$  so that we have  $\pi(w) = \tau(u)$ . Hence  $(w, u)$  is the extremal partial isometry in  $A$  which lifts  $u$  since we can view  $A$  as the pullback construction of  $A/I$  and  $M(I)$ .  $\square$

**Remark 2.6.** We do not know the result of Proposition 2.5 is true under the weaker hypothesis  $C(I)$  is extremally rich. In addition,  $I$  in Example 2.3 and Example 2.4 are (stable rank one)  $C^*$ -algebras such that  $M(I)$  is not extremally rich.

**Corollary 2.7.** *Let  $I$  be the  $C^*$ -algebra such that  $M(I)$  is extremally rich and any Busby invariant into  $C(I)$  is extreme point preserving map. Suppose a unital  $C^*$ -algebra  $A$  contains  $I$  as an ideal. Then any extremal partial isometry  $u$  in  $A/I$  can be lifted to an extremal partial isometry.*

*Proof.* If a map is extreme point preserving, it sends an extremal partial isometry to an extremal partial isometry. Thus the result follows from above diagram.  $\square$

We have seen an example of  $I$  such that any extremal partial isometry in  $A/I$  can't be lifted to a partial isometry in  $A$  for some  $C^*$ -algebra  $A$  which contains  $I$  as an ideal in Example 2.4. If this does not happen, we can show the lifting question (3) might be true for a certain class of idealizers of  $I$ .

**Proposition 2.8.** *Let  $I$  be the  $C^*$ -algebra such that any extremal partial isometry in  $A/I$  is lifted to a partial isometry whenever a  $C^*$ -algebra  $A$  contains  $I$  as an ideal. If  $A$  is an extremally rich  $C^*$ -algebra, then any extremal partial isometry in  $A/I$  can be lifted to an extremal partial isometry in  $A$ .*

*Proof.* Let  $q$  be the quotient map from  $A$  to  $A/I$ . If  $u$  is an extremal partial isometry in  $B = A/I$ , it can be lifted to a partial isometry  $v$  in  $A$ . Since  $A$  is an extremally rich  $C^*$ -algebra,  $v$  has an extremal extension  $w$  in  $\mathcal{E}(A)$  such that  $v = vw^*v = vv^*w$  [4, Proposition 2.6]. Now let  $I^+$  and  $I^-$  be the defect ideals of  $u$  and consider the map  $\pi^+ \oplus \pi^- : B \rightarrow B/I^+ \oplus B/I^-$  which is injective. Since  $\pi^+ \oplus \pi^-(q(v)) = \pi^+ \oplus \pi^-(q(w))$ ,  $q(v) = q(w) = u$ . Hence an extremal partial isometry  $w$  is an inverse image of the extremal partial isometry  $u$ .  $\square$

However, the following examples show us that affirmative examples for question (3) might be hard to get without putting a condition on  $I$ .

**Example 2.9.** Let

$$A = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in M_2(B(H)) \mid B, C \in K \right\}$$

Then  $I = M_2(K)$  is an ideal of  $A$ . And  $B = A/M_2(K)$  is isomorphic to  $Q(H) \oplus Q(H)$ . Now consider  $\begin{pmatrix} S & 0 \\ 0 & T^* \end{pmatrix}$  where  $S$  and  $T$  are isometries

such that  $1 - SS^*$  and  $1 - TT^*$  are not compacts. Then  $\pi \left( \begin{pmatrix} S & 0 \\ 0 & T^* \end{pmatrix} \right)$

becomes an extremal partial isometry in  $Q(H) \oplus Q(H)$  where  $\pi : I \rightarrow B$ . We can check, however, that there is no extremal partial isometry

which lifts  $\pi \left( \begin{pmatrix} S & 0 \\ 0 & T^* \end{pmatrix} \right)$ : Suppose there is  $R \in M_2(K)$  such that  $V =$

$\begin{pmatrix} S & 0 \\ 0 & T^* \end{pmatrix} + R$  is an extremal partial isometry. Since  $A$  contains  $M_2(K)$

as an ideal, either one of defect projections of  $V$  must be zero. But

note that  $1 - V^*V = \begin{pmatrix} 0 & 0 \\ 0 & 1 - TT^* \end{pmatrix} + \text{compact}$  is in  $M_2(K)$ . Therefore

$1 - VV^* = 0$  implies  $1 - TT^*$  is compact which is a contradiction. The

other case is also similar.

The following clever example is due to Larry Brown although it was slightly generalized by the author.

**Example 2.10.** Let  $A$  be a unital  $C^*$ -algebra such that  $RR(A) = 0$ ,  $\text{tsr}(A) = 1$  and  $K_1(A) = 0$ . Then it is known that  $\text{tsr}(C[0, 1] \otimes A) = 1$

[12]. Now let's consider  $I$  as  $C[0, 1] \otimes (A \otimes \mathcal{K})$  which has stable rank one too. Note that  $M(A \otimes \mathcal{K})_s$  is equipped with the strict topology. Then  $M(I) = C([0, 1], M(A \otimes \mathcal{K})_s)$ .

If we denote  $[\mathcal{F}_A]$  by the set of homotopy classes of Freehold elements in  $M(A \otimes \mathcal{K})$ , there is Mingo's index map  $\partial : [\mathcal{F}_A] \rightarrow K_0(A)$  which is actually an isomorphism [13]. Suppose  $K_0(A)$  is an ordered group and let  $e$  be an order unit. Then let  $f_0 \in M(I)$  such that  $f_0(1) = 1$  and  $f_0(t)$  is an isometry with  $\partial(f_0(t)) = -2e$  for  $t < 1$ . Let  $u$  be a co-isometry in  $M(A \otimes \mathcal{K})$  with the index  $e$  and  $f$  be  $f_0u$ . Then we can see that  $\pi(f)$  is an isometry in  $\mathcal{C}(I)$  where  $\pi : M(I) \rightarrow C(I)$ . Assume there is a  $k \in I$  such that  $f + k$  is an extremal partial isometry in  $M(I)$  which is a continuous path of extremal partial isometries. Since  $M(A \otimes \mathcal{K})$  is a prime  $C^*$ -algebra, using index theory,  $f(1) + k(1)$  must be co-isometry of index  $e$ . It follows that there is a unitary  $v$  such that  $u + k(1) = uv$  [6, Theorem 2.1]. Since  $uv - u$  is in  $A \otimes \mathcal{K}$ ,  $u$  is a Fredholm element in  $M(A \otimes \mathcal{K})$ , and  $v \in 1 + A \otimes \mathcal{K}$ .

Finally, let  $f_1 = fv$  and  $k_1$  be  $f + k - f_1 = f(1 - v) + k \in I$ . Note that  $k_1(1) = 0$  and  $f + k = f_1 + k_1$ . Thus  $f_1(t) + k_1(t)$  must be an isometry for  $t < 1$  since  $f$  has negative index for  $t < 1$  and  $f(t) + k(t)$  is an extremal partial isometry (isometry or co-isometry in this case) for each  $t$ . Since  $f_1$  has a non-trivial kernel which is the range of  $v^*(1 - u^*u)$ , we know that  $\|k(t)\| \geq 1$  for  $t < 1$  which is a contradiction.

### 3. ACKNOWLEDGEMENTS

This is a part of the author's Ph.D. thesis. He would like to thank his advisor Larry Brown for many helpful suggestions and discussions.

### REFERENCES

- [1] L. G. Brown, G. K. Pedersen, *C\*-algebras of real rank zero*, J. Func. Anal. 99 (1991), 131-149
- [2] L. G. Brown *Interpolation by projections in C\*-algebras of real rank zero*, J.Operator Theory 26 (1991), 383-387
- [3] L. G. Brown, G. K. Pedersen *On the geometry of Unit ball of a C\* - algebra*, J. Reine agnew. Math. 469 (1995), 113-147
- [4] L. G. Brown, G. K. Pedersen *Limits and C\*-algebras of lower rank or dimension*, 2006 preprint
- [5] L. G. Brown, G. K. Pedersen *Non-stable K-theory and extremally rich C\*-algebras*, 2007 preprint
- [6] L. G. Brown, G. K. Pedersen *Extremal K-theory and Index for C\*-algebras*, K-theory 20 (2000), 201-241
- [7] J. W. Calkin *Tow sided ideals and congruences in the ring of bounded operators in Hilbert space*, Ann. of Math 42 (1941), 839-873



- [8] J. Cuntz, *K-theory for certain  $C^*$ -algebras*, Annals of Math., 113 (1981), 181-197
- [9] G. A. Elliott *Derivations of matroid  $C^*$ -algebras II*, Ann. Of Math., 100 (1974), 407-422
- [10] R.V. Kadison *Isometries of operator algebras*, Annals of Math. 54 (1951), 325-338
- [11] H. Lee *Some examples of non-stable K-theory*, Ph.D. Thesis Purdue University 2008
- [12] Nagisa, Masaru; Osaka, Hiroyuki; Phillips, N. Christopher *Ranks of algebras of continuous  $C^*$ -algebra valued functions*, Canad. J. Math. 53 (2001), no. 5, 979–1030.
- [13] J. A. Mingo, *K-theory and Multifiers of stable  $C^*$ -algebras*, Trans. of the Amer. Math. Soc. 299 (1987), 39 7-411
- [14] M. A. Rieffel *Dimensions and stable rank in the K-theory of  $C^*$ -algebras*, Proc. London Math.Soc. (3) 46 (1983), 301-333

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE,  
INDIANA 47907

*E-mail address:* ylee@math.purdue.edu