

Predator-Prey Quasi-cycles from a Path Integral Formalism

Thomas Butler and David Reynolds

*Department of Physics and Institute for Genomic Biology,
University of Illinois at Urbana Champaign, 1110 West Green Street, Urbana, IL 61801 USA*

(Dated: June 21, 2024)

The existence of beyond mean field quasi-cycle oscillations in a simple spatial model of predator-prey interactions is derived from a path integral formalism. The results agree substantially with those obtained from analysis of similar models using system size expansions of the master equation. In all of these analyses, the discrete nature of predator-prey populations and finite size effects lead to persistent oscillations in time, but spatial patterns fail to form. The path integral formalism goes beyond mean field theory and provides a focus on individual realizations of the stochastic time evolution of population not captured in the standard master equation approach.


PACS numbers: 87.23.Cc, 87.10.Mn, 02.50.Ey, 05.40.-a

When constructing models of biological phenomenon, observations of stable, periodic behavior have generally been taken to imply that the model will contain a stable limit cycle. In the context of ecological modeling, both simple heuristic arguments and field observations support predator-prey oscillations in ecosystems. However, the simple differential equation (mean field) models of predator-prey dynamics do not exhibit limit cycles [1, 2]. Several authors have addressed this difficulty by developing spatial individual level models (ILMs) that incorporate the stochastic effects of individual predator-prey interactions as in, for example, [3, 4, 5, 6]. These models generally yield limit cycles [6] or stochastically induced cycles dependent on space [3, 4, 5]. However, recent work on a 0 dimensional model has shown that space is not an essential ingredient for generation of temporal oscillations in predator-prey populations, but can arise by considering only the corrections to mean field theory emerging from an ILM for well mixed predator-prey populations [7]. Generalization of this work to space continues to show oscillations in time, but fails to exhibit oscillations in space [8].

The purpose of the present work is to develop a version of the spatial ILM of predator-prey interactions in [8] with only soft constraints and analyze the oscillatory fluctuations using path integral techniques. Our model includes the motion of both predator and prey, does not have a hard constraint on the number of organisms that can be present in a patch and will be found to have oscillations at the global scale consistent with previous results [8]. We use the standard mapping of the master equation to a bosonic field theory [9, 10, 11, 12] to obtain a simple derivation of coupled Langevin equations for the fluctuations of predator-prey populations. The path integral approach has the formal advantage of directly manipulating the population variables themselves rather than their probability distributions as in the master equation approach.

DEFINITION OF THE MODEL AND MASTER EQUATION

Consider a single, well-mixed patch of volume V . Species A is a predator for species B . We then have the following reactions:



We give the rates of the two body reactions an inverse V dependence, which is interpreted as the volume scaling of the probability in a volume V that the two organisms will be close enough to interact. We assume that the physical size of the patch is much greater than that of the organisms, but note that crowding must have an impact on the birth and death rates of the prey. We include this effect by imposing a concentration dependence on the death rate. For biological realism, one ought to include concentration dependence in the birth rate as well. However, the effect is the same if concentration dependence is put into one term or both — the limitation of prey population growth. Previous work has included concentration dependence by including a hard constraint on the number of available spaces in a patch, N , leading to an “Urn Model” description [13]. The probability of reactions then depends on the combinatorics of picking the appropriate reactants in the patch [7, 8, 13]. The relationship between the concentration dependent death rate presented in this paper and the urn model approach is analogous to the relationship between the $T < T_c$ ϕ^4 model and the nonlinear σ model; i.e. a soft versus hard constraint that does not change the universality class of the system [14]. An advantage of the current approach is that it avoids nonlinear diffusive cross terms in spatial models that do not seem to change the dynamics substantially from versions without the cross terms [8]. Ad-

ditionally, the hard constraints lead to substantial complications in the interpretation of the model parameters at the mean field level and in the master equation. This is due to the fact that reaction rates in urn models must be combined with the joint probability for drawing the reactants from the urn prior to use in the master equation or mean field description leading to complex combinations of parameters [13]. With the soft constraint applied here, the reaction rates have similar, predictable meanings at every level of description resulting in mean field parameters that correspond exactly to the reaction probabilities specified in Eq. 1 up to the volume scaling.

We include the concentration dependence of the death rate by noting that $n_A = N_A/V$ is small and write

$$d_1(n_A) = d_1(0) + cn_A + O(n_A^2), \quad c = d'(0) > 0 \quad (2)$$

We can now write a master equation for the patch

$$\begin{aligned} \partial_t P(m, n) = & d(-nP(m, n) + (n+1)P(m, n+1)) \\ & + c(-n^2P(m, n) + (n+1)^2P(m, n+1)) \\ & + b_1(-nP(m, n) + (n-1)P(m, n-1)) \\ & + p_1(-mnP(m, n) + (n+1)mP(m, n+1)) + \\ & p_2(-mnP(m, n) + (m-1)(n+1)P(m-1, n+1)) + \\ & d_2(-mP(m, n) + (m+1)P(m+1, n)) \quad (3) \end{aligned}$$

Where m denotes the number of predators, and n denotes the number of prey. This master equation defines the time evolution of the probability distribution of the different population states.

MAPPING TO PATH INTEGRAL FORMULATION

To analyze the predator prey dynamics, we map Eq. 3 to a field theory. This is done using the standard Doi formalism to obtain a second quantized Hamiltonian [9] and bosonic coherent states to map the resulting theory to a path integral. For our approach and helpful reviews, see [15, 16]. The mapping is achieved by introducing the state vector

$$|\psi\rangle = \sum_{m,n} P(m, n) |m, n\rangle \quad (4)$$

and the operator pairs a, \hat{a}, b, \hat{b} such that

$$\begin{aligned} a|m, n\rangle &= m|m-1, n\rangle \\ \hat{a}|m, n\rangle &= |m+1, n\rangle \\ [a, \hat{a}] &= 1 \\ b|m, n\rangle &= n|m, n-1\rangle \\ \hat{b}|m, n\rangle &= |m, n+1\rangle \\ [b, \hat{b}] &= 1 \quad (5) \end{aligned}$$

Finally, all other commutators are zero. We can then rewrite the dynamics given by the master equation (Eq. 3) as a Schrodinger like equation.

$$\partial_t |\psi\rangle = -\hat{H}(a, \hat{a}, b, \hat{b}) |\psi\rangle \quad (6)$$

We now can now specify the Hamiltonian (more accurately Liouvillian [12]) operator by multiplying the master equation by the state vector $|m, n\rangle$ summing over m and n , and applying the algebra of Eq. 6 to replace m and n by various combinations of the operators a, \hat{a} and b, \hat{b} . From this algebra, working out the structure of the Hamiltonian is direct and simple. As an example, we work out the term corresponding to prey birth explicitly

$$\begin{aligned} & b_1 \sum_{m,n} (-nP(m, n) + (n-1)P(m, n-1)) |m, n\rangle \\ = & b_1 \sum_{m,n} (-\hat{b}bP(m, n) + (n-1)P(m, n-1)) |m, n\rangle \\ = & -b_1 \hat{b}b |\psi\rangle + \sum_{m,n} nP(m, n) |m, n+1\rangle \\ = & -b_1 \hat{b}b |\psi\rangle + b_1 \hat{b}\hat{b}b |\psi\rangle \quad (7) \end{aligned}$$

Other terms are treated analogously. With normal ordering, this leads to the Hamiltonian

$$\begin{aligned} \hat{H} = & b_1(\hat{b}b - \hat{b}^2b) + d_1(\hat{b}b - b) + \frac{c}{V}(\hat{b}^2b^2 - \hat{b}b^2) \\ & + \frac{p_1}{V}(\hat{a}a\hat{b}b - \hat{a}ab) + \frac{p_2}{V}(\hat{a}a\hat{b}b - \hat{a}^2ab) \\ & + d_2(\hat{a}a - a) \quad (8) \end{aligned}$$

Expectation values of functions of the random variables m and n are given by

$$\langle f \rangle = \langle 0, 0 | e^{a+b} f(\hat{a}, a, \hat{b}, b) e^{-H(\hat{a}, a, \hat{b}, b)t} | \psi(0) \rangle \quad (9)$$

Using bosonic coherent states, it is easy to write Eq. 9 as a path integral resulting in a Lagrangian description of the dynamics with generalization to space following standard procedures [10, 11]. Since we are interested in persistent oscillations around the only stable fixed point in the system, our choice of initial conditions is irrelevant and can be ignored for the purposes of studying persistent quasi-cycles. To add space, we add diffusion terms to the resulting Lagrangian. Such an addition of space does have an implicit continuum limit, but if nearest neighbor hopping probabilities are scaled appropriately, it is well behaved [11]. The resulting Lagrangian density is given by

$$\begin{aligned} \mathcal{L} = & a^* \partial_t a + b^* \partial_t b - D_1 a^* \nabla^2 a - D_2 b^* \nabla^2 b \\ & + H(\hat{b}, \hat{a}, b, a) \quad (10) \end{aligned}$$

With fields derived from boson operators, the Lagrangian form of the master equation is not simply interpreted. This is because the field variables in the Lagrangian are not simply related to the physical variables of population number. This proves to be the source of difficulties in deriving correlation functions that are physically meaningful. To address this difficulty, we use a standard semi canonical Cole-Hopf transformation [17] to transform the field variables to density variables

$$a = ze^{-\hat{z}}, \hat{a} = e^{\hat{z}} \quad (11)$$

$$b = \rho e^{-\hat{\rho}}, \hat{b} = e^{\hat{\rho}} \quad (12)$$

This formulation has the advantage that z and ρ can be directly interpreted as the density variables for predator and prey respectively, while $\hat{\rho}$ and \hat{z} generate noise terms at quadratic order. The transformed Lagrangian takes the form

$$\begin{aligned} \mathcal{L} = & \hat{z}\partial_t z + \hat{\rho}\partial_t \rho - D_1 \hat{z}\nabla^2 z - D_1 z(\nabla \hat{z})^2 \\ & - D_2 \rho(\nabla \hat{\rho})^2 - D_2 \hat{\rho}\nabla^2 \rho - b_1 \rho(1 - e^{\hat{\rho}}) \\ & + d_1 \rho(1 - e^{-\hat{\rho}}) + \frac{c}{V} \rho^2(1 - e^{-\hat{\rho}}) \\ & + \frac{p_1}{V} z \rho(1 - e^{-\hat{\rho}}) + \frac{p_2}{V} z \rho(1 - e^{\hat{z}-\hat{\rho}}) \\ & + d_2 z(1 - e^{-\hat{z}}) \end{aligned} \quad (13)$$

In this form, the Lagrangian has diffusive noise, and difficult to handle exponential terms. In the following section, we exploit the small parameter $1/V$ to resolve these difficulties and analyse the theory.

DERIVATION OF MEAN FIELD THEORY AND QUASI-CYCLE OSCILLATIONS THROUGH LARGE V EXPANSION

From the Lagrangian in Eq. 13, we can proceed directly by rewriting the fields as

$$\begin{aligned} \hat{z} & \rightarrow \frac{\hat{z}}{\sqrt{V}} \\ \hat{\rho} & \rightarrow \frac{\hat{\rho}}{\sqrt{V}} \\ z & = V\varphi + \sqrt{V}\eta \\ \rho & = V\phi + \sqrt{V}\xi \end{aligned} \quad (14)$$

and inserting them into the Lagrangian. These forms are intended to capture Gaussian fluctuations in the spirit of the traditional system size expansion of the master equation due to Van Kampen [18] while directly manipulating the population variables. The fields \hat{z} and $\hat{\rho}$ have a mean field value of 0 after the Cole-Hopf transformation due to

conservation of probability [16]. This means that within the Gaussian approximation, the leading order term in those fields is a small correction of order $1/\sqrt{V}$ as above.

To derive the mean field theory and the fluctuations, we then insert the rhs forms of the fields in Eq. 14 into the Lagrangian Eq. 13 and retain only leading and next to leading order, resulting in an effective Lagrangian of the form

$$\mathcal{L} = \sqrt{V}\mathcal{L}_1 + \mathcal{L}_2 + O(1/\sqrt{V}) \quad (15)$$

Deriving each of these terms is straightforward. For purposes of illustration, we will carry out the expansion for the prey birth term explicitly

$$\begin{aligned} & b_1 \rho(1 - e^{\hat{\rho}}) \\ = & b_1(V\phi + \sqrt{V}\xi)\left(-\frac{\hat{\rho}}{V} - \frac{\hat{\rho}^2}{2V}\right) \\ = & b_1(-\sqrt{V}\hat{\rho}\phi - \frac{\hat{\rho}^2\phi}{2} - \hat{\rho}\eta) \end{aligned} \quad (16)$$

Carrying this out for each term in the Lagrangian and collecting terms yields at order \sqrt{V}

$$\begin{aligned} \mathcal{L}_1 = & \hat{\rho}\partial_t \phi + \hat{z}\partial_t \varphi - D_1 \hat{z}\nabla^2 \varphi - D_2 \hat{\rho}\nabla^2 \phi \\ & - b_1 \phi \hat{\rho} + d_1 \varphi \hat{\rho} + c \hat{\rho} \phi^2 + p_1 \hat{\rho} \varphi \phi + p_2 \hat{\rho} \phi \varphi \\ & - p_2 \hat{z} \phi \varphi + d_2 \hat{z} \varphi \end{aligned} \quad (17)$$

Minimizing this term provides the mean field theory. For $V \rightarrow \infty$, this minimum is exact. The Euler-Lagrange equations applied to \mathcal{L}_1 provide the mean field equations

$$\begin{aligned} \frac{\delta \mathcal{L}_1}{\delta \hat{z}} & = \partial_t \varphi - D_1 \nabla^2 \varphi - p_2 \phi \varphi + d_2 \varphi = 0 \\ \frac{\delta \mathcal{L}_1}{\delta \hat{\rho}} & = \partial_t \phi - D_2 \nabla^2 \phi - b_1 \phi + d_1 \phi + c \phi^2 \\ & + p_1 \varphi \phi + p_2 \phi \varphi = 0 \end{aligned} \quad (18)$$

These are the standard Lotka-Volterra equations generalized to include space. They arise as the leading order contribution in large volume limit as the saddle point of the path integral and flow to stable, spatially uniform solutions with magnitudes given by the fixed points of the ordinary differential equations obtained by dropping the diffusion operator in Eqs. 18 above.

At next to leading order, we Fourier transform and switch to matrix notation, defining

$$\mathbf{x} = \begin{pmatrix} \eta \\ \xi \end{pmatrix}, \mathbf{y} = \begin{pmatrix} \hat{z} \\ \hat{\rho} \end{pmatrix} \quad (19)$$

By simply collecting terms as in Eq. 14 we can write down \mathcal{L}_2 as

$$\mathcal{L}_2 = i\omega \mathbf{y}^T \mathbf{x} + \mathbf{y}^T \mathbf{A} \mathbf{x} - \frac{1}{2} \mathbf{y}^T \mathbf{B} \mathbf{y} \quad (20)$$

The matrices are given by

$$\mathbf{A} = \begin{pmatrix} D_1 k^2 & -p_2 \varphi \\ (p_1 + p_2) \phi & D_2 k^2 + c\phi \end{pmatrix} \quad (21)$$

and

$$\mathbf{B} = \begin{pmatrix} 2(d_2 + D_1 k^2) \varphi & -p_2 \varphi \phi \\ -p_2 \varphi \phi & 2(b_1 + D_2 k^2) \phi \end{pmatrix} \quad (22)$$

We now can recognize that the vector \mathbf{y} is a response field in the Martin Siggia Rose response function formalism for Langevin equations [19, 20]. Thus the fluctuations around mean field in the path integral are coupled Langevin equations. The resulting Langevin equations with the appropriate noise and correlations are

$$\begin{aligned} -i\omega \mathbf{x} &= \mathbf{A} \mathbf{x} + \gamma(\omega) \\ \langle \gamma_i(\omega) \gamma_j(-\omega) \rangle &= B_{ij} \end{aligned} \quad (23)$$

With a couple of modifications due to the addition of space and the model differences discussed above, these equations are the same as the equations reported in [7, 21] and are easily solved for the power spectrum. We solve for the average power spectrum, which captures oscillations but is free of phase cancellations and is the appropriate quantity for studying oscillations in the system [7].

To obtain the power spectrum, we proceed with simple linear algebra manipulations [21]

$$\begin{aligned} \mathbf{x} &= -(\mathbf{A} + i\omega)^{-1} \gamma(\omega) \equiv \mathbf{D}(\omega)^{-1} \gamma(\omega) \\ \rightarrow x_1 = \eta &= -\det(\mathbf{D})^{-1} (\mathbf{D}_{11} \gamma_1 - \mathbf{D}_{12} \gamma_2) \\ x_2 = \xi &= -\det(\mathbf{D})^{-1} (\mathbf{D}_{21} \gamma_1 - \mathbf{D}_{22} \gamma_2) \end{aligned} \quad (24)$$

The power spectrum is then obtained by taking the amplitude squared and averaging. For the predator fluctuations this gives

$$\langle x_1 x_1^* \rangle = \frac{\alpha_k + \beta_k \omega^2}{(\omega^2 - \Omega_k^2)^2 + \Gamma_k^2 \omega^2} \quad (25)$$

with

$$\begin{aligned} \alpha_k &= B_{11}(k) A_{22}^2 + B_{22}(k) A_{12}^2 \\ \beta_k &= B_{11}(k) \\ \Omega_k^2 &= D_1 k^2 (D_2 k^2 + c\phi) + p_2 (p_1 + p_2) \phi \varphi > 0 \\ \Gamma &= -A_{11} - A_{22} \end{aligned} \quad (26)$$

The power spectrum contains a nontrivial peak in ω corresponding to the expected temporal oscillations. The peak in k is at 0 wavenumber as can be seen from the strictly increasing functions of k present in the spectrum. These results are in qualitative agreement with previous results from expansion of the master equation for spatially generalized Urn models of predator prey interactions [7, 8]. Further work is underway to apply the methods of this paper to problems involving fluctuating phage-host dynamics and other problems in population biology that require detailed analysis of fluctuations, and transparent connection of different levels of description in order to compare to experimental data.

We thank Nigel Goldenfeld for suggesting this problem and for helpful discussions. This work was partially supported by the National Science Foundation and the Department of Energy grants NSF-EF-0526747 and DOE-2005-05818.

-
- [1] M. A. Nowak, *Evolutionary Dynamics* (Belknap/Harvard Press, 2006).
 - [2] S. H. Strogatz, *Nonlinear Dynamics and Chaos* (Westview Press, 1994).
 - [3] M. Mobilia, I. T. Georgiev, and U. C. Tauber, Phys. Rev. E **73**, 040903 (2006).
 - [4] M. Mobilia, I. T. Georgiev, and U. C. Tauber, J. Stat. Phys. **128**, 447 (2007).
 - [5] N. Boccara, R. O., and M. Roger, Phys. Rev. E **50**, 4531 (1994).
 - [6] T. Antal and M. Droz, Phys. Rev. E **63**, 056119 (2001).
 - [7] A. J. McKane and T. J. Newman, Phys. Rev. Lett. **94**, 218102 (2005).
 - [8] C. Lugo and A. J. McKane, arXiv:0806.1287v1 [q-bio.PE] (2008).
 - [9] M. Doi, J. Phys. A. **9**, 1465 (1976).
 - [10] A. S. Mikhailov, Phys. Lett. **85**, 214 (1981).
 - [11] L. Peliti, P.J. Physique **46**, 1469 (1985).
 - [12] N. Goldenfeld, J. Phys. A **17**, 2807 (1984).
 - [13] A. J. McKane and T. J. Newman, Phys. Rev. E **70**, 041902 (2004).
 - [14] M. Stone, *The Physics of Quantum Fields* (Springer, 2000).
 - [15] D. Mattis and M. L. Glasser, Rev. Mod. Phys. **70**, 979 (1998).
 - [16] J. Cardy, arXiv:cond-mat/9607163v2 (1996).
 - [17] H. K. Janssen and U. C. Tauber, Annals of Physics **315**, 147 (2005).
 - [18] N. G. Van Kampen, *Stochastic Processes in Physics and Chemistry* (Elsevier, New York, 1992).
 - [19] P. C. Martin, E. D. Siggia, and H. A. Rose, Phys. Rev. A **8**, 423 (1973).
 - [20] R. Bausch, H. K. Janssen, and H. Wagner, Z. Phys. B. **24**, 113 (1976).
 - [21] A. McKane, J. D. Nagy, T. J. Newman, and M. O. Stefanini, J. Stat. Phys. **128**, 165 (2007).