# Dressed-particle approach in the nonrelativistic classical limit

I. Y. Dodin and N. J. Fisch

Department of Astrophysical Sciences, Princeton University, Princeton, New Jersey 08544, USA

(Dated: October 24, 2018)

For a nonrelativistic classical particle undergoing arbitrary oscillations, the generalized effective potential  $\Psi$  is derived from nonlinear eigenfrequencies of the particle-field system. Specifically, the ponderomotive potential is extended to a nonlinear oscillator, resulting in multiple branches near the primary resonance. For a pair of natural frequencies in a beat resonance,  $\Psi$  scales linearly with the internal actions and is analogous to the dipole potential for a two-level quantum system. Thus cold quantum particles and highly-excited quasiclassical objects permit uniform manipulation tools, particularly, one-way walls.

PACS numbers: 52.35.Mw, 05.45.-a, 45.20.Jj, 45.05.+x

#### I. INTRODUCTION

Multiscale adiabatic dynamics of classical particles in oscillating and static fields is simplified within the oscillation-center (OC) approach, which allows separating fast quiver motion of the particles from their slow translational motion [1, 2, 3]. Hence the average forces are embedded into the properties of the OC, yielding a quasiparticle with a variable effective mass  $m_{\text{eff}}$  [4, 5]. In each given case,  $m_{\text{eff}}$  can be Taylor-expanded at nonrelativistic energies so as to appear as an effective potential  $\Psi$  [4], e.g., ponderomotive [6, 7, 8, 9] or diamagnetic [10]. Yet the nonrelativistic limit must permit also an independent calculation of  $\Psi$ . For linear oscillations, the generalized effective potential was derived in Ref. [3]. However, a comprehensive method of finding  $\Psi$  for nonlinear quiver motion has not been proposed.

The purpose of this work is to calculate, from first principles, the generalized effective potential  $\Psi$  for a nonrelativistic classical particle undergoing arbitrary oscillations in high-frequency or static fields. We proceed by finding eigenmodes in the particle-field system; hence  $\Psi$  is obtained like in the dressed-atom approach [11, 12, 13, 14] but from nonlinear classical equations. Specifically, we show that the ponderomotive potential extended to a nonlinear oscillator has multiple branches near the primary resonance. Also, for a pair of natural frequencies in a beat resonance,  $\Psi$  scales linearly with the internal actions and is analogous to the dipole potential for a twolevel quantum system. Thus cold quantum particles and highly-excited quasiclassical objects permit uniform manipulation tools, particularly, stationary asymmetric barriers, or one-way walls [15, 16, 17, 18, 19, 20, 21, 22, 23].

The work is organized as follows. In Sec. II, we obtain the general form of the effective potential  $\Psi$ . In Sec. III, we derive the equations for oscillation modes. In Sec. IV, we calculate the ponderomotive potential from the infinitesimal frequency shift of the oscillating field coupled to a particle at a primary resonance, both linear and nonlinear; see also Appendix A. In Sec. V, we find  $\Psi$  near a beat resonance and show the analogy with the dipole potential. In Sec. VI, we explain how  $\Psi$  allows one-way walls. In Sec. VII, we summarize our main results.

#### II. GENERALIZED EFFECTIVE POTENTIAL

Consider a particle which exhibits slow dynamics in canonical variables  $(\mathbf{r}, \mathbf{P})$  superimposed on fast oscillations in angle-action variables  $(\boldsymbol{\vartheta}, \mathcal{J} = \text{const})$ , such that zero  $\mathcal{J}$  corresponds to purely translational motion. The OC Lagrangian is then written as [4]

$$\mathcal{L}_0 = -m_{\rm eff} c^2 \sqrt{1 - v^2/c^2},\tag{1}$$

where  $\mathbf{v} \equiv \dot{\mathbf{r}}$ , and  $m_{\text{eff}}(\mathbf{r}, \mathbf{v}; \boldsymbol{\mathcal{J}})$  is the effective mass, the dependence on  $\boldsymbol{\mathcal{J}}$  being parametric [24]. Thus the complete Lagrangian, which describes also the oscillations, equals  $\mathcal{L}_0 + \boldsymbol{\vartheta} \cdot \boldsymbol{\mathcal{J}}$ , so the corresponding Hamiltonian  $\mathcal{H}$  matches that of the oscillation center:  $\mathcal{H} = \mathbf{P} \cdot \mathbf{v} - \mathcal{L}_0$ .

Assume nonrelativistic dynamics, i.e.,  $v \ll c$  and  $\delta m \equiv m_{\text{eff}} - m \ll m$ , where *m* is the true mass. Hence

$$\mathcal{L}_0 = \frac{1}{2} m v^2 - \mathcal{U}, \quad \mathcal{U} = \delta m c^2, \tag{2}$$

so the Hamiltonian reads

$$\mathcal{H} = \frac{P^2}{2m} + \Psi, \quad \Psi = \mathcal{U} - \frac{1}{2m} (\partial_{\mathbf{v}} \mathcal{U})^2, \quad (3)$$

where  $\mathbf{P} = m\mathbf{v} - \partial_{\mathbf{v}}\mathcal{U}$ , and  $\Psi(\mathcal{J} = 0) = 0$ .

Following Ref. [4], one can find  $\Psi$  from the relativistic particle trajectory in given fields; however, a general nonrelativistic approach is also possible. Formally, field modes can be understood as particle degrees of freedom. Then, like in the dressed-atom approach [11, 12], the effective potential can be found from eigenfrequencies  $\boldsymbol{\varpi} \equiv \boldsymbol{\vartheta} = \partial_{\mathcal{J}} \mathcal{H}$  of the particle-field system:

$$\Psi = \int \boldsymbol{\varpi} \cdot d\boldsymbol{\mathcal{J}}.$$
 (4)

In this case,  $\Psi$  depends on  $\mathbf{r}$  and  $\mathbf{P}$  only parametrically, through  $\boldsymbol{\varpi}(\mathbf{r}, \mathbf{P}, \mathcal{J})$ .

The canonical frequencies can be redefined such that  $\varpi \to \varpi + \text{const}$ , adding a constant to  $\Psi$ . Albeit arbitrarily large, this contribution does not affect the motion equations, so we abandon the requirement that  $\Psi$  must

remain small compared to  $mc^2$ ; hence actual physical frequencies can be used for  $\boldsymbol{\varpi}$ . Particularly, for uncoupled modes one gets  $\Psi = \Psi_0$ ,

$$\Psi_0 = \mathbf{\Omega} \cdot \mathbf{J} + \boldsymbol{\omega} \cdot \mathbf{I},\tag{5}$$

where we used Eq. (4) and  $(\Omega, \mathbf{J})$ ,  $(\boldsymbol{\omega}, \mathbf{I})$  for the unperturbed frequencies and actions of the particle and the field, correspondingly. For an unbounded field  $(I \to \infty)$ , the second term in Eq. (4) is infinite. As it is fixed though, the force on a particle is determined only by  $\Omega \cdot \mathbf{J}$  and the finite modification of the effective potential due to coupling,

$$\Phi \equiv \Psi - \Psi_0, \tag{6}$$

where  $\Phi$  also can be found from Eq. (4), as shown below.

## **III. PARTIAL MODE DECOMPOSITION**

Suppose weakly nonlinear oscillations  $\boldsymbol{\xi}(t)$ , both of the particle [3] and of external fields [25], so their Lagrangian reads  $\tilde{\mathcal{L}}(\boldsymbol{\xi}, \dot{\boldsymbol{\xi}}) = \tilde{\mathcal{L}}_0 + \tilde{\mathcal{L}}_{int}$ , where  $\tilde{\mathcal{L}}_{int}$  is a perturbation to a bilinear form  $\tilde{\mathcal{L}}_0$  [26],

$$\tilde{\mathcal{L}}_0 = \frac{1}{2} \left( \dot{\boldsymbol{\xi}} \cdot \hat{M} \dot{\boldsymbol{\xi}} \right) - \left( \dot{\boldsymbol{\xi}} \cdot \hat{R} \boldsymbol{\xi} \right) - \frac{1}{2} \left( \boldsymbol{\xi} \cdot \hat{Q} \boldsymbol{\xi} \right).$$
(7)

Here  $\hat{M}$ ,  $\hat{R}$ ,  $\hat{Q}$  are  $N \times N$  real matrices;  $\hat{M}$  and  $\hat{Q}$  are symmetric,  $\hat{R}$  is antisymmetric, and rank  $\hat{M} = N \equiv \dim \boldsymbol{\xi}$ . At zero  $\hat{R}$ ,  $\tilde{\mathcal{L}}_0(\boldsymbol{\xi}, \boldsymbol{\xi})$  can be diagonalized to yield

$$\tilde{\mathcal{L}}_0 = \sum_{j=1}^N L_j, \quad L_j = \frac{1}{2} M_j \dot{\xi}_j^2 - \frac{1}{2} Q_j \xi_j^2,$$
 (8)

where  $L_i$  describe individual modes  $\xi_i$  [27]. Then

$$\hat{D}_j \xi_j = \delta_{\xi_j} \tilde{\mathcal{L}}_{\text{int}}, \quad \hat{D}_j = M_j \, d_t^2 + Q_j, \tag{9}$$

 $\delta$  and  $d_t$  standing for the variational and time derivatives. Yet, such decomposition does not hold in the general case, so we redefine eigenmodes, following Ref. [28].

Extend the configuration space by introducing

$$\langle \ell | = (-\pi \hat{M}^{-1}, \boldsymbol{\xi}), \quad |r\rangle = (\boldsymbol{\xi}, \hat{M}^{-1}\pi)^{\mathrm{T}}$$
(10)

as the new, "left" and "right", coordinate vectors, where  $\pi = \hat{M}\dot{\xi} - \hat{R}\xi$  is the old canonical momentum. Then

$$\tilde{\mathcal{L}}_{0} = \frac{1}{4} \left[ \langle \dot{\ell} | \hat{\mathfrak{M}} | r \rangle - \langle \ell | \hat{\mathfrak{M}} | \dot{r} \rangle \right] + \frac{1}{2} \langle \ell | \hat{\mathfrak{F}} | r \rangle , \qquad (11)$$

where we omitted a full time derivative and introduced

$$\hat{\mathfrak{M}} = \begin{pmatrix} \hat{M} & 0\\ 0 & \hat{M} \end{pmatrix}, \quad \hat{\mathfrak{F}} = \begin{pmatrix} \hat{R} & \hat{M}\\ \hat{F} & \hat{R} \end{pmatrix}, \quad (12)$$

with  $\hat{F} = \hat{R}\hat{M}^{-1}\hat{R} - \hat{Q}$ . Thus the resulting equations are

$$\langle \dot{\ell} | \, \hat{\mathfrak{M}} + \langle \ell | \, \hat{\mathfrak{F}} = 0, \quad \hat{\mathfrak{M}} \, | \dot{r} \rangle - \hat{\mathfrak{F}} \, | r \rangle = 0, \tag{13}$$

both equivalent to

$$\hat{M}\ddot{\boldsymbol{\xi}} - 2\hat{R}\dot{\boldsymbol{\xi}} + \hat{Q}\boldsymbol{\xi} = 0.$$
(14)

Eq. (14) has 2N eigenmodes  $\boldsymbol{\xi}_j = \bar{\boldsymbol{\xi}}_j e^{-i\nu_j t}$ , with  $\nu_j$  hence assumed real and nonzero; therefore, for each  $\boldsymbol{\xi}_j$ , there also exists a mode  $\boldsymbol{\xi}_{-j} = \boldsymbol{\xi}_j^*$ , and  $\bar{\boldsymbol{\xi}}_j$  are generally not orthogonal. The corresponding eigenmodes of Eqs. (13) are

$$\langle \ell_j | = e^{i\nu_j t} \langle \bar{\ell}_j |, \quad |r_j \rangle = e^{-i\nu_j t} |\bar{r}_j \rangle,$$
 (15)

with vector amplitudes

$$\langle \bar{\ell}_j | = (-i\nu_j \bar{\boldsymbol{\xi}}_j^* - \bar{\boldsymbol{\xi}}_j^* \hat{R} \hat{M}^{-1}, \bar{\boldsymbol{\xi}}_j^*), \qquad (16)$$

$$|\bar{r}_j\rangle = (\bar{\boldsymbol{\xi}}_j, -i\nu_j \bar{\boldsymbol{\xi}}_j - \hat{M}^{-1} \hat{R} \bar{\boldsymbol{\xi}}_j)^{\mathrm{T}}, \qquad (17)$$

and  $\mathfrak{M}_{jk} \equiv \langle \bar{\ell}_j | \hat{\mathfrak{M}} | \bar{r}_k \rangle = -2i\rho_{jk}$ , where

$$\rho_{jk} = \frac{1}{2} \, \bar{\boldsymbol{\xi}}_j^* \cdot \left[ (\nu_j + \nu_k) \hat{M} - 2i\hat{R} \right] \cdot \bar{\boldsymbol{\xi}}_k. \tag{18}$$

The matrix  $\hat{\rho}$  is diagonal for distinct  $\nu_j$ , as seen from Eq. (14), or can be diagonalized when some of the frequencies coincide [28]; thus,

$$\mathfrak{M}_{jk} = -2i\rho_j\delta_{jk}, \quad \rho_j \equiv \nu_j(\bar{\boldsymbol{\xi}}_j^* \cdot \hat{M}\bar{\boldsymbol{\xi}}_j), \qquad (19)$$

 $\rho_j = -\rho_{-j}$ . (Hence modes with  $\nu_j = 0$  are orthogonal to the others and can be considered separately, as implied below.) Therefore any  $\langle \ell |$  and  $|r \rangle$  are decomposed as

$$\langle \ell | = \sum_{j=-N}^{N} \ell_j \langle \bar{\ell}_j |, \quad |r\rangle = \sum_{j=-N}^{N} r_j |\bar{r}_j\rangle, \qquad (20)$$

where the primes stand for skipping j = 0, and

$$\ell_j = \frac{i}{2\rho_j} \left\langle \ell | \hat{\mathfrak{M}} | \bar{r}_j \right\rangle, \quad r_j = \frac{i}{2\rho_j} \left\langle \bar{\ell}_j | \hat{\mathfrak{M}} | r \right\rangle.$$
(21)

Since  $\langle \ell |$  and  $|r \rangle$  are real, one has  $r_j = \ell_j^* \equiv \psi_j \sqrt{2}$  and  $\psi_{-j} = \psi_j^*$ ; hence Eq. (8), but with

$$L_{j} = \frac{i\rho_{j}}{2} \left( \dot{\psi}_{j} \psi_{j}^{*} - \psi_{j} \dot{\psi}_{j}^{*} \right) - \rho_{j} \nu_{j} |\psi_{j}|^{2}.$$
(22)

The resulting equations for individual modes are

$$\hat{D}_{j}\psi_{j} = \delta_{\psi_{j}^{*}}\tilde{\mathcal{L}}_{\text{int}}, \quad \hat{D}_{j} = \rho_{j}\left(\nu_{j} - id_{t}\right), \qquad (23)$$

similar to reduced Eqs. (9). Particularly, at zero  $\tilde{\mathcal{L}}_{int}$ ,

$$\psi_j = \sqrt{2\mathcal{J}_j} e^{-i\vartheta_j}, \quad \dot{\vartheta}_j = \nu_j, \quad \mathcal{J}_j = \text{const.}$$
 (24)

On the other hand,  $L_j = (\dot{\vartheta}_j - \nu_j)\mathcal{J}_j$ ; thus  $\partial_{\dot{\vartheta}_j}L_j = \mathcal{J}_j$ is also the action corresponding to the angle  $\vartheta_j$ :

$$\mathcal{J}_j = \rho_j |\psi_j|^2, \qquad (25)$$

so  $\nu_j$  is the canonical frequency. Then the mode energy is  $\nu_j \mathcal{J}_j$  (thus  $\rho_j > 0$  for stable modes with  $\nu_j > 0$ , henceforth implied), and Eq. (5) is recovered.

In the next sections, we apply Eqs. (23) to find eigenmodes for nonzero  $\tilde{\mathcal{L}}_{int}$ , with  $\psi$  becoming partial oscillations. Hence the effective potential modification  $\Phi$ [Eq. (6)] is obtained from Eq. (4).

## IV. PRIMARY RESONANCE

#### A. Linear oscillator

First, we calculate  $\Psi$  for a linear coupling between a pair of modes  $\psi_1$  and  $\psi_2$ , say,

$$\tilde{\mathcal{L}}_{\text{int}} = \sigma \psi_1 \psi_2^* + \sigma^* \psi_1^* \psi_2, \qquad (26)$$

where  $\sigma = \text{const.}$  In this case, Eqs. (23) yields

$$\hat{D}_1\psi_1 = \sigma^*\psi_2, \quad \hat{D}_2\psi_2 = \sigma\psi_1;$$
 (27)

hence a quadratic equation for the eigenfrequencies  $\varpi$ ,

$$\rho_1 \rho_2 (\varpi - \nu_1) (\varpi - \nu_2) = |\sigma|^2, \tag{28}$$

from which  $\Psi = \varpi_1 \mathcal{J}_1 + \varpi_2 \mathcal{J}_2$  is obtained.

As a particular case, consider interaction of a particle internal mode having frequency  $\Omega$  and action  $J = \rho |\psi|^2$ with an external oscillating field  $E = \bar{E}e^{-i\omega t}$  having frequency  $\omega$  and action  $I = \rho_E |E|^2$ . Given that the field occupies a volume  $V \to \infty$ , the frequency shifts  $\delta\Omega$  and  $\delta\omega$  are infinitesimal, Eq. (28) yielding

$$\delta\Omega\rho = \frac{|\sigma|^2}{(\Omega - \omega)\rho_E}, \quad \delta\omega\rho_E = \frac{|\sigma|^2}{(\omega - \Omega)\rho}.$$
 (29)

Since  $\rho_E \propto V$ , one has  $\delta \Omega J \ll \delta \omega I$ , whereas

$$\delta\omega I = \frac{|\sigma|^2}{(\omega - \Omega)\rho} \, |\bar{E}|^2 \tag{30}$$

is nonvanishing. Then one gets

$$\Psi = \Omega J + \Phi_0, \quad \Phi_0 = -\frac{1}{4} \alpha |\bar{E}|^2,$$
 (31)

where  $\Phi_0$  is the so-called ponderomotive potential (for the general expression see Appendix A), an insignificant constant  $\omega I$  is removed, and  $\alpha = 4|\sigma|^2[(\Omega-\omega)\rho]^{-1}$ . Since

$$\Phi_0 = \frac{\kappa_\alpha^2 |\bar{E}|^2}{\omega - \Omega},\tag{32}$$

where  $\kappa_{\alpha}^2 \equiv |\sigma|^2/\rho > 0$ , the effective potential becomes infinite at the linear resonance [Fig. 1(a)]. However, nonlinear effects remove this singularity, as we show below.

### B. Nonlinear oscillator

Consider the effective potential near a nonlinear resonance, with a Duffing oscillator as a model system. Then

$$\tilde{\mathcal{L}}_{\text{int}} = \sigma \psi E^* + \sigma^* \psi^* E + \frac{1}{2} \beta |\psi|^4, \qquad (33)$$

where  $\beta = \text{const}$ , yielding

$$\hat{D}\psi = \sigma^* E + \beta |\psi|^2 \psi, \quad \hat{D}_E E = \sigma \psi.$$
 (34)

Separate the driven motion from free oscillations,  $\psi = Xe^{-i(\omega+\delta\omega)t} + Ye^{-i(\Omega+\delta\Omega)t}$ , so

$$-\delta\omega\,\rho_E\bar{E} = \sigma X,\tag{35}$$

$$(\Omega - \omega)\rho X = \sigma^* \bar{E} + 2\beta |Y|^2 X + \beta |X|^2 X, \qquad (36)$$

$$-\delta\Omega\,\rho Y = \beta |Y|^2 Y + 2\beta |X|^2 Y. \tag{37}$$

From Eq. (37), it follows that  $\delta\Omega J \sim \beta |XY|^2$ , which we assume, for simplicity, small compared to  $\Phi_0 \sim \delta\omega I$ ; hence  $\Psi \approx \Omega J + \Phi$ , where  $\Phi = \int_0^I \delta\omega \, dI$  is the modified ponderomotive potential. The field frequency shift is found from the cubic equation  $(1 + \zeta h^2)h = 1$  for  $h = \delta\omega I/\Phi_0$ ;  $\zeta = \beta \Phi_0/(\gamma \rho)^2$ ,  $\Phi_0 = |\sigma \bar{E}|^2/(\gamma \rho)$ ,  $\gamma = \omega - \Omega + \zeta(J)$ ,  $\zeta = 2\beta J/\rho^2$ , and the above condition of negligible  $\delta\Omega J$  reads  $\zeta/(\omega - \Omega) \ll 1$ . Thus,

$$\Phi = \frac{\gamma^2 \rho^2}{\beta} W(\zeta), \quad W(\zeta) = \int_0^{\zeta} h(\tilde{\zeta}) \, d\tilde{\zeta}, \qquad (38)$$

where  $h(\zeta)$  has three branches [Fig. 1(b)]. One of those is unstable [29]; hence two branches of the ponderomotive potential,  $\Phi_1$  and  $\Phi_2$ . Their asymptotic form flows from  $W_{1,2}(\zeta \to 0\pm) \approx \zeta$  and  $W_1(\zeta \to 0-) \approx 2\sqrt{-\zeta}$  [Fig. 1(c)]:

$$\Phi_{1,2}^{\pm} \approx \Phi_0 = \frac{\kappa_{\alpha}^2 |\bar{E}|^2}{\gamma}, \quad \Phi_1^- \approx 2\kappa_{\alpha}\rho |\bar{E}| \frac{\sqrt{|\gamma\beta|}}{\beta}, \quad (39)$$

and Eq. (32) is recovered from  $\Phi_{1,2}^{\pm}$  in the limit of small  $\zeta$ and  $\zeta = 0$  [Fig. 1(d)]. On the other hand,  $W_1(\zeta \to \infty) \approx \frac{3}{2} |\zeta|^{2/3}$ , so  $\Phi_1 \to 3 |\sigma \bar{E}|^{4/3} |\beta|^{2/3}/(2\beta)$  at the resonance, i.e., the singularity vanishes.

#### V. BEAT RESONANCE

Suppose that  $\mathcal{L}_{int}$  also contains a term cubic in  $\psi$ , say,

$$\delta \tilde{\mathcal{L}}_{\text{int}} = \epsilon \psi_1 \psi_2^* \psi_3^* + \epsilon^* \psi_1^* \psi_2 \psi_3, \qquad (40)$$

where  $\epsilon = \text{const.}$  That would normally yield a potential small compared to  $\Phi_0$ , yet maybe except when

$$\nu_1 \approx \nu_2 + \nu_3. \tag{41}$$

Below we study the beat resonance (41) neglecting the quadratic  $\tilde{\mathcal{L}}_{int}$ , so the envelopes  $\bar{\psi}_j = \psi_j e^{i\nu_j t}$  satisfy

$$-i\rho_1 \dot{\bar{\psi}}_1 = \epsilon^* \bar{\psi}_2 \bar{\psi}_3 e^{-i\Delta t}, \qquad (42)$$

$$-i\rho_2\bar{\psi}_2 = \epsilon\bar{\psi}_1\bar{\psi}_3^*e^{i\Delta t},\tag{43}$$

$$-i\rho_3\bar{\psi}_3 = \epsilon\bar{\psi}_1\bar{\psi}_2^*e^{i\Delta t},\tag{44}$$

where  $\Delta \equiv \nu_3 + \nu_2 - \nu_1$  is the detuning frequency.

## A. Linear coupling

Suppose that  $\Delta$  is large enough, so Eqs. (42)-(44) can be solved by averaging. Then split  $\bar{\psi}_i$  into driven and



FIG. 1: (a) Resonant ponderomotive potential  $\Phi_0$  [Eq. (32)] in units  $\Phi_{\alpha} \equiv \kappa_{\alpha}^2 |\vec{E}|^2 / \omega$  vs.  $\omega - \Omega$  in units  $\omega$ . (b) Solid – squared normalized frequency shift  $h = \delta \omega I / \Phi_0$  vs.  $\zeta = \beta \Phi_0 / (\gamma \rho)^2$ ,  $\zeta (J = 0) \propto (\omega - \Omega)^{-3}$ ; dashed – approximations  $h^2 = 1$ and  $h^2 = -\zeta^{-1}$ . (c)  $W(\zeta)$  [Eq. (38)], consists of branches  $W_1 > 0$  and  $W_2 < 0$ . (d) Effective potential  $\Phi$  in units  $\Phi_c \equiv \rho^2 \omega_c^2 / \beta = \frac{2}{3} \Phi(0)$  vs.  $\omega - \Omega$  in units  $\omega_c \equiv \beta^{1/3} |\sigma \vec{E}|^{2/3} / \rho$ , assuming J = 0 and  $\beta > 0$ : solid – Eq. (38) (consists of branches  $\Phi_1 > 0$  and  $\Phi_2 < 0$ ), dashed – Eqs. (39).

free motion:  $\bar{\psi}_j = X_j e^{-i\eta_j t} + Y_j e^{-i\delta\nu_j t}$ , where  $\delta\nu_j \ll \eta_j$  are the nonlinear frequency shifts, and

$$\eta_1 = \delta \nu_2 + \delta \nu_3 + \Delta, \tag{45}$$

$$\eta_2 = \delta \nu_1 - \delta \nu_3 - \Delta, \tag{46}$$

$$\eta_3 = \delta \nu_1 - \delta \nu_2 - \Delta. \tag{47}$$

Thus the corresponding equations read

$$\begin{aligned} &-\eta_1 \rho_1 X_1 = \epsilon^* Y_2 Y_3, \quad -\delta \nu_1 \rho_1 Y_1 = \epsilon^* (Y_2 X_3 + X_2 Y_3) \,, \\ &-\eta_2 \rho_2 X_2 = \epsilon Y_1 Y_3^*, \quad -\delta \nu_2 \rho_2 Y_2 = \epsilon (Y_1 X_3^* + X_1 Y_3^*) \,, \\ &-\eta_3 \rho_3 X_3 = \epsilon Y_1 Y_2^*, \quad -\delta \nu_3 \rho_3 Y_3 = \epsilon (Y_1 X_2^* + X_1 Y_2^*) \,, \end{aligned}$$

where insignificant oscillating terms are neglected. Since  $\delta \nu_j \ll \Delta$ , one has

$$\delta\nu_1 = -\frac{\mathcal{J}_2 + \mathcal{J}_3}{\Gamma\Delta}, \quad \delta\nu_2 = \frac{\mathcal{J}_3 - \mathcal{J}_1}{\Gamma\Delta}, \quad \delta\nu_3 = \frac{\mathcal{J}_2 - \mathcal{J}_1}{\Gamma\Delta},$$

where  $\Gamma \equiv \rho_1 \rho_2 \rho_3 / |\epsilon|^2 > 0$ . Then, from Eqs. (4), one gets  $\Psi = \Psi_0 + \Phi$ ,

$$\Phi = \frac{\mathcal{J}_1 \mathcal{J}_2 + \mathcal{J}_1 \mathcal{J}_3 - \mathcal{J}_2 \mathcal{J}_3}{(\nu_1 - \nu_2 - \nu_3) \,\Gamma}.$$
(48)

Particularly, if the third mode corresponds to a macroscopic oscillating field tuned close to the beat resonance

$$\omega \approx \Omega_1 - \Omega_2, \tag{49}$$

then  $J_{1,2} \ll I \propto |\bar{E}|^2$ . Hence the "hybrid" ponderomotive potential that is obtained is simultaneously proportional to  $|\bar{E}|^2$  and the internal actions  $J_i$ :

$$\Phi = \frac{\kappa_{\epsilon}^2}{\Delta} \left( J_2 - J_1 \right) |\bar{E}|^2, \tag{50}$$

where  $\Delta = \omega - (\Omega_1 - \Omega_2)$ , and  $\kappa_{\epsilon}^2 \equiv |\epsilon|^2 / (\rho_1 \rho_2) > 0$ .

#### B. Nonlinear coupling

At  $\Delta \lesssim \delta \nu_j$ , the internal mode equations

$$-i\rho_1 \dot{\bar{\psi}}_1 = \epsilon^* \bar{\psi}_2 \bar{E} e^{-i\Delta t}, \quad -i\rho_2 \dot{\bar{\psi}}_2 = \epsilon \bar{\psi}_1 \bar{E}^* e^{i\Delta t} \quad (51)$$

do not allow averaging, so the above derivation is modified as follows. First, consider strictly periodic oscillations, in which case  $\bar{\psi} \equiv (\bar{\psi}_1, \bar{\psi}_2)^{\mathrm{T}}$  rewrites as  $\bar{\psi} = \hat{T}_{\Delta} \hat{C}_{\psi}^{-1} \mathbf{x}$ , where  $\hat{T}_y = \text{diag}(e^{-iyt/2}, e^{iyt/2})$ , and  $\hat{C}_{\psi} = \hat{T}_{-\phi} \text{diag}(\rho_1, \rho_2)^{1/2}$ ,  $\phi = \arg(\epsilon \bar{E}^*) + \pi$ . Then

$$i\dot{x}_1 = \frac{\varepsilon}{2} x_2 - \frac{\Delta}{2} x_1, \quad i\dot{x}_2 = \frac{\varepsilon}{2} x_1 + \frac{\Delta}{2} x_2,$$
 (52)

where  $\varepsilon = 2\kappa_{\epsilon}|\bar{E}|$ . Eqs. (52) yield two eigenmodes at frequencies  $\pm \frac{1}{2}\Lambda$ ,  $\Lambda = (\varepsilon^2 + \Delta^2)^{1/2}$ , so  $\mathbf{x} = \hat{U}\hat{T}_{\Lambda}\bar{\mathbf{x}}$ , where

$$\hat{U} = \begin{pmatrix} \cos\Theta & -\sin\Theta\\ \sin\Theta & \cos\Theta \end{pmatrix}, \tag{53}$$

 $\bar{\mathbf{x}} \equiv (\bar{x}_+, \bar{x}_-)^{\mathrm{T}}$  is constant, and  $\Theta$  satisfies

$$\cos 2\Theta = -\Delta/\Lambda, \quad \sin 2\Theta = \varepsilon/\Lambda.$$
 (54)

Then

$$\boldsymbol{\psi} = \hat{T}_{\omega} \hat{S} \boldsymbol{\chi}, \tag{55}$$

where  $\hat{S} \equiv \hat{C}_{\psi}^{-1} \hat{U} \hat{C}_{\chi}$  is a constant matrix,  $\hat{C}_{\chi} = \text{diag}(\rho_+, \rho_-)^{1/2}, \ \rho_{\pm} \equiv \varpi_{\pm}$  (for uniformity),

$$\varpi_{\pm} = \frac{1}{2} \left( \Omega_1 + \Omega_2 \right) \pm \frac{1}{2} \Lambda, \tag{56}$$

 $\chi_{\pm} = \bar{\chi}_{\pm} e^{-i\vartheta_{\pm}}, \, \dot{\vartheta}_{\pm} = \varpi_{\pm}, \, \text{and} \, \bar{\chi} = \hat{C}_{\chi}^{-1} \bar{\mathbf{x}}.$ When the oscillation parameters evolve, treat Eq. (55)

When the oscillation parameters evolve, treat Eq. (55) as a formal change of variables for Eq. (22). Hence  $\tilde{\mathcal{L}} = L_+ + L_-$ , the field part being omitted, and

$$L_{\pm} = \frac{i\rho_{\pm}}{2} \left( \dot{\chi}_{\pm} \chi_{\pm}^* - \dot{\chi}_{\pm}^* \chi_{\pm} \right) - \rho_{\pm} \nu_{\pm} |\chi_{\pm}|^2.$$
(57)

Therefore  $\chi_{\pm}$  are independent linear modes with frequencies  $\varpi_{\pm}$  and conserved actions  $J_{\pm} = \rho_{\pm} |\chi_{\pm}|^2$  (Sec. III), and Eq. (4) yields

$$\Psi = \frac{1}{2} \left( \Omega_1 + \Omega_2 \right) \left( J_+ + J_- \right) + \frac{\Lambda}{2} \left( J_+ - J_- \right).$$
 (58)

At  $\Delta \gg \varepsilon$ ,  $J_{\pm} = J_{1,2}$  for  $\Delta < 0$  and  $J_{\pm} = J_{2,1}$  for  $\Delta > 0$ , so Eq. (50) is recovered by Taylor expansion of Eq. (58).

## C. Quantum analogy

The above classical particle is the limit of a quantum system with plentiful states coupled to the field simultaneously and  $\Omega_j$  being the unperturbed transition frequencies [Fig. 2(a)]. Yet, with  $\rho_j \rightarrow \hbar$ , the equations (51) are also equivalent to those describing a *two*-level system [13, 14, 30], with the unperturbed eigenfrequencies  $\Omega_1$ and  $\Omega_2$  and the Rabi frequency  $\Omega_R = \varepsilon$  [Fig. 2(b)]. Hence Eq. (58) yields as well the dipole potential for a two-level quantum object, e.g., a cold atom [11, pp. 454-461], [12]:

$$\Psi = \frac{\hbar}{2} \left( \Omega_1 + \Omega_2 \right) + \frac{\hbar}{2} \left( n_+ - n_- \right) \sqrt{\Delta^2 + \Omega_{\rm R}^2}, \qquad (59)$$

where  $n_j = |\chi_j|^2$  are the occupation numbers  $(J_j \rightarrow \hbar n_j)$ , satisfying  $n_+ + n_- = 1$ . Similarly, Eq. (50) is equivalent to the dipole potential at weak coupling [30, 31]:

$$\Phi = \frac{\hbar \Omega_{\rm R}^2}{4\Delta} \left( 1 - 2n_1 \right). \tag{60}$$

[At  $n_1 = 0$ , Eq. (60) is further analogous to Eq. (32), with the resonance at the transition frequency  $\Omega_1 - \Omega_2$ .]

This parallelism originates from Eq. (23) yielding Schrödinger-like equations for  $\tilde{\mathcal{L}}_{int}$  quadratic in internal mode amplitudes. Therefore, for any classical particle governed by a hybrid potential, an analogue is possible which is governed by a quantum dipole potential, and vice versa. Hence the two types of objects permit uniform manipulation techniques, as also discussed in Sec. VI.

### VI. ONE-WAY WALLS

As an *effective* potential,  $\Psi$  can have properties distinguishing it from true potentials. Particularly, it can yield asymmetric barriers, or one-way walls [15, 16, 17, 18, 19, 20, 21, 22, 23], allowing current drive [32, 33, 34] and translational cooling [17, 22, 35, 36]. We explain these barriers as follows.

Suppose a ponderomotive potential of the form (32) [or, similarly, (38)]. Given  $\Omega = \Omega(z)$ , with, say,  $\partial_z \Omega < 0$ , the average force  $F_z = -\partial_z \Psi$  is everywhere in +z direction, except at the exact resonance where the effective potential does not apply [Fig. 3(a)]. Hence particles can be transmitted when traveling in one direction but reflected otherwise, even assuming uniform  $\bar{E}(z)$  [32, 33, 34]. In



FIG. 2: (a) Schematic of a quasiclassical particle containing two harmonic oscillators with unperturbed frequencies  $\Omega_1$ and  $\Omega_2$ . The energy spectrum for each individual oscillator at large energies (close-up on the right) is equidistant. The coupling with an oscillating field  $\bar{E}$  near the beat resonance (49) is described by Eqs. (51). (b) A two-level system, with unperturbed eigenfrequencies  $\Omega_1$  and  $\Omega_2$ , formally described by the same Eqs. (51), assuming the Rabi frequency  $\Omega_{\rm R} = \varepsilon$ .

Ref. [15], such dynamics was confirmed for cyclotronresonant rf fields, and, in Ref. [16], a similar scheme employing abrupt  $\overline{E}(z)$  was proposed.

Hybrid potentials (Sec. V) permit yet another type of one-way walls. Assume uniform  $\Omega_j$  and  $\Delta > 0$ ; then  $\Psi$  [Eqs. (50), (58)] is repulsive for cold particles ( $J_1 < J_2$ ) but attractive for hot particles ( $J_1 > J_2$ ). Thus, if particles incident, say, from the left are preheated (via nonadiabatic interaction with another field), they will be transmitted, whereas those cold as incident from the right will be repelled [Fig. 3(b)]; hence the asymmetry.

In agreement with the parallelism shown in Sec. V C, similar one-way walls for cold atoms have been suggested [17, 18, 19, 20, 21, 37] and enjoyed experimental verification [22, 23]. However, the new result here is that quasiclassical particles like Rydberg atoms and molecules can, in principle, be manipulated in the same manner, despite their involved eigenspectrum is different (Fig. 2).

#### VII. CONCLUSIONS

We propose a method to calculate the generalized effective potential  $\Psi$  for a nonrelativistic classical particle undergoing arbitrary oscillations in high-frequency or static fields. We derive  $\Psi$  from the oscillation eigenfrequencies in the particle-field system [Eq. (4)], like in the dressed-atom approach [11, 12, 13, 14] but from nonlinear classical equations [Eqs. (23)]. Specifically, we show that the ponderomotive potential [Eqs. (32), (A6); Fig. 1(a)] extended to a nonlinear oscillator has multiple branches near the primary resonance [Eq. (38); Fig. 1(d)]. Also, for a pair of natural frequencies in a beat resonance,  $\Psi$ scales linearly with the internal actions [Eqs. (50), (58)]and is analogous to the dipole potential [Eqs. (59), (60)] for a two-level quantum system (Fig. 2). Thus cold quantum particles and highly-excited quasiclassical objects permit uniform manipulation tools, particularly, station $\Delta \ > 0$ 

FIG. 3: Schematic showing two types of one-way walls (arrows denote particle transmission and reflection). (a) The one-way wall is due to a ponderomotive potential near a primary resonance [Eqs. (31), (32)] with uniform  $\partial_z \Omega < 0$ , so  $F_z = -\partial_z \Psi > 0$  for all z, except at the exact resonance where the effective potential does not apply. [The inclined asymptote corresponds to  $\Psi = \Omega J + \text{const.}$ ] (b) The one-way wall is due to a hybrid potential [Eqs. (50), (58)] with uniform  $\Omega_j$  and  $\Delta > 0$ : particles incident from the left receive  $J_1 > J_2$ , so they see an attractive  $\Psi$ ; those incident from the right have  $J_1 < J_2$ , so they see a repulsive  $\Psi$ .

 $F_{\tau} > 0$ 

ary asymmetric barriers, or one-way walls (Fig. 3).

## VIII. ACKNOWLEDGMENTS

This work was supported by DOE Contracts No. DE-FG02-06ER54851, DE-FG02-05ER54838, and by the NNSA under the SSAA Program through DOE Research Grant No. DE-FG52-04NA00139.

## APPENDIX A: GENERAL EXPRESSION FOR THE PONDEROMOTIVE POTENTIAL

Consider generalization of the ponderomotive potential (31), (32) to multiple internal oscillations and vector field

$$\bar{\mathbf{E}} = \sum_{\mu} \bar{\mathbf{E}}_{\mu}, \quad \bar{\mathbf{E}}_{\mu} = \mathbf{e}_{\mu} \bar{E}_{\mu}, \quad (A1)$$

composed of modes  $\mu$  with polarizations  $\mathbf{e}_{\mu}$ . From Sec. IV A, it is known that the internal energy  $\delta \mathbf{\Omega} \cdot \mathbf{J}$ yields a negligible contribution to  $\Psi$ . Thus

$$\Phi_0 = \sum_{\mu} \delta \omega_{\mu} I_{\mu}, \qquad (A2)$$

where the infinitesimal frequency shifts  $\delta \omega_{\mu}$  of the field are found as follows. Use [3]

$$\tilde{\mathcal{L}}_{int} = \frac{1}{2} \operatorname{Re} \left[ \bar{\mathbf{E}}^* \cdot \bar{\mathbf{d}} \right], \tag{A3}$$

where insignificant oscillating terms are removed, and  $\bar{\mathbf{d}}$  is the particle induced dipole moment. From Eq. (23),

$$-\delta\omega_{\mu}\rho_{\mu}\bar{E}_{\mu} = \frac{1}{4}\,\mathbf{e}_{\mu}^{*}\cdot\bar{\mathbf{d}}.\tag{A4}$$

Multiply Eq. (A4) by  $\bar{E}^*_{\mu}$  and substitute  $\bar{\mathbf{d}} = \hat{\alpha} \bar{\mathbf{E}}$ , where  $\hat{\alpha}$  is the polarizability tensor; then

$$\delta\omega_{\mu}I_{\mu} = -\frac{1}{4}\,\bar{\mathbf{E}}_{\mu}^{*}\cdot\hat{\alpha}\bar{\mathbf{E}}.\tag{A5}$$

Hence summation over all field modes yields the known expression [38, 39]

$$\Phi_0 = -\frac{1}{4} \, (\bar{\mathbf{E}}^* \cdot \hat{\alpha} \bar{\mathbf{E}}), \tag{A6}$$

which holds for any internal modes contributing to  $\hat{\alpha}$  [3].

- [1] R. L. Dewar, Phys. Fluids 16, 1102 (1973).
- [2] P. L. Similon, A. N. Kaufman, and D. D. Holm, Phys. Fluids 29, 1908 (1986).
- [3] I. Y. Dodin and N. J. Fisch, Phys. Lett. A 349, 356 (2006).
- [4] I. Y. Dodin and N. J. Fisch, Phys. Rev. E 77, 036402 (2008).
- [5] I. Y. Dodin and N. J. Fisch, in Frontiers in Modern Plasma Physics (Trieste, Italy, 2008), also AIP Proc. 1061 (AIP, New York, 2008), p. 263.
- [6] A. V. Gaponov and M. A. Miller, Zh. Eksp. Teor. Fiz. 34, 242 (1958) [Sov. Phys. JETP 7, 168 (1958)].
- [7] H. Motz and C. J. H. Watson, Adv. Electron. 23, 153

(1967).

- [8] J. R. Cary and A. N. Kaufman, Phys. Rev. Lett. 39, 402 (1977).
- [9] T. Hatori and H. Washimi, Phys. Rev. Lett. 46, 240 (1981).
- [10] J. D. Jackson, *Classical electrodynamics* (Wiley, New York, 1975), Sec. 12.10.
- [11] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, Atom-photon interactions (Wiley, New York, 1992), Chap. VI.
- [12] J. Dalibard and C. Cohen-Tannoudji, J. Opt. Soc. Am. B 2, 1707 (1985).
- [13] P. R. Berman and R Salomaa, Phys. Rev. A 25, 2667

(1982).

- [14] E. Courtens and A. Szöke, Phys. Rev. A 15, 1588 (1977).
- [15] I. Y. Dodin, N. J. Fisch, and J. M. Rax, Phys. Plasmas 11, 5046 (2004).
- [16] I. Y. Dodin and N. J. Fisch, Phys. Rev. E 72, 046602 (2005).
- [17] M. G. Raizen, A. M. Dudarev, Q. Niu, and N. J. Fisch, Phys. Rev. Lett. 94, 053003 (2005).
- [18] A. Ruschhaupt and J. G. Muga, Phys. Rev. A 70, 061604(R) (2004).
- [19] A. Ruschhaupt and J. G. Muga, Phys. Rev. A 73, 013608 (2006).
- [20] A. Ruschhaupt, J. G. Muga, and M. G. Raizen, J. Phys.
   B: At. Mol. Opt. Phys. **39**, L133 (2006).
- [21] G. N. Price, S. T. Bannerman, E. Narevicius, and M. G. Raizen, Laser Phys. 17, 965 (2007).
- [22] G. N. Price, S. T. Bannerman, K. Viering, E. Narevicius, and M. G. Raizen, Phys. Rev. Lett. **100**, 093004 (2008).
- [23] J. J. Thorn, E. A. Schoene, T. Li, and D. A. Steck, Phys. Rev. Lett. **100**, 240407 (2008).
- [24] Slow dependence on time is also allowed here and further.
- [25] In addition to electromagnetic modes [40], any periodic forces can be included here, assuming that an extended phase space is used [41].
- [26] Sec. III corrects Appendix B of our Ref. [3].
- [27] H. Goldstein, *Classical mechanics* (Addison-Wesley, Reading, MA, 1950), Chap. 10.
- [28] A. K. Schenk, P. Arras, E. E. Flanagan, S. A. Teukolsky,

and I. Wasserman, Phys. Rev. D 65, 024001 (2001).

- [29] N. N. Bogoliubov and Y. A. Mitropolskii, Asymptotic methods in the theory of nonlinear oscillations (Gordon and Breach, New York, 1961), Sec. 15.
- [30] M. O. Scully and M. S. Zubairy, *Quantum optics* (Cambridge Univ. Press, Cambridge, 1997), Secs. 5.2, 17.1.4.
- [31] A. Ashkin, Phys. Rev. Lett. 40, 729 (1978).
- [32] E. V. Suvorov and M. D. Tokman, Fiz. Plazmy 14, 950 (1988) [Sov. J. Plasma Phys. 14, 557 (1988)].
- [33] A. G. Litvak, A. M. Sergeev, E. V. Suvorov, M. D. Tokman, and I. V. Khazanov, Phys. Fluids B 5, 4347 (1993).
- [34] N. J. Fisch, J. M. Rax, and I. Y. Dodin, Phys. Rev. Lett. **91**, 205004 (2003), Phys. Rev. Lett. **93**, 059902(E) (2004).
- [35] A. M. Dudarev, M. Marder, Q. Niu, N. J. Fisch, and M. G. Raizen, Europhys. Lett. **70**, 761 (2005).
- [36] A. Ruschhaupt, J. G. Muga, and M. G. Raizen, J. Phys.
   B: At. Mol. Opt. Phys. **39**, 3833 (2006).
- [37] E. Narevicius and M. G. Raizen, arXiv:0808.1383v1 (2008).
- [38] D. Auerbach, E. E. A. Bromberg, and L. Wharton, J. Chem. Phys. 45, 2160 (1966).
- [39] J. P. Gordon, Phys. Rev. A 8, 14 (1973).
- [40] G. Flores-Hidalgo and A. P. C. Malbouisson, Phys. Lett. A 337, 37 (2005).
- [41] I. Y. Dodin and N. J. Fisch, Phys. Lett. A 372, 6112 (2008).