

---

# Variation and Rough Path Properties of Local Times of Lévy Processes

Chunrong Feng<sup>1</sup>, Huaizhong Zhao<sup>2</sup>

<sup>1</sup> Department of Mathematics, Shanghai Jiao Tong University, Shanghai, 200240, China

<sup>2</sup> Department of Mathematical Sciences, Loughborough University, LE11 3TU, UK  
fcr@sjtu.edu.cn, H.Zhao@lboro.ac.uk

**Summary.** In this paper, we will prove that the local time of a Lévy process is of finite  $p$ -variation in the space variable in the classical sense, a.s. for any  $p > 2$ ,  $t \geq 0$ , and is a rough path of roughness  $p$  a.s. for any  $2 < p < 3$ . Then for any function  $g$  of finite  $q$ -variation ( $1 \leq q < 3$ ), we establish the integral  $\int_{-\infty}^{\infty} g(x) dL_t^x$  as a Young integral when  $1 \leq q < 2$  and a Lyons' rough path integral when  $2 \leq q < 3$ . We therefore apply these path integrals to extend the Tanaka-Meyer formula for a continuous function  $f$  if  $\nabla^- f$  exists and is of finite  $q$ -variation when  $1 \leq q < 3$ , for both continuous semi-martingales and a class of Lévy processes.

**Keywords:** semimartingale local time; geometric rough path; finite  $p$ -variation; Young integral; rough path integral; Lévy processes.

## 1 Introduction

The variation of a stochastic process is a classical problem of fundamental importance in probability theory. There are two kinds of variations, namely in the sense of probability and in the classical sense. The quadratic variation of a Brownian motion (a martingale) in the sense of probability made it possible to define Itô's stochastic integral of a square integrable progressive process with respect to the Brownian motion (the martingale) (Lévy [20], Itô [16], Kunita and Watanabe [18]). The classical  $p$ -variation of the Brownian motion and its Lévy area,  $p > 2$ , led to Lyons' pathwise approach to stochastic differential equations (Lyons [21], [22]). Local time is an important and useful stochastic process. The investigation of its variation and integration has attracted attentions of many mathematicians. Similar to the case of the Brownian motion, the variation of the local time of a semimartingale in the spatial variable is also fundamental in the construction of an integral with respect to the local time. There have been many works on the quadratic or  $p$ -variations (in the case of stable processes) of local times in the sense of probability. Bouleau and Yor ([3]), Perkins ([28]), first proved that, for the Brownian local time, and a sequence of partitions  $\{D_n\}$  of an interval  $[a, b]$ , with the mesh  $|D_n| \rightarrow 0$  when  $n \rightarrow \infty$ ,  $\lim_{n \rightarrow \infty} \sum_{D_n} (L_t^{x_{i+1}} - L_t^{x_i})^2 = 4 \int_a^b L_t^x dx$  in probability. Subsequently, the process  $x \rightarrow L_t^x$  can be regarded as a semimartingale (with appropriate filtration). This result allowed one to construct various stochastic integrals of the Brownian local time in the spatial variable. See also Rogers and Walsh [30]. Numerous important extensions on the variations, stochastic integrations of local times and Itô's formula have been done, e.g. Marcus and Rosen [24], [25], Eisenbaum [5], [6], Eisenbaum and Kyprianou [7], Flandoli, Rosso and Wolf [11], Föllmer, Protter and Shiryaev [12], Föllmer and Protter [13], Moret and Nualart

[27]. In their extensions of Itô's formula, the integrals of the local time are given as stochastic integrals in nature, for example as forward and backward stochastic integrals. What had been missing here in the literature mentioned above was the variation of the local time in the classical sense and whether or not the local time is a rough path (its meaning will be made precise later). In Feng and Zhao [9], we proved that for a continuous semi-martingale, its local time  $L_t^x$  is of finite  $p$ -variation in the classical sense in  $x$  for any  $t \geq 0$ , a.s. for any  $p > 2$ , i.e.

$$\sup_D \sum_{i=0}^{n-1} |L_t^{x_{i+1}} - L_t^{x_i}|^p < \infty, \quad (1)$$

where the supremum is taken over all finite partition  $D_{(-\infty, +\infty)} = \{-\infty < x_0 < x_1 < x_2 < \dots < x_n < \infty\}$ . This allowed us to define the path integral  $\int_{-\infty}^{\infty} g(x) d_x L_t^x$  as a Young integral, for any  $g$  being of a finite  $q$ -variation for a number  $q \in [1, 2)$ . The main purpose of this paper is to solve the rough path part of the problem, for the local times of both continuous semi-martingales and a class of Lévy processes.

As a first step, we consider the classical  $p$ -variation for the Lévy process which is represented by the following Lévy-Itô decomposition

$$X_t = X_0 + \sigma B_t + bt + \int_0^{t+} \int_{R \setminus \{0\}} y 1_{\{|y| \geq 1\}} N_p(dsdy) + \int_0^{t+} \int_{R \setminus \{0\}} y 1_{\{|y| < 1\}} \tilde{N}_p(dsdy). \quad (2)$$

This is a non-trivial problem as the jumps, especially the small jumps, create a lot of difficulties in estimating the increment  $L_t^{x_{i+1}} - L_t^{x_i}$ . Recall that for a general semimartingale  $X_t$ ,  $L = \{L_t^x; x \in R\}$  is defined from the following formula (Meyer [26]):

$$\int_0^t g(X_s) d \langle X^c \rangle_s = \int_{-\infty}^{\infty} g(x) L_t^x dx, \quad (3)$$

where  $X^c$  is the continuous part of  $X$ . There is a different notion of local time defined as the Radon-Nikodym derivative of the occupation measure of  $X$  with respect to the Lebesgue measure on  $R$  i.e.

$$\int_0^t g(X_s) ds = \int_{-\infty}^{\infty} g(x) \gamma_t^x dx, \quad (4)$$

for every Borel function  $g : R \rightarrow R^+$ . For the Lévy process (2), if  $\sigma \neq 0$ ,  $L_t^x$  and  $\gamma_t^x$  are the same (up to a multiple of a constant). In case  $\sigma = 0$  e.g. for a stable process, there is no diffusion part so these two definitions are different. In fact, in this case  $L_t^x = 0$ . The increment of  $\gamma_t^x$  for stable processes was considered by Boylan [4], Gettoor and Kesten [14] and Barlow [2], using potential theory approach, in order to establish the continuity of the local time in the space variable. The first main task of this paper is to prove that when  $\sigma \neq 0$ , and (12) is satisfied, for any  $p > 2$  and  $t \geq 0$ , the process  $x \mapsto L_t^x$ , is of finite  $p$ -variation in the classical sense almost surely. Both our result and our method are new in literature. Here the Tanaka formula is directly used in our approach. As a direct application, one can define the path integral  $\int_{-\infty}^{+\infty} g(x) dL_t^x$  as a Young integral for any  $g$  being of bounded  $q$ -variation for a  $q \in [1, 2)$ . It is noted when  $q \geq 2$ , Young's condition  $\frac{1}{p} + \frac{1}{q} > 1$  is broken.

The main task of this paper is to construct a geometric rough path over the processes  $Z(x) = (L_t^x, g(x))$ , for any deterministic function  $g$  being of finite  $q$ -variation when  $q \in [2, 3)$ . This implies establishing the path integrals  $\int_{-\infty}^{\infty} L_t^x d_x L_t^x$  and  $\int_{-\infty}^{\infty} g(x) d_x L_t^x$ . For these two integrals, all the classical integration theories of Riemann, Lebesgue and Young fail to work. To overcome the difficulty, we use

the rough path theory pioneered by Lyons, see [21], [22], [23], also [19]. However, our  $p$ -variation result of the local time does not automatically make the desired rough path exist or the integral well defined, though it is a crucial step to study first. Actually further hard analysis is needed to establish an iterated path integration theory for  $Z$ . First we introduce a piecewise curve of bounded variation as a generalized Wong-Zakai approximation to the stochastic process  $Z$ . Then we define a smooth rough path by defining the iterated integrals of the piecewise bounded variation process. We need to prove the smooth rough path converges to a geometric rough path  $\mathbf{Z} = (1, \mathbf{Z}^1, \mathbf{Z}^2)$  when  $1 \leq q < 3$ . For this, an important step is to compute  $E(L_t^{x_{i+1}} - L_t^{x_i})(L_t^{x_{j+1}} - L_t^{x_j})$ , and obtain the correct order in terms of the increments  $x_{i+1} - x_i$  and  $x_{j+1} - x_j$ , especially in disjoint intervals  $[x_i, x_{i+1}]$  and  $[x_j, x_{j+1}]$  when  $i \neq j$ . Using this key estimate, we can establish the geometric rough path  $\mathbf{Z} = (1, \mathbf{Z}^1, \mathbf{Z}^2)$ . Then from Chen's identity, we define the following two integrals

$$\int_a^b L_t^x dL_t^x = \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{1,1} + L_t(x_i)(L_t^{x_{i+1}} - L_t^{x_i})) \quad (5)$$

and

$$\int_a^b g(x) dL_t^x = \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1} + g(x_i)(L_t^{x_{i+1}} - L_t^{x_i})). \quad (6)$$

Note that the Riemann sum  $\sum_{i=0}^{r-1} L_t^{x_i}(L_t^{x_{i+1}} - L_t^{x_i})$  and  $\sum_{i=0}^{r-1} g(x_i)(L_t^{x_{i+1}} - L_t^{x_i})$  themselves may not have limits as the mesh  $m(D_{[a,b]}) \rightarrow 0$ . At least there are no integration theories, rather than Lyons' rough path theory, to guarantee the convergence of the Riemann sums for almost all  $\omega$ . Here it is essential to add Lévy areas to the Riemann sum. Furthermore, we can prove if a sequence of smooth functions  $g_j \rightarrow g$  as  $j \rightarrow \infty$ , then the Riemann integral  $\int_a^b g_j(x) dL_t^x$  converges to the rough path integral  $\int_a^b g(x) dL_t^x$  defined in (6).

Having established the path integration of local time and the corresponding convergence results, as a simple application, we can easily prove a useful extension of Itô's formula for the Lévy process when the function is less smooth: if  $f : R \rightarrow R$  is an absolutely continuous function and has left derivative  $\nabla^- f(x)$  being left continuous and of bounded  $q$ -variation, where  $1 \leq q < 3$ , then P-a.s.

$$\begin{aligned} f(X_t) &= f(X_0) + \int_0^t \nabla^- f(X_s) dX_s - \int_{-\infty}^{\infty} \nabla^- f(x) d_x L_t^x \\ &\quad + \sum_{0 \leq s \leq t} [f(X_s) - f(X_{s-}) - \Delta X_s \nabla^- f(X_{s-})], \quad 0 \leq t < \infty. \end{aligned} \quad (7)$$

Here the path integral  $\int_{-\infty}^{\infty} \nabla^- f(x) d_x L_t^x$  is a Lebesgue-Stieltjes integral when  $q = 1$ , a Young integral when  $1 < q < 2$  and a Lyons' rough path integral when  $2 \leq q < 3$  respectively. Needless to say that Tanaka's formula ([31]) and Meyer's formula ([26], [32]) are very special cases of our formula when  $q = 1$ . The investigation of Itô's formula to less smooth functions is crucial and useful in many problems e.g. studying partial differential equations with singularities, the free boundary problem in American options, and certain stochastic differential equations. Time dependent cases for a continuous semimartingale  $X_t$  were investigated recently by Elworthy, Truman and Zhao ([8]), Feng and Zhao ([9]), where two-parameter Lebesgue-Stieltjes integrals and two-parameter Young integral were used respectively. We would like to point out that a two-parameter rough path integration theory, which is important to the study of local times, and some other problems such as SPDEs, still remains open.

A part of the results about the rough path integral of local time for a continuous semimartingale was announced in Feng and Zhao [10]. In summary, we have obtained complete results of the variation and rough path of roughness  $p$  for any  $p \in (2, 3)$  of local times for any continuous semi-martingales and a class of Lévy processes satisfying (12) and  $\sigma \neq 0$ . We don't include the proof of the rough path result for continuous semimartingales in this paper as we believe the reader can draw a proof from the one that we present in this paper for the Lévy processes easily.

## 2 The $p$ -variation of the local time of Lévy processes

Let  $X_t$  be a one dimensional time homogeneous Lévy process, and  $(\mathcal{F}_t)_{t \geq 0}$  be generated by the sample paths  $X_t$ ,  $p(\cdot)$  be a stationary  $(\mathcal{F}_t)$ -Poisson process on  $R \setminus \{0\}$ . From the well-known Lévy-Itô decomposition theorem, we can write  $X_t$  as follows:

$$X_t := X_0 + \sigma B_t + V_t + \tilde{M}_t, \quad (8)$$

where

$$V_t := bt + \int_0^{t+} \int_{R \setminus \{0\}} y 1_{\{|y| \geq 1\}} N_p(dsdy),$$

$$\tilde{M}_t := \int_0^{t+} \int_{R \setminus \{0\}} y 1_{\{|y| < 1\}} \tilde{N}_p(dsdy).$$

Here,  $N_p$  is the Poisson random measure of  $p$ , the compensator of  $p$  is of the form  $\hat{N}(dsdy) = ds n(dy)$ , where  $n(dy)$  is the Lévy measure of process  $X$ . The compensated random measure  $\tilde{N}_p(t, U) = N_p(t, U) - \hat{N}_p(t, U)$  is an  $(\mathcal{F}_t)$ -martingale. Before we give our main theorem of this section, we first give a  $p$ -moment estimate formula. This will be used in later proofs.

**Lemma 1** *Assume  $f$  is  $(\mathcal{F}_t)$ -predictable, and satisfies*

$$E \left( \int_0^t \int_R |f(s, y, \omega)|^2 n(dy) ds \right) < \infty.$$

*Then we have the  $p$ -moment estimate formula: for any  $p \geq 1$*

$$E \left( \sum_{0 \leq s \leq t} |f(s, p_s(\omega), \omega)| \right)^p$$

$$\leq c_p E \left( \int_0^t \int_R |f(s, y, \omega)|^2 n(dy) ds \right)^{\frac{p}{2}} + p E \left( \int_0^t \int_R |f(s, y, \omega)| n(dy) ds \right)^p, \quad (9)$$

for constant  $c_p > 0$ .

**Proof:** From the definition of  $N_p$ ,  $\tilde{N}_p$ , the Burkholder-Davis-Gundy inequality, we can have the  $p$ -moment estimation:

$$E \left( \sum_{0 \leq s \leq t} |f(s, p_s(\omega), \omega)| \right)^p$$

$$= E \left( \int_0^{t+} \int_R |f(s, y, \omega)| N_p(dsdy) \right)^p$$

$$\begin{aligned}
&= E \left( \int_0^{t+} \int_R |f(s, y, \omega)| (N_p(dsdy) - n(dy)ds) + \int_0^t \int_R |f(s, y, \omega)| n(dy)ds \right)^p \\
&\leq pE \left( \int_0^{t+} \int_R |f(s, y, \omega)| \tilde{N}_p(dsdy) \right)^p + pE \left( \int_0^t \int_R |f(s, y, \omega)| n(dy)ds \right)^p \\
&\leq c_p E \left( \int_0^t \int_R |f(s, y, \omega)|^2 n(dy)ds \right)^{\frac{p}{2}} + pE \left( \int_0^t \int_R |f(s, y, \omega)| n(dy)ds \right)^p. \quad \diamond
\end{aligned}$$

Recall the Tanaka formula for the Lévy process  $X_t$  ([1]), we have

$$\begin{aligned}
L_t^a &= (X_t - a)^+ - (X_0 - a)^+ - \int_0^t \mathbf{1}_{\{X_{s-} > a\}} dX_s \\
&\quad + \sum_{0 \leq s \leq t} [(X_{s-} - a)^+ - (X_s - a)^+ + \mathbf{1}_{\{X_{s-} > a\}} \Delta X_s].
\end{aligned}$$

Since

$$\sum_{0 \leq s \leq t} \mathbf{1}_{\{X_{s-} > a\}} \Delta X_s \mathbf{1}_{\{|\Delta X_s| \geq 1\}} = \int_0^{t+} \int_{R \setminus \{0\}} \mathbf{1}_{\{X_{s-} > a\}} y \mathbf{1}_{\{|y| \geq 1\}} N_p(dsdy),$$

it turns out from the cancellation due to above that we have

$$L_t^a = \varphi_t(a) - bI_t(a) - \sigma \hat{B}_t^a - \hat{M}_t^a + K_1(t, a) + K_2(t, a), \quad (10)$$

where

$$\begin{aligned}
\varphi_t(a) &:= (X_t - a)^+ - (X_0 - a)^+, \\
I_t(a) &:= \int_0^t \mathbf{1}_{\{X_{s-} > a\}} ds, \\
\hat{B}_t^a &:= \int_0^t \mathbf{1}_{\{X_{s-} > a\}} dB_s, \\
\hat{M}_t^a &:= \int_0^t \mathbf{1}_{\{X_{s-} > a\}} d\tilde{M}_s, \\
K_1(t, a) &:= \sum_{0 \leq s \leq t} [(X_{s-} - a)^+ - (X_s - a)^+] \mathbf{1}_{\{|\Delta X_s| \geq 1\}}, \\
K_2(t, a) &:= \sum_{0 \leq s \leq t} [(X_{s-} - a)^+ - (X_s - a)^+ + \mathbf{1}_{\{X_{s-} > a\}} \Delta X_s] \mathbf{1}_{\{|\Delta X_s| < 1\}}.
\end{aligned}$$

Denote  $K(t, a) := K_1(t, a) + K_2(t, a)$ . For the convenience in what follows in later part, we denote

$$J_1(s, a) := (X_{s-} - a)^+ - (X_s - a)^+, \quad J_2(s, a) := \mathbf{1}_{\{X_{s-} > a\}} \Delta X_s.$$

Note we have the following important decomposition that we will use often: for any  $a_i < a_{i+1}$ ,

$$\begin{aligned}
&[J_1(s, a_{i+1}) - J_1(s, a_i)] + [J_2(s, a_{i+1}) - J_2(s, a_i)] \\
&= -(X_s - a_i) \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} - (a_{i+1} - a_i) \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{X_{s-} > a_{i+1}\}} \\
&\quad + (X_s - a_i) \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_{s-} \leq a_i\}} - (a_{i+1} - X_s) \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_{s-} > a_{i+1}\}} \\
&\quad + (a_{i+1} - a_i) \mathbf{1}_{\{X_s > a_{i+1}\}} \mathbf{1}_{\{X_{s-} \leq a_i\}} + (a_{i+1} - X_s) \mathbf{1}_{\{X_s > a_{i+1}\}} \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}}. \quad (11)
\end{aligned}$$

In the following, we assume

$$\int_{\mathbb{R} \setminus \{0\}} (|y| \wedge 1) n(dy) < \infty. \quad (12)$$

**Lemma 2** *Assume (12) is satisfied, then for any  $p \geq 2$ ,  $\sup_a E(L_t^a)^{\frac{p}{2}} < \infty$ ,  $\sup_a E(L_t^a)^p < \infty$ .*

**Proof:** Now from (10) and using the Burkholder-Davis-Gundy inequality again, we have

$$\begin{aligned} E(L_t^a)^{\frac{p}{2}} &\leq cE[|X_t - X_0|^{\frac{p}{2}} + t^{\frac{p}{2}} + \langle B \rangle_t^{\frac{p}{4}} + \langle \tilde{M} \rangle_t^{\frac{p}{4}} + |K(t, a)|^{\frac{p}{2}}] \\ &\leq cE \langle \tilde{M} \rangle_t^{\frac{p}{4}} + ct^{\frac{p}{2}} + t^{\frac{p}{4}} + cE \langle \tilde{M} \rangle_t^{\frac{p}{4}} + cE|K(t, a)|^{\frac{p}{2}}. \end{aligned}$$

Here  $c$  is a generic constant depending on  $p$ ,  $\sigma$  and  $b$ . First note

$$E \langle \tilde{M} \rangle_t^{\frac{p}{4}} = E \left[ \int_0^t \int_{\mathbb{R} \setminus \{0\}} y^2 \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right]^{\frac{p}{4}} < \infty.$$

To estimate the last term, first

$$\begin{aligned} E|K(t, a)|^{\frac{p}{2}} &\leq \frac{p}{2} E \left[ \sum_{0 \leq s \leq t} |J_1(s, a)| \mathbf{1}_{\{|\Delta X_s| \geq 1\}} \right]^{\frac{p}{2}} + \frac{p}{2} E \left[ \sum_{0 \leq s \leq t} |J_1(s, a) + J_2(s, a)| \mathbf{1}_{\{|\Delta X_s| < 1\}} \right]^{\frac{p}{2}} \\ &:= A + B. \end{aligned}$$

We need to prove  $A < \infty$  and  $B < \infty$ . The result  $A < \infty$  can be seen from

$$E \sum_{0 \leq s \leq t} |J_1(s, a)| \mathbf{1}_{\{|\Delta X_s| \geq 1\}} \leq E \sum_{0 \leq s \leq t} |\Delta X_s| \mathbf{1}_{\{|\Delta X_s| \geq 1\}} < \infty.$$

About  $B$ , from the  $p$ -moment estimate, we have

$$\begin{aligned} B &= \frac{p}{2} E \left( \int_0^{t^+} \int_{-\infty}^{\infty} |(X_s - a - y)^+ - (X_s - a)^+ + \mathbf{1}_{\{X_s - y > a\}} y| \mathbf{1}_{\{|y| < 1\}} N_p(dy ds) \right)^{\frac{p}{2}} \\ &\leq c(t, p) E \left( \int_0^t \int_{-\infty}^{\infty} |(X_s - a - y)^+ - (X_s - a)^+ + \mathbf{1}_{\{X_s - y > a\}} y|^2 \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{4}} \\ &\quad + c(t, p) E \left( \int_0^t \int_{-\infty}^{\infty} |(X_s - a - y)^+ - (X_s - a)^+ + \mathbf{1}_{\{X_s - y > a\}} y| \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{2}} \\ &= c(t, p) E \left( \int_0^t \int_{-\infty}^{\infty} |-(X_s - a)^- \mathbf{1}_{\{X_s - y > a\}} - (X_s - a)^+ \mathbf{1}_{\{X_s - y \leq a\}}|^2 \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{4}} \\ &\quad + c(t, p) E \left( \int_0^t \int_{-\infty}^{\infty} |-(X_s - a)^- \mathbf{1}_{\{X_s - y > a\}} - (X_s - a)^+ \mathbf{1}_{\{X_s - y \leq a\}}| \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{2}} \\ &\leq c(t, p) E \left( \int_0^t \int_{-\infty}^{\infty} |y|^2 \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{4}} + c(t, p) E \left( \int_0^t \int_{-\infty}^{\infty} |y| \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{2}} \\ &< \infty. \end{aligned}$$

Here  $c(t, p)$  is a generic constant depending on  $t$  and  $p$ . So we can get  $\sup_a E|K(t, a)|^{\frac{p}{2}} < \infty$  and therefore  $\sup_a E(L_t^a)^{\frac{p}{2}} < \infty$ . The assertion  $\sup_a E(L_t^a)^p < \infty$  can be proved similarly.  $\diamond$

The following theorem on the  $p$ -variation of local time in the spatial variable is the main result of this section. We need to consider the variation of each term in (10). As the terms related to the continuous part of the Lévy process were considered in [9], so the main difficulty is from the jumps related to Lévy process, especially small jumps. The main idea to deal with the jump part is to use the  $p$ -moment estimate formula for the jump part and change it to integration of  $X_s$ . Then we can use occupation times formula, Jensen's inequality and Fubini theorem to obtain the  $p$ -moment of the increment from  $a_i$  to  $a_{i+1}$ . The decomposition (11) of  $J_1(s, a_{i+1}) - J_1(s, a_i) + J_2(s, a_{i+1}) - J_2(s, a_i)$  plays an important role in our proof of the desired increment estimates. With the increment estimate, the  $p$ -variation can be proved by the classical approach of Lévy.

**Theorem 3** *If  $\sigma \neq 0$  and the Lévy measure  $n(dy)$  satisfies (12), then the local time  $L_t^a$  of time homogeneous Lévy process  $X_t$  given by (2) is of bounded  $p$ -variation in  $a$  for any  $t \geq 0$ , for any  $p > 2$ , almost surely, i.e.*

$$\sup_{D(-\infty, \infty)} \sum_i |L_t^{a_{i+1}} - L_t^{a_i}|^p < \infty \quad a.s.,$$

where the supremum is taken over all finite partition on  $R$ ,  $D(-\infty, \infty) := \{-\infty < a_0 < a_1 < \dots < a_n < \infty\}$ .

**Proof:** By the usual localization argument, we may first assume that there is a constant  $H$  for which in (8),  $\sup_{0 \leq s \leq t} |X_s|, \sup_{0 \leq s \leq t} |V_s|, \langle \tilde{M}, \tilde{M} \rangle_t \leq H$ . By Tanaka formula for such  $X_t$ , we have (10). First note that the function  $\varphi_t(a) := (X_t - a)^+ - (X_0 - a)^+$  is Lipschitz continuous in  $a$  with Lipschitz constant 2. This implies that for any  $p > 2$  and  $a_i < a_{i+1}$ ,

$$|\varphi_t(a_{i+1}) - \varphi_t(a_i)|^p \leq 2^p (a_{i+1} - a_i)^p. \quad (13)$$

Secondly, for the second term, by the occupation times formula, Jensen's inequality and Fubini theorem,

$$\begin{aligned} E|I_t(a_{i+1}) - I_t(a_i)|^p &= E \left( \int_0^t 1_{\{a_i < X_s \leq a_{i+1}\}} ds \right)^p \\ &= \frac{1}{\sigma^{2p}} (a_{i+1} - a_i)^p E \left( \frac{1}{a_{i+1} - a_i} \int_{a_i}^{a_{i+1}} L_t^x dx \right)^p \\ &\leq \frac{1}{\sigma^{2p}} (a_{i+1} - a_i)^p \sup_x E(L_t^x)^p. \end{aligned} \quad (14)$$

Hence it follows from Lemma 2 that

$$E|I_t(a_{i+1}) - I_t(a_i)|^p \leq c(t, \sigma, p)(a_{i+1} - a_i)^p. \quad (15)$$

Thirdly, for the term  $\hat{B}_t^a$ ,

$$\begin{aligned} E|\hat{B}_t^{a_{i+1}} - \hat{B}_t^{a_i}|^p &= E \left| \int_0^t 1_{\{a_i < X_s \leq a_{i+1}\}} dB_s \right|^p \\ &\leq c_p E \left( \int_0^t 1_{\{a_i < X_s \leq a_{i+1}\}} ds \right)^{\frac{p}{2}} \\ &\leq c(t, p, \sigma)(a_{i+1} - a_i)^{\frac{p}{2}}. \end{aligned} \quad (16)$$

The last estimate can be obtained similarly to (14). In the following we can use the Burkholder-Davis-Gundy inequality, occupation times formula, changing integration orders and Jensen's inequality to have

$$\begin{aligned}
E|\hat{M}_t^{a_{i+1}} - \hat{M}_t^{a_i}|^p &= E \left| \int_0^{t+} \int_{R \setminus \{0\}} \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} y \mathbf{1}_{\{|y| < 1\}} \tilde{N}_p(dsdy) \right|^p \\
&\leq c_p E \left| \int_0^t \int_{X_s - a_{i+1}}^{X_s - a_i} y^2 \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right|^{\frac{p}{2}} \\
&= \frac{c_p}{\sigma^p} E \left| \int_{-\infty}^{\infty} L_t^x \int_{x - a_{i+1}}^{x - a_i} y^2 \mathbf{1}_{\{|y| < 1\}} n(dy) dx \right|^{\frac{p}{2}} \\
&= c(p, \sigma) Q^{\frac{p}{2}} E \left| \frac{1}{Q} \int_{-\infty}^{\infty} \left( \int_{y+a_i}^{y+a_{i+1}} L_t^x dx \right) y^2 \mathbf{1}_{\{|y| < 1\}} n(dy) \right|^{\frac{p}{2}} \\
&\leq c(p, \sigma) Q^{\frac{p}{2}} E \left| \frac{1}{Q} \int_{-\infty}^{\infty} \left( \int_{y+a_i}^{y+a_{i+1}} (L_t^x)^{\frac{p}{2}} dx \right) y^2 \mathbf{1}_{\{|y| < 1\}} n(dy) \right|^{\frac{p}{2}} \\
&\leq c(p, \sigma) |a_{i+1} - a_i|^{\frac{p}{2}} \left( \int_{R \setminus \{0\}} y^2 \mathbf{1}_{\{|y| < 1\}} n(dy) \right)^{\frac{p}{2}} \sup_x E(L_t^x)^{\frac{p}{2}} \\
&\leq c(t, \sigma, p) |a_{i+1} - a_i|^{\frac{p}{2}}, \tag{17}
\end{aligned}$$

where

$$Q := (a_{i+1} - a_i) \int_{-\infty}^{\infty} y^2 \mathbf{1}_{\{|y| < 1\}} n(dy)$$

is the normalization constant to make  $\frac{1}{Q} y^2 \mathbf{1}_{\{|y| < 1\}} \mathbf{1}_{(y+a_i, y+a_{i+1})}(x) dx n(dy)$  a probability measure so that Jensen's inequality can be applied. About  $K_1(t, a)$ , it is easy to see that

$$\begin{aligned}
&|K_1(t, a_{i+1}) - K_1(t, a_i)| \\
&\leq \sum_{0 \leq s \leq t} |(X_{s-} - a_{i+1})^+ - (X_s - a_{i+1})^+ - (X_{s-} - a_i)^+ + (X_s - a_i)^+| \mathbf{1}_{\{|\Delta X_s| \geq 1\}} \\
&\leq 2(a_{i+1} - a_i) \sum_{0 \leq s \leq t} \mathbf{1}_{\{|\Delta X_s| \geq 1\}}.
\end{aligned}$$

So

$$E|K_1(t, a_{i+1}) - K_1(t, a_i)|^p \leq C(a_{i+1} - a_i)^p. \tag{18}$$

In the following, we will estimate  $E|K_2(t, a_{i+1}) - K_2(t, a_i)|^p$ . With the decomposition (11), we will estimate the sum of each term for jumps  $|\Delta X_s| < 1$ . There are six such terms.

1) For the first term in (11), by the  $p$ -moment estimate formula and occupation times formula,

$$\begin{aligned}
&E \left( \sum_{0 \leq s \leq t} |X_s - a_i| \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \\
&= E \left( \sum_{0 \leq s \leq t} |X_s - a_i| \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{a_i < X_s - \Delta X_s \leq a_{i+1}\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \\
&\leq c_p E \left( \int_0^t \int_{X_s - a_{i+1}}^{X_s - a_i} |X_s - a_i|^2 \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{2}}
\end{aligned}$$



$$\begin{aligned}
& + p E \left( \int_0^t \int_{X_s - a_{i+1}}^{X_s - a_i} |X_s - a_i| 1_{\{X_s \leq a_i\}} 1_{\{|y| < 1\}} n(dy) ds \right)^p \\
& = \frac{c_p}{\sigma^p} E \left( \int_{-\infty}^{a_i} L_t^x \int_{x - a_{i+1}}^{x - a_i} |x - a_i|^2 1_{\{|y| < 1\}} n(dy) dx \right)^{\frac{p}{2}} \\
& \quad + \frac{p}{\sigma^{2p}} E \left( \int_{-\infty}^{a_i} L_t^x \int_{x - a_{i+1}}^{x - a_i} |x - a_i| 1_{\{|y| < 1\}} n(dy) dx \right)^p.
\end{aligned}$$

Now we change the orders of the integration and use Jensen's inequality, and Lemma 2, we have

$$\begin{aligned}
& E \left( \sum_{0 \leq s \leq t} |X_s - a_i| 1_{\{X_s \leq a_i\}} 1_{\{a_i < X_s - \leq a_{i+1}\}} 1_{\{|\Delta X_s| < 1\}} \right)^p \\
& \leq \frac{c_p}{\sigma^p} E \left( \int_{-\infty}^{a_i - a_{i+1}} \int_{y + a_i}^{y + a_{i+1}} L_t^x |x - a_i|^2 1_{\{|y| < 1\}} dx n(dy) \right. \\
& \quad \left. + \int_{a_i - a_{i+1}}^0 \int_{y + a_i}^{a_i} L_t^x |x - a_i|^2 1_{\{|y| < 1\}} dx n(dy) \right)^{\frac{p}{2}} \\
& \quad + \frac{p}{\sigma^{2p}} E \left( \int_{-\infty}^{a_i - a_{i+1}} \int_{y + a_i}^{y + a_{i+1}} L_t^x |x - a_i| 1_{\{|y| < 1\}} dx n(dy) \right. \\
& \quad \left. + \int_{a_i - a_{i+1}}^0 \int_{y + a_i}^{a_i} L_t^x |x - a_i| 1_{\{|y| < 1\}} dx n(dy) \right)^p \\
& \leq c(\sigma, p) E \left( \int_{-\infty}^{a_i - a_{i+1}} \int_{y + a_i}^{y + a_{i+1}} L_t^x dx |y|^2 1_{\{|y| < 1\}} n(dy) \right. \\
& \quad \left. + \int_{a_i - a_{i+1}}^0 \int_{y + a_i}^{a_i} L_t^x |a_{i+1} - a_i|^2 dx 1_{\{|y| < 1\}} n(dy) \right)^{\frac{p}{2}} \\
& \quad + c(\sigma, p) E \left( \int_{-\infty}^{a_i - a_{i+1}} \int_{y + a_i}^{y + a_{i+1}} L_t^x dx |y| 1_{\{|y| < 1\}} n(dy) \right. \\
& \quad \left. + \int_{a_i - a_{i+1}}^0 \int_{y + a_i}^{a_i} L_t^x |a_{i+1} - a_i| dx 1_{\{|y| < 1\}} n(dy) \right)^p \\
& \leq c(\sigma, p) |a_{i+1} - a_i|^{\frac{p}{2}} \sup_x E(L_t^x)^{\frac{p}{2}} \int_{-\infty}^{a_i - a_{i+1}} |y|^2 1_{\{|y| < 1\}} n(dy) \\
& \quad + c(\sigma, p) |a_{i+1} - a_i|^p \sup_x E(L_t^x)^{\frac{p}{2}} \int_{a_i - a_{i+1}}^0 |y| 1_{\{|y| < 1\}} n(dy) \\
& \quad + c(\sigma, p) |a_{i+1} - a_i|^p \sup_x E(L_t^x)^p \int_{-\infty}^{a_i - a_{i+1}} |y| 1_{\{|y| < 1\}} n(dy) \\
& \quad + c(\sigma, p) |a_{i+1} - a_i|^p \sup_x E(L_t^x)^p \int_{a_i - a_{i+1}}^0 |y| 1_{\{|y| < 1\}} n(dy) \\
& \leq c(t, \sigma, p) |a_{i+1} - a_i|^{\frac{p}{2}}.
\end{aligned}$$

We can use the similar method to deal with other terms. In the following, we will only sketch the estimate without giving great details.

2) For the second term, we have

$$\begin{aligned}
& E \left( \sum_{0 \leq s \leq t} (a_{i+1} - a_i) \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{X_s > a_{i+1}\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \\
& \leq (a_{i+1} - a_i)^p \left[ c_p E \left( \int_0^t \int_{-\infty}^{\infty} \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{X_s - y > a_{i+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{2}} \right. \\
& \quad \left. + p E \left( \int_0^t \int_{-\infty}^{\infty} \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{X_s - y > a_{i+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^p \right] \\
& = (a_{i+1} - a_i)^p \left[ \frac{c_p}{\sigma^p} E \left( \int_{-\infty}^{a_i} L_t^x \int_{-\infty}^{x - a_{i+1}} \mathbf{1}_{\{|y| < 1\}} n(dy) dx \right)^{\frac{p}{2}} \right. \\
& \quad \left. + \frac{p}{\sigma^{2p}} E \left( \int_{-\infty}^{a_i} L_t^x \int_{-\infty}^{x - a_{i+1}} \mathbf{1}_{\{|y| < 1\}} n(dy) dx \right)^p \right] \\
& \leq c(\sigma, p) (a_{i+1} - a_i)^p \left[ E \left( \int_{-\infty}^{a_i - a_{i+1}} \left( \int_{y + a_{i+1}}^{a_i} L_t^x dx \right) \mathbf{1}_{\{|y| < 1\}} n(dy) \right)^{\frac{p}{2}} \right. \\
& \quad \left. + E \left( \int_{-\infty}^{a_i - a_{i+1}} \left( \int_{y + a_{i+1}}^{a_i} L_t^x dx \right) \mathbf{1}_{\{|y| < 1\}} n(dy) \right)^p \right] \\
& \leq c(\sigma, p) |a_{i+1} - a_i|^p \left( \sup_x E(L_t^x)^{\frac{p}{2}} + \sup_x E(L_t^x)^p \right) \int_{-\infty}^{a_i - a_{i+1}} |y| \mathbf{1}_{\{|y| < 1\}} n(dy) \\
& \leq c(t, \sigma, p) |a_{i+1} - a_i|^p.
\end{aligned}$$

3) For the third term, we have

$$\begin{aligned}
& E \left( \sum_{0 \leq s \leq t} |X_s - a_i| \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \\
& \leq c_p E \left( \int_0^t \int_{-\infty}^{\infty} |X_s - a_i|^2 \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_s - y \leq a_i\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^{\frac{p}{2}} \\
& \quad + p E \left( \int_0^t \int_{-\infty}^{\infty} |X_s - a_i| \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_s - y \leq a_i\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \right)^p \\
& = \frac{c_p}{\sigma^p} E \left( \int_{a_i}^{a_{i+1}} L_t^x \int_{x - a_i}^{\infty} |x - a_i|^2 \mathbf{1}_{\{|y| < 1\}} n(dy) dx \right)^{\frac{p}{2}} \\
& \quad + \frac{p}{\sigma^{2p}} E \left( \int_{a_i}^{a_{i+1}} L_t^x \int_{x - a_i}^{\infty} |x - a_i| \mathbf{1}_{\{|y| < 1\}} n(dy) dx \right)^p \\
& \leq c(\sigma, p) E \left( \int_0^{a_{i+1} - a_i} \int_{a_i}^{a_i + y} L_t^x |a_{i+1} - a_i|^2 dx \mathbf{1}_{\{|y| < 1\}} n(dy) \right. \\
& \quad \left. + \int_{a_{i+1} - a_i}^{\infty} \int_{a_i}^{a_{i+1}} L_t^x |a_{i+1} - a_i| dx |y| \mathbf{1}_{\{|y| < 1\}} n(dy) \right)^{\frac{p}{2}} \\
& \quad + E \left( \int_0^{a_{i+1} - a_i} \int_{a_i}^{a_i + y} L_t^x |a_{i+1} - a_i| dx \mathbf{1}_{\{|y| < 1\}} n(dy) \right)
\end{aligned}$$

$$\begin{aligned}
& + \int_{a_{i+1}-a_i}^{\infty} \int_{a_i}^{a_{i+1}} L_t^x dx |y| \mathbf{1}_{\{|y|<1\}} n(dy) \Big)^p \\
& \leq c(\sigma, p) |a_{i+1} - a_i|^p \left( \sup_x E(L_t^x)^{\frac{p}{2}} + \sup_x E(L_t^x)^p \right) \int_0^{\infty} |y| \mathbf{1}_{\{|y|<1\}} n(dy) \\
& \leq c(t, \sigma, p) |a_{i+1} - a_i|^p.
\end{aligned}$$

4) The fourth term is symmetric to the third term, so by a similar computation, we have

$$E \left( \sum_{0 \leq s \leq t} |a_{i+1} - X_s| \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_s > a_{i+1}\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \leq c(t, \sigma, p) |a_{i+1} - a_i|^p.$$

5) For the fifth term, as it is symmetric to the second term, so we can use a similar computation to have

$$E \left( \sum_{0 \leq s \leq t} (a_{i+1} - a_i) \mathbf{1}_{\{X_s > a_{i+1}\}} \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \leq c(t, \sigma, p) |a_{i+1} - a_i|^p.$$

6) The last term is symmetric to the first term, so by a similar computation, we have

$$E \left( \sum_{0 \leq s \leq t} |a_{i+1} - X_s| \mathbf{1}_{\{X_s > a_{i+1}\}} \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{|\Delta X_s| < 1\}} \right)^p \leq c(t, \sigma, p) |a_{i+1} - a_i|^{\frac{p}{2}}.$$

From the above six cases, we can derive that

$$E |K_2(t, a_{i+1}) - K_2(t, a_i)|^p \leq c(t, \sigma, p) |a_{i+1} - a_i|^{\frac{p}{2}}. \quad (19)$$

Now we use Proposition 4.1.1 in [23] ( $i = 1, \gamma > p - 1$ ), for any finite partition  $\{a_l\}$  of  $[a, b]$

$$\sup_D \sum_l |\hat{B}_t^{a_{l+1}} - \hat{B}_t^{a_l}|^p \leq c(p, \gamma) \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\hat{B}_t^{a_k^n} - \hat{B}_t^{a_{k-1}^n}|^p,$$

where

$$a_k^n = a + \frac{k}{2^n} (b - a), \quad k = 0, 1, \dots, 2^n.$$

The key point here is that the right hand side does not depend on the partition  $D$ . We take the expectation and use (16), it follows that

$$\begin{aligned}
E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\hat{B}_t^{a_k^n} - \hat{B}_t^{a_{k-1}^n}|^p &= \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} E |\hat{B}_t^{a_k^n} - \hat{B}_t^{a_{k-1}^n}|^p \\
&\leq c \sum_{n=1}^{\infty} n^\gamma \left( \frac{b-a}{2^n} \right)^{\frac{p}{2}-1} < \infty,
\end{aligned}$$

as  $p > 2$ . Therefore

$$\sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\hat{B}_t^{a_k^n} - \hat{B}_t^{a_{k-1}^n}|^p < \infty \text{ a.s.}$$

It turns out that for any interval  $[a, b] \subset R$

$$\sup_D \sum_l |\hat{B}_t^{a_{l+1}} - \hat{B}_t^{a_l}|^p < \infty \text{ a.s.} \quad (20)$$

But we know  $L_t^a$  has a compact support  $[-K, K]$  in  $a$ . So for the partition  $D := D_{-K, K} = \{-K = a_0 < a_1 < \dots < a_r = K\}$ , we obtain

$$\sup_D \sum_l |\hat{B}_t^{a_{l+1}} - \hat{B}_t^{a_l}|^p < \infty \text{ a.s.} \quad (21)$$

In the same way, from (15), (17), (18) and (19), we can prove that

$$\sup_D \sum_i |I_t^{a_{i+1}} - I_t^{a_i}|^p < \infty \text{ a.s.}, \quad (22)$$

$$\sup_D \sum_i |\hat{M}_t^{a_{i+1}} - \hat{M}_t^{a_i}|^p < \infty \text{ a.s.}, \quad (23)$$

$$\sup_D \sum_i |K_1(t, a_{i+1}) - K_1(t, a_i)|^p < \infty \text{ a.s.}, \quad (24)$$

$$\sup_D \sum_i |K_2(t, a_{i+1}) - K_2(t, a_i)|^p < \infty \text{ a.s.} \quad (25)$$

On the other hand, it is easy to see from (13) that

$$\sum_i |\varphi_t(a_{i+1}) - \varphi_t(a_i)|^p \leq 2^p \sum_i (a_{i+1} - a_i)^p \leq 2^p [\sum_i (a_{i+1} - a_i)]^p = 2^p (b - a)^p. \quad (26)$$

Then from (10), (21), (22), (23), (24), (25) and (26), it turns out that

$$\sup_D \sum_i |L_t^{a_{i+1}} - L_t^{a_i}|^p < \infty \text{ a.s.}$$

Finally we can use the usual localization procedure to remove the assumption that  $\sup_{0 \leq s \leq t} |X_s|, \sup_{0 \leq s \leq t} |V_s|, \langle \tilde{M}, \tilde{M} \rangle_t \leq K$ . For this, define a stopping time for an integer  $H > 0$ :  $\tau_H = \inf\{s : \min\{|X_s|, \int_0^s |dV_r|, \langle M, M \rangle_s\} > H\}$  if there exists  $s$  such that  $\min\{|X_s|, |V_s|, \langle \tilde{M}, \tilde{M} \rangle_s\} > H$  and  $\tau_H = +\infty$  otherwise. Then the above result shows that there exists  $\Omega_1 \subset \Omega$  with  $P(\Omega_1) = 1$  such that for each  $\omega \in \Omega_1$  and each given integer  $H > 0$ ,

$$\sup_D \sum_i |L_{t \wedge \tau_H}^{a_{i+1}} - L_{t \wedge \tau_H}^{a_i}|^p < \infty.$$

Since  $\sup_{0 \leq s \leq t} |X_s|(\omega), \sup_{0 \leq s \leq t} |V_s|(\omega)$  and  $\langle \tilde{M}, \tilde{M} \rangle_t(\omega)$  are finite almost surely so there exists  $\Omega_2 \subset \Omega$  with  $P(\Omega_2) = 1$  such that for each  $\omega \in \Omega_2$ , there exists an integer  $H(\omega) > 0$  such that  $\sup_{0 \leq s \leq t} |X_s|(\omega), \sup_{0 \leq s \leq t} |V_s|(\omega), \langle \tilde{M}, \tilde{M} \rangle_t(\omega) \leq H$ . This leads to  $\tau_H(\omega) > t$ . So for each  $\omega \in \Omega_1 \cap \Omega_2$ ,

$$\sup_D \sum_i |L_t^{a_{i+1}} - L_t^{a_i}|^p < \infty.$$

The result follows as  $P(\Omega_1 \cap \Omega_2) = 1$ .  $\diamond$

Recall (11) and denote

$$\begin{aligned} J^*(X_s, X_{s-}, a_i, a_{i+1}) &= J^*(X_s, X_s - \Delta X_s, a_i, a_{i+1}) \\ &:= \left[ J_1(s, a_{i+1}) - J_1(s, a_i) + J_2(s, a_{i+1}) - J_2(s, a_i) \right] 1_{\{|\Delta X_s| < 1\}}. \end{aligned}$$

From the proof of the above theorem, we can easily see that

**Corollary 4** *Assume (12) is satisfied, then for any  $p \geq 2$ ,*

$$E \left( \int_0^t \int_{-\infty}^{\infty} J^*(X_s, X_s - y, a_i, a_{i+1}) n(dy) ds \right)^p \leq c(t, \sigma, p) |a_{i+1} - a_i|^p.$$

The estimate will be used in the next section.

### 3 The local time rough path

The  $p$ -variation ( $p > 2$ ) result of the local time enables one to use Young's integration theory to define  $\int_{-\infty}^{\infty} g(x) dx L_t^x$  for  $g$  being of bounded  $q$ -variation when  $1 \leq q < 2$ . This is because in this case, for any  $q \in [1, 2)$ , one can always find a constant  $p > 2$  such that the condition  $\frac{1}{p} + \frac{1}{q} > 1$  for the existence of the Young integral is satisfied. However, when  $q \geq 2$ , Young integral is no longer well defined. We have to use a new integration theory. Lyons' integration of rough path provides a way to push the result further. But from [23], generally, we cannot expect to have an integration theory to define integrals such as  $\int_{-\infty}^{\infty} g(x) dx L_t^x$ . However, using the method in Chapter 6 in [23], we can treat  $Z_x := (L_t^x, g(x))$  as a process of variable  $x$  in  $R^2$ . Then it's easy to know that  $Z_x$  is of bounded  $\hat{q}$ -variation in  $x$ , where  $\hat{q} = q$ , if  $q > 2$ , and  $\hat{q} > 2$  can be taken any number when  $q = 2$ . Most of the analysis in this section works for  $2 \leq q < 4$ , especially we will establish the convergence of smooth rough path in the  $\theta$ -variation topology for any  $\theta \in (q, 4)$  so to obtain  $\mathbf{Z}_{a,b}^1$  and  $\mathbf{Z}_{a,b}^2$ . In particular, when  $2 \leq q < 3$ , we obtain the existence of the geometric rough path  $\mathbf{Z} = (1, \mathbf{Z}^1, \mathbf{Z}^2)$  associated to  $Z$ . In the following we consider  $2 \leq q < 4$ , otherwise we will explicitly say so. Recall the  $\theta$ -variation metric  $d_\theta$  on  $C_{0,\theta}(\Delta, T^{(\theta)}(R^2))$  defined in [23],

$$d_\theta(\mathbf{Z}, \mathbf{Y}) = \max_{1 \leq i \leq [\theta]} d_{i,\theta}(\mathbf{Z}^i, \mathbf{Y}^i) = \max_{1 \leq i \leq [\theta]} \sup_D \left( \sum_l |\mathbf{Z}_{x_{l-1}, x_l}^i - \mathbf{Y}_{x_{l-1}, x_l}^i|^{\frac{\theta}{i}} \right)^{\frac{i}{\theta}}.$$

Let  $[x', x'']$  be any interval in  $R$ . From the proof of Theorem 3, for any  $p \geq 2$ , we know there exists a constant  $c > 0$  such that

$$E |L_t^b - L_t^a|^p \leq c |b - a|^{\frac{p}{2}}, \quad (27)$$

i.e.  $L_t^x$  satisfies Hölder condition in [23] with exponent  $\frac{1}{2}$ . First we consider the case when  $g$  is continuous. Recall in [23], a control  $w$  is a continuous super-additive function on  $\Delta := \{(a, b) : x' \leq a < b \leq x''\}$  with values in  $[0, \infty)$  such that  $w(a, a) = 0$ . Therefore

$$w(a, b) + w(b, c) \leq w(a, c), \quad \text{for any } (a, b), (b, c) \in \Delta.$$

If  $g(x)$  is of bounded  $q$ -variation, we can find a control  $w$  s.t.

$$|g(b) - g(a)|^q \leq w(a, b),$$

for any  $(a, b) \in \Delta := \{(a, b) : x' \leq a < b \leq x''\}$ . It is obvious that  $w_1(a, b) := w(a, b) + (b - a)$  is also a control of  $g$ . Set  $h = \frac{1}{q}$ , it is trivial to see for any  $\theta > q$  (so  $h\theta > 1$ ) we have,

$$|g(b) - g(a)|^\theta \leq w_1(a, b)^{h\theta}, \text{ for any } (a, b) \in \Delta. \quad (28)$$

Considering (27), we can see  $Z_x$  satisfies, for such  $h = \frac{1}{q}$ , and any  $\theta > q$ , there exists a constant  $c$  such that

$$E|Z_b - Z_a|^\theta \leq cw_1(a, b)^{h\theta}, \text{ for any } (a, b) \in \Delta. \quad (29)$$

For any  $m \in N$ , define a continuous and bounded variation path  $Z(m)$  by

$$Z(m)_x := Z_{x_{l-1}^m} + \frac{w_1(x) - w_1(x_{l-1}^m)}{w_1(x_l^m) - w_1(x_{l-1}^m)} \Delta_l^m Z, \quad (30)$$

if  $x_{l-1}^m \leq x < x_l^m$ , for  $l = 1, \dots, 2^m$ , and  $\Delta_l^m Z = Z_{x_l^m} - Z_{x_{l-1}^m}$ . Here  $D_m := \{x' = x_0^m < x_1^m < \dots < x_{2^m}^m = x''\}$  is a partition of  $[x', x'']$  such that

$$w_1(x_l^m) - w_1(x_{l-1}^m) = \frac{1}{2^m} w_1(x', x''),$$

where  $w_1(x) := w_1(x', x)$ . It is obvious that  $x_l^m - x_{l-1}^m \leq \frac{1}{2^m} w_1(x', x'')$  and by the superadditivity of the control function  $w_1$ ,

$$w_1(x_{l-1}^m, x_l^m) \leq w_1(x_l^m) - w_1(x_{l-1}^m) = \frac{1}{2^m} w_1(x', x'').$$

The corresponding smooth rough path  $\mathbf{Z}(m)$  is built by taking its iterated path integrals, i.e. for any  $(a, b) \in \Delta$ ,

$$\mathbf{Z}(m)_{a,b}^j = \int_{a < x_1 < \dots < x_j < b} dZ(m)_{x_1} \otimes \dots \otimes dZ(m)_{x_j}. \quad (31)$$

In the following, we will prove  $\{\mathbf{Z}(m)\}_{m \in N}$  converges to a geometric rough path  $\mathbf{Z}$  in the  $\theta$ -variation topology when  $2 \leq q < 3$ . We call  $\mathbf{Z}$  the canonical geometric rough path associated to  $Z$ .

**Remark 5** *The bounded variation process  $Z(m)_x$  is a generalized Wong-Zakai approximation to the process  $Z$  of bounded  $\hat{q}$ -variation. Here we divide  $[x', x'']$  by equally partitioning the range of  $w_1$ . We then use (30) to form the piecewise curved approximation to  $Z$ . Note here Wong-Zakai's standard piecewise linear approximation does not work immediately.*

Let's first look at the first level path  $\mathbf{Z}(m)_{a,b}^1$ . The method is similar to Chapter 4 in [23] for Brownian motion. Similar to Proposition 4.2.1 in [23], we can prove that for all  $n \in N$ ,  $m \mapsto \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1|^\theta$  is increasing and for  $m \geq n$ ,

$$\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1 = \mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^1 = Z_{x_k^n} - Z_{x_{k-1}^n}. \quad (32)$$

Let  $\mathbf{Z}_{a,b}^1 = Z_b - Z_a$ . Then (29) implies  $E|\mathbf{Z}_{a,b}^1|^\theta \leq cw_1(a, b)^{h\theta}$ . For such points  $\{x_k^n\}$ ,  $k = 1, \dots, 2^n$ ,  $n = 1, 2, \dots$ , defined above we still have the inequality in Proposition 4.1.1 in [23],

$$\begin{aligned}
E \sup_D \sum_l |\mathbf{Z}_{x_{l-1}, x_l}^1|^\theta &\leq C(\theta, \gamma) E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta \\
&\leq C_1 \sum_{n=1}^{\infty} n^\gamma \left(\frac{1}{2^n}\right)^{h\theta-1} w_1(x', x'')^{h\theta},
\end{aligned} \tag{33}$$

for constant  $C_1 = C(\theta, \gamma)c$ . Since  $h\theta - 1 > 0$ , the series on the right-hand side of (33) is convergent, so that  $\sup_D \sum_l |\mathbf{Z}_{x_{l-1}, x_l}^1|^\theta < \infty$  almost surely. This shows that  $\mathbf{Z}^1$  has finite  $\theta$ -variation almost surely. Moreover, for any  $\gamma > \theta - 1$ , there exists a constant  $C_1(\theta, \gamma, c) > 0$  such that

$$\begin{aligned}
E \sup_m \sup_D \sum_l |\mathbf{Z}(m)_{x_{l-1}, x_l}^1|^\theta &\leq C(\theta, \gamma) E \sup_m \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1|^\theta \\
&\leq C(\theta, \gamma) E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta \\
&\leq C_1(\theta, \gamma, c) \sum_{n=1}^{\infty} n^\gamma \left(\frac{1}{2^n}\right)^{h\theta-1} w_1(x', x'')^{h\theta} \\
&< \infty.
\end{aligned} \tag{34}$$

So

$$\sup_m \sup_D \sum_l |\mathbf{Z}(m)_{x_{l-1}, x_l}^1|^\theta < \infty \quad a.s. \tag{35}$$

This means that  $\mathbf{Z}(m)_{a,b}^1$  have finite  $\theta$ -variation uniformly in  $m$ . And furthermore, from (32) and some standard arguments,

$$E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1 - \mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta \leq C \left(\frac{1}{2^m}\right)^{\frac{h\theta-1}{2}}, \tag{36}$$

where  $C$  depends on  $\theta$ ,  $h$ ,  $w_1(x', x'')$ , and  $c$  in (29). By Proposition 4.1.2 in [23], Jensen's inequality and (36),

$$\begin{aligned}
E \sum_{m=1}^{\infty} \sup_D \left( \sum_l |\mathbf{Z}(m)_{x_{l-1}, x_l}^1 - \mathbf{Z}_{x_{l-1}, x_l}^1|^\theta \right)^{\frac{1}{\theta}} &\leq E \sum_{m=1}^{\infty} \left( \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1 - \mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta \right)^{\frac{1}{\theta}} \\
&\leq \sum_{m=1}^{\infty} \left( E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1 - \mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta \right)^{\frac{1}{\theta}} \\
&\leq C \sum_{m=1}^{\infty} \left(\frac{1}{2^m}\right)^{\frac{h\theta-1}{2\theta}} \\
&< \infty,
\end{aligned} \tag{37}$$

for  $h\theta > 1$ . So we obtain

**Theorem 6** *Let  $L_t^x$  be the local time of the time homogeneous Lévy process  $X_t$  given by (2), and  $g$  be a continuous function of bounded  $q$ -variation. Assume  $q \geq 1$ ,  $\sigma \neq 0$  and the Lévy measure  $n(dy)$  satisfies (12). Then for any  $\theta > q$ , the continuous process  $Z_x = (L_t^x, g(x))$  satisfying (29), we have*

$$\sum_{m=1}^{\infty} \sup_D \left( \sum_l |\mathbf{Z}(m)_{x_{l-1}, x_l}^1 - \mathbf{Z}_{x_{l-1}, x_l}^1|^\theta \right)^{\frac{1}{\theta}} < \infty \text{ a.s.} \quad (38)$$

In particular,  $(\mathbf{Z}(m)_{a,b}^1)$  converges to  $(\mathbf{Z}_{a,b}^1)$  in the  $\theta$ -variation distance a.s. for any  $(a, b) \in \Delta$ .

We next consider the second level path  $\mathbf{Z}(m)_{a,b}^2$ . As in [23], we can also see that if  $m \leq n$ ,

$$\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2 = 2^{2(m-n)-1} (\Delta_l^m Z)^{\otimes 2}, \quad (39)$$

where  $l$  is chosen such that  $x_{l-1}^m \leq x_{k-1}^n < x_k^n \leq x_l^m$ ; if  $m > n$ ,

$$\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2 = \frac{1}{2} \Delta_k^n Z \otimes \Delta_k^n Z + \frac{1}{2} \sum_{\substack{r, l=2^{m-n}(k-1)+1 \\ r < l}}^{2^{m-n}k} (\Delta_r^m Z \otimes \Delta_l^m Z - \Delta_l^m Z \otimes \Delta_r^m Z),$$

so

$$\begin{aligned} & \mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2 \\ &= \frac{1}{2} \sum_{l=2^{m-n}(k-1)+1}^{2^{m-n}k} (\Delta_{2l-1}^{m+1} Z \otimes \Delta_{2l}^{m+1} Z - \Delta_{2l}^{m+1} Z \otimes \Delta_{2l-1}^{m+1} Z), \end{aligned} \quad (40)$$

$k = 1, \dots, 2^n$ . Similar to the proof of Proposition 4.3.3 in [23], we have

**Proposition 7** *Assume  $q \geq 2$ . Let  $\theta > q$ . Then for  $m \leq n$ ,*

$$\sum_{k=1}^{2^n} E |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^{\frac{\theta}{2}} \leq C \left( \frac{1}{2^{n+m}} \right)^{\frac{\theta h-1}{2}}, \quad (41)$$

where  $C$  depends on  $\theta$ ,  $h(= \frac{1}{q})$ ,  $w_1(x', x'')$ , and  $c$  in (29).

The main step to establish the geometric rough path integral over  $Z$  is the following estimate. The proof is rather complicated.

**Proposition 8** *Assume  $2 \leq q < 4$ . Let  $q < \theta < 4$ . Then for  $m > n$ , we have*

$$E |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^{\frac{\theta}{2}} \leq C \left[ \left( \frac{1}{2^n} \right)^{\frac{\theta}{4}} \left( \frac{1}{2^m} \right)^{\frac{1}{2} h \theta} + \left( \frac{1}{2^n} \right)^{\frac{\theta}{2}} \left( \frac{1}{2^m} \right)^{\frac{1}{2} h \theta - \frac{1}{8} \theta} \right], \quad (42)$$

where  $C$  is a generic constant and also depends on  $\theta$ ,  $h(= \frac{1}{q})$ ,  $w_1(x', x'')$ , and  $c$  in (29).

In order to prove the above proposition, we need the following key lemma about the correlation of  $\hat{M}_t^{a_{i+1}} - \hat{M}_t^{a_i}$  and  $K_2(t, a_{j+1}) - K_2(t, a_j)$ , and the correlation of  $K_2(t, a_{i+1}) - K_2(t, a_i)$  and  $K_2(t, a_{j+1}) - K_2(t, a_j)$ . Here  $\hat{M}$  and  $K_2$  were defined in the last section.

**Lemma 9** *For any  $a_0 < a_1 < \dots < a_m$ ,*

$$\begin{aligned} & \left| E \int_0^{t^+} \int_{|y| < 1} 1_{\{a_i < X_{s-} \leq a_{i+1}\}} y \tilde{N}_p(ds dy) (K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \\ & \leq \begin{cases} c(t, \sigma)(a_{i+1} - a_i), & \text{when } 0 \leq i = j \leq m, \\ c(t, \sigma)[(a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2], & \text{when } 0 \leq i \neq j \leq m, \end{cases} \end{aligned} \quad (43)$$

and

$$\begin{aligned} & \left| E (K_2(t, a_{i+1}) - K_2(t, a_i)) (K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \\ & \leq \begin{cases} c(t, \sigma)(a_{i+1} - a_i), & \text{when } 0 \leq i = j \leq m, \\ c(t, \sigma)[(a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2], & \text{when } 0 \leq i \neq j \leq m. \end{cases} \end{aligned} \quad (44)$$



**Proof:** When  $i = j$ , it is easy to see from (17) and (19), using the Cauchy-Schwarz inequality that

$$\begin{aligned} & \left| E \int_0^{t^+} \int_{|y|<1} 1_{\{a_i < X_{s-} \leq a_{i+1}\}} y \tilde{N}_p(dsdy) (K_2(t, a_{i+1}) - K_2(t, a_i)) \right| \\ & \leq \left( E \left| \int_0^{t^+} \int_{|y|<1} 1_{\{a_i < X_{s-} \leq a_{i+1}\}} y \tilde{N}_p(dsdy) \right|^2 \right)^{\frac{1}{2}} \left( E |K_2(t, a_{i+1}) - K_2(t, a_i)|^2 \right)^{\frac{1}{2}} \\ & \leq c(t, \sigma)(a_{i+1} - a_i). \end{aligned}$$

When  $i \neq j$ , without losing generality, we assume that  $i < j$ . Note  $K_2(t, a_{j+1}) - K_2(t, a_j)$  can be decomposed as the sum of the integral of random martingale measure and integral of compensator. The compensator part is of bounded variation. Now we use the quadratic variation formula for stochastic integrals with respect to martingales and the fact that  $E(P.M.) = 0$  if  $P$  is a continuous process of bounded variation and  $M$  is a martingale of mean zero. Then from (11), occupation times formula, Fubini theorem and Jensen's inequality, when  $a_{j+1} - a_j \geq a_{i+1} - a_i$ , we have

$$\begin{aligned} & \left| E \int_0^{t^+} \int_{|y|<1} 1_{\{a_i < X_{s-} \leq a_{i+1}\}} y \tilde{N}_p(dsdy) (K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \\ & \leq E \left[ \int_0^t \int_{|y|<1} 1_{\{a_i < X_{s-} \leq a_{i+1}\}} |y| 1_{\{a_j < X_s \leq a_{j+1}\}} (X_s - a_j) n(dy) ds \right. \\ & \quad \left. + \int_0^t \int_{|y|<1} 1_{\{a_i < X_{s-} \leq a_{i+1}\}} |y| 1_{\{X_s > a_{j+1}\}} (a_{j+1} - a_j) n(dy) ds \right] \\ & = \frac{1}{\sigma^2} E \left[ \int_{a_j}^{a_{j+1}} L_t^x \int_{x-a_{i+1}}^{x-a_i} |y| 1_{\{|y|<1\}} (x - a_j) n(dy) dx \right. \\ & \quad \left. + \int_{a_{j+1}}^{\infty} L_t^x \int_{x-a_{i+1}}^{x-a_i} |y| 1_{\{|y|<1\}} (a_{j+1} - a_j) n(dy) dx \right] \\ & = \frac{1}{\sigma^2} E \left[ \int_{a_j-a_{i+1}}^{a_j-a_i} \int_{a_j}^{y+a_{i+1}} L_t^x |y| 1_{\{|y|<1\}} (x - a_j) dx n(dy) \right. \\ & \quad \left. + \int_{a_j-a_i}^{a_{j+1}-a_{i+1}} \int_{y+a_i}^{y+a_{i+1}} L_t^x |y| 1_{\{|y|<1\}} (x - a_j) dx n(dy) \right. \\ & \quad \left. + \int_{a_{j+1}-a_{i+1}}^{a_{j+1}-a_i} \int_{y+a_i}^{a_{j+1}} L_t^x |y| 1_{\{|y|<1\}} (x - a_j) dx n(dy) \right] \\ & + \frac{1}{\sigma^2} E \left[ \int_{a_{j+1}-a_{i+1}}^{a_{j+1}-a_i} \int_{a_{j+1}}^{y+a_{i+1}} L_t^x |y| 1_{\{|y|<1\}} (a_{j+1} - a_j) dx n(dy) \right. \\ & \quad \left. + \int_{a_{j+1}-a_i}^{\infty} \int_{y+a_i}^{y+a_{i+1}} L_t^x |y| 1_{\{|y|<1\}} (a_{j+1} - a_j) dx n(dy) \right] \\ & \leq c(\sigma) [(a_{j+1} - a_j)^2 + (a_{i+1} - a_i)^2] \sup_x (EL_t^x) \int_{a_j-a_{i+1}}^{\infty} |y| 1_{\{|y|<1\}} n(dy) \\ & \leq c(t, \sigma) [(a_{j+1} - a_j)^2 + (a_{i+1} - a_i)^2]. \end{aligned}$$

When  $a_{j+1} - a_j < a_{i+1} - a_i$ , we also can get the same estimate as above, but the difference is only the integral

$$\begin{aligned}
& \int_{a_j}^{a_{j+1}} L_t^x \int_{x-a_{i+1}}^{x-a_i} |y| \mathbf{1}_{\{|y|<1\}}(x-a_j) n(dy) dx \\
&= \int_{a_j-a_{i+1}}^{a_{j+1}-a_{i+1}} \int_{a_j}^{y+a_{i+1}} L_t^x |y| \mathbf{1}_{\{|y|<1\}}(x-a_j) dx n(dy) \\
&\quad + \int_{a_{j+1}-a_{i+1}}^{a_j-a_i} \int_{a_j}^{a_{j+1}} L_t^x |y| \mathbf{1}_{\{|y|<1\}}(x-a_j) dx n(dy) \\
&\quad + \int_{a_j-a_i}^{a_{j+1}-a_i} \int_{y+a_i}^{a_{j+1}} L_t^x |y| \mathbf{1}_{\{|y|<1\}}(x-a_j) dx n(dy).
\end{aligned}$$

So we proved (43).

To estimate  $\left| E(K_2(t, a_{i+1}) - K_2(t, a_i)) \cdot (K_2(t, a_{j+1}) - K_2(t, a_j)) \right|$ , first when  $i = j$ , (44) follows from (19) directly. Now we consider the case when  $i \neq j$ . Without losing generality, we assume that  $i < j$ . From (11), it is easy to see that

$$\begin{aligned}
& (K_2(t, a_{i+1}) - K_2(t, a_i)) \cdot (K_2(t, a_{j+1}) - K_2(t, a_j)) \\
&= \int_0^{t+} \int_R J^*(X_s, X_s - y, a_i, a_{i+1}) N_p(dy ds) \cdot \int_0^{t+} \int_R J^*(X_s, X_s - y, a_j, a_{j+1}) N_p(dy ds),
\end{aligned}$$

so

$$\begin{aligned}
& E(K_2(t, a_{i+1}) - K_2(t, a_i)) \cdot (K_2(t, a_{j+1}) - K_2(t, a_j)) \\
&= E \int_0^t \int_R J^*(X_s, X_s - y, a_i, a_{i+1}) J^*(X_s, X_s - y, a_j, a_{j+1}) n(dy) ds \\
&\quad + E \int_0^t \int_R J^*(X_s, X_s - y, a_i, a_{i+1}) n(dy) ds \cdot \int_0^t \int_R J^*(X_s, X_s - y, a_j, a_{j+1}) n(dy) ds.
\end{aligned}$$

By using the Cauchy-Schwarz inequality and Corollary 4, we know,

$$\begin{aligned}
& \left| E \int_0^t \int_R J^*(X_s, X_s - y, a_i, a_{i+1}) n(dy) ds \cdot \int_0^t \int_R J^*(X_s, X_s - y, a_j, a_{j+1}) n(dy) ds \right| \\
&\leq \left( E \left( \int_0^t \int_R J^*(X_s, X_s - y, a_i, a_{i+1}) n(dy) ds \right)^2 \right)^{\frac{1}{2}} \left( E \left( \int_0^t \int_R J^*(X_s, X_s - y, a_j, a_{j+1}) n(dy) ds \right)^2 \right)^{\frac{1}{2}} \\
&\leq c(t, \sigma)(a_{i+1} - a_i)(a_{j+1} - a_j).
\end{aligned}$$

But from (11), we know

$$\int_0^t \int_R J^*(X_s, X_s - y, a_i, a_{i+1}) J^*(X_s, X_s - y, a_j, a_{j+1}) n(dy) ds = A_1 + A_2 + A_3 + A_4,$$

where

$$\begin{aligned}
A_1 &:= \int_0^t \int_R \left[ -(a_{i+1} - a_i)(a_j - X_s) \right] \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{a_j < X_s - \leq a_{j+1}\}} \mathbf{1}_{\{|y|<1\}} n(dy) ds \\
&\quad + \int_0^t \int_R \left[ (a_{i+1} - a_i)(a_{j+1} - a_j) \right] \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{X_s - > a_{j+1}\}} \mathbf{1}_{\{|y|<1\}} n(dy) ds, \\
A_2 &:= \int_0^t \int_R \left[ -(a_{i+1} - X_s)(a_j - X_s) \right] \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{a_j < X_s - \leq a_{j+1}\}} \mathbf{1}_{\{|y|<1\}} n(dy) ds \\
&\quad + \int_0^t \int_R (a_{i+1} - X_s)(a_{j+1} - a_j) \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_s - > a_{j+1}\}} \mathbf{1}_{\{|y|<1\}} n(dy) ds,
\end{aligned}$$

$$\begin{aligned}
A_3 &:= \int_0^t \int_R \left[ (a_{i+1} - a_i)(X_s - a_j) \right] \mathbf{1}_{\{X_{s-} \leq a_i\}} \mathbf{1}_{\{a_j < X_s \leq a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\quad + \int_0^t \int_R (a_{i+1} - a_i)(a_{j+1} - a_j) \mathbf{1}_{\{X_{s-} \leq a_i\}} \mathbf{1}_{\{X_s > a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds, \\
A_4 &:= \int_0^t \int_R \left[ -(X_s - a_{i+1})(X_s - a_j) \right] \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} \mathbf{1}_{\{a_j < X_s \leq a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\quad + \int_0^t \int_R \left[ -(X_s - a_{i+1})(a_{j+1} - a_j) \right] \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} \mathbf{1}_{\{X_s > a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds.
\end{aligned}$$

To estimate  $|EA_1|$ , we notice that

$$\begin{aligned}
|EA_1| &\leq E \int_0^t \int_{-\infty}^{\infty} (a_{i+1} - a_i)(a_j - X_s) \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{a_j < X_s - y \leq a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\quad + E \int_0^t \int_{-\infty}^{\infty} (a_{i+1} - a_i)(a_{j+1} - a_j) \mathbf{1}_{\{X_s \leq a_i\}} \mathbf{1}_{\{X_s - y > a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&:= A_{11} + A_{12}.
\end{aligned}$$

Let's estimate every term on the righthand side of the above inequality. By the occupation times formula, Fubini theorem, Jensen's inequality and Lemma 2, similar as before, we can do the following crucial computation:

$$\begin{aligned}
A_{11} &= \frac{1}{\sigma^2} E \int_{-\infty}^{a_i} L_t^x \int_{x-a_{j+1}}^{x-a_j} (a_{i+1} - a_i)(a_j - x) \mathbf{1}_{\{|y| < 1\}} n(dy) dx \\
&= \frac{1}{\sigma^2} (a_{i+1} - a_i) E \left[ \int_{-\infty}^{a_i - a_{j+1}} \int_{y+a_j}^{y+a_{j+1}} L_t^x (a_j - x) \mathbf{1}_{\{|y| < 1\}} dx n(dy) \right. \\
&\quad \left. + \int_{a_i - a_{j+1}}^{a_i - a_j} \int_{y+a_j}^{a_i} L_t^x (a_j - x) \mathbf{1}_{\{|y| < 1\}} dx n(dy) \right] \\
&\leq \frac{1}{\sigma^2} (a_{i+1} - a_i)(a_{j+1} - a_j) (\sup_x EL_t^x) \int_{-\infty}^{a_i - a_j} |y| \mathbf{1}_{\{|y| < 1\}} n(dy) \\
&\leq c(t, \sigma)(a_{i+1} - a_i)(a_{j+1} - a_j).
\end{aligned}$$

In the same way, we can have

$$A_{12} \leq \frac{1}{\sigma^2} (a_{i+1} - a_i)(a_{j+1} - a_j) \sup_x E(L_t^x) \int_{-\infty}^{a_i - a_{j+1}} |y| \mathbf{1}_{\{|y| < 1\}} n(dy) \leq c(t, \sigma)(a_{i+1} - a_i)(a_{j+1} - a_j).$$

Therefore, we get

$$|EA_1| \leq c(t, \sigma)((a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2).$$

Using the same method, we can have the similar estimation in the other cases:

$$\begin{aligned}
|EA_2| &\leq E \int_0^t \int_{-\infty}^{\infty} (a_{i+1} - X_s)(a_j - X_s) \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{a_j < X_s - y \leq a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\quad + E \int_0^t \int_{-\infty}^{\infty} (a_{i+1} - X_s)(a_{j+1} - a_j) \mathbf{1}_{\{a_i < X_s \leq a_{i+1}\}} \mathbf{1}_{\{X_s - y > a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\leq c(t, \sigma)((a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2);
\end{aligned}$$

and

$$\begin{aligned}
|EA_3| &\leq E \int_0^t \int_{-\infty}^{\infty} (a_{i+1} - a_i)(X_s - a_j) \mathbf{1}_{\{X_s - y \leq a_i\}} \mathbf{1}_{\{a_j < X_s \leq a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\quad + E \int_0^t \int_{-\infty}^{\infty} (a_{i+1} - a_i)(a_{j+1} - a_j) \mathbf{1}_{\{X_s - y \leq a_i\}} \mathbf{1}_{\{X_s > a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\leq c(t, \sigma)((a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2);
\end{aligned}$$

and

$$\begin{aligned}
|EA_4| &\leq E \int_0^t \int_{-\infty}^{\infty} (X_s - a_{i+1})(X_s - a_j) \mathbf{1}_{\{a_i < X_s - y \leq a_{i+1}\}} \mathbf{1}_{\{a_j < X_s \leq a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\quad + E \int_0^t \int_{-\infty}^{\infty} (X_s - a_{i+1})(a_{j+1} - a_j) \mathbf{1}_{\{a_i < X_s - y \leq a_{i+1}\}} \mathbf{1}_{\{X_s > a_{j+1}\}} \mathbf{1}_{\{|y| < 1\}} n(dy) ds \\
&\leq c(t, \sigma)((a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2).
\end{aligned}$$

So when  $i \neq j$ ,

$$|E(K_2(t, a_{i+1}) - K_2(t, a_i))(K_2(t, a_{j+1}) - K_2(t, a_j))| \leq c(t, \sigma)((a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2).$$

Therefore we proved (44).  $\diamond$

**Proof of Proposition 8:** For  $m > n$ , it is easy to see that

$$\begin{aligned}
&E|\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^2 \\
&= \frac{1}{4} E \left| \sum_{l=2^{m-n}(k-1)+1}^{2^{m-n}k} (\Delta_{2l-1}^{m+1} Z \otimes \Delta_{2l}^{m+1} Z - \Delta_{2l}^{m+1} Z \otimes \Delta_{2l-1}^{m+1} Z) \right|^2 \\
&= \frac{1}{4} \sum_{\substack{i,j=1 \\ i \neq j}}^2 E \sum_{l,r=2^{m-n}(k-1)+1}^{2^{m-n}k} (\Delta_{2l-1}^{m+1} Z^i \Delta_{2l}^{m+1} Z^j - \Delta_{2l}^{m+1} Z^i \Delta_{2l-1}^{m+1} Z^j) \\
&\quad \cdot (\Delta_{2r-1}^{m+1} Z^i \Delta_{2r}^{m+1} Z^j - \Delta_{2r}^{m+1} Z^i \Delta_{2r-1}^{m+1} Z^j) \\
&= \frac{1}{4} \sum_{l,r} \left[ E(\Delta_{2l-1}^{m+1} L_t^x \Delta_{2r-1}^{m+1} L_t^x)(\Delta_{2l}^{m+1} g(x) \Delta_{2r}^{m+1} g(x)) \right. \\
&\quad \left. + (\Delta_{2l-1}^{m+1} g(x) \Delta_{2r-1}^{m+1} g(x)) E(\Delta_{2l}^{m+1} L_t^x \Delta_{2r}^{m+1} L_t^x) \right] \\
&\quad - \frac{1}{4} \sum_{l,r} \left[ E(\Delta_{2l-1}^{m+1} L_t^x \Delta_{2r}^{m+1} L_t^x)(\Delta_{2l}^{m+1} g(x) \Delta_{2r-1}^{m+1} g(x)) \right. \\
&\quad \left. + (\Delta_{2l-1}^{m+1} g(x) \Delta_{2r}^{m+1} g(x)) E(\Delta_{2l}^{m+1} L_t^x \Delta_{2r-1}^{m+1} L_t^x) \right] \\
&\quad - \frac{1}{4} \sum_{l,r} \left[ E(\Delta_{2l}^{m+1} L_t^x \Delta_{2r-1}^{m+1} L_t^x)(\Delta_{2l-1}^{m+1} g(x) \Delta_{2r}^{m+1} g(x)) \right. \\
&\quad \left. + (\Delta_{2l}^{m+1} g(x) \Delta_{2r-1}^{m+1} g(x)) E(\Delta_{2l-1}^{m+1} L_t^x \Delta_{2r}^{m+1} L_t^x) \right] \\
&\quad + \frac{1}{4} \sum_{l,r} \left[ E(\Delta_{2l}^{m+1} L_t^x \Delta_{2r}^{m+1} L_t^x)(\Delta_{2l-1}^{m+1} g(x) \Delta_{2r-1}^{m+1} g(x)) \right. \\
&\quad \left. + (\Delta_{2l}^{m+1} g(x) \Delta_{2r}^{m+1} g(x)) E(\Delta_{2l-1}^{m+1} L_t^x \Delta_{2r-1}^{m+1} L_t^x) \right].
\end{aligned} \tag{45}$$

The main difficulty is to estimate the following expectation which can be derived from Tanaka's formula:

$$\begin{aligned}
& E \left[ \Delta_{2r-1}^{m+1} L_t^x \Delta_{2l-1}^{m+1} L_t^x \right] \\
&= E \left[ \left( L_t(x_{2r-1}^{m+1}) - L_t(x_{2r-2}^{m+1}) \right) \left( L_t(x_{2l-1}^{m+1}) - L_t(x_{2l-2}^{m+1}) \right) \right] \\
&= E \left[ \varphi_t(x_{2r-1}^{m+1}) - \varphi_t(x_{2r-2}^{m+1}) - b \int_0^t 1_{\{x_{2r-2}^{m+1} < X_{s-} \leq x_{2r-1}^{m+1}\}} ds \right. \\
&\quad \left. - \sigma \int_0^t 1_{\{x_{2r-2}^{m+1} < X_{s-} \leq x_{2r-1}^{m+1}\}} dB_s - \int_0^{t+} \int_{|y| < 1} 1_{\{x_{2r-2}^{m+1} < X_{s-} \leq x_{2r-1}^{m+1}\}} y \tilde{N}_p(dsdy) \right. \\
&\quad \left. + (K_1(t, x_{2r-1}^{m+1}) - K_1(t, x_{2r-2}^{m+1})) + (K_2(t, x_{2r-1}^{m+1}) - K_2(t, x_{2r-2}^{m+1})) \right] \\
&\cdot \left[ \varphi_t(x_{2l-1}^{m+1}) - \varphi_t(x_{2l-2}^{m+1}) - b \int_0^t 1_{\{x_{2l-2}^{m+1} < X_{s-} \leq x_{2l-1}^{m+1}\}} ds \right. \\
&\quad \left. - \sigma \int_0^t 1_{\{x_{2l-2}^{m+1} < X_{s-} \leq x_{2l-1}^{m+1}\}} dB_s - \int_0^{t+} \int_{|y| < 1} 1_{\{x_{2l-2}^{m+1} < X_{s-} \leq x_{2l-1}^{m+1}\}} y \tilde{N}_p(dsdy) \right. \\
&\quad \left. + (K_1(t, x_{2l-1}^{m+1}) - K_1(t, x_{2l-2}^{m+1})) + (K_2(t, x_{2l-1}^{m+1}) - K_2(t, x_{2l-2}^{m+1})) \right].
\end{aligned}$$

Firstly, from (13), (15), (16), the Cauchy-Schwarz inequality and the quadratic variation of stochastic integrals, we have

$$\begin{aligned}
& \left| E \left( \varphi_t(x_{2r-1}^{m+1}) - \varphi_t(x_{2r-2}^{m+1}) - b \int_0^t 1_{\{x_{2r-2}^{m+1} < X_{s-} \leq x_{2r-1}^{m+1}\}} ds - \sigma \int_0^t 1_{\{x_{2r-2}^{m+1} < X_{s-} \leq x_{2r-1}^{m+1}\}} dB_s \right) \right. \\
&\quad \left. \cdot \left( \varphi_t(x_{2l-1}^{m+1}) - \varphi_t(x_{2l-2}^{m+1}) - b \int_0^t 1_{\{x_{2l-2}^{m+1} < X_{s-} \leq x_{2l-1}^{m+1}\}} ds - \sigma \int_0^t 1_{\{x_{2l-2}^{m+1} < X_{s-} \leq x_{2l-1}^{m+1}\}} dB_s \right) \right| \\
&\leq c(t) \left[ (1 + 2b + b^2)(x_{2r-1}^{m+1} - x_{2r-2}^{m+1})(x_{2l-1}^{m+1} - x_{2l-2}^{m+1}) + \sigma(x_{2r-1}^{m+1} - x_{2r-2}^{m+1})(x_{2l-1}^{m+1} - x_{2l-2}^{m+1})^{\frac{1}{2}} \right. \\
&\quad \left. + \sigma(x_{2l-1}^{m+1} - x_{2l-2}^{m+1})(x_{2r-1}^{m+1} - x_{2r-2}^{m+1})^{\frac{1}{2}} \right] \\
&\quad + \sigma^2 E \left| \int_0^t 1_{\{x_{2r-2}^{m+1} \leq X_s < x_{2r-1}^{m+1}\}} 1_{\{x_{2l-2}^{m+1} \leq X_s < x_{2l-1}^{m+1}\}} ds \right| \\
&\leq C \left[ \left( \frac{1}{2^{m+1}} \right)^2 w_1(x', x'')^2 + \left( \frac{1}{2^{m+1}} \right)^{\frac{3}{2}} w_1(x', x'')^{\frac{3}{2}} \right] \\
&\quad + \sigma^2 E \left| \int_0^t 1_{\{x_{2r-2}^{m+1} \leq X_s < x_{2r-1}^{m+1}\}} 1_{\{x_{2l-2}^{m+1} \leq X_s < x_{2l-1}^{m+1}\}} ds \right| \\
&\leq \begin{cases} C \left( \frac{1}{2^{m+1}} \right)^{\frac{3}{2}}, & \text{if } r \neq l, \\ C \frac{1}{2^{m+1}}, & \text{if } r = l. \end{cases} \tag{46}
\end{aligned}$$

Here  $C$  is a generic constant and also depends on  $t, b, \sigma, w_1(x', x'')$ . Secondly, recall the fact that  $E(P.M.) = 0$ , if  $P$  is a process of bounded variation and  $M$  is a martingale with mean 0 and at least one of  $M$  and  $P$  is continuous. Note here  $K_1$  is a process of bounded variation. Recall also that the cross-variation of  $\int_0^t 1_{\{a_i < X_{s-} \leq a_{i+1}\}} dB_s$  and the jump parts such as  $\int_0^{t+} \int_{|y| < 1} 1_{\{a_j < X_{s-} \leq a_{j+1}\}} y \tilde{N}_p(dsdy)$  are zero. So using Itô product formula (c.f.[1]), we have

$$\begin{aligned}
& E \int_0^t 1_{\{a_i < X_{s-} \leq a_{i+1}\}} dB_s \cdot (K_1(t, a_{j+1}) - K_1(t, a_j)) = 0, \\
& E \int_0^t 1_{\{a_i < X_{s-} \leq a_{i+1}\}} dB_s \cdot \int_0^{t+} \int_{|y| < 1} 1_{\{a_j < X_{s-} \leq a_{j+1}\}} y \tilde{N}_p(dsdy) = 0,
\end{aligned}$$

$$E \int_0^t \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} ds \cdot \int_0^{t+} \int_{|y| < 1} \mathbf{1}_{\{a_j < X_{s-} \leq a_{j+1}\}} y \tilde{N}_p(dsdy) = 0,$$

$$E \int_0^t \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} dB_s \cdot (K_2(t, a_{j+1}) - K_2(t, a_j)) = 0.$$

Thirdly, by Lemma 9, we can see that

$$\left| E(K_2(t, a_{i+1}) - K_2(t, a_i))(K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2],$$

$$\left| E(K_2(t, a_{i+1}) - K_2(t, a_i)) \int_0^{t+} \int_{|y| < 1} \mathbf{1}_{\{a_j < X_{s-} \leq a_{j+1}\}} y \tilde{N}_p(dsdy) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)^2 + (a_{j+1} - a_j)^2].$$

For other terms, by the Cauchy-Schwarz inequality, it is easy to see that

$$\left| E(\varphi_t(a_{i+1}) - \varphi_t(a_i)) \int_0^{t+} \int_{|y| < 1} \mathbf{1}_{\{a_j < X_{s-} \leq a_{j+1}\}} y \tilde{N}_p(dsdy) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)(a_{j+1} - a_j)^{\frac{1}{2}}];$$

$$\left| E(\varphi_t(a_{i+1}) - \varphi_t(a_i))(K_1(t, a_{j+1}) - K_1(t, a_j)) \right| \leq c(t)[(a_{i+1} - a_i)(a_{j+1} - a_j)];$$

$$\left| E(\varphi_t(a_{i+1}) - \varphi_t(a_i))(K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)(a_{j+1} - a_j)^{\frac{1}{2}}];$$

$$\left| E \int_0^t \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} ds (K_1(t, a_{j+1}) - K_1(t, a_j)) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)(a_{j+1} - a_j)];$$

$$\left| E \int_0^t \mathbf{1}_{\{a_i < X_{s-} \leq a_{i+1}\}} ds (K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)(a_{j+1} - a_j)^{\frac{1}{2}}];$$

$$\left| E(K_1(t, a_{i+1}) - K_1(t, a_i)) \int_0^{t+} \int_{|y| < 1} \mathbf{1}_{\{a_j < X_{s-} \leq a_{j+1}\}} y \tilde{N}_p(dsdy) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)(a_{j+1} - a_j)^{\frac{1}{2}}];$$

$$\left| E(K_1(t, a_{i+1}) - K_1(t, a_i))(K_2(t, a_{j+1}) - K_2(t, a_j)) \right| \leq c(t, \sigma)[(a_{i+1} - a_i)(a_{j+1} - a_j)^{\frac{1}{2}}].$$

Thus

$$E \left[ \Delta_{2r-1}^{m+1} L_t^x \Delta_{2l-1}^{m+1} L_t^x \right] \leq \begin{cases} C \left( \frac{1}{2^{m+1}} \right)^{\frac{3}{2}}, & \text{if } r \neq l, \\ C \frac{1}{2^{m+1}}, & \text{if } r = l. \end{cases} \quad (47)$$

The other terms in (45) can be treated similarly, therefore

$$E |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^2 \leq C \left[ 2^{m-n} \left( \frac{1}{2^{m+1}} \right)^{1+2h} + 2^{2(m-n)} \left( \frac{1}{2^{m+1}} \right)^{\frac{3}{2}+2h} \right].$$

Hence, for  $2 \leq \theta < 4$ , by Jensen's inequality,

$$\begin{aligned} E |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^{\frac{\theta}{2}} &\leq \left( E |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^2 \right)^{\frac{\theta}{4}} \\ &\leq C \left[ 2^{m-n} \left( \frac{1}{2^{m+1}} \right)^{1+2h} + 2^{2(m-n)} \left( \frac{1}{2^{m+1}} \right)^{\frac{3}{2}+2h} \right]^{\frac{\theta}{4}} \\ &\leq C \left[ 2^{(m-n)\frac{\theta}{4}} \left( \frac{1}{2^{m+1}} \right)^{\frac{\theta}{4} + \frac{1}{2}h\theta} + 2^{(m-n)\frac{\theta}{2}} \left( \frac{1}{2^{m+1}} \right)^{\frac{3\theta}{8} + \frac{1}{2}h\theta} \right] \\ &\leq C \left[ \left( \frac{1}{2^n} \right)^{\frac{\theta}{4}} \left( \frac{1}{2^m} \right)^{\frac{1}{2}h\theta} + \left( \frac{1}{2^n} \right)^{\frac{\theta}{2}} \left( \frac{1}{2^m} \right)^{\frac{1}{2}h\theta - \frac{1}{8}\theta} \right], \end{aligned}$$

where  $C$  is a generic constant and also depends on  $\theta$ ,  $h$ ,  $w_1(x', x'')$ , and  $c$ .  $\diamond$

**Corollary 10** *Under the same assumption as in Proposition 8, we have*

$$\sup_m \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^{\frac{\theta}{2}} < \infty \quad a.s.$$

**Proof:** From the Minkowski inequality,

$$\begin{aligned}
& \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \\
& \leq \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n} - \mathbf{Z}(m-1)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} + \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m-1)_{x_{k-1}^n, x_k^n} - \mathbf{Z}(m-2)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \\
& \quad + \cdots + \left( \sum_{k=1}^{2^n} |\mathbf{Z}(1)_{x_{k-1}^n, x_k^n} - \mathbf{Z}(0)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} + \left( \sum_{k=1}^{2^n} |\mathbf{Z}(0)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \\
& \leq \sum_{m=1}^{\infty} \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n} - \mathbf{Z}(m-1)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} + \left( \sum_{k=1}^{2^n} |\mathbf{Z}(0)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}}. \tag{48}
\end{aligned}$$

Then it is easy to see from (48), Jensen's inequality, (41), (42) and (39)

$$\begin{aligned}
& E \sup_m \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \\
& \leq E \sum_{m=1}^{\infty} \left[ \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n} - \mathbf{Z}(m-1)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \right. \\
& \quad \left. + \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} E \left( \sum_{k=1}^{2^n} |\mathbf{Z}(0)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \right] \\
& \leq \sum_{m=1}^{\infty} \left[ \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \left( E \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n} - \mathbf{Z}(m-1)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \right. \\
& \quad \left. + \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} (2^{n+\frac{(-1-2n)\theta}{2}} w(x', x'') h\theta)^{\frac{2}{\theta}} \right] \\
& \leq C \sum_{n=m}^{\infty} n^{\frac{2}{\theta}\gamma} \sum_{m=1}^{\infty} \left( \frac{1}{2^{n+m}} \right)^{\left( \frac{h\theta-1}{2} \right) \frac{2}{\theta}} \\
& \quad + C \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \sum_{m=1}^{\infty} \left[ \left( \frac{1}{2^n} \right)^{\left( \frac{\theta}{4} - \frac{1}{2} \right) \frac{2}{\theta}} \left( \frac{1}{2^m} \right)^{\left( \frac{1}{2} h\theta - \frac{1}{2} \right) \frac{2}{\theta}} + \left( \frac{1}{2^n} \right)^{1-\frac{2}{\theta}} \left( \frac{1}{2^m} \right)^{\left( \frac{1}{2} h\theta - \frac{1}{8} \theta \right) \frac{2}{\theta}} \right] \\
& \quad + C \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \left( \frac{1}{2^n} \right)^{\frac{2(\theta-1)}{\theta}} \\
& < \infty,
\end{aligned}$$

as  $2 < \theta < 4$ ,  $h\theta > 1$ , where  $C$  is a generic constant and also depends on  $\theta$ ,  $h$ ,  $w_1(x', x'')$ , and  $c$ . Therefore,

$$\sup_m \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} < \infty \text{ a.s.}$$

However, it is easy to see as  $\theta > 2$ ,

$$\left( \sup_m \sum_{n=1}^{\infty} n^{\gamma} \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} \leq \sup_m \sum_{n=1}^{\infty} n^{\frac{2}{\theta}\gamma} \left( \sum_{k=1}^{2^n} |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}|^{\frac{\theta}{2}} \right)^{\frac{2}{\theta}} < \infty \text{ a.s.}$$

So the claim follows.  $\diamond$

**Theorem 11** *Let  $L_t^x$  be the local time of the time homogeneous Lévy process  $X_t$  given by (2). Assume  $2 \leq q < 4$ ,  $\sigma \neq 0$  and the Lévy measure  $n(dy)$  satisfies (12). Then for any  $\theta \in (q, 4)$ , the continuous process  $Z_x = (L_t^x, g(x))$  satisfying (29), there exists a unique  $\mathbf{Z}^i$  on  $\Delta$  taking values in  $(\mathbb{R}^2)^{\otimes i}$  ( $i = 1, 2$ ) such that*

$$\sum_{i=1}^2 \sup_D \left( \sum_l |\mathbf{Z}(m)_{x_{l-1}, x_l}^i - \mathbf{Z}_{x_{l-1}, x_l}^i|^{\frac{\theta}{4}} \right)^{\frac{4}{\theta}} \rightarrow 0,$$

both almost surely and in  $L^1(\Omega, \mathcal{F}, \mathcal{P})$  as  $m \rightarrow \infty$ . In particular, when  $2 \leq q < 3$ ,  $\mathbf{Z} = (1, \mathbf{Z}^1, \mathbf{Z}^2)$  is the canonical geometric rough path associated to  $Z$ . Moreover,  $\mathbf{Z}_{a,b}^1 = Z_b - Z_a$ .

**Proof:** The convergence of  $\mathbf{Z}(m)^1$  to  $\mathbf{Z}^1$  is actually the result of Theorem 6. In the following we will prove  $\mathbf{Z}(m)_{a,b}^2$  converges in the  $\theta$ -variation distance. By Proposition 4.1.2 in [23],

$$\begin{aligned} & E \sup_D \sum_l |\mathbf{Z}(m+1)_{x_{l-1}, x_l}^2 - \mathbf{Z}(m)_{x_{l-1}, x_l}^2|^{\frac{\theta}{2}} \\ & \leq C(\theta, \gamma) E \left( \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^1 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1|^\theta \right)^{\frac{1}{2}} \\ & \quad \cdot \left( \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} \left( |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^1|^\theta + |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1|^\theta \right) \right)^{\frac{1}{2}} \\ & \quad + C(\theta, \gamma) E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^2 - \mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^2|^{\frac{\theta}{2}} \\ & := A + B. \end{aligned}$$

We will estimate part A, B respectively. First from the Cauchy-Schwarz inequality, (34) and (36), we know

$$\begin{aligned} A & \leq C \left( E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} \left( |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^1 - \mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta + |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1 - \mathbf{Z}_{x_{k-1}^n, x_k^n}^1|^\theta \right) \right)^{\frac{1}{2}} \\ & \quad \cdot \left( E \sum_{n=1}^{\infty} n^\gamma \sum_{k=1}^{2^n} \left( |\mathbf{Z}(m+1)_{x_{k-1}^n, x_k^n}^1|^\theta + |\mathbf{Z}(m)_{x_{k-1}^n, x_k^n}^1|^\theta \right) \right)^{\frac{1}{2}} \\ & \leq C \left( \frac{1}{2^m} \right)^{\frac{h\theta-1}{4}} \left( \sum_{n=1}^{\infty} n^\gamma \left( \frac{1}{2^n} \right)^{h\theta-1} \right)^{\frac{1}{2}}. \end{aligned}$$

Secondly from Proposition 7 and Proposition 8, we know

$$\begin{aligned} B & \leq C \sum_{n=m}^{\infty} n^\gamma \left( \frac{1}{2^{m+n}} \right)^{\frac{h\theta-1}{2}} + C \left[ \sum_{n=1}^{m-1} n^\gamma \left( \frac{1}{2^n} \right)^{\frac{\theta}{4}-1} \left( \frac{1}{2^m} \right)^{\frac{h\theta}{2}} + \sum_{n=1}^{m-1} n^\gamma \left( \frac{1}{2^n} \right)^{\frac{\theta}{2}-1} \left( \frac{1}{2^m} \right)^{\frac{h\theta}{2} - \frac{\theta}{8}} \right] \\ & \leq C \left[ \left( \frac{1}{2^m} \right)^{\frac{h\theta-1}{2}} + \left( \frac{1}{2^m} \right)^{\frac{h\theta-1}{2}} + \left( \frac{1}{2^m} \right)^{\frac{h\theta}{2} - \frac{\theta}{8}} \right], \end{aligned}$$

as  $q < \theta < 4$ , and  $h\theta > 1$ . So



$$E \sup_D \sum_l |\mathbf{Z}(m+1)_{x_{l-1}, x_l}^2 - \mathbf{Z}(m)_{x_{l-1}, x_l}^2|^{\frac{\theta}{2}} \leq C \left[ \left( \frac{1}{2^m} \right)^{\frac{h\theta-1}{4}} + \left( \frac{1}{2^m} \right)^{\frac{h\theta}{2} - \frac{\theta}{8}} \right].$$

Similar to the proof of Theorem 6, we can easily deduce that  $(\mathbf{Z}(m)^2)_{m \in \mathbb{N}}$  is a Cauchy sequence in the  $\theta$ -variation distance. So when  $m \rightarrow \infty$ , it has a limit, denote it by  $\mathbf{Z}^2$ . And from the completeness under the  $\theta$ -variation distance (Lemma 3.3.3 in [23]),  $\mathbf{Z}^2$  is also of finite  $\theta$ -variation. The theorem is proved.  $\diamond$

**Remark 12** *We would like to point out that the above method does not seem to work for two arbitrary functions  $f$  of  $p$ -variation and  $g$  of  $q$ -variation ( $2 < p, q < 3$ ) to define a rough path  $Z_x = (f(x), g(x))$ . However the special property (47) of local times makes our analysis work. A similar method was used in [23] for fractional Brownian motion with the help of long-time memory. Here (47) serves a similar role of the long-time memory as in [23].*

In the following, we will only consider the case that  $2 \leq q < 3$  and take  $q < \theta < 3$ .

As local time  $L_t^x$  has a compact support in  $x$  for each  $\omega$  and  $t$ , so we can define integral of local time directly in  $R$ . For this, we take  $[x', x'']$  covering the support of  $L_t^x$ . From Chen's identity, it's easy to know that for any  $(a, b) \in \Delta$ ,

$$\mathbf{Z}_{a,b}^2 = \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} (\mathbf{Z}_{x_i, x_{i+1}}^2 + \mathbf{Z}_{a, x_i}^1 \otimes \mathbf{Z}_{x_i, x_{i+1}}^1).$$

In particular,

$$\begin{aligned} (\mathbf{Z}_{a,b}^2)_{2,1} &= \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1} + (\mathbf{Z}_{a, x_i}^1 \otimes \mathbf{Z}_{x_i, x_{i+1}}^1)_{2,1}) \\ &= \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1} + (g(x_i) - g(a))(L_t^{x_{i+1}} - L_t^{x_i})) \end{aligned}$$

exists. Here  $(\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1}$  means lower-left element of the  $2 \times 2$  matrix  $\mathbf{Z}_{x_i, x_{i+1}}^2$ . It turns out that

$$\begin{aligned} &\lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1} + g(x_i)(L_t^{x_{i+1}} - L_t^{x_i})) \\ &= \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1} + (g(x_i) - g(a))(L_t^{x_{i+1}} - L_t^{x_i})) + g(a)(L_t^b - L_t^a) \end{aligned}$$

exists. Denote this limit by  $\int_a^b g(x) dL_t^x$ . Similarly, we can define  $\int_a^b L_t^x dL_t^x$ . Therefore we have the following corollary.

**Corollary 13** *Assume all conditions of Theorem 11. Then the local time  $L_t^x$  is a geometrical rough path of roughness  $p$  in  $x$  for any  $t \geq 0$  a.s. for any  $p > 2$ , and  $(a, b) \in \Delta$ ,*

$$\int_a^b L_t^x dL_t^x = \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{1,1} + L(x_i)(L_t^{x_{i+1}} - L_t^{x_i})).$$

Moreover, if  $g$  is a continuous function with bounded  $q$ -variation,  $2 \leq q < 3$ , the integral  $\int_a^b g(x) dL_t^x$  is defined by

$$\int_a^b g(x) dL_t(x) = \lim_{m(D_{[a,b]}) \rightarrow 0} \sum_{i=0}^{r-1} ((\mathbf{Z}_{x_i, x_{i+1}}^2)_{2,1} + g(x_i)(L_t^{x_{i+1}} - L_t^{x_i})). \quad (49)$$

## 4 Convergence of the rough path integrals and applications to extensions of Itô's formula

In this section we will apply the Young integral and rough path integral of local time defined in sections 2 and 3 to prove a useful extension to Itô's formula. First we consider some convergence result of the rough path integrals.

Let  $Z_j(x) := (L_t^x, g_j(x))$ , where  $g_j(\cdot)$  is of bounded  $q$ -variation uniformly in  $j$  for  $2 \leq q < 3$ , and when  $j \rightarrow \infty$ ,  $g_j(x) \rightarrow g(x)$  for all  $x \in R$ . Repeating the above argument, for each  $j$ , we can find the canonical geometric rough path  $\mathbf{Z}_j = (1, \mathbf{Z}_j^1, \mathbf{Z}_j^2)$  associated to  $Z_j$ , and the smooth rough path  $\mathbf{Z}_j(m) = (1, \mathbf{Z}_j(m)^1, \mathbf{Z}_j(m)^2)$ . Actually,  $(\mathbf{Z}_j)_{a,b}^1 \rightarrow \mathbf{Z}_{a,b}^1$  in the sense of the uniform topology, and also in the sense of the  $\theta$ -variation topology. As for  $(\mathbf{Z}_j)_{a,b}^2$ , we can easily see that

$$d_{2,\theta}((\mathbf{Z}_j)^2, \mathbf{Z}^2) \leq d_{2,\theta}((\mathbf{Z}_j)^2, (\mathbf{Z}_j(m))^2) + d_{2,\theta}((\mathbf{Z}_j(m))^2, \mathbf{Z}(m)^2) + d_{2,\theta}(\mathbf{Z}(m)^2, \mathbf{Z}^2). \quad (50)$$

From Theorem 11, we know that  $d_{2,\theta}(\mathbf{Z}(m)^2, \mathbf{Z}^2) \rightarrow 0$  as  $m \rightarrow \infty$ . Moreover, it is not difficult to see from the proofs of Propositions 7, 8, and Theorem 11,  $d_{2,\theta}((\mathbf{Z}_j)^2, (\mathbf{Z}_j(m))^2) \rightarrow 0$  as  $m \rightarrow \infty$  uniformly in  $j$ . So for any given  $\varepsilon > 0$ , there exists an  $m_0$  such that when  $m \geq m_0$ ,  $d_{2,\theta}(\mathbf{Z}(m)^2, \mathbf{Z}^2) < \frac{\varepsilon}{3}$ ,  $d_{2,\theta}((\mathbf{Z}_j)^2, (\mathbf{Z}_j(m))^2) < \frac{\varepsilon}{3}$  for all  $j$ . In particular,  $d_{2,\theta}(\mathbf{Z}(m_0)^2, \mathbf{Z}^2) < \frac{\varepsilon}{3}$ ,  $d_{2,\theta}((\mathbf{Z}_j)^2, (\mathbf{Z}_j(m_0))^2) < \frac{\varepsilon}{3}$  for all  $j$ . It's easy to prove for such  $m_0$ ,  $d_{2,\theta}((\mathbf{Z}_j(m_0))^2, \mathbf{Z}(m_0)^2) < \frac{\varepsilon}{3}$  for sufficiently large  $j$ . Replacing  $m$  by  $m_0$  in (50), we can get  $d_{2,\theta}((\mathbf{Z}_j)^2, \mathbf{Z}^2) < \varepsilon$  for sufficiently large  $j$ . Then by (49) and the definition of  $\int_a^b g_j(x) dL_t^x$ , we know that  $\int_a^b g_j(x) dL_t^x \rightarrow \int_a^b g(x) dL_t^x$  as  $j \rightarrow \infty$ . Note now that the local time  $L_t^x$  has a compact support in  $x$  a.s. So it is easy to see from taking  $[x', x'']$  covering the support of  $L_t^x$  that the above construction of the integrals and the convergence can work for the integrals on  $R$ . Therefore we have

**Proposition 14** *Let  $Z_j(x) := (L_t^x, g_j(x))$ ,  $Z(x) := (L_t^x, g(x))$ , where  $g_j(\cdot)$ ,  $g(\cdot)$  are continuous and of bounded  $q$ -variation uniformly in  $j$ ,  $2 \leq q < 3$ . Assume  $g_j(x) \rightarrow g(x)$  as  $j \rightarrow \infty$  for all  $x \in R$ . Then as  $j \rightarrow \infty$ ,  $\mathbf{Z}_j(\cdot) \rightarrow \mathbf{Z}(\cdot)$  a.s. in the  $\theta$ -variation distance. In particular, as  $j \rightarrow \infty$ ,  $\int_{-\infty}^{\infty} g_j(x) dL_t^x \rightarrow \int_{-\infty}^{\infty} g(x) dL_t^x$  a.s.*

Now for any  $g$  being continuous and of bounded  $q$ -variation ( $2 \leq q < 3$ ), define

$$g_j(x) = \int_{-\infty}^{\infty} k^j(x-y)g(y)dy,$$

where  $k^j$  is the mollifier given by

$$k^j(x) = \begin{cases} cje^{\frac{1}{(jx-1)^2-1}}, & \text{if } x \in (0, \frac{2}{j}), \\ 0, & \text{otherwise.} \end{cases}$$

Here  $c$  is a constant such that  $\int_0^2 k^j(x)dx = 1$ . It is well known that  $g_j$  is a smooth function and  $g_j(x) \rightarrow g(x)$  as  $j \rightarrow \infty$  for each  $x$ . So the integral  $\int_{-\infty}^{\infty} g_j(x) dL_t^x$  is a Riemann integral for the smooth function  $g_j(x)$ . Moreover, Proposition 14 guarantees that  $\int_{-\infty}^{\infty} g_j(x) dL_t^x \rightarrow \int_{-\infty}^{\infty} g(x) dL_t^x$  a.s.

In the following, we will show that Proposition 14 is true for  $g$  being of bounded  $q$ -variation ( $2 \leq q < 3$ ) without assuming  $g$  being continuous. Note that a function with bounded  $q$ -variation ( $q \geq 1$ ) may have at most countable discontinuities. Using the method in [33], we will define the rough path integral  $\int_{x'}^{x''} L_t^x dg(x)$ . Here we assume  $g(x)$  is càdlàg in  $x$ .

First we can define a map

$$\tau_\delta(\cdot) : [x', x''] \rightarrow [x', x'' + \delta \sum_{n=1}^{\infty} |j(x_n)|^q],$$

in the following way:

$$\tau_\delta(x) = x + \delta \sum_{n=1}^{\infty} |j(x_n)|^q \mathbf{1}_{\{x_n \leq x\}}(x),$$

where  $j(x_i) := G(x_i) - G(x_i-)$ ,  $\{x_i\}_{i=1}^{\infty}$  are the discontinuous points of  $G$  inside  $[x', x'']$ ,  $\delta > 0$ . The map  $\tau_\delta(\cdot) : [x', x''] \rightarrow [x', \tau_\delta(x'')]$  extends the space interval into the one where we can define the continuous path  $G_\delta(y)$  from a càdlàg path  $G$  by:

$$G_\delta(y) = \begin{cases} G(x) & \text{if } y = \tau_\delta(x), \\ G(x_n-) + (y - \tau_\delta(x_n-))j(x_n)\delta^{-1}|j(x_n)|^{-q} & \text{if } y \in [\tau_\delta(x_n-), \tau_\delta(x_n)). \end{cases} \quad (51)$$

Take  $G$  to be  $g$  and  $L_t$ , we can define  $g_\delta$  and  $L_{t,\delta}$  respectively. As  $L_t^x$  is continuous, we can easily see that  $L_{t,\delta}(y) := L_{t,\delta}(\tau_\delta(x)) = L_t^x$ .

**Theorem 15** *Let  $g(x)$  be a càdlàg path with bounded  $q$ -variation ( $2 \leq q < 3$ ). Then*

$$\int_{x'}^{x''} L_t^x dg(x) = \int_{x'}^{\tau_\delta(x'')} L_{t,\delta}(y) dg_\delta(y). \quad (52)$$

**Proof:** First it is easy to see that the integral  $\int_{x'}^{\tau_\delta(x'')} L_{t,\delta}(y) dg_\delta(y)$  is a rough path integral that can be defined by the method of last section. Now note that at any discontinuous point  $x_r$ ,

$$\int_{x_r-}^{x_r} L_t^x dg(x) = L_t(x_r)(g(x_r) - g(x_r-))$$

and

$$\sum_r ((Z_\delta)_{\tau_\delta(x_r-), \tau_\delta(x_r)}^2)_{2,1} = \sum_r \int_{\tau_\delta(x_r-)}^{\tau_\delta(x_r)} (L_{t,\delta}(y) - L_{t,\delta}(\tau_\delta(x_r-))) dg_\delta(y) = 0,$$

where  $Z_\delta(y) := (L_{t,\delta}(y), g_\delta(y))$ . Thus

$$\begin{aligned} & \sum_r L_t^{x_r}(g(x_r) - g(x_r-)) \\ &= \sum_r L_{t,\delta}(\tau_\delta(x_r-))(g_\delta(\tau_\delta(x_r)) - g_\delta(\tau_\delta(x_r-))) \\ &= \sum_r \left[ L_{t,\delta}(\tau_\delta(x_r-))(g_\delta(\tau_\delta(x_r)) - g_\delta(\tau_\delta(x_r-))) + ((Z_\delta)_{\tau_\delta(x_r-), \tau_\delta(x_r)}^2)_{2,1} \right] \\ &< \infty, \end{aligned}$$

so

$$\begin{aligned} \int_{\tau_\delta(x_r-)}^{\tau_\delta(x_r)} L_{t,\delta}(y) dg_\delta(y) &= L_{t,\delta}(\tau_\delta(x_r-))(g_\delta(\tau_\delta(x_r)) - g_\delta(\tau_\delta(x_r-))) \\ &= L_t(x_r)(g(x_r) - g(x_r-)). \end{aligned}$$

Thus

$$\int_{x_r-}^{x_r} L_t^x dg(x) = \int_{\tau_\delta(x_r-)}^{\tau_\delta(x_r)} L_{t,\delta}(y) dg_\delta(y).$$

Now define  $g(x) = \tilde{g}(x) + h(x)$ , where  $h(x) = \sum_{x_r \leq x} (g(x_r) - g(x_{r-}))$ . Then  $\tilde{g}$  is the continuous part of  $g$  and  $h$  is the jump part of  $g$ . Moreover,  $\tilde{g}$  satisfies the  $q$ -variation condition. So  $\int_{x'}^{x''} L_t^x d\tilde{g}(x)$  can be well defined as in the last section. For  $h$ , we can define  $h_\delta$  by taking  $G = h$  in (51). So the integral  $\int_{x'}^{x''} L_t^x dh(x)$  can be well defined by the followings:

$$\begin{aligned} \int_{x'}^{\tau_\delta(x'')} L_{t,\delta}(y) dh_\delta(y) &= \sum_r \int_{\tau_\delta(x_{r-})}^{\tau_\delta(x_r)} L_{t,\delta}(y) dh_\delta(y) = \sum_r L_t(x_r)(h(x_r) - h(x_{r-})) \\ &= \sum_r L_t(x_r)(g(x_r) - g(x_{r-})) = \sum_r \int_{x_{r-}}^{x_r} L_t^x dh(x) = \int_{x'}^{x''} L_t^x dh(x). \end{aligned}$$

Therefore

$$\begin{aligned} \int_{x'}^{x''} L_t^x dg(x) &= \int_{x'}^{x''} L_t^x d\tilde{g}(x) + \int_{x'}^{x''} L_t^x dh(x) \\ &= \int_{x'}^{\tau_\delta(x'')} L_{t,\delta}(y) d\tilde{g}_\delta(y) + \int_{x'}^{\tau_\delta(x'')} L_{t,\delta}(y) dh_\delta(y) \\ &= \int_{x'}^{\tau_\delta(x'')} L_{t,\delta}(y) dg_\delta(y). \end{aligned}$$

◇

Similarly to Proposition 14, we have

**Proposition 16** *Under the condition of Proposition 14, as  $j \rightarrow \infty$ ,  $\int_{-\infty}^{\infty} L_t^x dg_j(x) \rightarrow \int_{-\infty}^{\infty} L_t^x dg(x)$  a.s. for such  $g$  with bounded  $q$ -variation ( $2 \leq q < 3$ ).*

**Proof:** Define  $F_j(x) := (g_j - g)(x)$ , so  $F_j(x) \rightarrow 0$  as  $j \rightarrow \infty$ , for all  $x$ . It's easy to see that  $F_{j,\delta}(x) \rightarrow 0$  as  $j \rightarrow \infty$ , for all  $x$ . From the above theorem and Proposition 14, we have

$$\int_{x'}^{x''} L_t^x d(g_j - g)(x) = \int_{x'}^{x''} L_t^x dF_j(x) = \int_{x'}^{\tau_\delta(x'')} L_{t,\delta} dF_{j,\delta}(y) \rightarrow 0, \quad \text{as } j \rightarrow \infty.$$

Then the proposition follows easily. ◇

With the definition of the integral of local time and the convergence results of the integrals, using standard smoothing procedure, we can prove the following theorem:

**Theorem 17** *Let  $X = (X_t)_{t \geq 0}$  be a time homogeneous Lévy process with  $\sigma \neq 0$  and Lévy measure  $n(dy)$  satisfying (12), and  $f : \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous function and have left derivative  $\nabla^- f(x)$  being left continuous and locally bounded. Assume  $\nabla^- f(x)$  is of bounded  $q$ -variation, where  $1 \leq q < 3$ , then  $P$ -a.s.*

$$\begin{aligned} f(X_t) &= f(X_0) + \int_0^t \nabla^- f(X_s) dX_s - \int_{-\infty}^{\infty} \nabla^- f(x) dx L_t^x \\ &\quad + \sum_{0 \leq s \leq t} [f(X_s) - f(X_{s-}) - \Delta X_s \nabla^- f(X_{s-})], \quad 0 \leq t < \infty. \end{aligned} \quad (53)$$

Here the integral  $\int_{-\infty}^{\infty} \nabla^- f(x) dx L_t^x$  is a Lebesgue-Stieltjes integral when  $q = 1$ , a Young integral when  $1 < q < 2$  and a Lyons' rough path integral when  $2 \leq q < 3$  respectively.

### Acknowledgements

It is our great pleasure to thank K.D. Elworthy, W.V. Li, T. Lyons, Z.M. Ma, S.G. Peng, Z.M. Qian for stimulating conversations. CF would like to acknowledge the support of National Basic Research Program of China (973 Program No. 2007CB814903) and National Natural Science Foundation of China (No. 70671069), Loughborough University and the London Mathematical Society that enabled her to visit Loughborough University.

### References

1. D. Applebaum, *Lévy processes and stochastic calculus*, Cambridge Studies in Advanced Mathematics No.93, Cambridge University Press (2004).
2. M. T. Barlow, *Necessary and sufficient conditions for the continuity of local time of Lévy processes*, Ann. Prob. 16 (1988), 1389-1427.
3. N. Bouleau and M. Yor, *Sur la variation quadratique des temps locaux de certaines semimartingales*, C.R.Acad. Sci. Paris, Ser.I Math 292 (1981), 491-494.
4. E. S. Boylan, *Local times for a class of Markov processes*, Illinois J. Math., 8: 19-39, 1694.
5. N. Eisenbaum, *Integration with respect to local time*, Potential analysis 13 (2000), 303-328.
6. N. Eisenbaum, *Local time-space stochastic calculus for Lévy processes*, Stochastic processes and their applications 116 (2006), 757-778.
7. N. Eisenbaum and A. Kyprianou, *On the parabolic generator of a general one-dimensional Lévy process*, Elect. Comm. in Probab. 13 (2008), 198-209.
8. K. D. Elworthy, A. Truman and H. Z. Zhao, *Generalized Itô Formulae and space-time Lebesgue-Stieltjes integrals of local times*, Séminaire de Probabilités, Vol XL, Lecture Notes in Mathematics 1899, Springer-Verlag, (2007), 117-136.
9. C. R. Feng and H. Z. Zhao, *Two-parameter  $p, q$ -variation Path and Integration of Local Times*, Potential Analysis, Vol 25 (2006), 165-204.
10. C.R. Feng and H.Z. Zhao, *Rough Path Integral of Local Time*, C.R. Acad. Sci. Paris, Ser. I 346 (2008), 431-434.
11. F. Flandoli, F. Russo and J. Wolf, *Some SDEs with distributional drift. Part II: Lyons-Zheng structure, Itô's formula and semimartingale characterization*, Random Oper. Stochastic Equations, Vol. 12, No. 2, (2004), 145-184.
12. H. Föllmer, P. Protter and A. N. Shiryaev, *Quadratic covariation and an extension of Itô's Formula*, Bernoulli 1 (1995), 149-169.
13. H. Föllmer and P. Protter, *On Itô's Formula for multidimensional Brownian motion*, Probability Theory and Related Fields 116 (2000), 1-20.
14. R. K. Gettoor and H. Kesten, *Continuous of local times for Markov processes*. Compositio Math., 24: 277-303, 1972.
15. N. Ikeda and S. Watanabe, *Stochastic Differential Equations and Diffusion Processes*, 2nd Edition, North-Holland Publ. Co., Amsterdam Oxford New York; Kodansha Ltd., Tokyo, 1981.
16. K. Itô *Stochastic Integral*, Proc. Imperial Acad. Tokyo 20, 519-524, 1944.
17. I. Karatzas and S. E. Shreve, *Brownian Motion and Stochastic Calculus, Second Edition*, Springer-Verlag: New York, 1998.
18. H. Kunita and S. Watanabe, *On square-integrable martingales*, Nagoya Math. J., Vol 30 (1967), 209-245.
19. A. Lejay, *An Introduction to Rough Paths*, Séminaire de probabilités XXXVII, Lecture Notes in Mathematics, Vol. 1832, 1-59, Springer-Verlag, 2003.
20. P. Lévy, *Processus Stochastiques et Mouvement Brownien*, Gauthier-Villars, Paris, (1948).
21. T. Lyons, *Differential equations driven by rough signals (I): An extension of an inequality of L. C. Young*, Math. Res. Lett., 1, 451-64, 1994.
22. T. Lyons, *Differential equations driven by rough signals*, Rev. Mat. Iberoamer., 14, 215-310, 1998.
23. T. Lyons and Z. Qian, *System Control and Rough Paths*, Clarendon Press Oxford, 2002.
24. M. B. Marcus and J. Rosen, *Sample path properties of the local times of strongly symmetric Markov processes via Gaussian processes*, Ann. Prob. Vol. 20 (1992), 1603-1684.
25. M. B. Marcus and J. Rosen,  *$p$ -variation of the local times of symmetric stable processes and of Gaussian processes with stationary increments*, Ann. Prob. Vol.20 (1992), 1685-1713.

26. P. A. Meyer, *Un cours sur les intégrales stochastiques*, Sémin. Probab 10, Lecture Notes in Math, No. 511, Springer-Verlag (1976), 245-400.
27. S. Moret and D. Nualart, *Generalization of Itô's formula for smooth nondegenerate martingales*, Stochastic Process. Appl. 91, 115-149, 2001.
28. E. Perkins, *Local time is a semimartingale*, Z. Wahrsch. Verw. Gebiete, Vol. 60 (1982), 79-117.
29. D. Revuz and M. Yor, *Continuous Martingales and Brownian Motion, Second Edition*, (Springer-Verlag: Berlin, Heidelberg, 1994).
30. L. C. G. Rogers and J. B. Walsh, *Local time and stochastic area integrals*, Annals of Probab. 19(2) (1991), 457-482.
31. H. Tanaka, *Note on continuous additive functionals of the 1-dimensional Brownian path*, Z. Wahrscheinlichkeitstheorie and Verw Gebiete 1 (1963), 251-257.
32. A. T. Wang, *Generalized Itô's formula and additive functionals of Brownian motion*, Z. Wahrscheinlichkeitstheorie and Verw Gebiete, 41(1977), 153-159.
33. D. R. E. Williams, *Path-wise solutions of stochastic differential equations driven by Levy processes*, Rev. Mat. Iberoam, 17 (2001), 295-329.
34. L. C. Young, *An inequality of Hölder type, connected with Stieltjes integration*, Acta Math., 67 (1936), 251-282.