Intrinsic Coupling between Current and Domain Wall Motion in (Ga,Mn)As

Kjetil Magne Dørheim Hals, Anh Kiet Nguyen, and Arne Brataas

Department of Physics, Norwegian University of Science and Technology, NO-7491, Trondheim, Norway

We consider current-induced domain wall motion and, the reciprocal process, moving domain wallinduced current. The associated Onsager coefficients are expressed in terms of scattering matrices. Uncommonly, in (Ga,Mn)As, the effective Gilbert damping coefficient α_w and the effective out-ofplane spin transfer torque parameter β_w are dominated by spin-orbit interaction in combination with scattering off the domain wall, and not scattering off extrinsic impurities. Numerical calculations give $\alpha_w \sim 0.01$ and $\beta_w \sim 1$ in dirty (Ga,Mn)As. The extraordinary large β_w parameter allows experimental detection of current or voltage induced by domain wall motion in (Ga,Mn)As.

The principle of giant magneto resistance is used to detect magnetic information. Large currents in magnetic nanostructures can manipulate the magnetization via spin transfer torques [1]. A deeper knowledge of the coupled out-of-equilibrium quasi-particle and magnetization dynamics is needed to precisely control and utilize current-induced spin transfer torques.

The magnetization relaxes towards its equilibrium configuration by releasing magnetic moments and energy into reservoirs. This friction process is usually described by the Gilbert damping constant α in the Landau-Lifshitz-Gilbert (LLG) equation. Spins traversing a magnetic domain wall exert an in-plane and an out-of-plane torque on the wall [2]. In *dirty* systems, when the domain wall is wider than the mean-free-path, the out-of-plane torque, often denoted the non-adiabatic torque, is parameterized by the so-called β -factor [2]. The Gilbert damping coefficient α , the in-plane spin-transfer torque, and the out-of-plane torque coefficient β determine how the magnetization is influenced by an applied current, e.g. the current-induced Walker domain wall drift velocity is proportional to β/α [2, 3, 4]. Scattering off impurities are important for α and β [2, 3, 4]. Additionally, domain wall scattering can contribute to α and β . In *ballistic* (Ga,Mn)As, intrinsic spin-orbit coupling causes significant hole reflection at the domain wall, even in the adiabatic limit when the wall is much thicker than the Fermi wavelength [5]. This grossly increases the out-ofplane spin-transfer torque, and consequently the currentdriven domain wall mobility. So far, there are no investigations on the effect of these domain wall induced hole reflections on the effective Gilbert damping constant α .

Experimental (Ga,Mn)As samples are dirty so that the effect of disorder on the effective Gilbert damping and the out-of-plane spin transfer torque should be taken into account. We find surprisingly that, in systems with a large intrinsic spin-orbit coupling, domain wall scattering contributes dominantly to α and β even in the dirty limit. Intrinsic current-domain wall motion coupling is robust against impurity scattering.

Current-induced domain-wall motion has been seen in many experiments [3]. The reciprocal effect, domain-wall motion induced current, is currently theoretically investigated [6, 7], and seen experimentally [8]. A precessing domain wall induces a charge current in ferromagnetic metals [6] similar to spin-pumping in layered ferromagnetnormal metal systems [9]. For rigid domain wall motion, the induced charge current is proportional to β/α [7]. We find that β and β/α in (Ga,Mn)As are so large that the current, or equivalently, the voltage induced by a moving domain wall is experimentally measurable.

Onsager's reciprocity relations dictate that response coefficients of domain wall motion induced current and current induced domain wall motion are related. In dirty systems, these relations have been discussed in Ref. [7]. Ref. [7] also used the scattering theory of adiabatic pumping to evaluate the non-adiabatic spin-transfer torque in ballistic systems without intrinsic spin-orbit interaction. We first extend the pumping approach to (Ga,Mn)As with strong intrinsic spin-orbit interaction, and second, also evaluate the Onsager coefficient as a function of sample disorder. In determining all Onsager coefficients, magnetization friction must be evaluated on the same footing. To this end, we generalize the energy pumping scattering theory of Gilbert damping [10] to domain wall motion. Our numerical calculation demonstrates, for the first time, that domain wall scattering is typically more important than impurity scattering for the effective domain wall motion friction in systems with a strong intrinsic spin-orbit interaction. Our novel Onsager scattering approach can also be used to compute the effective rigid domain wall motion α and β parameters in realistic materials like Fe, Ni, Co, and alloys thereof from first-principles.

Let us discuss in more detail the Onsager reciprocity relations in our system. The magnetic field is a thermodynamic force for the magnetization since it can move domain walls. The electric field is a thermodynamic force for the charges as it induces currents. In systems where charge carriers also carry spin, the magnetic and charge systems are coupled. Through this coupling, the electric field can move a domain wall and, vice versa, the magnetic field can induce a current. This phenomenon, where the thermodynamic force of one system can induce a flux in another system is well-known in thermodynamics [11]: Assume a system described by the quantities $\{q_i\}, X_i$ denotes the thermodynamic force, and J_i the flux associated with the quantity q_i . In linear response, $J_i = \sum_j L_{ij}X_j$, where L_{ij} are the Onsager coefficients. Onsager's reciprocity principle dictates $L_{ij} = \epsilon_i \epsilon_j L_{ji}$, where $\epsilon_i = 1$ ($\epsilon_i = -1$) if q_i is even (odd) under timereversal [11]. Fluxes and forces are not uniquely defined, but the Onsager reciprocity relations are valid when the entropy generation is $\dot{S} = \sum_i J_i X_i$ [11].

We first derive expressions for the Onsager coefficients and determine the Onsager reciprocity relations between a charge current and a moving domain wall in terms of the scattering matrix. Subsequently, we derive the relation between the Onsager coefficients and the effective Gilbert damping parameter α_w and the out-of-plane torque parameter β_w for domain wall motion. Finally, we numerically compute α_w and β_w for (Ga,Mn)As.

We start the derivation of the Onsager coefficients in terms of the scattering matrix by assuming the following free energy functional for the magnetic system

$$F[\mathbf{M}] = M_s \int d\mathbf{r} \left(\frac{J}{2} [(\nabla \theta)^2 + \sin^2(\theta) (\nabla \phi)^2] + \frac{K_\perp}{2} \sin^2(\theta) \sin^2(\phi) - \frac{K_z}{2} \cos^2(\theta) - H_{\text{ext}} \cos(\theta) \right), \quad (1)$$

where M_s , J and H_{ext} are the saturation magnetization, spin-stiffness and external magnetic field, respectively, and K_z and K_{\perp} are magnetic anisotropy constants. The local magnetization angles θ and ϕ are defined with respect to the z- and x-axis, respectively. The system contains a Bloch wall rotating in the (transverse) x-z plane, $\cos(\theta) = \tanh([y-r_w]/\lambda_w), \sin(\theta) = 1/\cosh([y-r_w]/\lambda_w),$ where r_w is the position of the wall, and λ_w is the wall width. We assume the external magnetic field is lower than the Walker threshold, so that the wall rigidly moves $(\dot{\phi} = 0)$ with a constant drift velocity. In this case r_w and ϕ completely characterize the magnetic system, and $\lambda_w = \sqrt{J/(K_z + K_{\perp} \sin^2(\phi))}$ [4]. The current is along the y-axis.

The heat dissipated per unit time from a charge current J_c is $\dot{Q} = J_c(V_L - V_R)$, where $V_L(V_R)$ is the voltage in the left (right) reservoir. Using the relation dS = dQ/T, this implies an entropy generation $\dot{S} = J_c(V_L - V_R)/T$. Thus, $X_c \equiv (V_L - V_R)/T$ is the thermodynamic force inducing the flux J_c . We assume the magnetic system to be at constant temperature, which means that the heat transported out of the magnetic system as the domain wall moves equals the loss of free energy. This implies an entropy generation $\dot{S} = \dot{Q}/T = -\dot{F}/T = (-\partial F[r_w, \phi]/T \partial r_w) \dot{r}_w = X_w J_w$, where we have defined the force $X_w \equiv -\partial F[r_w, \phi]/T \partial r_w$ and flux $J_w \equiv \dot{r}_w$. Using Eq. (1), we find $X_w = -2AM_s H_{\text{ext}}/T$, where A is the conductor's cross-section. Fluxes are related to forces by

$$J_w = L_{ww}X_w + L_{wc}X_c \tag{2}$$

$$J_c = L_{cc}X_c + L_{cw}X_w, (3)$$

where $L_{cc} = GT$ and G is the conductance. L_{ww} (L_{wc}) determine the induced domain wall velocity by an external magnetic field (a current). The induced current by a moving domain wall caused by an external magnetic field H_{ext} is controlled by L_{cw} . Both charge and r_w are even under time-reversal so that $L_{cw} = L_{wc}$ [12].

The current induced by a moving domain wall is parametric pumping in terms of the scattering matrix [9]:

$$J_{c,\alpha} = \frac{e\dot{r}_w}{2\pi} \sum_{\beta=1,2} \Im m \left\{ \operatorname{Tr} \left[\frac{\partial S_{\alpha\beta}}{\partial r_w} S_{\alpha\beta}^{\dagger} \right] \right\} \,, \qquad (4)$$

where $S_{\alpha\beta}$ is the scattering matrix between transverse modes in lead β to transverse modes in lead α . The system has two leads $(\alpha, \beta \in \{1, 2\})$. The trace is over all propagating modes at the Fermi energy E_F . From Eqs. (2) and (3) we find $J_c = L_{cw} \dot{r}_w / L_{ww}$.

We consider transport well below the critical transition temperature in (Ga,Mn)As, which is relatively low, and assume the energy loss in the magnetic system is transferred into the leads by holes. Generalizing Ref. [10] to domain wall motion, this energy-flux is related to the scattering matrix:

$$J_E = \frac{\hbar}{4\pi} \operatorname{Tr} \left\{ \frac{dS}{dt} \frac{dS^{\dagger}}{dt} \right\} = \frac{\hbar \dot{r}_w^2}{4\pi} \operatorname{Tr} \left\{ \frac{\partial S}{\partial r_w} \frac{\partial S^{\dagger}}{\partial r_w} \right\}.$$
 (5)

For a domain wall moved by an external magnetic field, we then find that $X_w J_w = J_w^2/L_{ww} = J_E/T$. In summary, the Onsager coefficients in Eq. (2) and Eq. (3) are

$$L_{ww} = \left(\frac{\hbar}{4\pi} \operatorname{Tr}\left\{\frac{\partial S}{\partial r_w}\frac{\partial S^{\dagger}}{\partial r_w}\right\}\right)^{-1}, \qquad (6)$$

$$L_{cw} = \frac{2e}{\hbar} \frac{\sum_{\beta=1,2} \Im m \left\{ \operatorname{Tr} \left[\frac{\partial S_{\alpha\beta}}{\partial r_w} S_{\alpha\beta}^{\dagger} \right] \right\}}{\operatorname{Tr} \left\{ \frac{\partial S}{\partial r_w} \frac{\partial S^{\dagger}}{\partial r_w} \right\}}, \qquad (7)$$

$$L_{cc} = \frac{e^2}{h} \operatorname{Tr} \left\{ t^{\dagger} t \right\} , \qquad (8)$$

where t is the transmission coefficient in the scattering matrix. We have omitted the temperature factor in the coefficients (6), (7), and (8) since it cancels with the temperature factor in the forces, *i.e.* we transform $L \rightarrow L/T$ and $X \rightarrow TX$. The Onsager coefficient expressions in terms of the scattering matrix are valid irrespective of impurity disorder and spin-orbit interaction in the band structures or during scattering events, and can treat transport both in ballistic and diffusive regimes.

Let us compare the global Onsager cofficients (6), (7), and (8) with the local Onsager coefficients in the dirty limit to gain additional understanding. In the dirty limit, all Onsager cofficients become local and the magnetization dynamics can be described by the following phenomenological local LLG equation [2, 3]:

$$\dot{\mathbf{m}} = -\gamma \mathbf{m} \times \mathbf{H}_{\text{eff}} + \alpha \mathbf{m} \times \dot{\mathbf{m}} \\ - (1 - \beta \mathbf{m} \times) (\mathbf{v}_s \cdot \nabla) \mathbf{m}, \qquad (9)$$

where **m** is the magnetization direction, \mathbf{H}_{eff} is the effective magnetic field, γ is the gyromagnetic ratio, $\mathbf{v}_s = -\hbar P \mathbf{j}/(eS_0)$, $S_0 = M_s/\gamma$, M_s the magnetization, α the Gilbert damping constant, P the spin-polarization along $-\mathbf{m}$ of the charge carriers [13], and β is the outof-plane spin-transfer torque parameter. Substituting a Walker ansatz into Eq. (9) gives below the Walker threshold [4]: $\alpha \dot{r}_w / \lambda_w = -\gamma H_{\text{ext}} - \hbar \beta P j / (eS_0 \lambda_w)$. In dirty, local, systems this equation determines the relation between the flux J_w and the forces X_w and X_c as $L_{ww} = \lambda_w / (2AS_0\alpha)$ and $L_{wc} = -\hbar \beta P G / (e\alpha S_0 A)$, where we have used $j = \sigma (V_L - V_R)/L$, and $G = \sigma A/L$. Here, L is the length of the conductor, e the electron charge, and σ the conductivity. This motivates defining the following dimensionless global coefficients:

$$\alpha_w \equiv \frac{\lambda_w}{2AS_0L_{ww}}, \quad \beta_w \equiv -\frac{\lambda_w e}{2\hbar PG}\frac{L_{wc}}{L_{ww}}.$$

 α_w is the effective Gilbert damping coefficient and β_w is the effective out-of-plane torque on the domain wall.

We will in the following investigate α_w and β_w for (Ga,Mn)As by calculating the scattering matrix expressions in Eq. (6) and Eq. (7). We use the following Hamiltonian to model quantum transport of itinerant holes:

$$H = H_{\rm L} + \mathbf{h}(\mathbf{r}) \cdot \mathbf{J} + V(\mathbf{r}) \,. \tag{10}$$

Here, $H_{\rm L}$ is the 4×4 Luttinger Hamiltonian (parameterized by γ_1 and γ_2) for zincblende semiconductors in the spherical approximation, while $\mathbf{h} \cdot \mathbf{J}$ describes the exchange interaction between the itinerant holes and the local magnetic moment of the Mn dopants. We introduce Anderson impurities as $V(\mathbf{r}) = \sum_i V_i \delta_{\mathbf{r},\mathbf{R}_i}$, where \mathbf{R}_i is the position of impurity i, V_i its impurity strength, and δ the Kronecker delta. More details about the model and the numerical method used can be found in Refs. [14, 15].

We consider a discrete conductor with transverse dimensions $L_x = 23nm$, $L_z = 17nm$ and length $L_y = 400nm$. The lattice constant is 1nm, much less than the typical Fermi wavelength $\lambda_F \sim 8nm$. The Fermi energy $E_F = 82meV$ is measured from the bottom of the lowest subband. $|\mathbf{h}| = 0.5 \times 10^{-20}J$ and $\gamma_1 = 7$. The typical mean-free path for the systems studied ranges from the diffusive to the ballistic regime $l \sim 23nm \to \infty$, and we are in the metallic regime $k_F l \gg 1$. The domain wall length is $\lambda_w = 40nm$. The spin-density S_0 from the local magnetic moments is $S_0 = 10\hbar x/a_{GaAs}^3$, a_{GaAs} the lattice constant for GaAs, and x = 0.05 the doping level[14].

Fig. 1a shows the computed effective Gilbert damping coefficient α_w versus λ_w/l for (Ga,Mn)As containing one Bloch wall. Note the relatively high $\alpha_w \sim 5 \times 10^{-3}$ in the ballistic limit. Additional impurities, in combination with the spin-orbit coupling, assist in releasing energy and angular momentum into the reservoirs and increase α_w . However, as shown in Fig. 1a, impurities contribute only about 20% to α_w even when the domain wall is two



free path when $\gamma_2 = 2.5$. Here, λ_w is kept fixed, and l is varied. Inset: α_w as a function of spin-orbit coupling γ_2 for a clean system, $l = \infty$. (b): β_w as a function of λ_w/l , where λ_w is the domain wall length and l is the mean free path when $\gamma_2 = 2.5$. Here, λ_w is kept fixed, and l is varied. Inset: β_w as a function of spin-orbit coupling γ_2 for a clean system, $l = \infty$. In all plots, line is guide to the eye.

times longer than the mean free path. Due to the strong spin-orbit coupling, ballistic domain walls have a large intrinsic resistance [5] that survives the adiabatic limit. When itinerant holes scatter off the domain wall their momentum changes and through the spin-orbit coupling their spin also changes. This is the dominate process for releasing energy and magnetization into the reservoirs. The saturated value $\alpha_w \sim 6 \times 10^{-3}$ is of the same order as the estimates in Ref. [16] for bulk (Ga,Mn)As. The inset in Fig. 1a shows the domain-wall contribution to α_w versus the spin-orbit coupling for a clean system with no impurities. α_w monotonically decreases for decreasing γ_2 and vanish for $\gamma_2 \to 0$. Since, $\lambda_w/\lambda_F \sim 5$, itinerant holes will, without spin-orbit coupling, traverse the do-

main wall adiabatically.

Fig. 1b shows β_w versus λ_w/l . β_w decreases with increasing disorder strength. This somewhat counter intuitive result stem from the fact that domain walls in systems with spin-orbit coupling have a large intrinsic domain wall resistance [5] which originates from the anisotropy in the distribution of conducting channels [5]. The reflected spins do not follow the magnetization of the domain wall, and thereby cause a large out-of-plane torque [2]. This causes the large β_w in the ballistic limit. Scalar, rotational symmetric impurities tend to reduce the anisotropy in the conducting channels, and thereby reduce the intrinsic domain wall resistance and consequently reduce β_w . Deeper into the diffusive regime, β saturates. Here, the domain wall resistance and β_w are kept at high levels due to the increase in the spin-flip rate caused by impurity scattering. The saturated value is $\beta \sim 1$. For even dirtier systems than a reasonable computing time allows, we expect a further increase in β_w . In comparison, simple microscopic theories for ferromagnetic metals where one disregards the spin-orbit coupling in the band structure predict $\beta \sim 0.001 - 0.01$ [2, 3, 4]. Similar to the Gilbert damping, in ballistic systems β_w increases with spin-orbit coupling because of the increased domain wall scattering [5], see Fig. 1b inset.

 β_w can be measured experimentally by the induced current or voltage from a domain wall moved by an external magnetic field as a function of the domain wall velocity [7]. From the Onsager relations we have that $J_c = L_{cw}X_w$. Using $X_w = J_w/L_{ww}$, the induced current and voltage are [7]:

$$J_c = -2\beta \frac{\hbar PG}{e\lambda_w} \dot{r}_w \Rightarrow V = -2\beta_w \frac{\hbar P}{e\lambda_w} \dot{r}_w \,. \tag{11}$$

An estimate of the maximum velocity of a domain wall moved by an external magnetic field below the Walker treshold is $\dot{r}_w \sim 10 \ m/s$ [17]. With $\lambda_w = 40 \ nm$ and P = 0.66 this indicates an experimentally measurable voltage $V \sim 0.2 \ \mu V$.

In conclusion, we have derived Onsager coefficients and reciprocity relations between current and domain wall motion in terms of scattering matrices. In (Ga,Mn)As, we find the effective Gilbert damping constant $\alpha_w \sim 0.01$ and out-of-plane spin transfer torque parameter $\beta_w \sim 1$. In contrast to ferromagnetic metals, the main contributions to α_w and β_w in (Ga,Mn)As are intrinsic, and induced by scattering off the domain wall, while impurity scattering is less important. The large β_w parameter implies a measurable moving domain wall induced voltage.

This work was supported in part by the Re-

search Council of Norway, Grants Nos. 158518/143 and 158547/431, computing time through the Notur project and EC Contract IST-033749 "DynaMax".

- J.C. Slonczewski, J. Magn. Magn. Mater. **159**, L1 (1996);
 L. Berger, Phys. Rev. B **54**, 9353 (1996).
- [2] G. Tatara, H. Kohno, Phys. Rev. Lett. 92,086601 (2004);
 S. Zhang, Z. Li, Phys. Rev. Lett. 93, 127204 (2004);
 S. E. Barnes, S. Maekawa, Phys. Rev. Lett. 95, 107204 (2005);
 A. Thiaville, Y. Nakatani, J. Miltat, Y. Suzuki, Europhys. Lett. 69, 990 (2005);
 Y. Tserkovnyak, H. J. Skadsem, A. Brataas, G. E. W. Bauer, Phys. Rev. B 74, 144405 (2006);
 A. K. Nguyen, H. J. Skadsem, A. Brataas, Phys. Rev. Lett. 98, 146602 (2007).
- [3] For reviews see *e.g.* D.C. Ralph, M.D. Stiles, J. Magn. Magn. Mater., **320** 1190 (2008).
- [4] Y. Tserkovnyak, A. Brataas, and G. E. W. Bauer, J. Magn. Magn. Mater., **320** 1282 (2008).
- [5] A. K. Nguyen, R. V. Shchelushkin and A. Brataas, Phys. Rev. Lett. **97**, 136603 (2006); R. Oszwaldowski, J. A. Majewski and T. Dietl, Phys. Rev. B **74**, 153310 (2006).
- [6] G. E. Volovik, J. Phys. C: Sol. State Phys. 20, L83 (1987);
 S. E. Barnes and S. Maekawa, Phys. Rev. Lett. 98, 246601 (2007);
 J.I. Ohe, A. Takeuchi, and G. Tatara, Phys. Rev. Lett. 99, 266603 (2007);
 S. A. Yang, D. Xiao, and Q. Niu, cond-mat/0709.1117.
- [7] R. A. Duine, Phys. Rev. B 77, 014409 (2008);
 arXiv:0809.2201v1; Y. Tserkovnyak and M. Mecklenburg, Phys. Rev. B 77, 134407 (2008).
- [8] S. A. Yang et al., Phys. Rev. Lett. 102, 067201 (2009).
- [9] Y. Tserkovnyak, A. Brataas, G. E. W. Bauer, and B. I. Halperin, Rev. Mod. Phys. 77, 1375 (2005).
- [10] A. Brataas, Y. Tserkovnyak and G.E.W Bauer, Phys. Rev. Lett. **101**, 037207 (2008).
- [11] S. R. de Groot, Thermodynamics of Irreversible Processes (North-Holland, Amsterdam, 1952).
- [12] The magnetic texture is inverted in this equation, i.e. $L_{cw} [\mathbf{m}] = L_{wc} [-\mathbf{m}].$
- [13] We define the spin-polarization $P = \langle J_z \rangle = \sum_j (\psi_j^{\dagger} J_z \psi_j) / N$ ($\mathbf{m} = -\hat{\mathbf{z}}$), where the sum is over N propagating modes, ψ_j are the corresponding spinor-valued wavefunctions, and J_z the dimensionless angular momentum operator.
- [14] T. Jungwirth, J. Sinova, J. Mašek, J. Kučera, and A. H. MacDonald, Rev. Mod. Phys. 78, 809 (2006).
- [15] A. K. Nguyen, and A. Brataas, Phys. Rev. Lett. 101, 016801 (2008).
- [16] J. Sinova et al., Phys. Rev. B 69, 085209 (2004); Y. Tserkovnyak, G. A. Fiete and B. I. Halperin, Appl. Phys. Lett. 84, 5234 (2004); I. Garate and A. MacDonald, arXiv:0808.3923v1.
- [17] A. Dourlat, V. Jeudy, A. Lemaitre and C. Gourdon, Phys. Rev. B 78, 161303(R) (2008).