

Counting closed geodesics in Moduli space.

Alex Eskin* and Maryam Mirzakhani†

February 18, 2019

1 Introduction.

Let \mathcal{M}_g denote the moduli space of Riemann surfaces of genus g . We may write $\mathcal{M}_g = \mathcal{T}_g/\Gamma$, where \mathcal{T}_g is the Teichmüller space of genus g surfaces, and Γ is the mapping class group. Let $N(R)$ denote the number of closed geodesics in \mathcal{M}_g of length at most R . Then $N(R)$ is also the number of conjugacy classes of pseudo-anosov elements of the mapping class group of translation length at most R . Our main result is the following:

Theorem 1.1 *As $R \rightarrow \infty$, we have*

$$N(R) \sim \frac{e^{hR}}{hR},$$

where $h = 6g - g$.

In the above theorem and below, the notation $A \sim B$ means that the ratio A/B tends to 1. Even though we assume that the surface has no punctures here, most of the results also hold on $\mathcal{M}_{g,n}$, provided h is replaced by $6g - 6 + 2n$.

In the proof of Theorem 1.1, the key is to estimate the number of closed geodesics which stay outside of compact sets.

Theorem 1.2 *Let $N_j(\delta, R)$ denote the number of closed geodesics γ of length at most R in \mathcal{M}_g such that for each point $X \in \gamma$, X has at least j curves of length less than δ . Then, for all $\epsilon > 0$ there exists $\delta > 0$ and $C = C(\epsilon)$ such that for all j , $1 \leq j \leq 3g - 3$, and all $R > 0$,*

$$N_j(\delta, R) \leq Ce^{(h-j+\epsilon)R}.$$

Let \mathcal{K} be a compact subset of \mathcal{M}_g . We let $N^{\mathcal{K}}(R)$ denote the number of geodesics in \mathcal{M}_g of length less than R which never intersect \mathcal{K} . Letting $j = 1$ in Theorem 1.2, we obtain the following:

Corollary 1.3 *For every $\epsilon > 0$ there exists a compact set \mathcal{K} and $C = C(\epsilon)$ such that for all $R > 0$,*

$$N^{\mathcal{K}}(R) \leq Ce^{(h-1+\epsilon)R}.$$

Corollary 1.3 is complimentary to the following result:

*partially supported by NSF grants DMS 0244542 and DMS 0604251 and the Clay foundation

†partially supported by the Clay foundation

Theorem 1.4 (Rafi, Hammenstadt) *For any compact \mathcal{K} , and sufficiently large R ,*

$$N^{\mathcal{K}}(R) \geq e^{(h-1)R}.$$

Previous Results. The first results on this problem are due to Veech [Ve]. He proved that there exists a constant c_2 such that

$$h \leq \liminf_{R \rightarrow \infty} \frac{\log N(R)}{R} \leq \limsup_{R \rightarrow \infty} \frac{\log N(R)}{R} \leq c_2$$

and conjectured that $c_2 = h$. In a remark in a paper by Ursula Hamenstadt [H1] (see also [H2]), in which the main focus is different, she proves that $c_2 \leq (6g - 6 + 2n)(6g - 5 + 2n)$.

Sasha Bufetov [Bu] proved the formula

$$\lim_{R \rightarrow \infty} \frac{\log \tilde{N}(R)}{R} = h, \tag{1}$$

where $\tilde{N}(R)$ is the number of periodic orbits of the Rauzy-Veech induction such that the log of the norm of the renormalization matrix is at most R . (This is a closely related problem; essentially $\tilde{N}(R)$ counts closed geodesics on a certain finite cover of \mathcal{M}_g). However the equation (1) does not easily imply

$$\lim_{R \rightarrow \infty} \frac{\log N(R)}{R} = h. \tag{2}$$

Very recently, Kasra Rafi [Ra1] proved Corollary 1.3 (which implies (2)) for the case of the five-punctured sphere.

Remarks. We note that (2) is an immediate consequence of Theorem 1.1, which is a bit more precise. In order to prove Theorem 1.1 one needs Corollary 1.3 and certain recurrence results for geodesics, which are based on [Ath].

Acknowledgements. We would like to thank Kasra Rafi who suggested a major simplification of the proof of Theorem 1.2. We would also like to thank Howard Masur for many useful discussions (including the proof of Lemma 2.2) and also William Cavendish whose lecture notes were used for part of this paper.

2 Geodesics in the thin part of moduli space

In this section we prove Theorem 1.2. The main idea, which is due to Margulis, is to prove a system of inequalities, which shows that the flow (or more precisely an associated random walk) is biased toward the thick part of Teichmüller space. Variations on this theme have been used in [EMM], [EM] and [Ath].

Notation. Let p denote the natural map from \mathcal{T}_g to $\mathcal{M}_g = \mathcal{T}_g/\Gamma$. Let \mathcal{Q}_g denote the Teichmüller space of unit area quadratic differentials on surfaces of genus g , and let $\pi : \mathcal{Q}_g \rightarrow \mathcal{T}_g$ denote the natural map. The space \mathcal{Q}_g carries a natural normalized smooth measure μ , preserved by the action of Γ . We set $\mathbf{m} = \pi_*\mu$.

2.1 A system of inequalities.

Suppose $0 < s < 1$ (in fact we will be using $s = 1/2$ only). Let $\tau \gg 1$ be a parameter to be chosen later. (In particular we will assume $e^{-(1-s)\tau} < 1/2$.) Let A_τ be the operator of averaging over a ball of radius τ in Teichmüller space. So if f is a real-valued function on Teichmüller space, then

$$(A_\tau f)(X) = \frac{1}{\mathbf{m}(B(X, \tau))} \int_{B(X, \tau)} f(y) d\mathbf{m}(y).$$

Remark. In [EM], [Ath] and [EMM] the average is over spheres. In this context, we use balls, since Minsky's product region theorem gives us much more precise information about balls than about spheres.

Let m be the maximal number of disjoint curves on a closed surface of genus g . Choose $K > e^{2m\tau}$, and pick constants $\epsilon_1 < \epsilon_2 < \dots < \epsilon_m < 1/K^3$ such that for all $1 \leq i \leq m-1$,

$$\epsilon_i < \frac{\epsilon_{i+1}}{K^2(2mK^2)^{2/s}} \quad (3)$$

Note that K and $\epsilon_1, \dots, \epsilon_m$ are constants which depend only on τ and the genus g .

For $1 \leq i \leq m$ and $X \in \mathcal{T}_g$ let $\ell_i(X)$ denote the extremal length of the i 'th shortest curve on X . Let $f_0 = 1$ and for $1 \leq j \leq m$ let

$$f_j(X) = \prod_{1 \leq i \leq j} \left(\frac{\epsilon_i}{\ell_i(X)} \right)^s.$$

Note that f_j is invariant under the action of the mapping class group, and thus descends to a function on \mathcal{M}_g . Let

$$u(X) = \sum_{k=1}^m f_k(X).$$

Let $\epsilon'_j = \epsilon_j/(mK^2)$. Let

$$W_j = \{X \in \mathcal{T}_g : \ell_{j+1}(X) > \epsilon'_j\}.$$

Note that W_0 is compact, and on W_j there are at most j short curves. If $X \notin W_{j-1}$ then X has at least j short curves, and thus if $X \in W_j \setminus W_{j-1}$ then X has exactly j short curves.

In this subsection, we prove the following:

Proposition 2.1 *Set $s = 1/2$. Then we may write*

$$(A_\tau u)(X) \leq c(X)u(X) + b(X), \quad (4)$$

where $b(X)$ is a bounded function which vanishes outside the compact set W_0 , and for all j and for all $X \notin W_{j-1}$,

$$c(X) \leq C'_j \tau^j e^{-j\tau},$$

where C'_j depends only on the genus.

We now begin the proof of Proposition 2.1. We recall the following:

Lemma 2.2 *There exists $L_0 > 0$ (depending only on g) and for every $L > L_0$ there exist constants $0 < c_1 < c_2$ such that for all $X \in \mathcal{T}_g$,*

$$c_1 \leq \mathbf{m}(B(X, L)) \leq c_2$$

The constants c_1 and c_2 depend on L and g (but not on X).

Proof. Suppose $X \in \mathcal{T}_g$. Let $\alpha_1, \dots, \alpha_k$ be the curves on X with hyperbolic length less than ϵ (where ϵ is the Bers constant). Let a_i denote the hyperbolic length of α_i . Then the extremal length $\text{Ext}_X(\alpha_i) \approx a_i$ (by e.g. [Mi]). It follows from the Kerchhoff formula for the Teichmüller distance CITE... that for all $Y \in B(X, L)$,

$$\text{Ext}_Y(\alpha_i) \leq L^2 \text{Ext}_X(\alpha_i) \leq CL^2 a_i,$$

where C depends only on the genus. Then, in view of the definition of extremal length, for any area 1 holomorphic quadratic differential q on $Y \in B(X, L)$,

$$\ell_q(\alpha_i)^2 \leq \text{Ext}_Y(\alpha_i) \leq CL^2 a_i, \tag{5}$$

where $\ell_q(\cdot)$ denotes length in the flat metric defined by q , and C depends only on the genus. Thus any flat metric in the conformal class of a surface in $B(X, L)$ has curves of flat length at most $L\sqrt{Ca_i}$.

Let \mathcal{F} be a fundamental domain for the action of Γ on \mathcal{T}_g . Then, in view of (5), and the definition of the measure $\mathbf{m}(\cdot)$, for any $\gamma \in \Gamma$,

$$\mathbf{m}(B(X, L) \cap \gamma\mathcal{F}) \leq C \prod_{i=1}^k CL^2 a_i \tag{6}$$

Let $I_{X,L}$ denote the set of elements $\gamma \in \Gamma$ such that $B(X, L) \cap \gamma\mathcal{F}$ is non-empty. We will now estimate the size of $I_{X,L}$.

Note that up to uniformly bounded index (the bound depending on g and L) $I_{X,L}$ consists only of twists around the α_i . By [Mi], the number of twists around α_i which one can take and still stay in $B(X, L)$ is $O(L^2/a_i)$. Thus,

$$|I_{X,L}| \approx C_L \prod_{i=1}^k \frac{1}{a_i}. \tag{7}$$

Now the upper bound of Lemma 2.2 follows from (6).

We now briefly outline the proof of the lower bound. Let $R \subset \mathcal{Q}_g/\Gamma$ be the set of flat structures such that each $S \in R$ has flat cylinders C_i with width (i.e. core curve) between $\sqrt{a_i}$ and $\sqrt{a_i}/2$ and height between $1/(g\sqrt{a_i})$ and $1/(2g\sqrt{a_i})$, and the rest of the arcs in a triangulation of $S \in R$ have length comparable to 1. It is easy to verify using the definition of the measures μ and \mathbf{m} that

$$\mathbf{m}(\pi(R)) \geq \mu(R) > c \prod_{i=1}^k a_i, \tag{8}$$

where c depends only on the genus. Note that by [Ra2] for any $Y \in \pi(R)$, the only short curves on Y (in the hyperbolic or extremal metric) are the core curves of the cylinders C_i , and the extremal

length of the core curve of C_i is within a constant multiple of a_i . Thus, there exists a constant L' depending only on the genus such that

$$\pi(R) \subset B(X, L').$$

Note that the above equation takes place in \mathcal{T}_g/Γ . We may think of it as taking place in \mathcal{T}_g if we identify $\pi(R)$ with a subset of the fundamental domain \mathcal{F} . Then, for any $\gamma \in I_{X, L'}$,

$$\gamma\pi(R) \subset B(X, 2L').$$

Thus, in view of (7) and (8),

$$\mathbf{m}(B(X, 2L')) \geq |I_{X, L'}| \mathbf{m}(\pi(R)) \geq c,$$

where c depends only on the genus. □

If $\alpha_1, \dots, \alpha_j$ are disjoint curves let $P(\alpha_1, \dots, \alpha_j)$ denote the product region where for all $1 \leq i \leq j$, the extremal length of α_i is at most $K\epsilon_i$.

Lemma 2.3 *Suppose $1 \leq j \leq m$, $1/2 \leq s < 1$, and suppose X and τ are such that $B(X, \tau)$ is completely contained in $P(\alpha_1, \dots, \alpha_j)$. Also suppose that for all $y \in B(X, \tau)$, the set $\{\alpha_1, \dots, \alpha_j\}$ coincides with the set of the j shortest curves on y . Then*

$$(A_\tau f_j)(X) \leq c_j f_j(X),$$

where for $1/2 < s < 1$,

$$c_j = C_j(s) e^{-2j(1-s)\tau}. \tag{9}$$

and if $s = 1/2$, $c_j = C_j \tau^j e^{-j\tau}$.

Sketch of proof. The product region theorem [Mi] states that $P(\alpha_1, \dots, \alpha_j)$ can be identified with a subset of $(\mathbb{H}^2)^j \times \mathcal{T}'$ (where \mathcal{T}' is the quotient Teichmüller space obtained by collapsing all the α_i), and the Teichmüller metric on $P(\alpha_1, \dots, \alpha_j)$ is within an additive constant of the supremum metric on $(\mathbb{H}^2)^j \times \mathcal{T}'$. Let \mathbf{m}' denote the product measure on $(\mathbb{H}^2)^j \times \mathcal{T}'$, and let A'_τ denote the averaging operator with respect to the product measure \mathbf{m}' , i.e. for a real-valued function f ,

$$(A'_\tau f)(X) = \frac{1}{\mathbf{m}'(B(X, \tau))} \int_{B(X, \tau)} f(y) d\mathbf{m}'(y).$$

We first establish the lemma with A_τ replaced by A'_τ .

We may write $X = (X_1, \dots, X_j, X')$ where X_k is in the k 'th copy of the hyperbolic plane and $X' \in \mathcal{T}'$. Because of the product region theorem $B(X, \tau)$ is essentially $B(X_1, \tau) \times \dots \times B(X_j, \tau) \times B'(X', \tau)$ where for $1 \leq k \leq j$, $B(X_k, \tau)$ is a ball of radius τ in the hyperbolic plane and $B'(X', \tau)$ a ball of radius τ in \mathcal{T}' . Also, by assumption, for any $Y \in B(X, \tau)$ the set of k shortest curves on Y is $\{\alpha_1, \dots, \alpha_k\}$. Thus, for $Y \in B(X, \tau)$, with $Y = (Y_1, \dots, Y_j, Y')$, we have

$$f_k(Y) = (\epsilon_1 \dots \epsilon_j)^s \prod_{k=1}^j \ell(Y_j)^{-2s}$$

where for $Y_j \in \mathbb{H}^2$, $\ell(Y_j)$ is the flat length of the shortest curve in the torus parametrized by Y_j . (The exponent is $2s$ instead of s since on the torus extremal length is the square of flat length). Hence,

$$(A'_\tau f_j)(X) \approx (\epsilon_1 \dots \epsilon_j)^s \prod_{k=1}^j \left(\frac{1}{\text{Vol}(B(X_k, \tau))} \int_{B(X_k, \tau)} \ell(Y)^{-2s} d\text{Vol}(Y) \right), \quad (10)$$

where Vol is the standard volume form on \mathbb{H}^2 , and the notation $A \approx B$ means that the ratio A/B is bounded from above and below in terms of the genus. Now the integral in the parenthesis, i.e. an average of ℓ^{-s} over a ball in a hyperbolic plane is essentially done in [EM, Lemma 7.4] (except that there the average is over spheres, but to get the average over balls one just makes an extra integral over the radius). One gets for $1/2 < s < 1$,

$$\frac{1}{\text{Vol}(B(X, \tau))} \int_{B(X, \tau)} \ell(Y)^{-2s} d\text{Vol}(Y) \leq c(s) e^{-2(1-s)\tau} \ell(X)$$

and for $s = 1/2$,

$$\frac{1}{\text{Vol}(B(X, \tau))} \int_{B(X, \tau)} \ell(Y)^{-1} d\text{Vol}(Y) \leq c' \tau e^\tau \ell(X).$$

Substituting these expressions into (10) completes the proof of Lemma 2.3 with A'_τ instead of A_τ .

Recall that a set \mathcal{N} is an $(c, 2c)$ separated net on a metric space \mathcal{X} if $\mathcal{N} \subset \mathcal{X}$, every point of \mathcal{X} is within $2L$ of a net point, and the minimal distance between net points is at least L . In view of Lemma 2.2, for any $(2, 2c)$ separated net \mathcal{N} in any Teichmüller space (including \mathbb{H}^2), for $\tau \gg 1$ and any X ,

$$c_1 |B(X, \tau) \cap \mathcal{N}| \leq \mathbf{m}(B(X, \tau)) \leq c_2 |B(X, \tau) \cap \mathcal{N}|, \quad (11)$$

where c_1, c_2 depend only on c and the genus. Similarly, in view of the form of the function f_k ,

$$c_1 \sum_{B(X, \tau) \cap \mathcal{N}} f_k(Y) \leq \int_{B(X, \tau)} f_k(Y) d\mathbf{m}(Y) \leq c_2 \sum_{B(X, \tau) \cap \mathcal{N}} f_k(Y). \quad (12)$$

Let $d(\cdot, \cdot)$ denote the Teichmüller metric and let $d'(\cdot, \cdot)$ denote the supremum metric on $(\mathbb{H}^2)^j \times \mathcal{T}'$. Minsky's product region theorem states that there exists a constant $\beta > 0$ depending only on the genus such that for all $X, Y \in P(\alpha_1, \dots, \alpha_j)$, $|d(X, Y) - d'(X, Y)| < \beta$. Choose $L \gg \beta$, and choose a $(L, 2L)$ -separated net \mathcal{N}_k in each factor. Let \mathcal{N} be the product of the \mathcal{N}_k . Then \mathcal{N} is an $(L - \beta, 2L + \beta)$ -separated net in $P(\alpha_1, \dots, \alpha_j)$. Now in view of (11),

$$\mathbf{m}(B(X, \tau)) \approx |B(X, \tau) \cap \mathcal{N}| \approx \prod_{k=1}^j |B(X_k, \tau) \cap \mathcal{N}_k| \approx \prod_{k=1}^j \mathbf{m}'(B(X_k, \tau)) \approx \mathbf{m}'(B(X, \tau))$$

(In the above equation, $A \approx B$ means that the ratio A/B is bounded by two constants depending only on β, L and g , and thus ultimately only on g). Similarly, using (12), we can show that

$$\int_{B(X, \tau)} f_k(Y) d\mathbf{m}(Y) \approx \int_{B(X, \tau)} f_k(Y) d\mathbf{m}'(Y).$$

Thus $(A'_\tau f_k)(X) \approx (A_\tau f_k)(X)$. □

Remark. The proof works even if at some point $Y \in B(X, \tau)$ there are short curves other than $\{\alpha_1, \dots, \alpha_j\}$ (but these other curves are longer than the maximum of the lengths of the α_j at Y). This is used in the next lemma.

Lemma 2.4 For $1 \leq j \leq m$, let $u_j(X) = \sum_{k=j}^m f_k(X)$. Suppose $\ell_j(X) < \epsilon_j$. Then (assuming τ is large enough),

$$(A_\tau u_j)(X) \leq \left(c_j + \frac{1}{2K} \right) u_j(X), \quad (13)$$

where c_j are as in (9). In particular, letting $j = 1$, and noting that the set $\{X \in \mathcal{M}_g : \ell_1(X) > \epsilon_1\}$ is compact, we have for all $X \in \mathcal{T}_g$,

$$(A_\tau u_1)(X) \leq \left(c_1 + \frac{1}{2K} \right) u_1(X) + b(\tau).$$

Proof. Note that for any $1 \leq i \leq m$, and any $X \in \mathcal{T}_g$,

$$\frac{1}{K} f_i(X) \leq (A_\tau f_i)(X) \leq K f_i(X) \quad (14)$$

(this is because in $B(X, \tau)$ the extremal length of any curve cannot change by more than $e^{2\tau}$).

We divide the set $\{j, j+1, \dots, m\}$ into two disjoint subsets: Let I_1 be the set of $k \in \{j, j+1, \dots, m\}$ such that

$$f_k(X) \leq \frac{u_j(X)}{2mK^2}, \quad (15)$$

and let I_2 be the set of $k \in \{j, j+1, \dots, m\}$ such that the opposite inequality to (15) holds. Suppose $k \in I_1$. Then, by (14), $(A_\tau f_k)(X) \leq \frac{1}{2mk} f_k(X)$, and thus

$$\sum_{k \in I_1} (A_\tau f_k)(X) \leq \frac{1}{2K} u_j(X). \quad (16)$$

Now suppose $k \in I_2$. We claim that

$$\ell_k(X) < (2mK^2)^{1/s} \epsilon_k. \quad (17)$$

Indeed, if $k = j$ then (17) is true by assumption. If $k > j$ then

$$u_j(X) \geq f_{k-1}(X) = f_k(X) \left(\frac{\ell_k(X)}{\epsilon_k} \right)^s \geq \frac{u_j(X)}{2mK^2} \left(\frac{\ell_k(X)}{\epsilon_k} \right)^s,$$

where we have used the inequality opposite to (15) in the last estimate. Thus (17) follows.

We now claim that under the assumption that $k \in I_2$ we have

$$\ell_{k+1}(X) \geq K^2 \ell_k(X) \quad (18)$$

If $k = m$ this is clear from (17) (since in the case where $\ell_m(X)$ is small, there are no other short curves on X). Now if $k < m$, then

$$u_j(X) \geq f_{k+1}(X) = f_k(X) \left(\frac{\epsilon_{k+1}}{\ell_{k+1}(X)} \right)^s \geq \frac{u_j(X)}{2mK^2} \left(\frac{\epsilon_{k+1}}{\ell_{k+1}(X)} \right)^s$$

where again we used the inequality opposite to (15) in the last estimate. Thus,

$$\ell_{k+1}(X) \geq \frac{\epsilon_{k+1}}{(2mK^2)^{1/s}}. \quad (19)$$

Now (18) follows from (19), (17) and (3).

Now in view of (17) and (18), Lemma 2.3 can be applied to f_k . Thus, for $k \in I_2$,

$$(A_\tau f_k)(X) \leq c_k f_k(X) \leq c_j f_k(X) \quad (20)$$

where for the last inequality we assumed that τ was large enough so that $c_k < c_j$ for $k > j$. Now (13) follows from (16) and (20). \square

Proof of Proposition 2.1. Suppose $X \notin W_{j-1}$. We may write

$$u(X) = u_j(X) + \sum_{k=1}^{j-1} f_k(X).$$

Note that for $X \notin W_{j-1}$ and $k < j$, $f_k(X) \leq u_j(X)/(mK^2)$. Hence, (4) follows from (13) and (14). \square

2.2 A uniform estimate for the measure of a ball.

Proposition 2.5 *There exists a constant C_2 such that for any X , any $\delta'' > 0$ and any sufficiently large τ , the volume of any $B(X, \tau)$ is bounded by $C_2 e^{(h+\delta'')\tau}$.*

Proof. See Appendix. \square

2.3 Proof of Theorem 1.2

We discretize Teichmüller space by fixing a $(1, 2)$ separated net $\mathcal{N} \subset \mathcal{T}_g$; this means that there exists constants $0 < \eta_1 < \eta_2$ such that the distance between any two net points is at least 1, and any point in Teichmüller space is within distance 2 of a net point.

Let

$$G(X) = \prod_{1 \leq i \leq m} \ell_i(X)^{-1/2}.$$

We note that there exist constants κ_1 and κ_2 such that for all $X \in \mathcal{T}_g$,

$$\kappa_1 u(X) \leq G(X) \leq \kappa_2 u(X), \quad (21)$$

where the κ_i depend on τ and the ϵ_i (and thus ultimately only on τ and the genus).

Trajectories of the random walk. Suppose $R \gg \tau$ and let n be the integer part of R/τ . By a trajectory of the random walk we mean a map $\lambda : \{0, n-1\} \rightarrow \mathcal{T}_g$ such that for all $0 < k \leq n-1$ we have $d(\lambda_k, \lambda_{k-1}) \leq \tau$ and also λ_k belongs to the net \mathcal{N} we are using in \mathcal{T}_g . Let $\mathcal{P}(X, R)$ denote the set of all trajectories for which $d(\lambda_0, X) = O(1)$. It is a corollary of Proposition 2.5 that

$$|\mathcal{P}(X, R)| \leq C_2 e^{(h+\delta'')R}, \quad (22)$$

where $|\cdot|$ denotes the cardinality of a set.

We say that a trajectory is *almost closed in the quotient* if the distance in \mathcal{M}_g between the projection to \mathcal{M}_g of $\lambda(0)$ and the projection to \mathcal{M}_g of λ_{n-1} is $O(1)$.

Let $\delta > 0$ be a constant to be chosen later. (We will have $\delta < \epsilon'_j$ for $1 \leq j \leq m$, where the ϵ'_j are as in §2.1). For $j \in \mathbb{N}$, let $\mathcal{P}_j(X, \delta, R)$ denote the set of all trajectories starting within $O(1)$ of X for which at any point, there are at least j curves of length at most δ . Let $\tilde{\mathcal{P}}_j(X, \delta, R)$ denote the subset of these trajectories which is almost closed in the quotient.

Lemma 2.6 *For any $j \in \mathbb{N}$, and any $\epsilon' > 0$ there exists $C > 0$ such that for τ large enough (depending on ϵ' and the genus g), and $\delta > 0$ small enough (depending on τ , ϵ' and g),*

$$|\tilde{\mathcal{P}}_j(X, \delta, R)| \leq C e^{(h-j+\epsilon')R}. \quad (23)$$

Here C depends on τ , δ , ϵ' and g .

Proof. Let $R = n\tau$, and let

$$q_j(X, R) = \sum_{\lambda \in \mathcal{P}_j(X, \delta, R)} u(\lambda_{n-1})$$

Note that if $\lambda \in \tilde{\mathcal{P}}_j(X, \delta, R)$ then $G(X) \approx G(\lambda_{n-1}) \leq \kappa_2 u(\lambda_{n-1})$. Therefore,

$$|\tilde{\mathcal{P}}_j(X, \delta, R)| \leq \frac{C_1}{G(X)} \sum_{\lambda \in \tilde{\mathcal{P}}_j(X, \delta, R)} \kappa_2 u(\lambda_{n-1}) = \frac{C_1 \kappa_2}{G(X)} q_j(X, R), \quad (24)$$

where $C_1 = O(1)$. For $0 < r = k\tau < R$ let

$$q_j(X, R, r) = \sum_{\lambda \in \mathcal{P}_j(X, \delta, R, r)} u(\lambda_{k-1}),$$

where the elements of $\mathcal{P}_j(X, \delta, R, r)$ are the trajectories λ belonging to $\mathcal{P}_j(X, \delta, R)$ but truncated after $k = r/\tau$ steps. Then

$$\begin{aligned} q_j(X, R, r + \tau) &= \sum_{\lambda \in \mathcal{P}_j(X, \delta, R, r + \tau)} u(\lambda_k) \\ &= \sum_{\lambda \in \mathcal{P}_j(X, \delta, R, r)} \sum_{\lambda_k \in \mathcal{N} \cap B(\lambda_{k-1}, \tau)} u(\lambda_k) \\ &\leq C \sum_{\lambda \in \mathcal{P}_j(X, \delta, R, r)} \int_{B(\lambda_{k-1}, \tau)} u(y) d\mathbf{m}(y) \\ &= C \sum_{\lambda \in \mathcal{P}_j(X, \delta, R, r)} \mathbf{m}(B(\lambda_{k-1}, \tau)) (A_\tau u)(\lambda_{k-1}) \end{aligned}$$

where in the next to last line we estimated a sum over $\mathcal{N} \cap B(\lambda_{k-1}, \tau)$ by a constant C times an integral over $B(\lambda_{k-1}, \tau)$.

Note that for $\lambda \in \mathcal{P}_j(X, \delta, R)$, the number of curves shorter than δ on λ_{k-1} is at least j . Thus, if δ is small enough, (depending on the ϵ'_j and thus ultimately only on τ and the genus),

$\lambda_{k-1} \notin W_{j-1}$. Then, from Proposition 2.1, and assuming τ is large enough so that Proposition 2.5 holds with $\delta'' < \epsilon'/2$, we get

$$q_j(X, R, r + \tau) \leq C \sum_{\lambda \in \mathcal{P}_j(X, \delta, R, r)} e^{(h+\epsilon'/2)\tau} C'_j \tau e^{-j\tau} u(\lambda_{k-1}) = CC'_j \tau e^{(h+(\epsilon'/2)-j)\tau} q_j(X, R, r). \quad (25)$$

Now iterating (25) $n = R/\tau$ times we get

$$q_j(X, R) \leq u(X) (CC'_j)^n \tau^n e^{(h-j+\epsilon'/2)n\tau} = u(X) e^{(h-j+\epsilon'/2+\frac{\log(CC'_j\tau)}{\tau})R} \quad (26)$$

We now choose τ so that $\frac{\log(CC'_j\tau)}{\tau} < \epsilon'/2$. Now the lemma follows from (21), (24), and (26). \square

Let $N_j(X, \delta, R)$ be the number of conjugacy classes of closed geodesics of length at most R which pass within $O(1)$ of the point X and always have at least j curves of length at most δ .

Lemma 2.7 *For any $\epsilon' > 0$ we may choose τ large enough (depending only on ϵ') so that for any $X \in \mathcal{T}_g$, any $\delta < 1/2$ and any sufficiently large R (depending only on ϵ', τ) we have*

$$N_j(X, \delta, (1 - \epsilon')R) < C |\tilde{\mathcal{P}}_j(X, 2\delta, R)|. \quad (27)$$

Proof. Let I_X denote the subset of the mapping class group which moves X by at most $O(1)$. Then up to uniformly bounded index, I_X consists only of Dehn twists around curves which are short on X .

Now consider a closed geodesic λ in \mathcal{M}_g which passes within $O(1)$ of $p(X)$ (recall that p denotes the natural map from $\mathcal{T}_g \rightarrow \mathcal{M}_g$). Let $[\lambda]$ denote the corresponding conjugacy class in Γ . Then there are approximately $|I_X|$ lifts of $[\lambda]$ to \mathcal{T}_g which start within $O(1)$ of X . Each lift γ is a geodesic segment of length equal to the length of λ .

We can mark points distance τ apart on γ , and replace these points by the nearest net points. (This replacement is the cause of the ϵ'). This gives a map Ψ from lifts of geodesics to trajectories. If the original geodesic λ has length at most $(1 - \epsilon')R$ and always has j curves shorter than δ , then the resulting trajectory belongs to $\tilde{\mathcal{P}}_j(X, 2\delta, R)$.

If two geodesic segments map to the same trajectory, then the segments fellow travel within $O(1)$ of each other. In particular if g_1 and g_2 are the pseudo-anosov elements corresponding to the two geodesics, then $d(g_2^{-1}g_1X, X) = O(1)$, thus $g_2^{-1}g_1 \in I_X$.

We now consider all possible geodesics contributing to $N_j(X, \delta, (1 - \epsilon')R)$; for each of these we consider all the possible lifts which pass near X , and then for each lift consider the associated random walk trajectory. We get:

$$N_j(X, \delta, (1 - \epsilon')R) |I_X| \leq C |I_X| |\tilde{\mathcal{P}}_j(X, 2\delta, R)|$$

(the factor of $|I_X|$ on the left hand side is due to the fact that we are considering all possible lifts which pass near X , and the factor of $|I_X|$ on the right is the maximum possible number of times a given random walk trajectory can occur as a result of this process). Thus, the factor of $|I_X|$ cancels, and the lemma follows. \square

The following lemma is due to Veech [Ve].

Lemma 2.8 *Suppose $\lambda \in \mathcal{M}_g$ is a closed geodesic of length at most R . Then for any $X \in \lambda$, $\ell_1(X) \geq \epsilon'_0 e^{-(6g-6)R}$, where ϵ'_0 depends only on g .*

Proof. We reproduce the proof for completeness. Let \tilde{X} be some point in \mathcal{T}_g with $p(\tilde{X}) = X$. Suppose the estimate is false, and let α be a curve on \tilde{X} with hyperbolic length less than $\epsilon'_0 e^{-(6g-4)R}$. Let γ be the element of the mapping class group associated to the lift of λ passing through \tilde{X} .

Recall that for very short curves, the hyperbolic and the extremal lengths are comparable. Therefore $\text{Ext}_\alpha(\tilde{X}) \leq \epsilon''_0 e^{-(6g-4)R}$. Therefore, by Kerchhoff's formula, for $j \in \mathbb{N}$,

$$\text{Ext}_{\gamma^j \alpha}(\tilde{X}) = \text{Ext}_\alpha(\gamma^j \tilde{X}) \leq \epsilon''_0 e^{-(6g-4+2j)R}$$

In particular, for $1 \leq j \leq 3g-2$, $\text{Ext}_{\gamma^j \alpha}(\tilde{X}) \leq \epsilon''_0$. Therefore (assuming that ϵ'_0 is small enough), for $1 \leq j \leq 3g-2$, the curves $\gamma^j \alpha$ are disjoint. This is a contradiction. \square

Proof of Theorem 1.2. Let $\epsilon' = \epsilon/8$. By Lemma 2.7 and Lemma 2.6 we can choose τ and δ so that (27) holds and also (23) holds with δ replaced by 2δ . We get, for sufficiently large R ,

$$N_j(X, \delta, R) < C' e^{(h-j+\epsilon/2)R}. \quad (28)$$

Finally $N_j(\delta, R)$ is at most $\sum_X N_j(X, \delta, R)$, where we have to let X vary over a net points within distance 1 of a fundamental domain for the action of the mapping class group. In view of Lemma 2.8, the number of relevant points in the net is at most polynomial in R . Thus Theorem 1.2 follows. \square

3 Recurrence of Geodesics.

For a compact subset \mathcal{K} of \mathcal{M}_g and a number $\theta > 0$ let $N^\mathcal{K}(\theta, R)$ denote the number of closed geodesics γ of length at most R such that γ spends at least θ -fraction of the time outside \mathcal{K} .

Theorem 3.1 *Suppose $\theta > 0$. Then there exists a compact subset \mathcal{K} of \mathcal{M}_g and $\delta > 0$ such that for sufficiently large R ,*

$$N^\mathcal{K}(\theta, R) \leq e^{(h-\delta)R}.$$

Proof. In view of Corollary 1.3, there exists a compact set \mathcal{K}_1 , such that the number of geodesics which do not return to \mathcal{K}_1 is $O(e^{(h-0.99)R})$. But then using Proposition 2.1 and [Ath, Corollary 2.7] (cf. [Ath, Theorem 2.3]), there exists a compact set \mathcal{K} depending on \mathcal{K}_1 , θ and there exists $\delta' > 0$, such that the number of random walk trajectories which start in \mathcal{K}_1 spend at least θ -fraction of the time outside of \mathcal{K} is at most $e^{(h-\delta')R}$. (Note that even though [Ath, Corollary 2.7] is not stated with an exponential bound, the proof does in fact imply this, as is done in the statement and proof of [Ath, Theorem 2.3]). It follows that the same kind estimate is true for the number of geodesics (see the proof of Lemma 2.7). \square

A Teichmüller geodesic γ is in fact a path in the unit tangent bundle of \mathcal{M}_g , i.e. the space \mathcal{Q}_g/Γ , where \mathcal{Q}_g is the space of unit area holomorphic quadratic differentials on surfaces of genus g . Let $\mathcal{P}(1, \dots, 1) \subset \mathcal{Q}_g$ denote the *principal stratum*, i.e. the set of pairs (M, q) where q is a holomorphic quadratic differential on M with simple zeroes. As above, for a compact subset \mathcal{K} of $\mathcal{P}(1, \dots, 1)$ and $\theta > 0$ we denote by $N^\mathcal{K}(\theta, R)$ denote the number of closed geodesics γ of length at most R such that γ spends at least θ -fraction of the time outside \mathcal{K} .

Theorem 3.2 *Suppose $\theta > 0$. Then there exists a compact subset $\mathcal{K} \subset \mathcal{P}(1, \dots, 1)$ and $\delta > 0$ such that for sufficiently large R ,*

$$N^{\mathcal{K}}(\theta, R) \leq e^{(h-\delta)R}.$$

The rest of this subsection will consist of the proof of Theorem 3.2.

Notation. If $S \in \mathcal{Q}_g$ is a pair (M, q) where M is a genus g surface and q is a quadratic differential on M , then we let $\ell(S)$ denote the length of the shortest saddle connection on S (in the flat metric defined by q). Let $d_{\mathcal{T}}(\cdot, \cdot)$ denote the Teichmüller metric.

Lemma 3.3 *Suppose $\mathcal{K}_1 \subset \mathcal{M}_g$ is a compact set, and $\epsilon > 0$. Suppose $\gamma : [0, L] \rightarrow \mathcal{Q}_g$ is a geodesic segment (parametrized by arclength), such that $(p \circ \pi)(\gamma(0)) \in \mathcal{K}_1$, $(p \circ \pi)(\gamma(L)) \in \mathcal{K}_1$ and*

$$|\{t \in [0, L] : \ell(\gamma(t)) > \epsilon\}| > (1/2)L.$$

Then there exists a constants c_0, L_0 , and ϵ' , depending only on ϵ and \mathcal{K}_1 such that if $L > L_0$ and if for any points p_1, p_2 with $d_{\mathcal{T}}(p_1, \pi(\gamma(0))) < c_0$, $d_{\mathcal{T}}(p_2, \pi(\gamma(L))) < c_0$, if we let $\hat{\gamma}(t)$ be the geodesic segment connecting p_1 and p_2 , then

$$|\{t \in [0, L] : \ell(\hat{\gamma}(t)) > \epsilon'\}| > (1/3)L.$$

Proof. For each $t \in [0, L]$, let \mathcal{S}_t be a short basis for on the orienting double cover of $\gamma(t)$ (see [ABEM, §8.4] for the definition). By assumption, for each $t \in [0, L]$, $\ell(\gamma(t)) \geq \epsilon e^{-L}$. Note that since $\pi(\gamma(L))$ belongs to $\Gamma\mathcal{K}_1$, the flat metric on $\gamma(L)$ cannot have any short flat cylinders. By [ABEM, Theorem 4.1], this implies that for each $\lambda \in \mathcal{S}_L$, (and assuming $L \gg 1$),

$$\|*c_{\lambda}\|_L \leq C(\epsilon)L,$$

where $*c_{\lambda}$ is as in [ABEM, §4.1], and $\|\cdot\|_t$ denotes the Hodge norm on the surface $\pi(\gamma(t))$. Similarly, for each $\lambda \in \mathcal{S}_0$,

$$\|*c_{\lambda}\|_0 \leq C(\epsilon)L,$$

Now suppose $\lambda \in \mathcal{S}_0$ and $\eta \in \mathcal{S}_L$. Then there exists $\alpha > 0$ depending only on ϵ and the genus such that

$$|\langle *c_{\lambda}, *c_{\eta} \rangle_L| \leq \|*c_{\lambda}\|_L \|*c_{\eta}\|_L \leq C e^{(1-\alpha)L} \|*c_{\lambda}\|_0 \|*c_{\eta}\|_L \leq C_1(\epsilon) L^2 e^{(1-\alpha)L}, \quad (29)$$

where $\langle \cdot, \cdot \rangle_L$ denotes the hodge inner product on the surface $\gamma(L)$, and we used the inequality $\|*c_{\lambda}\|_L \leq C e^{(1-\alpha)t} \|*c_{\lambda}\|_0$ which follows from a theorem of Forni [Fo], restated as [ABEM, Theorem 4.3].

Now let A_t be the linear transformation which takes the basis \mathcal{S}_0 to the basis \mathcal{S}_t . Then, in view of (29),

$$\max(\|A\|_t, \|A\|_t^{-1}) \leq C L e^{(1-\alpha)t}. \quad (30)$$

Write $A = A_L$. Then, assuming L is sufficiently large,

$$\max(\|A\|, \|A\|^{-1}) \leq C L e^{(1-\alpha)L} \leq C e^{(1-(\alpha/2))L}. \quad (31)$$

As in [ABEM, §2], we denote the strong unstable, unstable, stable and strong stable foliations of the geodesic flow by \mathcal{F}^{uu} , \mathcal{F}^u , \mathcal{F}^s and \mathcal{F}^{ss} respectively. Let \mathcal{F}_t^u (resp. \mathcal{F}_t^{uu}) be the leaf of \mathcal{F}^u (resp. \mathcal{F}^{uu}) passing through $\gamma(t)$. Let $d_E(\cdot, \cdot)$ be the Euclidean metric as defined in [ABEM, §8.4], and let




Figure 1. Proof of Lemma 3.3.

$B_E(x, r)$ denote the radius r ball with center x in the Euclidean metric. Since \mathcal{K}_1 is compact, there is a number $\rho_1 > 0$ depending only on \mathcal{K}_1 and ϵ such that $B_E(q, \rho_1)$ is contained in one fundamental domain for the action of the mapping class group Γ .

Now let $S_0 = S(\pi(\gamma(0)))$ denote the sphere at $\pi(\gamma(0))$, i.e. the set of unit area holomorphic quadratic differentials on the surface $\pi(\gamma(0))$. For $q \in S_0$ near $\gamma(0)$, let $f(q) \in \mathcal{Q}_g$ and $t(q) \in \mathbb{R}$ be such that $f(q) \in \mathcal{F}_0^{uu}$ and $g_{t(q)}q$ and $f(q)$ are on the same leaf of \mathcal{F}^{ss} . Then clearly $f(\gamma(0)) = \gamma(0)$, and there exists a number $\rho_3 > 0$ depending only on \mathcal{K}_1 such that the restriction of f to $B_E(\gamma(0), \rho_3) \cap S_0$ is a homeomorphism onto a neighborhood of $\gamma(0)$ in \mathcal{F}_0^{uu} . Let

$$U = \bigcup_{|t| < \rho_3} \{g_t q : q \in B_E(\gamma(0), \rho_3/2) \cap S_0\}.$$

We now claim that there exists $\rho_0 > 0$ depending only on ρ_3 and \mathcal{K}_1 such

$$\pi(g_t U) \supset B_{\mathcal{T}}(\pi(\gamma(L)), \rho_0). \quad (32)$$

Assume for the moment that (32) holds. Let p_2 be as in the statement of the lemma, and let $\hat{\gamma}$ denote the geodesic segment connecting $\pi(\gamma(0))$ and p_2 . Then by (32), $\hat{\gamma}(0) \in S_0 \cap U$ and the length L' of $\hat{\gamma}$ satisfies $|L' - L| < \rho_3$. Then, in particular,

$$d_E(\hat{\gamma}(0), f(\hat{\gamma}(0))) \leq \rho_3.$$

Then, in view of (30), for L sufficiently large (depending only on \mathcal{K}_1 and ϵ) and $t \in [L/10, L]$,

$$d_E(\hat{\gamma}(t), f(\hat{\gamma}(t))) \leq \epsilon/4.$$

(Indeed if λ is a path in \mathcal{F}^s connecting $\hat{\gamma}(0)$ and $f(\hat{\gamma}(0))$ and realizing the Euclidean distance between these points, then the Euclidean length of $\lambda_t = g_t\lambda$ is bounded by $e^{-t}\|A_t\|$ times the Euclidean length of λ). In particular,

$$d_E(\hat{\gamma}(L), f(\hat{\gamma}(L))) \leq \epsilon/4.$$

Then,

$$d_E(\gamma(L), f(\hat{\gamma}(L))) < C,$$

where C depends only on ϵ and c_0 . Since $\gamma(L)$ and $f(\hat{\gamma}(L))$ are on the same leaf of \mathcal{F}^{uu} , it follows from (30) that for $t \in [0, 9L/10]$, if L is sufficiently large,

$$d_E(\gamma(t), f(\hat{\gamma}(t))) < \epsilon/4,$$

Thus, for L sufficiently large (depending only on \mathcal{K}_1 and ϵ) and $t \in [L/10, 9L/10]$,

$$d_E(\tilde{\gamma}(t), \gamma(t)) \leq \epsilon/2$$

Therefore, by the assumption of $\gamma(t)$, there exists a subset $W \subset [0, L]$ with $|W| \geq (5/12)L$ such that for $t \in W$, $\ell(\tilde{\gamma}(t)) \geq (\epsilon/2)$. Thus the lemma is true if $p_1 = \gamma(0)$. Now to handle the general case, we repeat the above argument with γ replaced by $\tilde{\gamma}$ in the opposite orientation. This implies the lemma with $c_0 = \rho_0/2$.

It remains to verify (32). This is straightforward (in view of the hyperbolicity (31)) but somewhat tedious argument. Let

$$V = \bigcup_{|t| < \rho_3} \{g_{t+t(q)}f(q) : q \in B_E(\gamma(0), \rho_3/2) \cap S_0\}.$$

Then V is relatively open as a subset of \mathcal{F}_0^u . Let ∂V denote the boundary of V viewed as a subset of \mathcal{F}_0^u . Therefore we can choose $\rho_2 > 0$ depending only on \mathcal{K}_1 such that for all $q' \in \partial V$,

$$d_E(q', \gamma(0)) \geq \rho_2$$

By (31) and the fact that $\partial V \subset \mathcal{F}_0^u$, this implies that for some constant ρ'_2 , (depending only on \mathcal{K}_1 and ϵ), all $q' \in \partial V$ and all $L > 0$,

$$d_E(g_L q', \gamma(L)) \geq \rho'_2. \tag{33}$$

Note that $U \subset \bigcup_{t \in \mathbb{R}} g_t S_0 \cong \mathbb{R} \times S_0$. Let ∂U denote the boundary of U viewed as a subset of $\mathbb{R} \times S_0$. Suppose $q_1 \in \partial U$. We may write $q_1 = g_t q$ for some $q \in S_0$. Then the fact that $q_1 \in \partial U$ implies that either $d_E(q, \gamma(0)) = \rho_3/2$ or $|t| = \rho_3$. In either case, let $q_2 = g_{t+t(q)}f(q)$. Then $q_2 \in \partial V$, q_1 and q_2 are on the same leaf of \mathcal{F}^{ss} , and

$$d_E(q_1, q_2) \leq C,$$

where C depends only on \mathcal{K}_1 . Then by (31) there exists L_0 (depending only on \mathcal{K}_1 and ϵ) such that for $L > L_0$,

$$d_E(g_L q_1, g_L q_2) \leq \rho'_2/2.$$

Then, in view of (33), for all $q_1 \in \partial U$ and all $L > L_0$,

$$d_E(q_L q_1, \gamma(L)) \geq \rho'_2/2.$$

Then by [ABEM, Lemma 8.6], there exists ρ_0 depending only on \mathcal{K}_1 such that for $L > L_0$, and all $q_1 \in \partial U$,

$$d_{\mathcal{T}}(\pi(g_L q_1), \pi(\gamma(L))) \geq \rho_0. \quad (34)$$

Let $\phi : U \rightarrow \mathcal{T}_g$ denote the map $\phi(q) = \pi(g_L(q))$. Then ϕ is continuous, and $\phi(\gamma(0)) = \pi(\gamma(L))$. Now in view of (34), (32) holds. \square

Proof of Theorem 3.2. Choose $\theta_1 > 0$. Let $\mathcal{K}_1 \subset \mathcal{M}_g$ be such that Theorem 3.1 holds for $\mathcal{K} = \mathcal{K}_1$, and $\theta = \theta_1$. Let \mathcal{K}_2 be a compact subset of $\mathcal{P}(1, \dots, 1)/\Gamma$ such that $\mathcal{K}_2 \subset p^{-1}(\mathcal{K}_1)$, and let \mathcal{K}_3 be a subset of the interior of \mathcal{K}_2 . We may choose these sets so that $\mu(\mathcal{K}_3) > (1/2)$, where μ is the Lebesgue measure on Q_g/Γ normalized so that $\mu(Q_g/\Gamma) = 1$. We also choose \mathcal{K}_2 and \mathcal{K}_3 to be symmetric, i.e if $q \in \mathcal{K}_2$ then $-q \in \mathcal{K}_2$ (and same for \mathcal{K}_3). Then there exists $\epsilon > 0$ such that for $X \in \mathcal{K}_3$, $\ell(X) > \epsilon$. Let c_0 be as in Lemma 3.3. We now choose a (c_1, c_2) separated net \mathcal{N} on \mathcal{T}_g , which $c_1 < c_0$, $c_2 < c_0$. We may assume that $\mathcal{N} \cap p^{-1}(\mathcal{K}_1)$ is invariant under the action of the mapping class group.

Suppose $X \in \mathcal{T}_g$, and let $S(X)$ denote the unit sphere at X , i.e. the set of area 1 holomorphic quadratic differentials on X . Let

$$U(X, T) = \{q \in S(X) : |\{t \in [0, T] : g_t q \in \mathcal{K}_2^c\}| > (1/2)T\}.$$

Let

$$V(X, T) = \bigcup_{0 \leq t \leq T} \pi(g_t U(X, t)),$$

so that $V(X, T)$ is the subset of $B(X, T)$ consisting of points $Y \in B(X, T)$ such that the geodesic from X to Y spends more than half the time outside \mathcal{K}_2 .

By [ABEM, Theorem 6.2], for any $\theta_1 > 0$, there exists $T > 0$ such that for any $\tau > T$ and any $X \in \mathcal{N} \cap p^{-1}(\mathcal{K}_1)$,

$$m(\text{Nbhd}_{c_2}(V(X, \tau) \cap p^{-1}(\mathcal{K}_1))) \leq \theta_1 e^{h\tau},$$

where $\text{Nbhd}_a(A)$ denotes the set of points within Teichmüller distance a of the set A . Then, since \mathcal{K}_1 is compact and θ_1 is arbitrary, this implies that for any $\theta_2 > 0$ there exists $T > 0$ such that for any $\tau > T$ and any $X \in \mathcal{N} \cap p^{-1}(\mathcal{K}_1)$,

$$|\mathcal{N} \cap V(X, \tau) \cap p^{-1}(\mathcal{K}_1)| \leq \theta_2 e^{h\tau}.$$

By the compactness of \mathcal{K}_1 and [ABEM, Theorem 1.2 and Theorem 10.1] there exists $C_1 > 1$ such that for τ sufficiently large and any $X \in \mathcal{N} \cap p^{-1}(\mathcal{K}_1)$,

$$C_1^{-1} e^{h\tau} \leq |\mathcal{N} \cap B(X, \tau)| \leq C_1 e^{h\tau}.$$

Thus, for any $\theta_3 > 0$ there exists $T > 0$ such that for $\tau > T$,

$$|\mathcal{N} \cap V(X, \tau) \cap p^{-1}(\mathcal{K}_1)| < \theta_3 |\mathcal{N} \cap B(X, \tau)|. \quad (35)$$

From now on we assume that τ is sufficiently large so that (35) holds.

Let $\mathcal{K}'_1 = \text{Nbhd}_{c_2}(\mathcal{K}_1)$, and let $\mathcal{G}(R)$ denote the set of closed geodesics in \mathcal{M}_g of length at most R , and let $\mathcal{G}^{\mathcal{K}'_1}(\theta_3, R) \subset \mathcal{G}(R)$ denote the subset which contributes to $N^{\mathcal{K}'_1}(\theta_3, R)$. In view of Theorem 3.1, it is enough to show that there exists $\delta_0 > 0$ such that for R sufficiently large,

$$|\mathcal{G}(R) \setminus \mathcal{G}^{\mathcal{K}'_1}(\theta_3, R)| \leq e^{(h-\delta_0)R} \quad (36)$$

As in §2, we associate a random walk trajectory $\Phi(\gamma)$ to each closed geodesic $\gamma \in \mathcal{G}(R)$. Let $\mathcal{P}_1(R) = \Phi(\mathcal{G}(R) \setminus \mathcal{G}^{\mathcal{K}'_1}(\theta_3, R))$ denote the set of resulting trajectories. Note that by construction, every trajectory in \mathcal{P}_1 spends at most θ_3 fraction of the time outside \mathcal{K}_1 .

Suppose $\Lambda = (\lambda_1, \dots, \lambda_n)$ be a trajectory in $\mathcal{P}_1(R)$. Let $J(\Lambda)$ denote the number of j , $1 \leq j \leq n$ such that

$$\lambda_j \in p^{-1}(\mathcal{K}'_1) \quad \text{or} \quad \lambda_{j+1} \in V(\lambda_j, \tau) \cup p^{-1}(\mathcal{K}'_1). \quad (37)$$

Let $\mathcal{P}_2(R) \subset \mathcal{P}_1(R)$ denote the trajectories for which $J(\Lambda) < (\theta/2)n$. Then, as long as θ_3 is chosen sufficiently small, the Law of Large Numbers implies that there exists $\delta_1 > 0$ such that for $n = R/\tau$ sufficiently large,

$$|\mathcal{P}_1(R) \setminus \mathcal{P}_2(R)| \leq e^{(h-\delta_1)R}.$$

Since every trajectory in $\mathcal{P}_1(R)$ intersects the compact set \mathcal{K}_1 , for any $\Lambda \in \mathcal{P}_1(R)$, the cardinality of $\Phi^{-1}(\Lambda)$ is bounded by a constant C depending only on \mathcal{K}_1 . Thus,

$$|\Phi^{-1}(\mathcal{P}_1(R) \setminus \mathcal{P}_2(R))| \leq Ce^{(h-\delta_1)R}.$$

To complete the proof we will show that there exists a compact set $\mathcal{K} \subset \mathcal{P}(1, \dots, 1)$ such that any geodesic $\gamma \in \Phi^{-1}(\mathcal{P}_2(R))$ spends at least $1 - \theta$ fraction of the time in \mathcal{K} . Indeed, suppose $\Lambda = (\lambda_1, \dots, \lambda_n) \in \mathcal{P}_2(R)$. and $\gamma \in \Phi^{-1}(\Lambda)$. Then there exist points $q_j = \gamma(t_j) \in \mathcal{Q}_g$ such that $q_j \in \gamma$ and for $1 \leq j \leq n$, $d_{\mathcal{T}}(\pi(q_j), \lambda_j) \leq c_2 < c_0$. Now suppose $j \notin J(\Lambda)$. Then by Lemma 3.3, there exists $\epsilon' > 0$ depending only on \mathcal{K}_1 and ϵ , and a point $t_j < t' < t_{j+1}$ such that $\ell(\gamma(t')) > \epsilon'$. But then for all $t_j < t < t_{j+1}$, $\ell(\gamma(t)) \geq e^{-\tau}\epsilon'$, so the entire geodesic γ spends at least $(1 - \theta)$ -fraction of the time in the compact set $\{q \in \mathcal{Q}_g : \ell(q) > e^{-\tau}\epsilon'\}$. \square

4 A Closing Lemma.

Lemma 4.1 (Closing Lemma) *Suppose $\mathcal{K}_1 \subset \mathcal{M}_g$ is a compact set, and $\epsilon > 0$. Suppose $\gamma : [0, L] \rightarrow \mathcal{Q}_g$ is a geodesic segment (parametrized by arclength), such that $(p \circ \pi)(\gamma(0)) \in \mathcal{K}_1$, $(p \circ \pi)(\gamma(L)) \in \mathcal{K}_1$ and*

$$|\{t \in [0, L] : \ell(\gamma(t)) > \epsilon\}| > (1/2)L.$$

Then there exists constants c_0, c_1, L_0 , and ϵ' , depending only on ϵ and \mathcal{K}_1 such that if $L > L_0$ and if

$$d_{\mathcal{T}}((p \circ \pi)(\gamma(0)), (p \circ \pi)(\gamma(L))) < c_0,$$

then there exists a closed geodesic γ' with the following properties:

- (a) γ' is homotopic to the union of γ and the geodesic segment from $\gamma(L)$ to $\gamma(0)$.
- (b) The length of γ' is within ϵ of L .
- (c) γ' passes within c_1 of $\gamma(0)$ (and $\gamma(L)$).

Outline of Proof. This is very similar to the proof of Lemma 3.3. In view of the hyperbolicity statement (31), there is a neighborhood of $\gamma(0)$ (of size at most c_0 where c_0 is as in Lemma 3.3) such that the time L geodesic flow restricted to the neighborhood expands along the leaves of \mathcal{F}^{uu} and contracts along the leaves of \mathcal{F}^{ss} . Then the contraction mapping principle (applied first to the map on \mathcal{F}^{ss} and then to the inverse of the map on \mathcal{F}^{uu}) allows us to find a fixed point for the geodesic flow near $\gamma(0)$. \square

5 Proof of Theorem 1.1.

In view of Theorem 3.2, the proof essentially reduces to the now standard hyperbolic dynamics argument of Margulis [Mar], see also [KH]. The argument below is not rigorous: its aim is to recall some of the key ideas with emphasis on what is different in this setting. We refer the reader to e.g. [KH, §20.6] for the missing details of the argument.

Choose any $\theta > 0$. Fix a compact set $\mathcal{K} \subset \mathcal{P}(1, \dots, 1)$ such that Theorem 3.2 holds, and $\mu(\mathcal{K}) > (1 - \theta/2)$. Let $\epsilon' > 0$ be such that $\ell(q) > \epsilon'$ for all $q \in \mathcal{K}$. Let $\mathcal{K}_1 = \pi^{-1}(\pi(\mathcal{K}))$. Let $U \subset \mathcal{K}_1$ be a small box in the tangent space \mathcal{Q}_g . We assume that the Teichmüller diameter of U is at most c_0 , where c_0 is as in Lemma 3.3 (with ϵ' instead of ϵ).

Recall that the geodesic flow on \mathcal{Q}_g is mixing, i.e. for $A, B \subset (\mathcal{Q}_g)$, $\mu(g_t(A) \cap B) \approx \mu(A)\mu(B)$ for large values of t . We now want to consider what happens to U under the action of the flow. We can think of the action of the geodesic flow as stretching U along \mathcal{F}^{uu} and contracting U along \mathcal{F}^{ss} . Because of the mixing property of the flow, we see the image of U under the flow intersecting U many times.

Let R be large enough that $g_R(U) \cap U$ has more than one component. We say that a connected component is “regular” if for some $q \in U$,

$$|\{t \in [0, R] : p(g_t q) \in \mathcal{K}\}| \geq (1/2)R \quad (38)$$

(so that the geodesic segment spends at least half the time in \mathcal{K}). Note that in view of Lemma 3.3, if (38) holds for some $q \in U$ then for all $q \in U$,

$$|\{t \in [0, R] : p(g_t q) \in \mathcal{K}'\}| \geq (1/3)R \quad (39)$$

where $\mathcal{K}' \supset \mathcal{K}$ is also a compact subset of $\mathcal{P}(1, \dots, 1)$.

It follows from the closing lemma (Lemma 4.1) that each regular component connected component of the intersection $g_R(U) \cap U$ contains exactly one closed geodesic of length between R and $R + \epsilon$, where ϵ is related to the diameter of U . Conversely, each closed geodesic of length between R and $R + \epsilon$ which spends at least half the time in \mathcal{K} belongs to a regular connected component of $g_R(U) \cap U$. Thus,

$$N_1(U, R, R + \epsilon) \leq C(U, R) \leq N_2(U, R, R + \epsilon), \quad (40)$$

where $C(U, R)$ is the number of regular connected components of the intersection $g_R(U) \cap U$, $N_1(U, R, R + \epsilon)$ is the number of closed geodesics of length between R and $R + \epsilon$ which intersect U and spend at least $(1 - \theta)$ -fraction of the time in \mathcal{K} , and $N_2(U, R, R + \epsilon)$ is the number of closed geodesics of length between R and $R + \epsilon$ which intersect U and spend at least $(1 - 2\theta)$ -fraction of the time in \mathcal{K}' .

Note that the irregular connected components are all contained in the set

$$E_R = \{q : |\{t \in [0, R] \text{ with } p(g_t q) \in \mathcal{K}^c > \theta R\}|\}$$

By the ergodic theorem, $\mu(E_R) \rightarrow 0$ as $R \rightarrow \infty$. Thus, for R sufficiently large, most of the measure of $g_R U \cap U$ is contained in regular connected components.

Now we can estimate,

$$C(U, R, R) \approx \frac{\mu(U \cap g_R(U))}{\mu(A)}$$

where A is a single connected component. The flow contracts the measure along \mathcal{F}^{ss} by e^{-hR} , so the area of each component of the intersection has area about $\mu(U)e^{-hR}$. By the mixing property, we know that the numerator in the preceding equation is about $\mu(U)^2$, so we get that

$$C(U, R) \sim \mu(U)e^{hR}. \quad (41)$$

(In the above we made the simplifying assumption that all regular intersections have the same area. This can be removed with a bit of care: see [KH, §20.6] for the details). Combining (41) with (40) and Theorem 3.2, we get

$$N_1(U, R, R + \epsilon) \sim \mu(U)e^{hR}.$$

Let $N_1(R, R + \epsilon)$ denote the number of closed geodesics of length between R and $R + \epsilon$ which spend at least θ -fraction of the time in \mathcal{K} . Of course not all geodesics contributing to $N_1(R, R + \epsilon)$ will pass through U . To catch all of them, we create a tiling of \mathcal{K}_1 by boxes U_i . Naively, the number of such geodesics of length R passing through U_i is about $\mu(U_i)e^{hR}$, so the total number of geodesics is $\sum \mu(U_i)e^{hR} = e^{hR} \sum \mu(U_i) = e^{hR} \mu(\mathcal{K}_1) \approx e^{hR}$. This is wrong, of course, because a geodesic will pass through many boxes. If the width of the boxes is ϵ , and the length of the geodesic is about R , and it spends at least $1 - \theta$ fraction of the time in \mathcal{K}_1 , we get that each of our paths passed through about R/ϵ boxes. This shows that our counting was off by a factor of $(1 + O(\theta))R/\epsilon$, so we get

$$N_1(R, R + \epsilon) \sim (1 + O(\theta))e^{hR}/(R/\epsilon) = (1 + O(\theta))\epsilon \cdot e^{hR}/R.$$

Thus the total number $N_1(R)$ of geodesics of length less than R which stay at least half the time in \mathcal{K} can be estimated as follows:

$$N_1(R) = \sum_{k=1}^{1/\epsilon} N_1(k\epsilon R, (k+1)\epsilon R) \sim (1+O(\theta)) \sum_{k=1}^{1/\epsilon} \epsilon \frac{e^{hk\epsilon R}}{kR} = (1+O(\theta)) \sum_{k=1}^{1/\epsilon} \frac{e^{h(k+1)\epsilon R}}{h(k+1)\epsilon R} - \frac{e^{hk\epsilon R}}{h\epsilon R} + O(\epsilon e^{hR}/R)$$

The sum on the right telescopes, and we get

$$N_1(R) \sim (1 + O(\theta))(1 + O(\epsilon)) \frac{e^{hR}}{hR}.$$

Since $\epsilon > 0$ and $\theta > 0$ are arbitrary, the theorem follows. \square

A Appendix: Proof of Proposition 2.5

Let $\tilde{\Delta}$ be a net in \mathcal{M}_g , and let $\Delta \subset \pi^{-1}(\tilde{\Delta})$ be a net in \mathcal{T}_g ; in other words, there are constants $c_1, c_2 > 0$ such that 1) : given $X \in \mathcal{T}_g$ there exists $Z \in \Delta$ such that $d_T(X, Z) \leq c_1$, and 2) : for any $Z_1 \neq Z_2 \in \Delta$, we have $d_T(Z_1, Z_2) \geq c_2$. In this case $\pi(\Delta) = \tilde{\Delta}$. Here $\pi : \mathcal{T}_g \rightarrow \mathcal{M}_g$ is the natural projection to the moduli space.

We recall that there exists a constant $C_2 > 0$ such that for any $X \in \mathcal{T}_g$

$$|\pi(B(X, \tau)) \cap \tilde{\Delta}| \leq C_2 \tau^{6g-6}. \quad (42)$$

Given X and Y in \mathcal{T}_g and $\tau > 0$, let $F_\tau(X, Y)$ denote the intersection of $B(X, \tau)$ with ΓY . Proposition 2.5 will follow from the following:

Proposition A.1 *Given $\epsilon > 0$, there exists $\tau_0 > 0$ such that for any $\tau > \tau_0$ and $X, Y \in \mathcal{T}_g$ we have*

$$|F_\tau(X, Y)| \leq e^{(h+\epsilon)\tau} G(Y)^2.$$

Proof of Proposition 2.5 assuming Proposition A.1. Suppose $X \in \mathcal{T}_g$ is arbitrary, and suppose $Y \in \tilde{\Delta}$. Then, by Proposition A.1, for τ sufficiently large,

$$|B(X, \tau) \cap \Gamma Y| \leq e^{(h+\epsilon/2)\tau} G(Y)^2.$$

Note that

$$|\Gamma Y \cap B(Y, c_2)| \geq cG(Y)^2. \quad (43)$$

Thus, since distinct points in Δ are at least c_2 apart,

$$|B(X, \tau) \cap \Gamma Y \cap \Delta| \leq ce^{(h+\epsilon/2)\tau}.$$

In view of (42), this implies that

$$|B(X, \tau) \cap \Delta| \leq c' \tau^{6g-6} e^{(h+\epsilon/2)\tau} \leq e^{(h+\epsilon)\tau}.$$

Now this implies Proposition 2.5 in view of Lemma 2.2. \square

Proof of Proposition A.1. Given $X \in \mathcal{T}_g$, let $\ell_\gamma(X) = \sqrt{\text{Ext}_\gamma(X)}$. Fix a very small ϵ_0 . Here we say γ is short on X if $\ell_\gamma(X) \leq \epsilon_0$. Let \mathcal{A}_X denote the set of all short simple closed curves on X . Given $X, Y \in \mathcal{T}_g$, and $\mathcal{B} \subset \mathcal{A}_X$ let

$$F_\tau(X, Y, \mathcal{B}) = \{gY \mid g \in \Gamma, d_T(X, gY) \leq \tau, \mathcal{A}_X \cap \mathcal{A}_{gY} = \mathcal{B}\} \subset F_\tau(X, Y).$$

Using the method in [ABEM, §10], we have

$$|F_\tau(X, Y, \mathcal{B})| \leq C_1 e^{(h-|\mathcal{B}|\tau)\tau} G(X)G(Y). \quad (44)$$

Then, in view of (43), given $Z \in \mathcal{T}_g$, $\mathcal{B} \subset \mathcal{A}_Z$, and $W \in \tilde{\Delta}$, the inequality (44) implies that:

$$\begin{aligned} |B(Z, \tau) \cap \pi^{-1}(W) \cap \Delta_{\mathcal{B}}| &\leq C_1 e^{(h-|\mathcal{B}|\tau)\tau} G(Z)G(W)/G(W)^2 \\ &\leq C_1 e^{(h-|\mathcal{B}|\tau)\tau} G(Z)/G(W), \end{aligned} \quad (45)$$

where $\Delta_{\mathcal{B}} = \{Z_1 \in \Delta \mid \forall \gamma \in \mathcal{B}, \ell_{\gamma}(Z_1) \leq \epsilon_0\} \subset \Delta$.

Let $\mathcal{A}_X = \{\gamma_1, \dots, \gamma_k\}$ be the set of short simple closed curves on X such that

$$\ell_{\gamma_1}(X) \geq \dots \geq \ell_{\gamma_s} \geq e^{-\tau} > \dots \ell_{\gamma_k}(X).$$

Then $3g-3 \geq k \geq s$. Let $\tau_0 = 0$, and for $i \geq 1$, $\tau_i = -\log(\ell_{\gamma_i}(X))$. Also for $1 \leq i \leq s$, $d_i = \tau_i - \tau_{i-1}$, $m_i = h - k + i - 1$, $d_{s+1} = \tau - \tau_s$, and $m_{s+1} = h - k$. Then it is easy to verify that

1.

$$\sum_{i=1}^{s+1} d_i = \tau, \quad \sum_{i=1}^{s+1} m_i d_i = (h - k + s)\tau - \tau_1 \dots - \tau_s, \quad G(X) = e^{\tau_1 + \dots + \tau_k};$$

2. By Kerckhoff's theorem, for $1 \leq i \leq s$, $\gamma_i, \dots, \gamma_k$ are all short in $B(X, \tau_i)$.

3. Since $\gamma_{s+1}, \dots, \gamma_k$ stay short in $B(X, \tau)$, again by Kerckhoff's theorem,

$$e^{\tau_{s+1} + \dots + \tau_k} \leq G(Y) e^{(k-s)\tau}, \quad (46)$$

for any $Y \in B(X, \tau)$.

Consider

$$\mathcal{P} = \{(X = Z_0, Z_1, \dots, Z_s, gY = Z_{s+1}) \mid Z_1, \dots, Z_s \in \Delta, \forall i, d_T(Z_i, Z_{i-1}) \leq d_i, g \in \Gamma_g\},$$

$$Z = \{(W_1, \dots, W_s) \mid W_i \in \tilde{\Delta} \cap \pi(B(X, \tau))\},$$

and finally

$$\mathcal{P}(W_1, \dots, W_s) = \{(X, Z_1, \dots, Z_s, gY) \in \mathcal{P} \mid \pi(Z_i) = W_i\}.$$

Note that for any $(X, Z_1, \dots, Z_s, Z_{s+1} = gY) \in \mathcal{P}$, $Z_i \in B(X, \tau_i)$.

On the other hand, since we can approximate a geodesic by points in the net Δ , we have

$$|F_{\tau}(X, Y)| \leq |\mathcal{P}|,$$

also by the definition,

$$\mathcal{P} \subset \bigcup_{W \in Z} \mathcal{P}(W).$$

By equation (42) we have

$$|Z| = O(\tau^{c_g}),$$

where $c_g \leq (12g - 12)s < (12g - 12)^2$.

Let $Z_0 = X$. then for a given $Z = Z_{i-1} \in \Delta \cap B(X, \tau_{i-1})$, $B(Z, d_i) \subset B(X, \tau_i)$. This implies that $h - m_i$ curves (i.e, the curves in $\mathcal{B}_i = \{\gamma_i, \dots, \gamma_k\}$) are short on $B(Z, d_i)$.

Hence equation (45) for $\mathcal{B} = \mathcal{B}_i$, $W = W_i$, and $Z = W_{i-1}$ implies that

$$|\pi^{-1}(W_i) \cap \Delta \cap B(Z_{i-1}, d_i)| = |\pi^{-1}(W_i) \cap \Delta_{\mathcal{B}_i} \cap B(Z_{i-1}, d_i)| \leq e^{m_i d_i} G(W_{i-1}) / G(W_i). \quad (47)$$

Now given $(Z_1, \dots, Z_s) \in Z$, we can apply (47) for $(X, W_1), \dots, (W_{s-1}, W_s)$. , and apply (44) for (W_s, Y) . We get

$$\begin{aligned}
|\mathcal{P}(W_1, \dots, W_s)| &\leq e^{m_1 d_1} G(X)/G(W_1) \dots e^{m_s d_s} G(W_{s-1})/G(W_s) \times \\
&\times e^{m_{s+1} d_{s+1}} G(W_s) G(Y) = \frac{e^{(h-k+s)\tau}}{e^{\tau_1 + \dots + \tau_s}} G(X) G(Y) = e^{(h-k+s)\tau} \times e^{\tau_{s+1} + \dots + \tau_k} G(Y) \leq \\
&\leq e^{h\tau} G(Y)^2.
\end{aligned}$$

We are using (46) to obtain the last inequality. Now we have,

$$|\mathcal{P}| \leq |Z| e^{h\tau} G(Y)^2 \leq \tau^{c_g} e^{h\tau} G(Y)^2,$$

where $c_g = O(g^2)$. □

References

- [Ath] J. Athreya. *Quantitative recurrence and large deviations for Teichmüller geodesic flow*. *Geom. Dedicata* 119 (2006), 121–140.
- [ABEM] J. Athreya, A. Bufetov, A. Eskin and M. Mirzakhani. *Lattice Point Asymptotics and Volume Growth on Teichmüller space*. Preprint
- [Bu] A. Bufetov. *Logarithmic asymptotics for the number of periodic orbits of the Teichmüller flow on Veech’s space of zippered rectangles*. Preprint.
- [EMM] A. Eskin, G. Margulis, S. Mozes. *Upper bounds and asymptotics in a quantitative version of the Oppenheim conjecture*. *Ann. of Math. (2)* **147**(1998), no. 1, 93–141
- [EM] A. Eskin and H. Masur. Asymptotic formulas on flat surfaces. *Ergodic Theory Dynam. Systems*, **21** (2001), 443–478.
- [Fo] G. Forni, Deviation of ergodic averages for area-preserving flows on surfaces of higher genus. *Ann. of Math. (2)*, 155 (2002), no. 1, 1–103.
- [H1] U. Hamenstädt. *Bernoulli Measures for the Teichmüller Flow*. Preprint.
- [H2] U. Hamenstädt. *Dynamics of the Teichmüller flow on compact invariant sets*. Preprint.
- [KH] A. Katok, B. Hasselblat Introduction to the Modern Theory of Dynamical Systems. Cambridge University Press, 1995.
- [Ker] S. Kerckhoff. The asymptotic geometry of Teichmüller space. *Topology* **19**(1980), 23–41.
- [Mar] G.A. Margulis, On some aspects of the theory of Anosov flows, Ph.D. Thesis, 1970, Springer, 2003.
- [Mi] Y. Minsky. Extremal length estimates and product regions in Teichmüller space. *Duke Math. J.* 83 (1996), no. 2, 249–286.
- [Ra1] K. Rafi. *Closed geodesics in the thin part of moduli space*. In preparation.
- [Ra2] K. Rafi. *Thick-thin decomposition of quadratic differentials*. preprint, 2005.
- [Ve] W. Veech, *The Teichmüller geodesic flow*, *Annals of Mathematics*, 1986.