

**SOLUTION OF PARTIAL DIFFERENTIAL EQUATIONS BY
METHOD OF HYPERHOLOMORPHIC FUNCTIONS**

ANATOLIY A. POGORUI

ABSTRACT. It is well known that the real and imaginary parts of any holomorphic function are harmonic functions of two variables. In this paper we generalize this property to finite-dimensional commutative algebras. We prove that if some basis of a subspace of a commutative algebra satisfies a polynomial equation then components of a hyperholomorphic function on the subspace are solutions of the respective partial differential equation.

Let \mathbf{A} be a finite-dimensional commutative unitary algebra over $K = \mathbb{R}$ (or \mathbb{C}), a set of vectors $\vec{e}_0, \vec{e}_1, \dots, \vec{e}_n$ be a basis of \mathbf{A} , and \vec{e}_0 be the unit of the algebra. Suppose \mathbf{B} is m -dimensional subspace of \mathbf{A} with the basis $\vec{e}_0, \vec{e}_1, \dots, \vec{e}_m$, $m \leq n$. Consider a function $\vec{f}: \mathbf{B} \rightarrow \mathbf{A}$ of the following form

$$\vec{f}(\vec{x}) = \sum_{k=0}^n \vec{e}_k u_k(\vec{x}),$$

where $u_k(\vec{x}) = u_k(x_0, x_1, \dots, x_m)$ are real (or complex) functions of $m + 1$ arguments.

Definition 1. $\vec{f}(\vec{x})$ is called differentiable at a point $\vec{x}_0 \in \mathbf{B}$ if there exists the function $\vec{f}': \mathbf{B} \rightarrow \mathbf{A}$ such that for any $\vec{h} \in \mathbf{B}$

$$\vec{h}\vec{f}'(\vec{x}_0) = \lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x}_0 + \varepsilon\vec{h}) - \vec{f}(\vec{x}_0)}{\varepsilon}, \quad (1)$$

where \vec{f}' doesn't depend on \vec{h} .

A function \vec{f} is said to be hyperholomorphic if \vec{f} is differentiable at every point of \mathbf{B} .

Theorem 1. A function $\vec{f}(\vec{x}) = \sum_{k=0}^n \vec{e}_k u_k(\vec{x})$ is hyperholomorphic if and only if there exists the function $\vec{f}': \mathbf{B} \rightarrow \mathbf{A}$ such that for all $k = 0, 1, \dots, m$, and $\forall \vec{x} \in \mathbf{B}$

$$\vec{e}_k \vec{f}'(\vec{x}) = \lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x} + \varepsilon \vec{e}_k) - \vec{f}(\vec{x})}{\varepsilon}, \quad (2)$$

where \vec{f}' doesn't depend on \vec{e}_k .

Date: November 16, 2008.

2000 Mathematics Subject Classification. Primary 35C99; Secondary 32W50.

Key words and phrases. Hyperholomorphic function, commutative algebra, partial differential equation.

Proof. Suppose that (2) is fulfilled, then it is easily verified that

$$\begin{aligned}
\vec{f}' &= \lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x} + \varepsilon \vec{e}_0) - \vec{f}(\vec{x})}{\varepsilon} = \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}, \\
\vec{e}_1 \vec{f}' &= \lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x} + \varepsilon \vec{e}_1) - \vec{f}(\vec{x})}{\varepsilon} = \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_1} = \vec{e}_1 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}, \\
&\vdots \\
\vec{e}_m \vec{f}' &= \lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x} + \varepsilon \vec{e}_m) - \vec{f}(\vec{x})}{\varepsilon} = \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_m} = \vec{e}_m \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}.
\end{aligned} \tag{3}$$

Consider $\vec{h} = \sum_{k=0}^m h_k \vec{e}_k$. It follows from Eqs.(3) that

$$\begin{aligned}
h_0 \vec{f}' &= h_0 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}, \\
h_1 \vec{e}_1 \vec{f}' &= h_1 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_1}, \\
&\vdots \\
h_m \vec{e}_m \vec{f}' &= h_m \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_m}.
\end{aligned}$$

This implies that

$$\begin{aligned}
\vec{h} \vec{f}' &= h_0 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0} + h_1 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_1} + \dots + h_m \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_m} = \\
&\lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x} + \varepsilon \vec{h}) - \vec{f}(\vec{x})}{\varepsilon}.
\end{aligned}$$

Furthermore, it follows from Eqs.(3) that

$$\begin{aligned}
&h_0 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0} + h_1 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_1} + \dots + h_m \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_m} = \\
&h_0 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0} + h_1 \vec{e}_1 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0} + \dots + h_m \vec{e}_m \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}.
\end{aligned}$$

Therefore, for every $\vec{h} \in \mathbf{B}$

$$\vec{h} \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0} = \lim_{\varepsilon \rightarrow 0} \frac{\vec{f}(\vec{x} + \varepsilon \vec{h}) - \vec{f}(\vec{x})}{\varepsilon}$$

or

$$\vec{f}' = \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0} \tag{4}$$

□

By using Eq.(3), we have

$$\begin{aligned}
 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_1} &= \vec{e}_1 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}, \\
 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_2} &= \vec{e}_2 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}, \\
 &\vdots \\
 \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_m} &= \vec{e}_m \sum_{k=0}^n \vec{e}_k \frac{\partial u_k}{\partial x_0}.
 \end{aligned} \tag{5}$$

Eqs.(5) will be called the Cauchy-Riemann type conditions. In paper [1] we studied hyperholomorphic functions $\vec{f}: \mathbf{A} \rightarrow \mathbf{A}$.

1. SOLUTION OF SOME DIFFERENTIAL EQUATIONS

Let us consider the following partial differential equation

$$\sum_{i_0+i_1+\dots+i_m=r} C_{i_0 i_1 \dots i_m} \frac{\partial^r}{\partial x_0^{i_0} \partial x_1^{i_1} \dots \partial x_m^{i_m}} u(x_0, x_1, \dots, x_m) = 0, \tag{6}$$

where $C_{i_0 i_1 \dots i_m} \in K$ are constant coefficients.

Theorem 2. *Suppose that some subspace \mathbf{B} of algebra \mathbf{A} has a basis $\vec{e}_0, \vec{e}_1, \dots, \vec{e}_m$ such that*

$$\sum_{i_0+i_1+\dots+i_m=r} C_{i_0 i_1 \dots i_m} (\vec{e}_0)^{i_0} (\vec{e}_1)^{i_1} \dots (\vec{e}_m)^{i_m} = 0,$$

and a function $\vec{f}(\vec{x}) = \sum_{k=0}^n \vec{e}_k u_k(\vec{x})$ ($\vec{f}: \mathbf{B} \rightarrow \mathbf{A}$) is hyperholomorphic.

Then functions $u_k(\vec{x})$, $k = 0, 1, \dots, n$ are solutions of Eq.(6).

Proof. It follows from the Cauchy-Riemann condition (5) that

$$\frac{\partial \vec{f}}{\partial x_k} = \vec{e}_k \frac{\partial \vec{f}}{\partial x_0}, \quad k = 0, 1, \dots, m.$$

This implies

$$\frac{\partial^i \vec{f}}{\partial x_k^i} = (\vec{e}_k)^i \frac{\partial^i \vec{f}}{\partial x_0^i}.$$

Therefore, we obtain

$$\begin{aligned}
 \sum_{i_0+i_1+\dots+i_m=r} C_{i_0 i_1 \dots i_m} \frac{\partial^r}{\partial x_0^{i_0} \partial x_1^{i_1} \dots \partial x_m^{i_m}} \vec{f}(x_0, x_1, \dots, x_m) = \\
 \frac{\partial^r \vec{f}(x_0, x_1, \dots, x_m)}{\partial x_0^r} \sum_{i_0+i_1+\dots+i_m=r} C_{i_0 i_1 \dots i_m} (\vec{e}_0)^{i_0} (\vec{e}_1)^{i_1} \dots (\vec{e}_m)^{i_m} = 0,
 \end{aligned}$$

□

Example 1. In case where \mathbf{A} and \mathbf{B} are complex numbers a differentiable function is holomorphic. Basis $\vec{e}_0 = 1, \vec{e}_1 = i$ satisfies the equation $(\vec{e}_0)^2 + (\vec{e}_1)^2 = 0$. Therefore, the real and imaginary parts of any holomorphic function satisfy the Laplace equation

$$\frac{\partial^2}{\partial x^2} u(x, y) + \frac{\partial^2}{\partial y^2} u(x, y) = 0.$$

In paper [2] it is considered algebras which differentiable functions have harmonic components that satisfy the three-dimensional Laplace equation.

REFERENCES

1. POGORUI A.A. (2007) Hyperholomorphic functions on commutative algebras *Complex Variables and Elliptic Equations*. Vol. 52, No. 12, 1155-1159.
2. KETCHUM P.W. (1929) A complete solution of Laplace's equation by an infinite hypervariable. *American Journal of Mathematics*. 51, 179-188.

DEPARTMENT OF MATHEMATICS, ZHYTOMYR STATE UNIVERSITY, ZHYTOMYR, UKRAINE 10008
E-mail address: `pogor@zu.edu.ua`