A Z-BASIS FOR THE CLUSTER ALGEBRA ASSOCIATED TO AN AFFINE QUIVER

MING DING, JIE XIAO AND FAN XU

ABSTRACT. The canonical bases of cluster algebras of finite types and rank 2 are given explicitly in [\[4\]](#page-19-0) and [\[14\]](#page-20-0) respectively. In this paper, we will deduce Z-bases for cluster algebras for affine types $\widetilde{A}_{n,n}$, \widetilde{D} and \widetilde{E} . Moreover, we give an inductive formula for computing the multiplication between two generalized cluster variables associated to objects in a tube.

1. INTRODUCTION

Cluster algebras were introduced by S. Fomin and A. Zelevinsky [\[11\]](#page-20-1) in order to develop a combinatorial approach to study problems of total positivity. By the work of [\[13\]](#page-20-2), the link between acyclic cluster algebras and representation theory of quivers found its general framework in [\[1\]](#page-19-1) where the authors introduced the cluster category. Let Q be an acyclic quiver with vertex set $Q_0 = \{1, 2, \dots, n\}$. Let $A = \mathbb{C}Q$ be the path algebra of Q and we denote by P_i the indecomposable projective $\mathbb{C} Q$ -module with the simple top S_i corresponding to $i \in Q_0$ and I_i the indecomposable injective $\mathbb{C} Q$ -module with the simple socle S_i . Let $\mathcal{D}^b(Q)$ be the bounded derived category of mod $\mathbb{C}Q$ with the shift functor T and the AR-translation τ . The cluster category associated to Q is the orbit category $\mathcal{C}(Q) := \mathcal{D}^b(Q)/F$ with $F = T \circ \tau^{-1}$. Let $\mathbb{Q}(x_1, \dots, x_n)$ be a transcendental extension of \mathbb{Q} . The Caldero-Chapton map of an acyclic quiver Q is the map

$$
X_?^Q: \mathrm{obj}(\mathcal{C}(Q)) \to \mathbb{Q}(x_1, \cdots, x_n)
$$

defined in [\[3\]](#page-19-2) by the following rules:

(1) if M is an indecomposable $\mathbb{C}Q$ -module, then

$$
X_M^Q = \sum_{\mathbf{e}} \chi(\text{Gr}_{\underline{e}}(M)) \prod_{i \in Q_0} x_i^{-\langle \underline{e}, s_i \rangle - \langle s_i, \underline{\text{dim}} M - \underline{e} \rangle};
$$

(2) if $M = TP_i$ is the shift of the projective module associated to $i \in Q_0$, then

$$
X_M^Q = x_i;
$$

(3) for any two objects M, N of C_Q , we have

$$
X_{M \oplus N}^{Q} = X_M^Q X_N^Q.
$$

Here, we denote by $\langle -, - \rangle$ the Euler form on $\mathbb{C}Q$ -mod and $Gr_e(M)$ is the e -Grassmannian of M, i.e. the variety of submodules of M with dimension vector \underline{e} . For any object $M \in \mathcal{C}(Q)$, X_M^Q will be called the generalized cluster variable for M.

We note that the indecomposable $\mathbb{C}Q$ -modules and TP_i for $i \in Q_0$ exhaust the indecomposable objects of the cluster category $\mathcal{C}(Q)$:

$$
ind - C(Q) = ind - CQ \sqcup \{TP_i : i \in Q_0\}.
$$

Key words and phrases: Z-basis, cluster algebra.

Each object M in $C(Q)$ can be uniquely decomposed in the following way:

$$
M=M_0\oplus TP_M
$$

where M_0 is a module and P_M is a projective module. The module M_0 can be recovered using the following homological functor:

$$
H^0 = \text{Hom}_{\mathcal{C}(Q)}(\mathbb{C}Q, -): \mathcal{C}(Q) \longrightarrow \mathbb{C}Q - \text{mod}
$$

Hence we can rewrite $M = M_0 \oplus TP_M$ as:

$$
M = H^0(M) \oplus TP_M.
$$

Let $R = (r_{ij})$ be a matrix of size $|Q_0| \times |Q_0|$ satisfying

$$
r_{ij} = \dim_{\mathbb{C}} \operatorname{Ext}^1(S_i, S_j)
$$

for any $i, j \in Q_0$. The Caldero-Chapton map can be reformulated by the following rules [\[16\]](#page-20-3)[\[12\]](#page-20-4):

(1)

(2)

$$
X_{\tau P} = X_{TP} = x^{\text{dim}P/radP}, X_{\tau^{-1}I} = X_{T^{-1}I} = x^{\text{dim}socI}
$$

for any projective $\mathbb{C} Q$ -module P and any injective $\mathbb{C} Q$ -module I;

$$
X_M = \sum_{\underline{e}} \chi(\text{Gr}_{\underline{e}}(M)) x^{\underline{e}R + (\underline{\text{dim}}M - \underline{e})R^{tr} - \underline{\text{dim}}M}
$$

where M is a $\mathbb{C}Q$ -module, R^{tr} is the transpose of the matrix R and $x^v =$ $x_1^{v_1} \cdots x_n^{v_n}$ for $v = (v_1, \dots, v_n) \in \mathbb{Z}^n$.

Let $\mathcal{AH}(Q)$ be the subalgebra of $\mathbb{Q}(x_1, \dots, x_n)$ generated by

 $\{X_M, X_{\tau P} \mid M, P \in \text{mod} - \mathbb{C}Q, P \text{ is projective module}\}.$

Let $\mathcal{E}\mathcal{H}(Q)$ be the subalgebra of $\mathcal{A}\mathcal{H}(Q)$ generated by

$$
\{X_M \mid M \in \text{ind} - C(Q), \text{Ext}^1_{C(Q)}(M, M) = 0 \}.
$$

In [\[5\]](#page-19-3), the authors showed that the Caldero-Chapton map induces a one to one correspondence between indecomposable objects in $\mathcal{C}(Q)$ without self-extension and the cluster variables of the cluster algebra $\mathcal{A}(Q)$. Hence, one can view $\mathcal{E}\mathcal{H}(Q)$ as the cluster algebra for a quiver Q. When Q is a simply laced Dynkin quiver, $\mathcal{AH}(Q)$ is coincides with $\mathcal{E}\mathcal{H}(Q)$. In [\[4\]](#page-19-0), the authors showed a Z-basis of $\mathcal{AH}(Q)$. When Q is the Kronecker quiver, $\mathcal{AH}(Q)$ is also coincides with $\mathcal{EH}(Q)$. In [\[6\]](#page-19-4), the authors gave a Z-basis of $\mathcal{AH}(Q)$ called the semicanonical basis. When Q is the quiver of type \tilde{D}_4 , $\mathcal{AH}(Q)$ is still equal to $\mathcal{EH}(Q)$ and a Z-basis is given in [\[8\]](#page-19-5)(also see Section [6.3\)](#page-14-0).

Now let Q be an affine quiver of type \widetilde{A}_n , \widetilde{D} or \widetilde{E} . There are many references about the theory of representations of affine quivers, for example, see [\[2\]](#page-19-6) and [\[7\]](#page-19-7). The main goal of this paper is to give a \mathbb{Z} -basis of $\mathcal{AH}(Q)$ for an affine quiver Q. We will prove the following theorem.

Theorem 1.1. Let Q be an affine quiver. A \mathbb{Z} -basis of $\mathcal{AH}(Q)$ is the following set: ${X_L, X_{T \oplus R} | \underline{\dim}(T_1 \oplus R_1) \neq \underline{\dim}(T_2 \oplus R_2), Ext^1_{\mathcal{C}(Q)}(T, R) = 0, Ext^1_{\mathcal{C}(Q)}(L, L) = 0}$ where L is any non-regular exceptional object, R is 0 or any regular exceptional module and T is 0 or any indecomposable regular module with self-extension and in addition if $Q \neq A_{1,1}$ and $\dim(T \oplus R) = m\delta$ for some $m \in \mathbb{N}$, then $R = 0$ and T is an indecomposable module of dimension vector mδ in a non-homogeneous tube.

Moreover, we give an inductive formula for computing the multiplication between two generalized cluster variables associated to objects in a tube in Section 6. Our tools are the following cluster multiplication theorems.

Theorem 1.2. [\[15\]](#page-20-5) (1) For any A-modules $V_{\xi'}$, $V_{\eta'}$ we have

$$
d^1(\xi',\eta')X_{V_{\xi'}}X_{V_{\eta'}} = \int_{\lambda \neq \xi' \oplus \eta'} \chi(\mathbb{P}Ext^1_A(V_{\xi'},V_{\eta'})_{V_{\lambda}})X_{V_{\lambda}}
$$

$$
+ \int_{\gamma,\beta} \chi(\mathbb{P}Hom_A(V_{\eta'}, \tau V_{\xi'})_{V_{\beta}[1] \oplus V_{\gamma}})X_{V_{\beta}}X_{\tau^{-1}V_{\gamma}}
$$

where $V_{\xi'}$ has no projective direct summand and $d^1(\xi', \eta') = \dim_{\mathbb{C}} Ext_A(V_{\xi'}, V_{\eta'}).$ (2) For any A-module $V_{\xi'}$ and $P \in \rho$ is projective Then

$$
d(\rho, \xi') X_{V_{\xi'}} x^{\underline{\dim} P/radP} = \int_{\delta, \iota'} \chi(\mathbb{P} \text{Hom}_{A}(V_{\xi'}, I)_{V_{\delta}[1] \oplus I'}) X_{V_{\delta}} x^{\underline{\dim} \text{soc} I'}
$$

+
$$
\int_{\gamma, \rho'} \chi(\mathbb{P} \text{Hom}_{A}(P, V_{\xi'})_{P'[1] \oplus V_{\gamma}}) X_{V_{\gamma}} x^{\underline{\dim} P'/radP'}
$$

where $I = DHom_A(P, A)$, $I' \in \iota'$ injective, $P' \in \rho'$ projective, and $d(\rho, \xi') =$ dim_CHom_A $(V_\rho, V_{\xi'})$.

Theorem 1.3. [\[3\]](#page-19-2) Let Q be an acyclic quiver and M any indecomposable nonprojective A-module, then

$$
X_M X_{\tau M} = 1 + X_E
$$

where E is the middle term of the Auslander-Reiten sequence ending in M .

Theorem 1.4. [\[5\]](#page-19-3) Let Q be an acyclic quiver and M,N be any two objects in $\mathcal{C}(Q)$ such that dim $Ext^1_{\mathcal{C}(Q)}(M,N) = 1$, then

$$
X_M X_N = X_B + X_{B'}
$$

where B and B' are the unique objects such that there exists non-split triangles

$$
M \longrightarrow B \longrightarrow N \longrightarrow M[1]
$$
 and $N \longrightarrow B' \longrightarrow M \longrightarrow N[1]$

In [\[10\]](#page-20-6), the author construct a Z-basis for a cluster algebra of type \widetilde{A} referred as the semicanonical basis. It is interesting to compare it to the Z-bases in this paper.

2. Numerators of Laurent expansions in generalized cluster **VARIABLES**

In the following, we will suppose that Q is one of $\widetilde{A}_{n,n}$, \widetilde{D} and \widetilde{E} with an orientation where every vertex is a sink or a source. For any object $M = M_0 \oplus (\oplus_i s_i T P_i) \in$ $\mathcal{C}(Q)$ where M_0 is a module, we extend the dimension vector to the objects in the cluster category by setting

$$
\underline{\dim}(M) = \underline{\dim}(M_0) - (s_1, \cdots, s_n).
$$

Let $E_i[n]$ be the indecomposable regular module with quasi-socle E_i and quasilength n and $X_0 = 1$. For any CQ-module M, we denoted by $\dim_{\mathbb{C}} M(i)$ the *i*-th component of dimM.

Definition 2.1. For $M, N \in \mathcal{C}(Q)$ with $\dim(M) = (m_1, \dots, m_n)$ and $\dim(N) =$ (r_1, \dots, r_n) , we write $\underline{\dim}(M) \preceq \underline{\dim}(N)$ if $m_i \leq r_i$ for $1 \leq i \leq n$. Moreover, if there exists some i such that $m_i < r_i$, then we write $\dim(M) \prec \dim(N)$.

Remark 2.2. Note that for the cluster multiplication formula in the Theorem $1.2(1)$ $1.2(1)$, we have the following exact sequences:

$$
0 \longrightarrow V_{\eta'} \longrightarrow V_{\lambda} \longrightarrow V_{\xi'} \longrightarrow 0
$$

and

$$
0 \longrightarrow V_{\beta} \longrightarrow V_{\eta'} \longrightarrow \tau V_{\xi'} \longrightarrow V_{\gamma} \longrightarrow 0.
$$

If $\dim_{\mathbb{C}} Ext_A(V_{\xi'}, V_{\eta'}) \neq 0 \implies \underline{\dim} V_{\beta} + \underline{\dim} \tau^{-1} V_{\gamma} \prec \underline{\dim} V_{\eta'} + \underline{\dim} V_{\xi'} = \underline{\dim} V_{\lambda}$ because $\tau V_{\xi'}$ has no injective summand.

According to the definition of the Caldero-Chapton map, we consider the Laurent expansions in generalized cluster variables $X_M = \frac{P(x)}{\prod_{x \in \mathcal{X}} P(x)}$ $\frac{F(x)}{\prod_{1 \leq i \leq n} x_i^{m_i}}$ for $M \in \mathcal{C}(Q)$ such that the integral polynomial $P(x)$ in the variables x_i is not divisible by any x_i . We define the denominator vector of X_M as (m_1, \dots, m_n) [\[10\]](#page-20-6). The following theorem is called as the denominator theorem.

Theorem 2.3. [\[4\]](#page-19-0) Let Q be an acyclic quiver. Then for any object M in $\mathcal{C}(Q)$, the denominator vector of X_M is $\underline{\dim}(M)$.

According to the denominator Theorem [2.3,](#page-3-0) we can prove the following propositions.

Proposition 2.4. If M is P_i or I_i for $1 \leq i \leq n$, then $X_M = \frac{P(x)}{x^{\dim(M)}}$ where the constant term of $P(x)$ is 1.

Proof. 1) If i is a sink point, we have the following short exact sequence:

$$
0 \longrightarrow P_i \longrightarrow I_i \longrightarrow I' \longrightarrow 0
$$

Then by the cluster multiplication theorem in Theorem [1.2,](#page-2-0) we have:

$$
X_{\tau P_i} X_{P_i} = x^{\underline{dim} socI'} + 1
$$

Thus the constant term of numerator in X_{P_i} as an irreducible fraction of integral polynomials in the variables x_i is 1 because of $X_{\tau P_i} = x_i$.

If i is a source point, we have the following short exact sequence:

$$
0 \longrightarrow P' \longrightarrow P_i \longrightarrow I_i \longrightarrow 0
$$

Similarly we have:

$$
X_{\tau P_i} X_{P_i} = X_{P'} + 1
$$

Thus we can finish it by induction on P' .

2) For X_{I_i} , it is totally similar.

Note that $X_{\tau P_i} = x_i = \frac{1}{\tau_i}$ $\frac{1}{x_i^{-1}}$ and $\underline{\dim}(\tau P_i) = (0, \cdots, 0, -1, 0, \cdots, 0)$ with i-th component 1 and others 0. Hence we denote the denominator of $X_{\tau P_i}$ by x_i^{-1} , and assert the constant term of numerator in $X_{\tau P_i}$ is 1. With these notations, we have the following Proposition [2.5.](#page-3-1)

Proposition 2.5. For any object $M \in \mathcal{C}(Q)$, then $X_M = \frac{P(x)}{x^{\dim(M)}}$ where the constant term of $P(x)$ is 1.

Proof. It is enough to consider the case that M is an indecomposable module.

1) When M is an indecomposable preprojective module, then by exchange relation in Thereom [1.3](#page-2-1) we have

$$
X_M X_{\tau M} = \prod_i X_{B_i} + 1.
$$

Thus by Proposition [2.4,](#page-3-2) we can prove that $X_M = \frac{P(x)}{x^{\dim(M)}}$ where the constant term of $P(x)$ is 1 by induction with the help of the directness of AR-quiver in the preprojective component of mod $\mathbb{C}Q$. The discussion is similar for any indecomposable preinjective module.

2) When M is an indecomposable regular module, we only need to prove that the proposition holds for any regular simple module according to the exchange relations.

First, suppose M is in some homogeneous tube with dimension vector δ . Note that there exists a point e such that $\dim \delta(e) = 1$. Thus we have

$$
\dim Ext^1_{\mathbb{C} Q}(M, P(e)) = \dim Hom_{\mathbb{C} Q}(P(e), M) = 1.
$$

Then we obtain the following two non-split exact sequences:

$$
0 \longrightarrow P(e) \longrightarrow L \longrightarrow M \longrightarrow 0
$$

and

$$
0 \longrightarrow L' \longrightarrow P(e) \longrightarrow M \longrightarrow L'' \longrightarrow 0
$$

where L and L' are preprojective modules and L'' is a preinjective module. Using Theorem [1.2](#page-2-0) or Theorem [1.4,](#page-2-2) we have

$$
X_M X_{P(e)} = X_L + X_{L'} X_{\tau^{-1} L''}
$$

where $\underline{\dim}(L' \oplus \tau^{-1}L'') \prec \underline{\dim}(P(e) \oplus M)$.

We have already known that the constant term of the numerator in $X_{P(e)}$, X_L as an irreducible fraction of integral polynomials in the variables x_i is 1 by 1), then the constant term of the numerator in X_M must be 1.

Now we consider these non-homogeneous tubes. Note that by the AR formula in Theorem [1.3,](#page-2-1) we only need to prove the constant term of the numerator in X_{E_i} is 1 for $1 \leq i \leq r$. Suppose M is a regular simple module such that $\dim M(e)$ $\dim \delta(e) = 1$. We denote M by E_1 , thus $\dim E_1(e) = 1$, and $\dim E_i(e) = 0$ for $2 \leq i \leq r$. Therefore $\dim Ext^1(E_2, P(e)) = 1$, then we have the following non-split exact sequences combining the relation $\tau E_2 = E_1$

$$
\quad\text{and}\quad
$$

$$
0 \longrightarrow L' \longrightarrow P(e) \longrightarrow E_1 \longrightarrow L'' \longrightarrow 0
$$

 $0 \longrightarrow P(e) \longrightarrow L \longrightarrow E_2 \longrightarrow 0$

where L and L' are preprojective modules and L'' is a preinjective module. Then we have

$$
X_{E_2}X_{P(e)} = X_L + X_{L'}X_{\tau^{-1}L''}
$$

where $\underline{\dim}(L' \oplus \tau^{-1}L'') \prec \underline{\dim}(P(e) \oplus E_2)$.

We have already known that the constant term of the numerator in $X_{P(e)}$, X_L as an irreducible fraction of integral polynomials in the variables x_i is 1 by 1), then the constant term of the numerator in X_{E_2} must be 1.

Note that $\dim E_1[2](e) = \dim E_1(e) + \dim E_2(e) = 1$, by similar discussions, we can obtain the constant term of the numerator in $X_{E_1[2]}$ must be 1. Thus by $X_{E_1}X_{E_2} = X_{E_1[2]} + 1$, we obtain that the constant term of the numerator in X_{E_1} must be 1. Using the same method, we can prove the constant term of the numerator in X_{E_i} must be 1 for $3 \leq i \leq r$.

3. Generalized cluster variables on tubes

Let Q be an affine quiver with the minimal imaginary root $\delta = (\delta_i)_{i \in Q_0}$. Then tubes of indecomposable regular CQ-modules are indexed by the projective line \mathbb{P}^1 . Let λ be the index of a homogeneous tube and $M(\lambda)$ be the regular simple $\mathbb{C}Q$ -module with dimension vector δ in this homogeneous tube. Let M[i] be the regular module with quasi-socle M and quasi-regular length i for any $i \in \mathbb{N}$. Let X_M be the generalized cluster variable associated to M by the reformulation of the Caldero-Chapton map. Then we have

Proposition 3.1. Let λ and μ be in \mathbb{P}^1 such that $M(\lambda)$ and $M(\mu)$ are two regular simple modules of dimension vector δ . Then $X_{M(\lambda)} = X_{M(\mu)}$.

Proof. Choose a vertex $p \in Q_0$ such that $\delta_p = 1$. We assume that p is a sink. Let Q have the underlying graph not of type A_n . Then there is unique edge $\alpha \in Q_1$ with the head p and tail p'. It is easy to check $\delta_{p'} = 2$. Let P_p be the indecomposable projective module corresponding p and $I(\delta - p)$ be the indecomposable preinjective module of dimension vector $\delta - \underline{\dim} S_p$. Then $\dim_{\mathbb{C}} Ext_{\mathbb{C}Q}^1(I(\delta - p), P_p) = 2$. Given

any $\epsilon \in Ext^1_{\mathbb{C}Q}(I(\delta - p), P_p)$, we have a short exact sequence whose equivalence class is ϵ as follows:

$$
\varepsilon: \quad 0 \longrightarrow P_p \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} M_{\epsilon} \xrightarrow{\begin{pmatrix} 0 & 1 \end{pmatrix}} I(\delta - p) \longrightarrow 0
$$

where $(M_{\epsilon})_i = (P_p)_i \oplus I(\delta - p)_i$ for any $i \in Q_0$, $(M_{\epsilon})_{\beta} = I(\delta - p)_{\beta}$ for $\beta \neq \alpha$ and $(M_{\epsilon})_{\alpha}$ is

$$
(M_\epsilon)_\alpha=\left(\begin{array}{cc} 0 & m(\epsilon,\alpha) \\ 0 & 0 \end{array}\right)
$$

where $m(\epsilon, \alpha) \in Hom_{\mathbb{C}}(I(\delta - p)'_p, P_p)$. For any $\epsilon, \epsilon' \in Ext^1_{\mathbb{C}Q}(I(\delta - p), P_p), M_{\epsilon} \cong M_{\epsilon'}$ if and only if $m(\epsilon, \alpha) = tm(\epsilon', \alpha)$ for some $t \in \mathbb{C}$. The regular simple $\mathbb{C}Q$ -modules (denoted by $M(\lambda)$) with dimension vector δ satisfy that $M(\lambda)_{\alpha}$ is as follows

$$
M(\lambda)_{p'} = \mathbb{C}^2 \xrightarrow{\begin{pmatrix} 1 & \lambda \end{pmatrix}} M(\lambda)_p = \mathbb{C}
$$

where $\lambda \in \mathbb{C}^* \setminus \{1\}$. Let $M(\lambda)$ and $M(\lambda')$ be any two regular simple $\mathbb{C}Q$ -modules with dimension vector δ . Let P is an indecomposable projective module such that $P \subseteq M(\lambda)$ and $(\underline{\dim} P)_p = 0$. Then P is also a submodule of $M(\lambda')$ and $(\underline{\dim} P)_{p'} =$ 0. Let P be an indecomposable projective module such that $P \subseteq M(\lambda)$ and $(\underline{\dim} P)_p = 1$. Assume that P_α is $\mathbb{C} \xrightarrow{a+b\lambda} \mathbb{C}$, then there exists $P' \in Gr_{\underline{e}}(M(\lambda'))$ such that $P' \cong P$ and P'_α is $\mathbb{C} \xrightarrow{a+b\lambda'} \mathbb{C}$. Since $\tau M(\lambda) = M(\lambda)$, we know $\tau^{-i}P$ and $\tau^{-1}P'$ are the submodules of $M(\lambda)$ and $M(\lambda')$ for any $i \in \mathbb{N}$, respectively. Hence, any preprojective submodule X of $M(\lambda)$ corresponds to a preprojective submodule X' of $M(\lambda')$ and $X \cong X'$. Let Q be of type \widetilde{A}_n . Then there are two adjacent edge α and β. Any regular simple module $M(\lambda)$ satisfies that $M(\lambda)$ _α is as follows:

$$
\mathbb{C}\overset{1}{\longrightarrow}\mathbb{C}\overset{\lambda}{\longleftarrow}\mathbb{C}~.
$$

The discussion is similar as above. If p is a source, the discussion is also similar. Therefore, there is a homeomorphism between $Gr_{\underline{e}}(M(\lambda))$ and $Gr_{\underline{e}}(M(\lambda'))$ for any dimension vector <u>e</u>. By definition, $X_{M(\lambda)} = X_{M(\mu)}$. .

We note that there is an alternative proof of Proposition [3.1](#page-4-0) in [\[10,](#page-20-6) Lemma 3.14].

Proposition 3.2. For any $m, n \in \mathbb{N}$ and $m \geq n$, we have

$$
X_{M[m]}X_{M[n]} = X_{M[m+n]} + X_{M[m+n-2]} + \cdots + X_{M[m-n+2]} + X_{M[m-n]}.
$$

Proof. When $n = 1$, we know $\dim_{\mathbb{C}} \mathrm{Ext}^1(M[m], M) = \dim_{\mathbb{C}} \mathrm{Hom}(M, M[m]) = 1$. The involving non-split short exact sequences are

$$
0 \longrightarrow M \longrightarrow M[m+1] \longrightarrow M[m] \longrightarrow 0
$$

and

$$
0 \longrightarrow M \longrightarrow M[m] \longrightarrow M[m-1] \longrightarrow 0.
$$

Thus by the cluster multiplication theorem in Theorem [1.2](#page-2-0) or Theorem [1.4](#page-2-2) and the fact $\tau M[k] = M[k]$ for any $k \in \mathbb{N}$, we obtain the equation

$$
X_{M[m]}X_M = X_{M[m+1]} + X_{M[m-1]}
$$

Suppose that it is right for $n \leq k$. When $n = k + 1$, we have

$$
X_{M[m]}X_{M[k+1]} = X_{M[m]}(X_{M[k]}X_{M} - X_{M[k-1]}) = X_{M[m]}X_{M[k]}X_{M} - X_{M[m]}X_{M[k-1]}
$$

=
$$
\sum_{i=0}^{k} X_{M[m+k-2i]}X_{M} - \sum_{i=0}^{k-1} X_{M[m+k-1-2i]}
$$

A Z-BASIS FOR THE CLUSTER ALGEBRA ASSOCIATED TO AN AFFINE QUIVER $\,$ 7 $\,$

$$
= \sum_{i=0}^{k} (X_{M[m+k+1-2i]} + X_{M[m+k-1-2i]}) - \sum_{i=0}^{k-1} X_{M[m+k-1-2i]}
$$

=
$$
\sum_{i=0}^{k+1} X_{M[m+k+1-2i]}.
$$

By Proposition [3.1](#page-4-0) and Proposition [3.2,](#page-5-0) we can define $X_{n\delta} := X_{M[n]}$ for $n \in \mathbb{N}$.

Now consider the non-homogeneous tubes $T(k)$ with rank r_k for $1 \leq k \leq m$. The corresponding regular simple modules are $E_{k,1}, \dots, E_{k,r_k}$ with $\tau E_{k,i+1} = E_{k,i}$. In fact $m = 2$ or 3 in our conditions. If we restrict the discussion to one tube, we will omit the index k for convenience. Set $q.soc(E_{k,i}[n]) = E_{k,i}$ and $X_{n\delta_{k,i}} = X_{E_{k,i}[nr_k]}$ for $n \in \mathbb{N}$.

Proposition 3.3. Let E_i and E_j be two regular simples in a non-homogeneous tube with rank r. Then we have

$$
X_{E_i[mr]} = X_{E_j[mr]} + X_{E_{i+1}[mr-2]} - X_{E_{j+1}[mr-2]}
$$

where $1 \leq i < j \leq r$ and $m \in \mathbb{N}$.

Proof. It is easy to prove that

$$
X_{E_i} X_{E_{i+1}[mr-1]} = X_{E_i[mr]} + X_{E_{i+2}[mr-2]},
$$

\n
$$
X_{E_{i+1}[mr-1]} X_{E_i} = X_{E_{i+1}[mr]} + X_{E_{i+1}[mr-2]}
$$

Hence, we have

$$
X_{E_i[mr]} = X_{E_{i+1}[mr]} + X_{E_{i+1}[mr-2]} - X_{E_{i+2}[mr-2]}.
$$

Similarly we have

$$
X_{E_{i+1}[mr]} = X_{E_{i+2}[mr]} + X_{E_{i+2}[mr-2]} - X_{E_{i+3}[mr-2]},
$$

\n
$$
X_{E_{i+2}[mr]} = X_{E_{i+3}[mr]} + X_{E_{i+3}[mr-2]} - X_{E_{i+4}[mr-2]},
$$

\n
$$
\vdots
$$

\n
$$
X_{E_{j-1}[mr]} = X_{E_j[mr]} + X_{E_j[mr-2]} - X_{E_{j+1}[mr-2]}.
$$

Thus

$$
X_{E_i[mr]} + X_{E_{i+1}[mr]} + \cdots + X_{E_{j-1}[mr]}
$$

= $(X_{E_{i+1}[mr]} + X_{E_{i+1}[mr-2]} - X_{E_{i+2}[mr-2]}) + (X_{E_{i+2}[mr]} + X_{E_{i+2}[mr-2]} - X_{E_{i+3}[mr-2]})$
+ $\cdots + (X_{E_j[mr]} + X_{E_j[mr-2]} - X_{E_{j+1}[mr-2]})$

 $= X_{E_{i+1}[mr]} + X_{E_{i+2}[mr]} + \cdots + X_{E_j[mr]} + X_{E_{i+1}[mr-2]} - X_{E_{j+1}[mr-2]}.$ Therefore

$$
X_{E_i[mr]} = X_{E_j[mr]} + X_{E_{i+1}[mr-2]} - X_{E_{j+1}[mr-2]}.
$$

 \Box

Example 3.4. Consider $r=3$, we have

$$
X_{E_1}X_{E_2}X_{E_3} = (X_{E_1[2]} + 1)X_{E_3} = X_{E_1[3]} + X_{E_1} + X_{E_3}.
$$

Similarly we have

$$
X_{E_2}X_{E_3}X_{E_1} = X_{E_2[3]} + X_{E_2} + X_{E_1},
$$

and

$$
X_{E_3}X_{E_1}X_{E_2} = X_{E_2[3]} + X_{E_3} + X_{E_2}.
$$

Therefore

$$
X_{E_1[3]} = X_{E_2[3]} + X_{E_2} - X_{E_3} = X_{E_3[3]} + X_{E_2} - X_{E_1}.
$$

Proposition 3.5. $X_{n\delta_{i,1}} = X_{n\delta_{j,1}} + \sum_{\underline{\dim}L \prec n\delta} a_L X_L$ for $i \neq j$ in different nonhomogeneous tubes where $a_L \in \mathbb{Q}$.

 \sum *Proof.* Denote $\delta = (v_1, v_2, \dots, v_n)$ and $\underline{\dim}(E_{i,j}) = (v_{j1}, v_{j2}, \dots, v_{jn})$, then $\delta =$ $_{1\leq j\leq r_i}$ $\underline{\dim}(E_{i,j})$. Thus by the cluster multiplication theorem in Theorem [1.2,](#page-2-0) and the fact that for any dimension vector there is at most one exceptional module up to isomorphism, we have

$$
X_{S_1}^{v_1} X_{S_2}^{v_2} \cdots X_{S_n}^{v_n}
$$

= $(X_{S_1}^{v_{11}} X_{S_2}^{v_{12}} \cdots X_{S_n}^{v_{1n}}) \cdots (X_{S_1}^{v_{r_i 1}} X_{S_2}^{v_{r_i 2}} \cdots X_{S_n}^{v_{r_i n}})$
= $(a_1 X_{E_{i,1}} + \sum_{\substack{\dim L' \prec \dim E_{i,1}}} a_{L'} X_{L'}) \cdots (a_{r_i} X_{E_{i,r_i}} + \sum_{\substack{\dim L'' \prec \dim E_{i,r_i}}} a_{L''} X_{L''})$
= $a_1 \cdots a_{r_i} X_{\delta_{i,1}} + \sum_{\substack{\dim M \prec \delta}} a_M X_M$.

Similarly

$$
X_{S_1}^{v_1} X_{S_2}^{v_2} \cdots X_{S_n}^{v_n}
$$

= $(b_1 X_{E_{j,1}} + \sum_{\substack{\dim T' \prec \dim E_{j,1}}} b_{T'} X_{T'}) \cdots (b_{r_j} X_{E_{j,r_j}} + \sum_{\substack{\dim T'' \prec \dim E_{j,r_j}}} b_{T''} X_{T''})$
= $b_1 \cdots b_{r_j} X_{\delta_{j,1}} + \sum_{\substack{\dim N \prec \delta}} b_N X_N.$

Thus we have

$$
a_1\cdots a_nX_{\delta_{i,1}}=b_1\cdots b_nX_{\delta_{j,1}}+\sum_{\underline{\dim}N\prec\delta}b_NX_N-\sum_{\underline{\dim}M\prec\delta}a_MX_M.
$$

Therefore by Proposition [2.5](#page-3-1) and the denominator theorem in Theorem [2.3,](#page-3-0) we have:

$$
X_{\delta_{i,1}} = X_{\delta_{j,1}} + \sum_{\underline{\dim} N' \prec \delta} b_{N'} X_{N'}.
$$

Now, suppose the proposition holds for $k \leq n$, then on the one hand

$$
X_{n\delta_{i,1}} X_{\delta_{i,1}} = X_{(n+1)\delta_{i,1}} + \sum_{\underline{\dim} L' \prec (n+1)\delta} b_{L'} X_{L'}.
$$

On the other hand by induction

$$
X_{n\delta_{i,1}} X_{\delta_{i,1}}
$$
\n
$$
= (X_{n\delta_{j,1}} + \sum_{\substack{\dim L \prec n\delta}} a_L X_L) X_{\delta_{i,1}}
$$
\n
$$
= (X_{n\delta_{j,1}} + \sum_{\substack{\dim L \prec n\delta}} a_L X_L) (X_{\delta_{j,1}} + \sum_{\substack{\dim N' \prec \delta}} b_{N'} X_{N'})
$$
\n
$$
= X_{(n+1)\delta_{j,1}} + \sum_{\substack{\dim L'' \prec (n+1)\delta}} b_{L''} X_{L''}.
$$

Therefore

$$
X_{(n+1)\delta_{i,1}} = X_{(n+1)\delta_{j,1}} + \sum_{\underline{\dim}L'' \prec (n+1)\delta} b_{L''} X_{L''} - \sum_{\underline{\dim}L' \prec (n+1)\delta} b_{L'} X_{L'}.
$$

Thus the proof is finished.

Proposition 3.6. $X_{n\delta} = X_{n\delta_{i,1}} + \sum_{\text{dim}L \prec n\delta} a_L X_L$, where $a_L \in \mathbb{Q}$.

Proof. Suppose Q be of type \tilde{D} or \tilde{E} and $\delta = (v_1, v_2, \dots, v_n)$, using the same method in Proposition [3.5,](#page-7-0) we have

$$
X_{S_1}^{v_1} X_{S_2}^{v_2} \cdots X_{S_n}^{v_n}
$$

= $(a_1 X_{E_{i,1}} + \sum_{\substack{\dim L' \prec \dim E_{i,1}}} a_{L'} X_{L'}) \cdots (a_{r_i} X_{E_{i,r_i}} + \sum_{\substack{\dim L'' \prec \dim E_{i,r_i}}} a_{L''} X_{L''})$
= $a_1 \cdots a_{r_i} X_{\delta_{i,1}} + \sum_{\substack{\dim M \prec \delta}} a_M X_M.$

We note that there exists a submodule L_1 satisfying the following short exact sequence:

$$
0 \longrightarrow L_1 \longrightarrow \delta \longrightarrow L_2 \longrightarrow 0
$$

where L_1, L_2 are preprojective, preinjective modules respectively. Thus by Proposition [3.3](#page-6-0) and Proposition [3.5,](#page-7-0) we have

$$
X_{S_1}^{v_1} X_{S_2}^{v_2} \cdots X_{S_n}^{v_n} = (b_1 X_{L_1} + \sum_{\substack{\dim L' \prec \dim L_1}} b_{L'} X_{L'}) (b_2 X_{L_2} + \sum_{\substack{\dim L'' \prec \dim L_2}} b_{L''} X_{L''})
$$

= $b_1 b_2 X_{\delta} + \sum_{k=1}^m b_{k,1} X_{\delta_{k,1}} + \sum_{\substack{\dim N \prec \delta}} b_N X_N$
= $b_1 b_2 X_{\delta} + b X_{\delta_{i,1}} + \sum_{\substack{\dim N' \prec \delta}} b_{N'} X_{N'}.$

Thus

$$
a_1\cdots a_nX_{\delta_{i,1}}+\sum_{\underline{\dim}M\prec\delta}a_{M}X_{M}=b_1b_2X_{\delta}+bX_{\delta_{i,1}}+\sum_{\underline{\dim}N'\prec\delta}b_{N'}X_{N'}.
$$

Therefore by Proposition [2.5](#page-3-1) and the denominator theorem in [\[12\]](#page-20-4), we have

$$
X_{\delta} = X_{\delta_{i,1}} + \sum_{\underline{\dim} M' \prec \delta} a_{M'} X_{M'}.
$$

Then we can finish the proof by induction as in the proof of Proposition [3.5.](#page-7-0)

Now we assume Q is of the form \tilde{A} . We give an alternative proof of the difference property in $[10]$. Let Q be a quiver as follows $[7]$:

Let $\lambda \in \mathbb{C}^*$ and $M(\lambda)$ be the regular simple module as follows

Its proper submodules $M_0(\lambda)$ are of the forms as follows:

Let $M(0)$ be the regular module as follows

Its proper submodules $M_0(0)$ are of the following two forms:

and

The proper submodules with the second form lie in the non-homogeneous tube indexed by 0 with quasi socle S_{d_q} . Hence, there is a bijection between the submodules of $M(\lambda)$ and the submodules of $M(0)$ of the first form. It is easy to conclude the difference property in [\[10\]](#page-20-6):

$$
X_{M(0)} = X_{M(\lambda)} + X_{q,radM(0)/S_{d_q}}.
$$

Proposition 3.7. If $\underline{\dim}(T_1 \oplus R_1) = \underline{\dim}(T_2 \oplus R_2)$ where R_i are 0 or any regular exceptional modules, T_i are 0 or any indecomposable regular modules with selfextension in non-homogeneous tubes and there are no extension between R_i and T_i , then

$$
X_{T_1 \oplus R_1} = X_{T_2 \oplus R_2} + \sum_{\underline{\dim} R \prec \underline{\dim}(T_2 \oplus R_2)} a_R X_R
$$

where $a_R \in \mathbb{Q}$.

Proof. Suppose $\underline{\dim}(T_1 \oplus R_1) = (d_1, d_2, \cdots, d_n)$, using the same method in Proposition [3.5,](#page-7-0) we have

$$
X_{S_1}^{d_1} X_{S_2}^{d_2} \cdots X_{S_n}^{d_n} = (a_1 X_{E_{i,1}} + \sum_{\substack{\dim L' \prec \dim E_{i,1}}} a_{L'} X_{L'}) \cdots (a_s X_{E_{i,s}} + \sum_{\substack{\dim L'' \prec \dim E_{i,s} \\ \dim L'' \prec \dim R_1}} a_{L''} X_{L''})
$$

$$
\times (a_{R_1} X_{R_1} + \sum_{\substack{\dim L'' \prec \dim R_1 \\ \dim L \prec \dim(T_1 \oplus R_1)}} a_L X_{L'})
$$

$$
X_{S_1}^{d_1} X_{S_2}^{d_2} \cdots X_{S_n}^{d_n} = (b_1 X_{E_{j,1}} + \sum_{\substack{\dim M' \prec \dim E_{j,1} \\ \dim M'' \prec \dim E_{j,1}}} b_{M'} X_{M'}) \cdots (b_t X_{E_{j,t}} + \sum_{\substack{\dim M'' \prec \dim E_{i,t} \\ \dim M'' \prec \dim E_{i,t}}} b_{M''} X_{M''})
$$

$$
= b X_{T_2 \oplus R_2} + \sum_{\substack{\dim L \prec \dim(T_2 \oplus R_2) \\ \dim(T_2 \oplus R_2)}} b_M X_{M}.
$$

Thus

$$
aX_{T_1\oplus R_1} + \sum_{\underline{\dim}L \prec \underline{\dim}(T_1\oplus R_1)} a_L X_L = bX_{T_2\oplus R_2} + \sum_{\underline{\dim}L \prec \underline{\dim}(T_2\oplus R_2)} b_M X_M.
$$

Therefore by Proposition [2.5](#page-3-1) and the denominator theorem in Theorem [2.3,](#page-3-0) we have

$$
X_{T_1 \oplus R_1} = X_{T_2 \oplus R_2} + \sum_{\underline{\dim} R \prec \underline{\dim}(T_2 \oplus R_2)} a_R X_R.
$$

4. A Z-basis for the cluster algebra of the alternating quiver of $A_{n,n}$, D or E

Recall that for an acyclic quiver, the matrix B associated to Q is the antisymmetric matrix given by

$$
b_{ij} = |i \longrightarrow j \in Q_1| - |j \longrightarrow i \in Q_1|
$$

where $1 \leq i, j \leq n$.

Definition 4.1. [\[4\]](#page-19-0)[\[9\]](#page-19-8) Let Q be an acyclic quiver with associated matrix B . Q is called graded if there exists a linear form ϵ on \mathbb{Z}^n such that $\epsilon(B\alpha_i) < 0$ for any $1 \leq i \leq n$ where α_i denotes the *i*-th vector of the canonical basis of \mathbb{Z}^n .

Theorem 4.2. [\[4\]](#page-19-0) Let Q be a graded quiver and $\{M_1, \dots, M_r\}$ a family objects in $\mathcal{C}(Q)$ such that $\underline{\dim}(M_i) \neq \underline{\dim}(M_j)$ for $i \neq j$, then X_{M_1}, \cdots, X_{M_r} are linearly independent over Q.

In the section, we still suppose that Q is one of $\widetilde{A}_{n,n}$, \widetilde{D} and \widetilde{E} with an orientation where every vertex is a sink or a source. Note that the quiver Q we consider is graded. We will prove the following theorem.

Theorem 4.3. A Z-basis for $\mathcal{AH}(Q)$ is the following set denoted by $\mathcal{S}(Q)$:

 ${X_L, X_{T \oplus R} | \underline{\dim}(T_1 \oplus R_1) \neq \underline{\dim}(T_2 \oplus R_2), Ext^1_{\mathcal{C}(Q)}(T, R) = 0, Ext^1_{\mathcal{C}(Q)}(L, L) = 0}$

where L is any non-regular exceptional object, R is 0 or any regular exceptional module and T is 0 or any indecomposable regular module with self-extension.

Proof. Note that for any objects $M, N \in \mathcal{C}(Q)$, the final results of their multiplication $X_M X_N$ must be Q−combinatorics of

$$
\{X_L, X_{T \oplus R} | Ext^1_{\mathcal{C}(Q)}(T, R) = 0, Ext^1_{\mathcal{C}(Q)}(L, L) = 0\}
$$

where L is any non-regular exceptional object, R is 0 or any regular exceptional module and T is 0 or any indecomposable regular module with self-extension.

Hence from those Propositions in Section 3, we can easily find that $X_M X_N$ is a \mathbb{Q} −combinatorics of $\mathcal{S}(Q)$. Then the proof is finished by the following Proposition [4.7](#page-11-0) and Proposition [4.8.](#page-12-0)

Remark 4.4. By Proposition [4.9](#page-12-1) in the following, we can also rewrite the \mathbb{Z} -basis as $\{X_{M(d_1,\dots,d_n)}:(d_1,\dots,d_n)\in\mathbb{Z}^n\}$, where $M(d_1,\dots,d_n)=L$ or $T\oplus R$ associated to the element in $\mathcal{S}(Q)$.

Corollary 4.5. $S(Q)$ is a Z-basis of the cluster algebra $\mathcal{E}\mathcal{H}(Q)$.

Proof. It is obvious that $X_{E_i} \in \mathcal{EH}(Q)$. Then by Proposition 6.2 in [\[9\]](#page-19-8), one can show that $X_{E_i[n]} \in \mathcal{EH}(Q)$ for any $n \in \mathbb{N}$. Thus by Proposition [3.6,](#page-7-1) we can prove that $X_{m\delta} \in \mathcal{EH}(Q)$. By definition, $X_L \in \mathcal{EH}(Q)$ for L satisfying $\text{Ext}^1_{\mathcal{C}(Q)}(L,L) = 0$. Thus $\mathcal{S}(Q) \subset \mathcal{E}\mathcal{H}(Q)$. It follows that $\mathcal{S}(Q)$ is a Z-basis of the cluster algebra $\mathcal{E}\mathcal{H}(Q)$ by Theorem [4.3.](#page-10-0)

According to Theorem [4.3](#page-10-0) and Corollary [4.5,](#page-11-1) we have

Corollary 4.6. $\mathcal{E}\mathcal{H}(Q) = \mathcal{AH}(Q)$.

Firstly, by Theorem [4.2](#page-10-1) we need to prove the dimension vectors of these objects associated to the corresponding elements in $\mathcal{S}(Q)$ are different.

Proposition 4.7. Let M be a regular module associated to some element in $S(Q)$ and L be a non-regular exceptional object in $\mathcal{C}(Q)$. Then $\dim(M) \neq \dim(L)$.

Proof. If L contains some τP_i as its direct summand, we know that

$$
\underline{\dim}(\tau P_i) = (0, \cdots, 0, -1, 0, \cdots, 0)
$$

where the *i*-th component is -1 . Suppose $L = \tau P_i \oplus \tau P_{i_1} \cdots \tau P_{i_r} \oplus N$ where N is an exceptional module. Because L is an exceptional object, $X_{\tau P_i} X_N = X_{\tau P_i \oplus N}$ i.e. $\dim_{\mathbb{C}} \text{Hom}(P_i, N) = 0.$ Thus we have $\dim_{\mathbb{C}} N(i) = 0$ and $\dim_{\mathbb{C}} (\tau P_i \oplus \tau P_{i_1} \cdots \tau P_{i_r} \oplus$ $N(i) \leq -1$. However, $\underline{\dim}(M) \geq 0$. Therefore, $\underline{\dim}(M) \neq \underline{\dim}(L)$.

If L is a module. Suppose $\underline{\dim}(M) = \underline{\dim}(L)$. Because L is an exceptional module, we know that M belongs to the orbit of L and then M is a degeneration of L. Hence, there exists some $\mathbb{C}Q$ -module U such that

$$
0 \longrightarrow M \longrightarrow L \oplus U \longrightarrow U \longrightarrow 0
$$

is an exact sequence. Choose minimal U so that we cannot separate the following exact sequence

$$
0 \longrightarrow 0 \longrightarrow U_1 \longrightarrow U_1 \longrightarrow 0
$$

from the above short exact sequence. Thus M has a non-zero map to every direct summand of L. Therefore L has no preprojective modules as direct summand because M is a regular module.

Dually there exists a $\mathbb{C}Q$ -module V such that

$$
0 \longrightarrow V \longrightarrow V \oplus L \longrightarrow M \longrightarrow 0
$$

is an exact sequence. We can choose minimal V so that one cannot separate the following exact sequence

$$
0 \longrightarrow V_1 \longrightarrow V_1 \longrightarrow 0 \longrightarrow 0
$$

from the above short exact sequence. Thus every direct summand of L has a non-zero map to M . Therefore L has no preinjective modules as direct summand because M is a regular module.

Therefore L is a regular exceptional module, it is a contradiction.

$$
\Box
$$

Secondly, we need to prove that $\mathcal{S}(Q)$ is a Z−basis.

Proposition 4.8. $X_M X_N$ belongs to $\mathbb{Z}S(Q)$ for any $M, N \in \mathcal{C}(Q)$.

Proof. According to these above discussions, for any objects $M, N \in \mathcal{C}(Q)$, we have

$$
X_M X_N = b_L X_L + \sum_{\underline{\dim}L' \prec \underline{\dim}(M \oplus N)} b_{L'} X_{L'}
$$

where $\underline{\dim}(L) = \underline{\dim}(M \oplus N), X_L, X_{L'} \in \mathcal{S}(Q)$ and $b_L, b_{L'} \in \mathbb{Q}$.

Therefore by Proposition [2.5](#page-3-1) and the denominator theorem in Theorem [2.3,](#page-3-0) we have $b_L = 1$. Note that there exists a partial order on these dimension vectors by Definition [2.1.](#page-2-3) Thus in these remained L' , we choose these maximal elements denoted by L'_1, \dots, L'_s . Then by $b_L = 1$ and the coefficients of Laurent expansions in generalized cluster variables are integers, we obtain that $a_{L'_1}, \cdots, a_{L'_s}$ are integers. Using the same method, we have $b_{L'} \in \mathbb{Z}$.

We denote $X_{S_i}^{-d} = X_{\tau P_i}^d$ for $d \in \mathbb{N}$. Then we have the following result.

Proposition 4.9.

$$
X_{S_1}^{d_1} X_{S_2}^{d_2} \cdots X_{S_n}^{d_n} = X_{M(d_1, \cdots, d_n)} + \sum_{\underline{dim} L \prec (d_1, \cdots, d_n)} b_L X_L
$$

where $X_{M(d_1,\dots,d_n)}, X_L \in \mathcal{S}(Q)$, $\underline{\dim}M = (d_1,\dots,d_n) \in \mathbb{Z}^n$ and $b_L \in \mathbb{Z}$.

Proof. By Proposition [2.5,](#page-3-1) Theorem [2.3](#page-3-0) and Theorem [4.3.](#page-10-0)

Note that $\{X_{M(d_1,\dots,d_n)}:(d_1,\dots,d_n)\in\mathbb{Z}^n\}$ is a Z-basis of $\mathcal{AH}(Q)$, then we have the following Corollary [4.10](#page-12-2) by Proposition [4.9.](#page-12-1)

Corollary 4.10. $\{X_{S_1}^{d_1}X_{S_2}^{d_2}\cdots X_{S_n}^{d_n} : (d_1,\cdots,d_n) \in \mathbb{Z}^n\}$ is a \mathbb{Z} -basis of $\mathcal{AH}(Q)$, further, there is an isomorphism of Z−algebras:

$$
\mathcal{AH}(Q) \simeq \mathbb{Z}[X_{S_1}, \cdots, X_{S_n}, X_{\tau P_1}, \cdots, X_{\tau P_n}].
$$

Remark 4.11. In [\[9\]](#page-19-8), G.Dupont has proved the following isomorphism of $\mathbb{Z}-algebras:$

$$
\mathcal{A}(\overrightarrow{A_n}) \simeq \mathbb{Z}[X_{S_1}, \cdots, X_{S_n}, X_{\tau P_1}, \cdots, X_{\tau P_n}]
$$

where $n > 0$.

5. A Z-BASIS FOR THE CLUSTER ALGEBRA OF TYPE $\widetilde{A}_{n,n}, \widetilde{D}$ or \widetilde{E}

In this section, we will deduce a Z-basis for the cluster algebra of $\widetilde{A}_{n,n}$, \widetilde{D} or \widetilde{E} which is independent of their orientations.

Define the reflected quiver $\sigma_i(Q)$ by reversing all the arrows ending at i. The mutations can be viewed as generalizations of reflections i.e. if i is a sink or a source in Q_0 , then $\mu_i(Q) = \sigma_i(Q)$ where μ_i denotes the mutation in the direction i. Thus there is a natural isomorphism of cluster algebras

$$
\Phi : \mathcal{A}(Q) \longrightarrow \mathcal{A}(Q')
$$

where Q' is a quiver mutation equivalent to Q , and Φ is called the canonical cluster algebras isomorphism in [\[10\]](#page-20-6).

Now suppose Q is an acyclic quiver and i is a sink in Q_0 . Let $Q' = \sigma_i(Q)$ and R_i^+ : $\mathcal{C}(Q) \longrightarrow \mathcal{C}(Q')$ be the extended BGP-reflection functor defined in [\[17\]](#page-20-7).

Denote by $X_?^Q$ (resp. by $X_?^{\sigma_i Q}$) the Caldero-Chapton map associated to Q (resp. to $\sigma_i Q$).

Then the following hold.

Lemma 5.1. [\[17\]](#page-20-7) Let Q be an acyclic quiver and i be a sink in Q. Then R_i^+ induces a triangle equivalence

$$
R_i^+:\mathcal{C}(Q)\longrightarrow \mathcal{C}(\sigma_i Q)
$$

Lemma 5.2. [\[10\]](#page-20-6) Let Q be an affine quiver and i be a sink in Q and M be an indecomposable regular CQ-module with quasi-composition series

$$
0 = M_0 \subset M_1 \subset \cdots \subset M_s = M
$$

then R_i^+M is a regular $\mathbb{C}\sigma_iQ$ -module with quasi-composition series

$$
0 = R_i^+ M_0 \subset R_i^+ M_1 \subset \cdots \subset R_i^+ M_s = R_i^+ M
$$

In particular, R_i^+ send quasi-socles to quasi-socles, quasi-radicals to quasi-radicals and preserves quasi-lengths.

Lemma 5.3. [\[10\]](#page-20-6) Let Q be an affine quiver and i be a sink in Q . Denote by $\Phi_i\,:\,\mathcal{A}(Q)\,\longrightarrow\, \mathcal{A}(\sigma_iQ)\,$ the canonical cluster algebra isomorphism and by R_i^+ : $\mathcal{C}(Q) \longrightarrow \mathcal{C}(\sigma_i Q)$ the extended BGP-reflection functor. Then

$$
\Phi_i(X_M^Q) = X_{R_i^+M}^{\sigma_i Q}
$$

where M is any rigid object in $\mathcal{C}(Q)$ or any regular module in non-homogeneous tubes of rank $r > 1$.

In the following, we will suppose that Q is one of $\widetilde{A}_{n,n}$, \widetilde{D} and \widetilde{E} with an orientation where every vertex is a sink or a source, and Q' is an acyclic quiver of $\widetilde{A}_{n,n}, \widetilde{D}$ or \widetilde{E} . Then there exists an admissible sequence of sinks (i_1, \dots, i_s) such that $Q' = \sigma_{i_s} \cdots \sigma_{i_1}(Q)$. Denote by $\Phi : \mathcal{A}(Q) \longrightarrow \mathcal{A}(Q')$ the canonical cluster algebra isomorphism and by $R^+ = R^+_{i_s} \cdots R^+_{i_1} : C(Q) \longrightarrow C(Q')$. Note that R^+ is an equivalence of triangulated categories by Lemma [5.1.](#page-13-0) Then we have the following main theorem.

Theorem 5.4. A \mathbb{Z} -basis for the cluster algebra of Q' is the following set denoted by $\mathcal{B}(Q')$:

 ${X_{L', X_{T'\oplus R'}\mid dim(T'_1\oplus R'_1)\neq \underline{dim}(T'_2\oplus R'_2), Ext^1_{C(Q')}(T', R')=0, Ext^1_{C(Q')}(L', L')=0}}$

where L' is any non-regular exceptional object, R' is 0 or any regular exceptional module, T' is 0 or any indecomposable regular module with self-extension and in addition if $Q' \neq A_{1,1}$ and $\underline{\dim}(T' \oplus R') = m\delta$ for some $m \in \mathbb{N}$, then $R' = 0$ and T' is an indecomposable module of dimension vector mδ in a non-homogeneous tube.

Proof. If $Q' = \tilde{A}_{1,1}$, it is obvious that

$$
\{X_M, X_{n\delta} : M \in obj\mathcal{C}(Q), Ext^1(M, M) = 0\}
$$

is a Z-basis for cluster algebra of $A_{1,1}$, which is called the semicanonical basis in [\[6\]](#page-19-4).

If $Q' \neq \widetilde{A}_{1,1}$, then by Theorem [4.3,](#page-10-0) we have obtained a Z-basis for cluster algebra of Q:

 ${X_L, X_{T \oplus R} | \underline{\dim}(T_1 \oplus R_1) \neq \underline{\dim}(T_2 \oplus R_2), Ext^1_{\mathcal{C}(Q)}(T, R) = 0, Ext^1_{\mathcal{C}(Q)}(L, L) = 0}$ where L is any non-regular exceptional object, R is 0 or any regular exceptional module and T is 0 or any indecomposable regular module with self-extension.

If $\underline{\dim}(T \oplus R) = m\delta$ for some $m \in \mathbb{N}$, then we take $R = 0$ and T an indecomposable module of dimension vector $m\delta$ in a non-homogeneous tube. We denote this Z-basis by $\mathcal{B}(Q)$. Thus $\Phi(\mathcal{B}(Q))$ is a Z-basis for the cluster algebra of Q' because $\Phi : \mathcal{A}(Q) \longrightarrow \mathcal{A}(Q')$ is the canonical cluster algebras isomorphism. Then by Lemma [5.1,](#page-13-0) Lemma [5.2](#page-13-1) and Lemma [5.3,](#page-13-2) we know that $\Phi(\mathcal{B}(Q))$ is exactly the basis $\mathcal{B}(Q')$ in Theorem [5.4.](#page-13-3)

6. some special cases

In this section, we will give some examples for special cases to explain the Z-basis explicitly.

6.1. Z-basis for finite type. For finite type, we know that there are no regular parts in $\mathcal{B}(Q)$. Thus the Z-bases are exactly

$$
\{X_M : M \in obj\mathcal{C}(Q), Ext^1(M, M) = 0\}
$$

which coincides with the canonical basis for a cluster algebra of finite type in [\[4\]](#page-19-0).

6.2. Z-basis for the Kronecker quiver. Consider the Kronecker quiver, we note that $R = 0$ and $T = n\delta$ for $n \ge 1$. Thus a Z-basis is

$$
\{X_M, X_{n\delta} : M \in obj\mathcal{C}(Q), Ext^1(M, M) = 0\}
$$

which is called the semicanonical basis in [\[6\]](#page-19-4). If we modify:

$$
z_1 := X_\delta, \ z_n := X_{n\delta} - X_{(n-2)\delta}.
$$

Then $\{X_M, z_n : M \in obj\mathcal{C}(Q), Ext^1(M, M) = 0\}$ is the canonical basis for cluster algebra of Kronecker quiver in [\[14\]](#page-20-0).

6.3. **Z-basis for** \widetilde{D}_4 . Let Q be the tame quiver of type \widetilde{D}_4 as follows

We denote the minimal imaginary root by $\delta = (2, 1, 1, 1, 1)$. The regular simple modules of dimension vector δ are

with linear maps

$$
\alpha_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \alpha_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \alpha_3 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \alpha_4 = \begin{pmatrix} \lambda \\ \mu \end{pmatrix}
$$

where $\lambda/\mu \in \mathbb{P}^1, \lambda/\mu \neq 0, 1, \infty$. Let M be any regular simple $\mathbb{C}Q$ -module of dimension vector δ and $M[i]$ be the regular module with top M and regular length i for any $i \in \mathbb{N}$. Let X_M be the generalized cluster variable associated to M by the reformulation of the Caldero-Chapton map. Then we have

Proposition 6.1. [8]
$$
X_M = \frac{1}{x_1^2 x_2 x_3 x_4 x_5} + \frac{4}{x_1 x_2 x_3 x_4 x_5} + \frac{x_1^2 + 4x_1 + 6}{x_2 x_3 x_4 x_5} + \frac{x_2 x_3 x_4 x_5 + 2}{x_1^2} + \frac{4}{x_1}.
$$

We define $X_{n\delta} := X_{M[n]}$ for $n \in \mathbb{N}$. Now, we consider three non-homogeneous tubes labelled by the subset $\{0,1,\infty\}$ of \mathbb{P}^1 . Let $X_{\delta_1}, X_{\delta_2}, X_{\delta_3}, X_{\delta_4}, X_{\delta_5}, X_{\delta_6}$ be generalized cluster variables associated to regular modules in non-homogeneous tubes of dimension vector δ . The regular simple modules in non-homogeneous tubes are denoted by $E_1, E_2, E_3, E_4, E_5, E_6$, where

$$
\underline{\dim}(E_1) = (1, 1, 1, 0, 0), \ \underline{\dim}(E_2) = (1, 0, 0, 1, 1), \ \underline{\dim}(E_3) = (1, 1, 0, 1, 0),
$$

 $\underline{\dim}(E_4) = (1, 0, 1, 0, 1), \ \underline{\dim}(E_5) = (1, 0, 1, 1, 0), \ \underline{\dim}(E_6) = (1, 1, 0, 0, 1).$

We note that ${E_1, E_2}, {E_3, E_4}$ and ${E_5, E_6}$ are pairs of the regular simple modules in the bottom of non-homogeneous tubes labelled by $1, \infty$ and 0, respectively.

Let $E_i[n]$ be the indecomposable regular module with top E_i and regular length *n* for $1 \leq i \leq 6$. We set $X_{n\delta_i} := X_{E_i[2n]}$ for $1 \leq i \leq 6$.

Proposition 6.2. [\[8\]](#page-19-5) For any $n \in \mathbb{N}$, we have $X_{n\delta_1} = X_{n\delta_2} = X_{n\delta_3} = X_{n\delta_4} = X_{n\delta_5}$ $X_{n\delta_5} = X_{n\delta_6}$ and $X_{n\delta_1} = X_{n\delta} + X_{(n-1)\delta}$ where $X_0 = 1$.

Theorem 6.3. [\[8\]](#page-19-5) A Z-basis for cluster algebra of \widetilde{D}_4 is the following set denoted by $\mathcal{B}(Q)$:

$$
\{X_L, X_{m\delta}, X_{E_i[2k+1]\oplus R}|\underline{\dim}(E_i[2k+1]\oplus R_1) \neq \underline{\dim}(E_j[2l+1]\oplus R_2),
$$

$$
\text{Ext}^1_{\mathcal{C}(Q)}(L, L) = 0, \text{Ext}^1_{\mathcal{C}(Q)}(E_i[2k+1], R) = 0, m, k, l \ge 0, 1 \le i, j \le 6
$$

where L is any non-regular exceptional object, R is 0 or any regular exceptional module.

7. the inductive multiplication formula for a tube

Now we fix a tube with rank r and these regular simple modules are E_1, \dots, E_r with $\tau E_{i+1} = E_i$ where $E_i = E_{i+mr}$ for $1 \leq i \leq r$ and $m \in \mathbb{N}$. Let X_{E_i} be the corresponding generalized cluster variable for $i = 1, \dots, r$. With these notations, we have the following inductive cluster multiplication formula.

Theorem 7.1. Let i, j, k, l, m and r be in Z such that $1 \leq k \leq mr + l, 0 \leq l \leq$ $r-1, 1 \leq i, j \leq r, m \geq 0.$

$$
(1)
$$
 When $j \leq i$, then

1)for $k + i \geq r + j$, we have $X_{E_i[k]}X_{E_j[mr+l]} = X_{E_i[(m+1)r+l+j-i]}X_{E_j[k+i-r-j]} +$ $X_{E_i[r+j-i-1]}X_{E_{k+i+1}[(m+1)r+l+j-k-i-1]},$ 2)for $k + i < r + j$ and $i \leq l + j \leq k + i - 1$, we have $X_{E_i[k]}X_{E_j[mr+l]} =$ $X_{E_j[mr+k+i-j]}X_{E_i[l+j-i]} + X_{E_j[mr+i-j-1]}X_{E_{l+j+1}[k+i-l-j-1]},$

3)for other conditions, we have $X_{E_i[k]}X_{E_j[mr+l]} = X_{E_i[k] \oplus E_j[mr+l]}$. (2) When $j > i$, then

1)for $k \geq j - i$, we have $X_{E_i[k]}X_{E_i[mr+l]} = X_{E_i[i-i-1]}X_{E_{k+i+1}[mr+l+j-k-i-1]} +$ $X_{E_i[mr+l+j-i]}X_{E_j[k+i-j]},$

2) for
$$
k < j - i
$$
 and $i \leq l + j - r \leq k + i - 1$, we have $X_{E_i[k]}X_{E_j[mr+l]} = X_{E_j[(m+1)r+k+i-j]}X_{E_i[l+j-r-i]} + X_{E_j[(m+1)r+i-j-1]}X_{E_{l+j+1}[k+r+i-l-j-1]},$
3) for other conditions, we have $X_{E_i[k]}X_{E_j[mr+l]} = X_{E_i[k] \oplus E_j[mr+l]}.$

Proof. We only prove (1) and (2) is totally similar to (1).

1) When $k = 1$, by $k + i \ge r + j$ and $1 \le j \le i \le r \Longrightarrow i = r$ and $j = 1$.

Then by the cluster multiplication theorem in Theorem [1.2](#page-2-0) or Theorem [1.4,](#page-2-2) we have

 $X_{E_r} X_{E_1[mr+l]} = X_{E_r[mr+l+1]} + X_{E_2[mr+l-1]}.$

When $k = 2$, by $k + i \ge r + j$ and $1 \le j \le i \le r \Longrightarrow i = r$ or $i = r - 1$. For $i = r \Longrightarrow j = 1$ or $j = 2$:

The case for $i = r$ and $j = 1$, we have

$$
X_{E_r[2]}X_{E_1[mr+l]} = (X_{E_r}X_{E_1} - 1)X_{E_1[mr+l]}
$$

\n
$$
= X_{E_1}(X_{E_r[mr+l+1]} + X_{E_2[mr+l-1]}) - X_{E_1[mr+l]}
$$

\n
$$
= X_{E_1}X_{E_r[mr+l+1]} + (X_{E_1[mr+l]} + X_{E_3[mr+l-2]}) - X_{E_1[mr+l]}
$$

\n
$$
= X_{E_1}X_{E_r[mr+l+1]} + X_{E_3[mr+l-2]}.
$$

The case for $i = r$ and $j = 2$, we have

$$
X_{E_r[2]}X_{E_2[mr+l]} = (X_{E_r}X_{E_1} - 1)X_{E_2[mr+l]}
$$

\n
$$
= X_{E_r}(X_{E_1[mr+l+1]} + X_{E_3[mr+l-1]}) - X_{E_2[mr+l]}
$$

\n
$$
= X_{E_r[mr+l+2]} + (X_{E_2[mr+l]} + X_{E_r}X_{E_3[mr+l-1]}) - X_{E_2[mr+l]}
$$

\n
$$
= X_{E_r[mr+l+2]} + X_{E_r}X_{E_3[mr+l-1]}.
$$

For $i = r - 1 \Longrightarrow j = 1$:

$$
X_{E_{r-1}[2]}X_{E_1[mr+l]} = (X_{E_{r-1}}X_{E_r} - 1)X_{E_1[mr+l]}
$$

\n
$$
= X_{E_{r-1}}(X_{E_r[mr+l+1]} + X_{E_2[mr+l-1]}) - X_{E_1[mr+l]}
$$

\n
$$
= (X_{E_{r-1}[mr+l+2]} + X_{E_1[mr+l]}) + X_{E_{r-1}}X_{E_2[mr+l-1]} - X_{E_1[mr+l]}
$$

\n
$$
= X_{E_{r-1}[mr+l+2]} + X_{E_{r-1}}X_{E_2[mr+l-1]}.
$$

Now, suppose it holds for $k \leq n$, then by induction we have

$$
X_{E_i[n+1]}X_{E_j[mr+l]}
$$

= $(X_{E_i[n]}X_{E_{i+n}} - X_{E_i[n-1]})X_{E_j[mr+l]}$
= $X_{E_{i+n}}(X_{E_i[n]}X_{E_j[mr+l]}) - X_{E_i[n-1]}X_{E_j[mr+l]}$
= $X_{E_{i+n}}(X_{E_i[(m+1)r+l+j-i]}X_{E_j[n+i-r-j]} + X_{E_i[r+j-i-1]}X_{E_{n+i+1}[(m+1)r+l+j-n-i-1]})$
 $-(X_{E_i[(m+1)r+l+j-i]}X_{E_j[n+i-r-j-1]} + X_{E_i[r+j-i-1]}X_{E_{n+i}[(m+1)r+l+j-n-i]})$
= $X_{E_i[(m+1)r+l+j-i]}(X_{E_j[n+i+1-r-j]} + X_{E_j[n+i-r-j-1]})$
+ $X_{E_i[r+j-i-1]}(X_{E_{n+i}[(m+1)r+l+j-n-i]} + X_{E_{n+i+2}[(m+1)r+l+j-n-i-2]})$
 $-(X_{E_i[(m+1)r+l+j-i]}X_{E_j[n+i-r-j-1]} + X_{E_i[r+j-i-1]}X_{E_{n+i}[(m+1)r+l+j-n-i-2]})$
= $X_{E_i[(m+1)r+l+j-i]}X_{E_j[n+i+1-r-j]} + X_{E_i[r+j-i-1]}X_{E_{n+i+2}[(m+1)r+l+j-n-i-2]}.$
2) When $k = 1$, by $i \le l + j \le k + i - 1 \Longrightarrow i \le l + j \le i \Longrightarrow i = l + j$.

Then by the cluster multiplication theorem in Theorem [1.2](#page-2-0) or Theorem [1.4,](#page-2-2) we have

 $X_{E_{i}}X_{E_{j}[mr+l]} = X_{E_{l+j}}X_{E_{j}[mr+l]} = X_{E_{j}[mr+l+1]} + X_{E_{j}[mr+l-1]}$ When $k = 2$, by $i \leq l+j \leq k+i-1 \Longrightarrow i \leq l+j \leq i+1 \Longrightarrow i = l+j$ or $i+1 = l+j$: For $i = l + j$, we have

$$
X_{E_i[2]}X_{E_j[mr+l]} = X_{E_{l+j}[2]}X_{E_j[mr+l]} = (X_{E_{l+j}}X_{E_{l+j+1}} - 1)X_{E_j[mr+l]}
$$

= $(X_{E_j[mr+l+1]} + X_{E_j[mr+l-1]})X_{E_{l+j+1}} - X_{E_j[mr+l]}$
= $X_{E_j[mr+l+2]} + X_{E_j[mr+l]} + X_{E_{l+j+1}}X_{E_j[mr+l-1]} - X_{E_j[mr+l]}$
= $X_{E_j[mr+j+2]} + X_{E_{l+j+1}}X_{E_j[mr+l-1]}.$

For $i + 1 = l + j$, we have

$$
X_{E_i[2]}X_{E_j[mr+l]} = X_{E_{l+j-1}[2]}X_{E_j[mr+l]} = (X_{E_{l+j-1}}X_{E_{l+j}} - 1)X_{E_j[mr+l]}
$$

= $(X_{E_j[mr+l+1]} + X_{E_j[mr+l-1]})X_{E_{l+j-1}} - X_{E_j[mr+l]}$
= $X_{E_j[mr+l+1]}X_{E_{l+j-1}} + (X_{E_j[mr+l]} + X_{E_j[mr+l-2]}) - X_{E_j[mr+l]}$
= $X_{E_j[mr+l+1]}X_{E_{l+j-1}} + X_{E_j[mr+l-2]}.$

Suppose it holds for $k \leq n$, then by induction we have

$$
X_{E_i[n+1]}X_{E_j[mr+l]}
$$

= $(X_{E_i[n]}X_{E_{i+n}} - X_{E_i[n-1]})X_{E_j[mr+l]}$
= $(X_{E_i[n]}X_{E_j[mr+l]})X_{E_{i+n}} - X_{E_i[n-1]}X_{E_j[mr+l]}$
= $(X_{E_j[mr+n+i-j]}X_{E_i[l+j-i]} + X_{E_j[mr+i-j-1]}X_{E_{l+j+1}[n+i-l-j-1]})X_{E_{i+n}}$
 $-(X_{E_j[mr+n+i-j-1]}X_{E_i[l+j-i]} + X_{E_j[mr+i-j-1]}X_{E_{l+j+1}[n+i-l-j-2]})$
= $(X_{E_j[mr+n+i+1-j]} + X_{E_j[mr+n+i-j-1]})X_{E_i[l+j-i]}$
 $+(X_{E_{l+j+1}[n+i-l-j]} + X_{E_{l+j+1}[n+i-l-j-2]})X_{E_j[mr+i-j-1]}$
 $-(X_{E_j[mr+n+i-j-1]}X_{E_i[l+j-i]} + X_{E_j[mr+i-j-1]}X_{E_{l+j+1}[n+i-l-j-2]})$
= $X_{E_j[mr+n+i+1-j]}X_{E_i[l+j-i]} + X_{E_j[mr+i-j-1]}X_{E_{l+j+1}[n+i-l-j]}.$
3) It is trivial by the definition of the Calder-Chapton map.

Note that in the above section, we have already obtained a \mathbb{Z} -basis for the cluster algebra of $\widetilde{A}_{n,n}$, \widetilde{D} or \widetilde{E} . Now by using Theorem [7.1](#page-15-0) and Proposition [3.3](#page-6-0) we can easily express $X_{E_i[m]}X_{E_j[n]}$ as a $\mathbb{Z}-$ combinatorics of the basis for the cluster algebra of $A_{n,n}, D$ or E where $E_i[m]$ and $E_j[n]$ are regular modules in one fixed tube. In the following section, we will explain it by an example.

8. an example

We consider a tube with rank 3. By Theorem [7.1,](#page-15-0) we can easily obtain the following proposition.

Proposition 8.1. (1) For $n > 3m + 1$, then

$$
X_{E_2[3m+1]}X_{E_1[n]} = X_{E_2}X_{E_1[n+3m]} + X_{E_1[n+3m-3]} + X_{E_2}X_{E_1[n+3m-6]} + X_{E_1[n+3m-9]} + \cdots + X_{E_2}X_{E_1[n-3m+6]} + X_{E_1[n-3m+3]} + X_{E_2}X_{E_1[n-3m]},
$$

(2) For $n \ge 3m+2$, then

 $X_{E_2[3m+2]}X_{E_1[n]} = X_{E_2[n+3m+2]} + X_{E_2}X_{E_2[n+3m-1]} + X_{E_2[n+3m-4]} + X_{E_2}X_{E_2[n+3m-7]}$ $+ \cdots + X_{E_2[n-3m+2]} + X_{E_2}X_{E_2[n-3m-1]},$

(3) For $n \geq 3m + 3$, then

$$
X_{E_2[3m+3]}X_{E_1[n]} = X_{E_1[n+3m+3]} + X_{E_3[n+3m+1]} + X_{E_2[n+3m-1]} + X_{E_1[n+3m-3]} + X_{E_3[n+3m-5]} + X_{E_2[n+3m-7]} + \cdots + X_{E_1[n-3m+3]} + X_{E_3[n-3m+1]} + X_{E_2[n-3m-1]} + X_{E_1[n-3m-3]}.
$$

Proof. (1) By Theorem [7.1,](#page-15-0) then

 $X_{E_2[3m+1]}X_{E_1[n]}$ $= X_{E_2[n+2]}X_{E_1[3m-1]} + X_{E_2}X_{E_1[n-3m]}$

 $= X_{E_2[3m-2]}X_{E_1[n+3]} + X_{E_1[n-3m+3]} + X_{E_2}X_{E_1[n-3m]}$

 $= X_{E_2[n+5]}X_{E_1[3m-4]} + X_{E_2}X_{E_1[n-3m+6]} + X_{E_1[n-3m+3]} + X_{E_2}X_{E_1[n-3m]}$

$$
= X_{E_2[3m-5]} X_{E_1[n+6]} + X_{E_1[n-3m+9]} + X_{E_2} X_{E_1[n-3m+6]} + X_{E_1[n-3m+3]} + X_{E_2} X_{E_1[n-3m]}
$$

. .

 $= X_{E_2}X_{E_1[n+3m]} + X_{E_1[n+3m-3]} + X_{E_2}X_{E_1[n+3m-6]} + X_{E_1[n+3m-9]}$ $+ \cdots + X_{E_2} X_{E_1[n-3m+6]} + X_{E_1[n-3m+3]} + X_{E_2} X_{E_1[n-3m]}.$ (2) By Theorem [7.1,](#page-15-0) then $X_{E_2[3m+2]}X_{E_1[n]}$ $= X_{E_2[n+2]}X_{E_1[3m]} + X_{E_2}X_{E_2[n-3m-1]}$

A Z-BASIS FOR THE CLUSTER ALGEBRA ASSOCIATED TO AN AFFINE QUIVER 19

- $= X_{E_2[3m-1]} X_{E_1[n+3]} + X_{E_2[n-3m+2]} + X_{E_2} X_{E_2[n-3m-1]}$
- $= X_{E_2[n+5]}X_{E_1[3m-3]} + X_{E_2}X_{E_2[n-3m+5]} + X_{E_2[n-3m+2]} + X_{E_2}X_{E_2[n-3m-1]}$

. .

- $= X_{E_2[3m-4]}X_{E_1[n+6]}+X_{E_2[n-3m+8]}+X_{E_2}X_{E_2[n-3m+5]}+X_{E_2[n-3m+2]}+X_{E_2}X_{E_2[n-3m-1]}$
- . $= X_{E_2[n+3m+2]} + X_{E_2}X_{E_2[n+3m-1]} + X_{E_2[n+3m-4]} + X_{E_2}X_{E_2[n+3m-7]}$ $+ \cdots + X_{E_2[n-3m+2]} + X_{E_2}X_{E_2[n-3m-1]}.$
- (3) By Theorem [7.1,](#page-15-0) then
	- $X_{E_2[3m+3]}X_{E_1[n]}$
	- $= X_{E_2[n+2]}X_{E_1[3m+1]} + X_{E_2}X_{E_3[n-3m-2]}$
	- $= X_{E_2[n+2]}X_{E_1[3m+1]} + X_{E_2[n-3m-1]} + X_{E_1[n-3m-3]}$
	- $= X_{E_2[3m]} X_{E_1[n+3]} + X_{E_3[n-3m+1]} + X_{E_2[n-3m-1]} + X_{E_1[n-3m-3]}$
	- $= X_{E_1[n+3m+3]} + X_{E_3[n+3m+1]} + X_{E_2[n+3m-1]}$ $+X_{E_1[n+3m-3]}$ + $X_{E_3[n+3m-5]}$ + $X_{E_2[n+3m-7]}$ $+ \cdots + X_{E_1[n-3m+3]} + X_{E_3[n-3m+1]} + X_{E_2[n-3m-1]} + X_{E_1[n-3m-3]}.$

. . .

 \Box

Corollary 8.2. When $n = 3k + 1$, we can rewrite Proposition [8.1](#page-17-0) as following: (1) For $n \geq 3m + 1$, then

 $X_{E_2[3m+1]}X_{E_1[n]} = X_{E_1[n+3m+1]} + X_{E_1[n+3m-1]} + X_{E_1[n+3m-3]} + \cdots$ $+X_{E_1[n-3m+5]} + X_{E_1[n-3m+3]} + X_{E_1[n-3m+1]} + X_{E_1[n-3m-1]},$ (2) For $n \geq 3m + 2$, then $X_{E_2[3m+2]}X_{E_1[n]} = X_{E_2[n+3m+2]} + X_{E_2[n+3m]} + X_{E_2[n+3m-2]} + X_{E_2[n+3m-4]}$

 $+ \cdots + X_{E_2[n-3m+2]} + X_{E_2[n-3m]} + X_{E_2[n-3m-2]},$

(3) For $n > 3m + 3$, then

 $X_{E_2[3m+3]}X_{E_1[n]} = X_{E_1[n+3m+3]} + X_{E_3[n+3m+1]} + X_{E_2[n+3m-1]}$ $+X_{E_1[n+3m-3]} + X_{E_3[n+3m-5]} + X_{E_2[n+3m-7]} + \cdots$ $+X_{E_1[n-3m+3]} + X_{E_3[n-3m+1]} + X_{E_2[n-3m-1]} + X_{E_1[n-3m-3]}$

Proof. When $n = 3k + 1$, we have the following equations:

$$
X_{E_2} X_{E_1[n]} = X_{E_1[n+1]} + X_{E_1[n-1]}
$$

$$
X_{E_2} X_{E_2[n-1]} = X_{E_2[n]} + X_{E_2[n-2]}
$$

Then the proof is immediately finished from Proposition [8.1.](#page-17-0) \Box

In the same way by using Theorem [7.1,](#page-15-0) we have the following proposition.

Proposition 8.3. (1) For $n \geq 3m + 1$, then

$$
X_{E_1[3m+1]}X_{E_1[n]} = X_{E_1}X_{E_1[n+3m]} + X_{E_1[n+3m-3]} + X_{E_1}X_{E_1[n+3m-6]} + X_{E_1[n+3m-9]} + \cdots + X_{E_1}X_{E_1[n-3m+6]} + X_{E_1[n-3m+3]} + X_{E_1}X_{E_1[n-3m]},
$$
\n(2) For $n \ge 3m+2$, then
\n
$$
X_{E_1[3m+2]}X_{E_1[n]} = X_{E_1[2]}(X_{E_1[n+3m]} + X_{E_1[n+3m-6]} + \cdots + X_{E_1[n-3m]}),
$$
\n(3) For $n \ge 3m+3$, then
\n
$$
X_{E_1[3m+3]}X_{E_1[n]} = X_{E_1[n+3m+3]} + X_{E_2}X_{E_1[n+3m]} + X_{E_1[n+3m-3]} + X_{E_2}X_{E_1[n+3m-6]} + \cdots + X_{E_1[n-3m+3]} + X_{E_2}X_{E_1[n-3m]} + X_{E_1[n-3m-3]}.
$$

Remark 8.4. By the same method in Corollary [8.2,](#page-18-0) we can also rewrite Proposition [8.3](#page-18-1) if we consider these different n. Here we omit it.

Remark 8.5. In fact, we can also check Proposition [8.1](#page-17-0) Corollary [8.2](#page-18-0) and Proposition [8.3](#page-18-1) by induction.

If we consider the quiver is one of $\widetilde{A}_{n,n}$, \widetilde{D} and \widetilde{E} . Then from the above discus-sions and Proposition [3.3,](#page-6-0) we can easily express $X_{E_i[m]}X_{E_i[n]}$ as a Z-combinatorics of the basis in this tube with rank 3.

9. A Z-basis for the cluster algebra of affine type

In [\[10\]](#page-20-6) Theorem 4.21, G. Dupont asserts that the following set is a \mathbb{Z} -basis for the cluster algebra of type A :

 ${cluster \, monomials} \cup {X_{M_\lambda^{\oplus n} \oplus E}} : n \geq 1, E \, \, are \, \, rigid \, \,regular \, \,modules).$

We will use Theorem [4.3](#page-10-0) and the above result in $[10]$ to give a \mathbb{Z} -basis for the cluster algebra of affine type.

Proof of Theorem [1.1.](#page-1-0) By Theorem [5.4,](#page-13-3) we only prove that the set $\mathcal{B}(Q')$ is a \mathbb{Z} basis for the cluster algebra of $A_{r,s}$ for $r \neq s$. By Theorem 3.25 in [\[10\]](#page-20-6) and Theorem [7.1,](#page-15-0) we have

$$
X_{M_{\lambda}^{\oplus n} \oplus E} = (X_{\delta_{i,1}} - X_{q,radM_{E_{i,1}}/E_{i,1}})^n X_E
$$

\n
$$
= (X_{\delta_{i,1}}^n + \sum_{\substack{\dim(T' \oplus R') \prec n\delta}} a_{T' \oplus R'} X_{T' \oplus R'}) X_E
$$

\n
$$
= X_{T \oplus R} + \sum_{\substack{\dim(T'' \oplus R'') \prec \dim(T \oplus R)}} a_{T'' \oplus R''} X_{T'' \oplus R''}
$$

where $a_{T'\oplus R'}, a_{T''\oplus R''} \in \mathbb{Z}$, $\underline{\dim}(T \oplus R) = \underline{\dim}(M_{\lambda}^{\oplus n} \oplus E)$ and $X_{T \oplus R}, X_{T'' \oplus R''} \in$ $\mathcal{B}(Q')$. Thus from the above upper triangle equations and these dimension vectors of $M_{\lambda}^{\oplus n} \oplus E$ in [\[10\]](#page-20-6) are different, we have

$$
\{X_{M_{\lambda}^{\oplus n} \oplus E} : n \ge 1, E \text{ are rigid regular modules}\} = \{X_{T \oplus R} | \underline{\dim}(T_1 \oplus R_1) \ne \underline{\dim}(T_2 \oplus R_2)\}\
$$

where R is 0 or any regular exceptional module, T is any indecomposable regular module with self-extension and there are no extension between R and T .

Also it is easy to see that

$$
{cluster\ monomials} = {X_L|Ext^1_{\mathcal{C}(Q)}(L,L) = 0}
$$

Therefore $\mathcal{B}(Q')$ is a Z-basis for the cluster algebra of $\widetilde{A}_{r,s}$.

REFERENCES

- [1] A. Buan, R. Marsh, M. Reineke, I. Reiten and G. Todorov, Tilting theory and cluster combinatorics. Advances in Math. 204 (2006), 572-618.
- [2] W. Crawley-Boevy, *Lectures on representations of quivers*, 1992.
- [3] P. Caldero and F. Chapoton, *Cluster algebras as Hall algebras of quiver representations*, Comm. Math. Helvetici, 81 (2006), 596-616.
- [4] P. Caldero and B. Keller, From triangulated categories to cluster algebras, Invent. math. 172 (2008), no. 1, 169-211.
- [5] P. Caldero and B. Keller, From triangulated categories to cluster algebras II, Annales Scientifiques de l'Ecole Normale Superieure, 39 (4) (2006), 83-100.
- [6] P. Caldero and A. Zelevinsky, Laurent expansions in cluster algebras via quiver representations, Moscow Math. J. 6 (2006), no. 2, 411-429.
- [7] V. Dlab and C. M. Ringel, Indecomposable representations of graphs and algebras, Memories of the AMS, 173 (1976), 1-57.
- [8] M.Ding and F.Xu, A Z-basis for the cluster algebra of type \widetilde{D}_4 , preprint.
- [9] G. Dupont,Cluster multiplication in stable tubes via generalized Chebyshev polynomials, arXiv:math.RT/0801.3964.
- [10] G. Dupont, Generic variables in Acyclic cluster algebras, arXiv:math.RT/0811.2909v1.
- [11] S. Fomin and A. Zelevinsky, Cluster algebras. I. Foundations. J. Amer. Math. Soc. 15 (2002), no. 2, 497-529.
- [12] A. Hubery, Acyclic cluster algebras via Ringel-Hall algebras, preprint.
- [13] R. Marsh, M. Reineke, and A. Zelevinsky generalized associahedra via quiver representations. Trans.AMS, 355(1) (2003), 4171-4186.
- [14] P. Sherman and A. Zelevinsky, Positivity and canonical bases in rank 2 cluster algebras of finite and affine types, Moscow Math. J. 4 (2004), no.4, 947–974.
- [15] J. Xiao and F. Xu, Green's formula with \mathbb{C}^* -action and Caldero-Keller's formula, arXiv:math.RT/0707.1175.
- [16] F. Xu, On the cluster multiplication theorem for acyclic cluster algebras, arXiv:math.RT/ 0711.3255.
- [17] B. Zhu, Equivalence between cluster categories, Journal of algebra, 304:832-850,2006.

Department of Mathematical Sciences, Tsinghua University, Beijing 10084, P. R. China E-mail address: m -ding04@mails.tsinghua.edu.cn (M.Ding), jxiao@math.tsinghua.edu.cn (J.Xiao), f-xu04@mails.tsinghua.edu.cn (F.Xu)