

Wellposedness of Cauchy problem for the Fourth Order Nonlinear Schrödinger Equations in Multi-dimensional Spaces

Chengchun Hao ^{a,*}, Ling Hsiao ^{a,2}, Baoxiang Wang ^b

^aAcademy of Mathematics and Systems Science, CAS, Beijing 100080, P.R. China

^bSchool of Mathematical Sciences, Peking University, Beijing 100871, P.R. China

Abstract

We study the wellposedness of Cauchy problem for the fourth order nonlinear Schrödinger equations

$$i\partial_t u = -\varepsilon\Delta u + \Delta^2 u + P((\partial_x^\alpha u)_{|\alpha|\leq 2}, (\partial_x^\alpha \bar{u})_{|\alpha|\leq 2}), \quad t \in \mathbb{R}, x \in \mathbb{R}^n,$$

where $\varepsilon \in \{-1, 0, 1\}$, $n \geq 2$ denotes the spatial dimension and $P(\cdot)$ is a polynomial excluding constant and linear terms.

2000 Mathematics Subject Classification. 35Q55, 35G25, 35A07.

Key words: Fourth order nonlinear Schrödinger equations, Cauchy problem, wellposedness, smoothing effects

1 Introduction

We consider the Cauchy problem for the fourth order nonlinear Schrödinger equations

$$i\partial_t u = -\varepsilon\Delta u + \Delta^2 u + P((\partial_x^\alpha u)_{|\alpha|\leq 2}, (\partial_x^\alpha \bar{u})_{|\alpha|\leq 2}), \quad t \in \mathbb{R}, x \in \mathbb{R}^n, \quad (1.1)$$

$$u(0, x) = u_0(x), \quad (1.2)$$

* Corresponding author.

Email addresses: hcc@amss.ac.cn (Chengchun Hao), hsiaol@amss.ac.cn (Ling Hsiao), wbx@math.pku.edu.cn (Baoxiang Wang).

¹ Supported partially by the Innovation Funds of AMSS, CAS of China.

² Supported partially by NSFC(Grant No.10431060), NSAF(Grant No.10276036).

where $\varepsilon \in \{-1, 0, 1\}$, $\alpha \in \mathbb{Z}^n$, $u = u(t, x)$ is a complex valued wave function of $(t, x) \in \mathbb{R}^{1+n}$ and Δ is the Laplace operator on \mathbb{R}^n with $n \geq 2$. $P(\cdot)$ is a complex valued polynomial defined in \mathbb{C}^{n^2+3n+2} such that

$$P(\vec{z}) = P(z_1, z_2, \dots, z_{n^2+3n+2}) = \sum_{\substack{l \leq |\beta| \leq h \\ \beta \in \mathbb{Z}^{n^2+3n+2}}} a_\beta z^\beta, \quad \text{for } l, h \in \mathbb{N}, \quad (1.3)$$

and there exists $a_{\beta_0} \neq 0$ for some $\beta_0 \in \mathbb{Z}^{n^2+3n+2}$ with $|\beta_0| = l \geq 2$.

This class of nonlinear Schrödinger equation has been widely applied in many branches in applied science such as deep water wave dynamics, plasma physics, optical communications and so on [6]. A large amount of interesting work has been devoted to the study of Cauchy problem to dispersive equations, such as [1,3–5,7–24,26] and references cited therein.

In order to study the influence of higher order dispersion on solitary waves, instability and the collapse phenomena, V. I. Karpman [12] introduced a class of nonlinear Schrödinger equations

$$i\Psi_t + \frac{1}{2}\Delta\Psi + \frac{\gamma}{2}\Delta^2\Psi + f(|\Psi|^2)\Psi = 0, \quad x \in \mathbb{R}^n, \quad t \in \mathbb{R}.$$

This system with different nonlinearities was discussed by lots of authors. In [22], by using the method of Fourier restriction norm, J. Segata studied a special fourth order nonlinear Schrödinger equation in one dimensional space and considered the three-dimensional motion of an isolated vortex filament, which was introduced by Da Rios, embedded in inviscid incompressible fluid fulfilled in an infinite region. And the results have been improved in [11,23].

In [1], M. Ben-Artzi, H. Koch and J. C. Saut discussed the sharp space-time decay properties of fundamental solutions to the linear equation

$$i\Psi_t - \varepsilon\Delta\Psi + \Delta^2\Psi = 0, \quad \varepsilon \in \{-1, 0, 1\}.$$

In [8], B. L. Guo and B. X. Wang considered the existence and scattering theory for the Cauchy problem of nonlinear Schrödinger equations with the form

$$iu_t + (-\Delta)^m u + f(u) = 0, \quad (1.4)$$

$$u(0, x) = \varphi(x), \quad (1.5)$$

where $m \geq 1$ is an integer. H. Pecher and W. von Wahl in [21] proved the existence of classical global solutions of (1.4)–(1.5) for the space dimensions $n \leq 7m$ for the case $m \geq 1$. In [9], we discussed the local wellposedness of

the Cauchy problem (1.4)–(1.5) without smallness of data for $m = 2$ and $f(u) = P((\partial_x^j u)_{j \leq 2}, (\partial_x^j \bar{u})_{j \leq 2})$ in one dimension.

In the present paper we deal with the equation (1.1) in which the difficulty arises from the operator semigroup itself and the fact that the nonlinearity of P involves the first and the second derivatives. This could cause the so-called loss of derivatives so long as we make direct use of the standard methods, such as the energy estimates, the space-time estimates, etc. In addition, compared to the one-dimensional case, it is quite different in the multi-dimensional case, since we have to separate the spatial space \mathbb{R}^n into a family of non-overlapping cubes of bounded size.

In [17], C. E. Kenig, G. Ponce and L. Vega made a great progress on the nonlinear Schrödinger equation of the form

$$\partial_t u = i\Delta u + P(u, \bar{u}, \nabla_x u, \nabla_x \bar{u}), \quad t \in \mathbb{R}, \quad x \in \mathbb{R}^n$$

and proved that the corresponding Cauchy problem is locally wellposed for small data in the Sobolev space $H^s(\mathbb{R}^n)$ and in its weighted version by pushing forward the linear estimates associated with the Schrödinger group $\{e^{it\Delta}\}_{-\infty}^{\infty}$ and by introducing suitable function spaces where these estimates act naturally. They also studied generalized nonlinear Schrödinger equations in [18] and quasi-linear Schrödinger equations in [19]. In the one dimensional case, $n = 1$, the smallness assumption on the size of the data was removed by N. Hayashi and T. Ozawa [10] by using a change of variable to obtain an equivalent system with a nonlinear term independent of $\partial_x u$, where the new system could be treated by the standard energy method. And H. Chihara [4], was able to remove the size restriction on the data in any dimension by using an invertible classical pseudo-differential operator of order zero.

T. Kato showed, in [13], that solutions of the Korteweg-de Vries equation

$$\partial_t u + \partial_x^3 u + u\partial_x u = 0, \quad t, x \in \mathbb{R},$$

satisfy that

$$\int_{-T}^T \int_{-R}^R |\partial_x u(t, x)|^2 dx dt \leq C(T, R, \|u(0, x)\|_2).$$

And in [20], S. N. Kruzkov and A. V. Faminskiĭ, for $u(0, x) = u_0(x) \in L^2$ such that $x^\alpha u_0(x) \in L^2((0; +\infty))$, proved that the weak solution of the Korteweg-de Vries equation constructed there has l -continuous space derivatives for all $t > 0$ if $l < 2\alpha$. The corresponding version of the above estimate for the Schrödinger group $\{e^{it\Delta}\}_{-\infty}^{\infty}$

$$\int_{-T}^T \int_{-R}^R |(1 - \Delta)^{1/4} e^{it\Delta} u_0|^2 dx dt \leq C(T, R, \|u_0\|_2)$$

was simultaneously established by P. Constantin and J.-C. Saut [5], P. Sjölin [24] and L. Vega [26] and others.

In the present paper, we will develop the arguments in [17] to study the Cauchy problem (1.1)–(1.2). We first discuss the local smoothing effects of the unit group $\{S(t)\}_{-\infty}^{\infty}$ ($S(t)$ is defined as below) in order to overcome the loss of derivatives in Sec.2. To construct the work space, we have to study, in Sec.3, the boundedness properties of the maximal function $\sup_{[0,T]} |S(t)\cdot|$. This idea is implicit in the splitting argument introduced by J. Ginibre and Y. Tsutsumi [7] to deal with uniqueness for the generalized KdV equation. Finally, we will consider some special cases in Sec.4 to apply the estimates we have obtained in the previous sections.

For convenience, we introduce some notations. $S(t) := e^{it(\varepsilon\Delta - \Delta^2)}$ denotes the unitary group generated by $i(\varepsilon\Delta - \Delta^2)$ in $L^2(\mathbb{R}^n)$. \bar{z} denotes the conjugate of the complex number z . $\mathcal{F}u$ or \hat{u} ($\mathcal{F}^{-1}u$, respectively) denotes the Fourier (inverse, respectively) transform of u with respect to all variables except the special announcement. \mathcal{S} denotes the space of Schwartz' functions. $\{Q_\alpha\}_{\alpha \in \mathbb{Z}^n}$ is the family of non-overlapping cubes of size R such that $\mathbb{R}^n = \bigcup_{\alpha \in \mathbb{Z}^n} Q_\alpha$.

$D_x^\gamma f(x) = \mathcal{F}^{-1} |\xi|^\gamma \mathcal{F} f(x)$ is the homogeneous derivative of order γ of f . H^s (\dot{H}^s , respectively) denotes the usual inhomogeneous (homogeneous, respectively) Sobolev space. We denote $\|\cdot\|_{s,2} = \|\cdot\|_{H^s}$, $\|\cdot\|_p = \|\cdot\|_{L^p}$ for $1 \leq p \leq \infty$ and

$$\|f\|_{l,2,j} = \|f\|_{H^l(\mathbb{R}; |x|^j dx)} = \sum_{|\gamma| \leq l} \left(\int_{\mathbb{R}^n} |\partial_x^\gamma f(x)|^2 |x|^j dx \right)^{1/2}. \quad (1.6)$$

Throughout the paper, the constant C might be different from each other and $\lfloor \frac{n}{2} \rfloor$ denotes the greatest integer that is less than or equal to $\frac{n}{2}$.

Now we state the main results of this paper.

Theorem 1 (Case $l \geq 3$) *Let $n \geq 2$. Given any polynomial P as in (1.3) with $l \geq 3$, then, for any $u_0 \in H^s(\mathbb{R}^n)$ with $s \geq s_0 = n + 2 + \frac{1}{2}$, there exists a $T = T(\|u_0\|_{s,2}) > 0$ such that the Cauchy problem (1.1)–(1.2) has a unique solution $u(t)$ defined in the time interval $[0, T]$ and satisfying*

$$u \in C([0, T]; H^s(\mathbb{R}^n)),$$

and

$$u \in L^2([0, T]; \dot{H}^{s+\frac{1}{2}}(Q)),$$

for any cube Q of unit size in \mathbb{R}^n .

Theorem 2 (Case $l = 2$) *Let $n \geq 2$. Given any polynomial P as in (1.3) with $l = 2$, then, for any $u_0 \in H^s(\mathbb{R}^n) \cap H^{n+4[\frac{n}{2}]+8}(\mathbb{R}^n; |x|^{2[\frac{n}{2}]+2} dx)$ with $s \geq$*

$s_0 = 2n + 3 \left[\frac{n}{2} \right] + 15 + \frac{1}{2}$, there exists a $T = T(\|u_0\|_{s,2}, \|u_0\|_{n+4[\frac{n}{2}]+8,2, [\frac{n}{2}]+2}) > 0$ such that the Cauchy problem (1.1)-(1.2) has a unique solution $u(t)$ defined in the time interval $[0, T]$ and satisfying

$$u \in C \left([0, T]; H^s(\mathbb{R}^n) \cap H^{n+[\frac{n}{2}]+8}(\mathbb{R}^n; |x|^{2[\frac{n}{2}]+2} dx) \right),$$

and

$$u \in L^2([0, T]; \dot{H}^{s+\frac{1}{2}}(Q)),$$

for any cube Q of unit size in \mathbb{R}^n .

2 Local Smoothing Effects

We will prove the local smoothing effects exhibited by the semigroup $\{S(t)\}_{-\infty}^{\infty}$ in this section.

Lemma 3 (Local smoothing effect: homogeneous case) *We have the following estimate*

$$\sup_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_{-\infty}^{\infty} |D_x^{3/2} S(t) u_0(x)|^2 dt dx \right)^{1/2} \leq CR^{1/2} \|u_0\|_2, \quad (2.1)$$

and the corresponding dual version

$$\left\| D_x^{3/2} \int_0^T S(-\tau) f(\tau, \cdot) d\tau \right\|_2 \leq CR^{1/2} \sum_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_0^T |f(x, t)|^2 dt dx \right)^{1/2}. \quad (2.2)$$

Proof. (2.1) and (2.2) can be derived from [15, Theorem 4.1] for which we omit the details. \square

More precisely, (2.2) yields, for $t \in [0, T]$, that

$$\left\| D_x^{3/2} \int_0^t S(t - \tau) f(\tau, \cdot) d\tau \right\|_2 \leq CR^{1/2} \sum_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_0^T |f(x, t)|^2 dt dx \right)^{1/2}. \quad (2.3)$$

Moreover, with the help of the Fubini theorem, (2.1) implies that

$$\sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |S(t) u_0(x)|^2 dx dt \right)^{1/2} \leq CR^{1/2} \|u_0\|_{\dot{H}^{-\frac{3}{2}}}. \quad (2.4)$$

In addition, we have

$$\begin{aligned}
& \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |S(t)u_0(x)|^2 dx dt \right)^{1/2} \\
& \leq T^{\frac{1}{2}} \left(\sup_t \int_{\mathbb{R}^n} |S(t)u_0(x)|^2 dx \right)^{1/2} \\
& \leq T^{\frac{1}{2}} \|u_0\|_2.
\end{aligned} \tag{2.5}$$

Interpolating both (2.4) and (2.5), we can obtain that

$$\sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |S(t)u_0(x)|^2 dx dt \right)^{1/2} \leq CR^{\frac{1}{6}} T^{\frac{1}{3}} \|u_0\|_{\dot{H}^{-\frac{1}{2}}}. \tag{2.6}$$

Now we turn to consider the inhomogeneous Cauchy problem:

$$i\partial_t u = -\varepsilon \Delta u + \Delta^2 u + F(t, x), \quad t \in \mathbb{R}, x \in \mathbb{R}^n, \tag{2.7}$$

$$u(0, x) = 0, \tag{2.8}$$

with $F \in \mathcal{S}(\mathbb{R} \times \mathbb{R}^n)$. We have the following estimate on the local smoothing effect in this inhomogeneous case.

Lemma 4 (Local smoothing effect: inhomogeneous case) *The solution $u(t, x)$ of the Cauchy problem (2.7)-(2.8) satisfies*

$$\sup_{\alpha \in \mathbb{Z}^n} \left\| D_x^2 u(t, x) \right\|_{L_x^2(Q_\alpha; L_t^2([0, T]))} \leq CRT^{\frac{1}{4}} \sum_{\alpha \in \mathbb{Z}^n} \|F\|_{L_x^2(Q_\alpha; L_t^2([0, T]))}. \tag{2.9}$$

Proof. Separating

$$F = \sum_{\alpha \in \mathbb{Z}^n} F \chi_{Q_\alpha} = \sum_{\alpha \in \mathbb{Z}^n} F_\alpha,$$

and

$$u = \sum_{\alpha \in \mathbb{Z}^n} u_\alpha,$$

where $u_\alpha(t, x)$ is the corresponding solution of the Cauchy problem

$$i\partial_t u_\alpha = -\varepsilon \Delta u_\alpha + \Delta^2 u_\alpha + F_\alpha(t, x), \quad t \in \mathbb{R}, x \in \mathbb{R}^n, \tag{2.10}$$

$$u_\alpha(0, x) = 0, \tag{2.11}$$

we formally take Fourier transform in both variables t and x in the equation (2.10) and obtain

$$\hat{u}_\alpha(\tau, \xi) = \frac{\hat{F}_\alpha(\tau, \xi)}{\tau - \varepsilon |\xi|^2 - |\xi|^4}, \quad \text{for each } \alpha \in \mathbb{Z}^n.$$

By the Plancherel theorem in the time variable, we can get

$$\begin{aligned}
& \sup_{\alpha \in \mathbb{Z}^n} \left\| D_x^2 u_\beta(t, x) \right\|_{L_x^2(Q_\alpha; L_t^2([0, T]))} = \sup_{\alpha \in \mathbb{Z}^n} \left\| D_x^2 \mathcal{F}_t u_\beta(\tau, x) \right\|_{L_x^2(Q_\alpha; L_\tau^2(\mathbb{R}))} \\
&= \sup_{\alpha \in \mathbb{Z}^n} \left\| D_x^2 \mathcal{F}_\xi^{-1} \left(\frac{\hat{F}_\beta(\tau, \xi)}{\tau - \varepsilon |\xi|^2 - |\xi|^4} \right) \right\|_{L_x^2(Q_\alpha; L_\tau^2(\mathbb{R}))} \\
&= \sup_{\alpha \in \mathbb{Z}^n} \left\| \mathcal{F}_\xi^{-1} \left(\frac{|\xi|^2}{\tau - \varepsilon |\xi|^2 - |\xi|^4} \hat{F}_\beta(\tau, \xi) \right) \right\|_{L_x^2(Q_\alpha; L_\tau^2(\mathbb{R}))} \\
&= \sup_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_{-\infty}^{\infty} \left| \mathcal{F}_\xi^{-1} \left(\frac{|\xi|^2}{\tau - \varepsilon |\xi|^2 - |\xi|^4} \hat{F}_\beta(\tau, \xi) \right) \right|^2 d\tau dx \right)^{1/2}. \quad (2.12)
\end{aligned}$$

In order to continue the above estimate, we introduce the following estimate.

Lemma 5 *Let $\mathcal{M}f = \mathcal{F}^{-1}m(\xi)\mathcal{F}f$ and $m(\xi) = \frac{|\xi|^2}{1 - \varepsilon\tau^{-\frac{1}{2}}|\xi|^2 - |\xi|^4}$ for $\tau > 0$ and $\varepsilon \in \{-1, 0, 1\}$, where $\mathcal{F}(\mathcal{F}^{-1})$ denotes the Fourier (inverse, respectively) transform in x only. Then, we have*

$$\sup_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} |\mathcal{M}(g\chi_{Q_\beta})|^2 dx \right)^{1/2} \leq CR \left(\int_{Q_\beta} |g|^2 dx \right)^{1/2}. \quad (2.13)$$

Proof. We have

$$\begin{aligned}
& 1 - \varepsilon\tau^{-\frac{1}{2}}|\xi|^2 - |\xi|^4 \\
&= \left(1 - \frac{\varepsilon}{2\sqrt{\tau}}|\xi|^2 \right)^2 - \left(\frac{\varepsilon^2}{4\tau} + 1 \right) |\xi|^4 \\
&= \left[1 + \left(\sqrt{\frac{\varepsilon^2}{4\tau} + 1} - \frac{\varepsilon}{2\sqrt{\tau}} \right) |\xi|^2 \right] \left[1 - \left(\sqrt{\frac{\varepsilon^2}{4\tau} + 1} + \frac{\varepsilon}{2\sqrt{\tau}} \right) |\xi|^2 \right] \\
&= \left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2 + 4\tau} - \varepsilon} + |\xi|^2 \right) \left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2 + 4\tau} + \varepsilon} - |\xi|^2 \right).
\end{aligned}$$

Let $\varphi \in C_0^\infty(\mathbb{R})$ with $\text{supp}\varphi \subset [-1, 1]$, $\varphi \equiv 1$ in $[-1/2, 1/2]$ and $0 \leq \varphi \leq 1$. Denote $a = \sqrt{\frac{2\sqrt{\tau}}{\varepsilon^2 + 4\tau + \varepsilon}}$. Let $\varphi_1(\xi) = \varphi(2(|\xi| - a))$ and choose $\varphi_2(\xi)$ such that $\varphi_1(\xi) + \varphi_2(\xi) = 1$.

Thus, we have $\text{supp}\varphi_1 \subset \{\xi : a - \frac{1}{2} \leq |\xi| \leq a + \frac{1}{2}\}$ and $\varphi_1(\xi) = 1$ for $a - \frac{1}{4} \leq |\xi| \leq a + \frac{1}{4}$.

Define

$$\mathcal{M}_j f(x) = \mathcal{F}^{-1}m_j(\xi)\mathcal{F}f(x), \quad j = 1, 2,$$

where $m_j(\xi) = \varphi_j(\xi)m(\xi)$.

First, we shall establish (2.13) for the operator \mathcal{M}_2 whose symbol $m_2(\xi)$ has no singularities. For p, p' such that

$$\frac{1}{p} + \frac{1}{p'} = 1 \text{ and } \frac{1}{p'} - \frac{1}{p} = \frac{1}{n},$$

it follows, by Hölder's inequality, Sobolev's embedding theorem and the Mihlin theorem, that

$$\begin{aligned} \left(\int_{Q_\alpha} |\mathcal{M}_2(g\chi_{Q_\beta})|^2 dx \right)^{\frac{1}{2}} &\leq C \|1\|_{L^{\frac{1}{1/2-1/p}}(Q_\alpha)} \|\mathcal{M}_2(g\chi_{Q_\beta})\|_{L^p(Q_\alpha)} \\ &\leq CR^{n(\frac{1}{2}-\frac{1}{p})} \|\mathcal{M}_2(g\chi_{Q_\beta})\|_{L^p(Q_\alpha)} \\ &\leq CR^{n(\frac{1}{2}-\frac{1}{p})} \|D_x \mathcal{M}_2(g\chi_{Q_\beta})\|_{L^{p'}(Q_\alpha)} \\ &\leq CR^{n(\frac{1}{2}-\frac{1}{p})} \|g\chi_{Q_\beta}\|_{L^{p'}(Q_\alpha)} \\ &\leq CR \|g\chi_{Q_\beta}\|_{L^2(Q_\alpha)}, \end{aligned}$$

where we have used the fact

$$\|D_x \mathcal{M}_2 f\|_p \leq C \|f\|_p, \text{ for } 1 < p < \infty. \quad (2.14)$$

In fact,

$$\begin{aligned} D_x \mathcal{M}_2 f &= \mathcal{F}^{-1} |\xi| m_2(\xi) \mathcal{F} f \\ &= \mathcal{F}^{-1} \frac{|\xi|^3 [1 - \varphi(2(|\xi| - a))]}{\left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2 + 4\tau - \varepsilon}} + |\xi|^2\right) (a^2 - |\xi|^2)} \mathcal{F} f. \end{aligned}$$

Denote $\rho(\xi) = \frac{|\xi|^3 [1 - \varphi(2(|\xi| - a))]}{\left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2 + 4\tau - \varepsilon}} + |\xi|^2\right) (a^2 - |\xi|^2)}$. In the case $|\xi| \geq a + \frac{1}{4}$, we have

$$|\rho(\xi)| \leq \frac{1 - \varphi(2(|\xi| - a))}{4} \leq \frac{1}{4}.$$

In the case $|\xi| \leq a - \frac{1}{4}$ where we assume that $a > \frac{1}{4}$, we have the same estimate. Thus, $|\rho(\xi)| \leq C$. By calculating the derivatives of ρ with respect to the variable ξ , we are able to get

$$\left| \frac{\partial^\alpha}{\partial \xi^\alpha} \rho(\xi) \right| \leq C |\xi|^{-|\alpha|}, \quad |\alpha| \leq L,$$

for some integer $L > \frac{n}{2}$. Therefore, we have the desired estimate (2.14) in view of the Mihlin multiplier theorem.

To estimate \mathcal{M}_1 , we split its symbol $m_1(\xi)$ into a finite number of pieces (depending only on the dimension n). Let $\theta \in C_0^\infty(\mathbb{R})$ with $\text{supp}\theta \subset [-a, a]$ where a is the same as above, define

$$m_{1,1}(\xi) = m_1(\xi)\theta(4|\bar{\xi}|),$$

and

$$\mathcal{M}_{1,1}f = \mathcal{F}^{-1}m_{1,1}(\xi)\mathcal{F}f,$$

where $\xi = (\bar{\xi}, \xi_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. By a rotation argument, $m_1(\cdot)$ can be expressed as a finite sum of $m_{1,1}$'s. Notice that

$$\begin{aligned} \text{supp}m_{1,1} &\subset \left\{ \xi = (\bar{\xi}, \xi_n) \in \mathbb{R}^{n-1} \times \mathbb{R} : |\bar{\xi}| \leq \frac{a}{4} \text{ and } a - \frac{1}{2} \leq |\xi| \leq a + \frac{1}{2} \right\}, \\ Q_\alpha &\subset \mathbb{R}^{n-1} \times [a_\alpha, a_\alpha + R] \text{ and } Q_\beta \subset \mathbb{R}^{n-1} \times [a_\beta, a_\beta + R] \end{aligned}$$

for appropriate constants a_α and a_β .

Thus, by the Plancherel theorem in the \bar{x} variable, we have

$$\begin{aligned} &\int_{Q_\alpha} |\mathcal{M}_{1,1}(g\chi_{Q_\beta})|^2 dx \\ &\leq \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} |\mathcal{M}_{1,1}(g\chi_{Q_\beta})|^2 d\bar{x} dx_n \\ &= \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} \left| \int_{-\infty}^{\infty} e^{ix_n\xi_n} m_{1,1}(\xi) \widehat{g\chi_{Q_\beta}}(\bar{\xi}, \xi_n) d\xi_n \right|^2 d\bar{\xi} dx_n \\ &= \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} \left| \int_{-\infty}^{\infty} e^{ix_n\xi_n} m_{1,1}(\xi) \int_{a_\beta}^{a_\beta+R} \int_{\mathbb{R}^{n-1}} e^{-iy\xi} (g\chi_{Q_\beta})(\bar{y}, y_n) d\bar{y} dy_n d\xi_n \right|^2 d\bar{\xi} dx_n \\ &= \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} \left| \int_{a_\beta}^{a_\beta+R} \int_{\mathbb{R}^{n-1}} e^{-i\bar{y}\bar{\xi}} (g\chi_{Q_\beta})(\bar{y}, y_n) \int_{-\infty}^{\infty} e^{i(x_n-y_n)\xi_n} m_{1,1}(\xi) d\xi_n d\bar{y} dy_n \right|^2 d\bar{\xi} dx_n \\ &= \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} \left| \int_{a_\beta}^{a_\beta+R} \int_{\mathbb{R}^{n-1}} e^{-i\bar{y}\bar{\xi}} (g\chi_{Q_\beta})(\bar{y}, y_n) b(x_n, y_n, \bar{\xi}) d\bar{y} dy_n \right|^2 d\bar{\xi} dx_n \\ &=: E, \end{aligned}$$

where

$$\begin{aligned} b(x_n, y_n, \bar{\xi}) &= \int_{-\infty}^{\infty} e^{i(x_n-y_n)\xi_n} m_{1,1}(\xi) d\xi_n \\ &= \int_{-\infty}^{\infty} e^{i(x_n-y_n)\xi_n} \frac{|\xi|^2 \varphi(2(|\xi| - a))}{\left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2+4\tau-\varepsilon}} + |\xi|^2\right) (a^2 - |\xi|^2)} \theta(4|\bar{\xi}|) d\xi_n \\ &= \theta(4|\bar{\xi}|) \int_{-\infty}^{\infty} e^{i\lambda\xi_n} \frac{|\xi|^2}{\left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2+4\tau-\varepsilon}} + |\xi|^2\right) (a^2 - |\xi|^2)} \varphi(2(|\xi| - a)) d\xi_n \\ &= K(\lambda, \bar{\xi}), \end{aligned}$$

here $\lambda = x_n - y_n$ and the support of the integrand is contained in

$$A = \left\{ \xi = (\bar{\xi}, \xi_n) \in \mathbb{R}^{n-1} \times \mathbb{R} : |\bar{\xi}| \leq \frac{a}{4} \text{ and } a - \frac{1}{2} \leq |\xi| \leq a + \frac{1}{2} \right\}.$$

For $(\bar{\xi}, \xi_n) \in A$, we have $|\xi|^2 - a^2 = \xi_n^2 + |\bar{\xi}|^2 - a^2 = \xi_n^2 - \mu^2$, where $\mu^2 = a^2 - |\bar{\xi}|^2 > \frac{15}{16}a^2$. Then, we separate K into two parts, *i.e.*, $K = K_+ + K_-$ with

$$\begin{aligned} K_+(\lambda, \bar{\xi}) &= \theta(4|\bar{\xi}|) \int_0^\infty e^{i\lambda\xi_n} \frac{1}{\xi_n - \mu} \cdot \frac{|\xi|^2}{\xi_n + \mu} \cdot \frac{\varphi(2(|\xi| - a))}{\left(\frac{2\sqrt{\tau}}{\sqrt{\varepsilon^2 + 4\tau - \varepsilon}} + |\xi|^2\right)} d\xi_n \\ &= \theta(4|\bar{\xi}|) \mathcal{F}_{\xi_n}^{-1} \left(\frac{1}{\xi_n - \mu} \psi(\bar{\xi}, \xi_n) \right) (\lambda) \\ &= \theta(4|\bar{\xi}|) \mathcal{F}_{\xi_n}^{-1} \left(\frac{1}{\xi_n - \mu} \right) (\lambda) * \mathcal{F}_{\xi_n}^{-1} \psi(\bar{\xi}, \xi_n) (\lambda) \\ &= \theta(4|\bar{\xi}|) e^{i\lambda\mu} \text{sgn}(\lambda) * \mathcal{F}_{\xi_n}^{-1} (\psi(\bar{\xi}, \xi_n)) (\lambda) \\ &= \theta(4|\bar{\xi}|) \int_{-\infty}^\infty e^{iy_n\mu} \text{sgn}(y_n) \mathcal{F}_{\xi_n}^{-1} \psi(\bar{\xi}, \xi_n) (\lambda - y_n) dy_n, \end{aligned}$$

where $\psi(\bar{\xi}, \xi_n) \in C_0^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^+)$.

Thus, it follows that

$$|K_+(\lambda, \bar{\xi})| \leq \int_{-\infty}^\infty \left| \mathcal{F}_{\xi_n}^{-1} \psi(\bar{\xi}, \xi_n)(y_n) \right| dy_n \leq C. \quad (2.15)$$

In the same way, we have

$$|K_-(\lambda, \bar{\xi})| \leq C. \quad (2.16)$$

(2.15) and (2.16) yield that $|b(x_n, y_n, \bar{\xi})| \leq C$. Now, we can turn to the estimate of $\int_{Q_\alpha} \left| \mathcal{M}_{1,1}(g\chi_{Q_\beta}) \right|^2 dx$ and get, by the Schwartz inequality in y_n variable and the Plancherel theorem in \bar{y} variable, that

$$\begin{aligned} E &\leq R \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} \int_{a_\beta}^{a_\beta+R} \left| \int_{\mathbb{R}^{n-1}} e^{-i\bar{y}\bar{\xi}} (g\chi_{Q_\beta})(\bar{y}, y_n) d\bar{y} \right|^2 b^2(x_n, y_n, \bar{\xi}) dy_n d\bar{\xi} dx_n \\ &\leq CR \int_{a_\alpha}^{a_\alpha+R} \int_{\mathbb{R}^{n-1}} \int_{a_\beta}^{a_\beta+R} \left| \int_{\mathbb{R}^{n-1}} e^{-i\bar{y}\bar{\xi}} (g\chi_{Q_\beta})(\bar{y}, y_n) d\bar{y} \right|^2 dy_n d\bar{\xi} dx_n \\ &= CR \int_{a_\alpha}^{a_\alpha+R} \int_{a_\beta}^{a_\beta+R} \int_{\mathbb{R}^{n-1}} \left| \int_{\mathbb{R}^{n-1}} e^{-i\bar{y}\bar{\xi}} (g\chi_{Q_\beta})(\bar{y}, y_n) d\bar{y} \right|^2 d\bar{\xi} dy_n dx_n \\ &= CR \int_{a_\alpha}^{a_\alpha+R} \int_{a_\beta}^{a_\beta+R} \int_{\mathbb{R}^{n-1}} |(g\chi_{Q_\beta})(\bar{y}, y_n)|^2 d\bar{y} dy_n dx_n \end{aligned}$$

$$= CR^2 \int_{\mathbb{R}^n} |(g\chi_{Q_\beta})(y)|^2 dy.$$

Therefore, we obtain

$$\sup_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} |\mathcal{M}(g\chi_{Q_\beta})|^2 dx \right)^{1/2} \leq CR \|g\chi_{Q_\beta}\|_{L^2(\mathbb{R}^n)} = CR \|g\|_{L^2(Q_\beta)}$$

as desired. \square

Now, we go back to the proof of Lemma 4. We first consider the part when $\tau > 0$ in (2.12), *i.e.*,

$$\begin{aligned} & \sup_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_0^\infty \left| \mathcal{F}_\xi^{-1} \left(\frac{|\xi|^2}{\tau - \varepsilon |\xi|^2 - |\xi|^4} \hat{F}_\beta(\tau, \xi) \right) \right|^2 d\tau dx \right)^{1/2} \\ &= \left(\int_0^\infty \tau^{\frac{n}{4}-1} \sup_{\alpha \in \mathbb{Z}^n} \int_{\tau^{\frac{1}{4}} Q_\alpha} \left| \int_{\mathbb{R}^n} e^{iy\eta} \frac{|\eta|^2}{1 - \varepsilon \tau^{-\frac{1}{2}} |\eta|^2 - |\eta|^4} \hat{F}_\beta(\tau, \tau^{\frac{1}{4}} \eta) d\eta \right|^2 dy d\tau \right)^{1/2} \\ &= \left(\int_0^\infty \tau^{-\frac{n}{4}-1} \sup_{\alpha \in \mathbb{Z}^n} \int_{\tau^{\frac{1}{4}} Q_\alpha} |\mathcal{M} \mathcal{F}_t F_\beta(\tau, \tau^{-\frac{1}{4}} y)|^2 dy d\tau \right)^{1/2} \\ &\leq CR \left(\int_0^\infty \tau^{-\frac{n}{4}-\frac{1}{2}} \int_{\tau^{\frac{1}{4}} Q_\beta} |\mathcal{F}_t F_\beta(\tau, \tau^{-\frac{1}{4}} y)|^2 dy d\tau \right)^{1/2} \\ &\leq CR \left(\int_0^\infty \tau^{-\frac{1}{2}} \int_{Q_\beta} |\mathcal{F}_t F(\tau, x)|^2 dx d\tau \right)^{1/2} \end{aligned}$$

where we have used the changes of variables, Lemma 5 and the identity

$$(\mathcal{F}_\eta^{-1} \hat{F}_\beta(\tau, \tau^{\frac{1}{4}} \eta))(\tau, y) = \tau^{-\frac{n}{4}} \mathcal{F}_t F_\beta(\tau, \tau^{-\frac{1}{4}} y).$$

For the part when $\tau \in (-\infty, 0)$ in (2.12), we are able to split it into two cases. First, it is easier to handle the case $\varepsilon \in \{0, 1\}$ since this corresponds to the symbol $\frac{|\eta|^2}{1 + \varepsilon |\tau|^{-\frac{1}{2}} |\eta|^2 + |\eta|^4}$ which has no singularity. Next, for the case $\varepsilon = -1$, we have

$$\frac{|\eta|^2}{1 + \varepsilon |\tau|^{-\frac{1}{2}} |\eta|^2 + |\eta|^4} = \frac{|\eta|^2}{\left(1 - \frac{1}{2|\tau|^{\frac{1}{2}}}\right)^2 + \left(1 - \frac{1}{4|\tau|}\right) |\eta|^4} =: B.$$

It is obvious that B has no singularity when $|\tau| > \frac{1}{4}$. Consequently, we have to deal with the case $|\tau| < \frac{1}{4}$. In fact, we have

$$B = \frac{|\eta|^2}{\left(\frac{2|\tau|^{\frac{1}{2}}}{1 - \sqrt{1-4|\tau|}} - |\eta|^2\right) \left(\frac{2|\tau|^{\frac{1}{2}}}{1 + \sqrt{1-4|\tau|}} - |\eta|^2\right)}.$$

Thus, we can use the same argument as in the proof of Lemma 5 step by step for the cases $|\eta|^2 \leq \frac{1}{2\sqrt{|\tau|}}$ and $|\eta|^2 > \frac{1}{2\sqrt{|\tau|}}$ respectively, and get that

$$\sup_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \left| \mathcal{F}_\eta^{-1} \frac{|\eta|^2}{1 - |\tau|^{-\frac{1}{2}} |\eta|^2 + |\eta|^4} \mathcal{F}_x(g\chi_{Q_\beta}) \right|^2 dx \right)^{\frac{1}{2}} \leq CR \left(\int_{Q_\beta} |g|^2 dx \right)^{\frac{1}{2}}.$$

Therefore, we obtain, by the Plancherel theorem, the Sobolev embedding theorem and the Hölder inequality, that

$$\begin{aligned} & \sup_{\alpha \in \mathbb{Z}^n} \|D_x^2 u_\beta(t, x)\|_{L_x^2(Q_\alpha; L_t^2([0, T]))} \\ & \leq CR \left(\int_{-\infty}^{\infty} |\tau|^{-\frac{1}{2}} \int_{Q_\beta} |\mathcal{F}_t F(\tau, x)|^2 dx d\tau \right)^{1/2} \\ & \leq CR \left(\int_{Q_\beta} \int_{-\infty}^{\infty} \left| |\tau|^{-\frac{1}{4}} \mathcal{F}_t F(\tau, x) \right|^2 d\tau dx \right)^{1/2} \\ & \leq CR \left(\int_{Q_\beta} \|F(\cdot, x)\|_{\dot{H}^{-\frac{1}{4}}([0, T])}^2 dx \right)^{1/2} \\ & \leq CR \left(\int_{Q_\beta} \|F(\cdot, x)\|_{L^{\frac{4}{3}}([0, T])}^2 dx \right)^{1/2} \\ & \leq CR \left(\int_{Q_\beta} \|1\|_{L^4([0, T])}^2 \|F(\cdot, x)\|_{L^2([0, T])}^2 dx \right)^{1/2} \\ & \leq CRT^{\frac{1}{4}} \|F\|_{L_x^2(Q_\beta; L_{t \in [0, T]}^2)}, \end{aligned}$$

which implies the desired result (2.9). \square

3 Estimates for the Maximal Function

For simplicity, let $\{Q_\alpha\}_{\alpha \in \mathbb{Z}^n}$ denote the mesh of dyadic cubes of unit size. We start with an L^2 -continuity result for the maximal function $\sup_{[0, T]} |S(t) \cdot|$.

Lemma 6 *For any $s > n + \frac{1}{2}$ and $T \in (0, 1]$*

$$\left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{\alpha \in \mathbb{Z}^n} |S(t)u_0(x)|^2 \right)^{1/2} \leq C \|u_0\|_{s, 2}. \quad (3.1)$$

In particular,

$$\left(\int_{\mathbb{R}^n} \sup_{t \in [0, T]} |S(t)u_0(x)|^2 dx \right)^{1/2} \leq C \|u_0\|_{s,2}. \quad (3.2)$$

Proof. It is clear that it suffices to prove the case $t = 1$. Let $\{\psi_k\}_{k=0}^\infty$ be a smooth partition of unity in \mathbb{R}^n such that the ψ_k 's are radial with $\text{supp}\psi_0 \subset \{\xi : |\xi| \leq 1\}$, $\text{supp}\psi_k \subset \{\xi : 2^{k-1} \leq |\xi| \leq 2^{k+1}\}$, $\psi_k(x) \in [0, 1]$ and $|\psi'_k(x)| \leq C2^{-k}$ for $k \in \mathbb{Z}^+$. For $k \geq 1$ and $t \in [0, 2]$, define

$$I(t, r) = \int_{-\infty}^{\infty} e^{i\phi_r(s)} \psi_k(s) ds,$$

where the phase function $\phi_r(s) = -\varepsilon t s^2 - t s^4 + r s$. Consequently, we have $\phi'_r(s) = -2\varepsilon t s - 4t s^3 + r$, $\phi''_r(s) = -2\varepsilon t - 12t s^2$ and $\phi'''_r(s) = -24t$.

Denote

$$\Omega = \left\{ s \in \mathbb{R}^+ : |\phi'_r(s)| < \frac{r}{2} \right\}$$

and

$$I_k = [2^{k-1}, 2^{k+1}].$$

For $r \in (0, 1)$, it is obvious that $|I(t, r)| \leq \frac{3}{2} \cdot 2^k$.

For $r > 1$, we divide it into three cases:

- i) Ω is located to the left of I_k ;
- ii) $\Omega \cap I_k \neq \emptyset$;
- iii) Ω is located to the right of I_k .

In the cases i) and ii), we have $t \geq C \frac{r}{2^{3k}}$ and $|\phi'''_r(s)| \geq C \frac{r}{2^{3k}}$. Thus, with the help of the Van der Corput lemma, we have $|I(t, r)| \leq C \left(\frac{2^{3k}}{r} \right)^{1/4}$, where we have used $|\psi'_k(x)| \leq C2^{-k}$.

In the case iii), we have $|\phi'_r(s)| \leq \frac{r}{2}$ and $\frac{r}{t} \leq C2^{3k}$. Integration by parts gives that

$$\begin{aligned} |I(t, r)| &\leq \int_0^\infty \left[\left| \frac{\phi''_r(s)}{(\phi'_r(s))^2} \right| \psi_k(s) + \frac{1}{|\phi'_r(s)|} |\psi'_k(s)| \right] ds \\ &\leq \int_{I_k} \left[\frac{4(2t + 12ts^2)}{r^2} \psi_k(s) + \frac{2}{r} |\psi'_k(s)| \right] ds \end{aligned}$$

$$\begin{aligned}
&\leq C \int_{I_k} \left[\frac{4(2 + 12 \cdot 2^{2(k+1)})}{r \cdot 2^{3k}} \psi_k(s) + \frac{2}{r} 2^{-k} \right] ds \\
&\leq \frac{C}{r} \leq C \left(\frac{2^{3k}}{r} \right)^{1/4},
\end{aligned}$$

where we have used the condition $r > 1$ in the last step.

Finally, we always have the case in which Ω is located to the right of I_k when $r \geq C2^{3k}$. Hence, we can integrate by parts as in the case iii) N -times and get that $|I(t, r)| \leq C_N/r^N$ for any $N \in \mathbb{Z}^+$.

Therefore, we have obtained

$$|I(t, r)| \leq \begin{cases} C2^k, & \text{for } r \in (0, 1), \\ C \left(\frac{2^{3k}}{r} \right)^{1/4}, & \text{for } r \in (1, C2^{3k}), \\ \frac{C_N}{r^N}, & \text{for } r \geq C2^{3k}. \end{cases}$$

In order to continue the proof, we have to introduce some estimates. The Fourier transform of a radial function $f(|x|) = f(s)$ is given by the formula (cf. [25, p.154, Theorem 3.3])

$$\hat{f}(r) = \hat{f}(|\xi|) = r^{-\frac{n-2}{2}} \int_0^\infty f(s) J_{\frac{n-2}{2}}(rs) s^{\frac{n}{2}} ds,$$

and the Bessel function is defined as

$$J_m(r) = \frac{\left(\frac{r}{2}\right)^m}{\Gamma(m + \frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{irs} (1 - s^2)^{\frac{2m-1}{2}} ds,$$

where $\Gamma(\cdot)$ is Gamma functions.

Lemma 7 ([17, Lemma 3.6])

$$\begin{aligned}
&J_m(r) = O(r^m) \quad \text{as } r \rightarrow 0, \\
&J_m(r) = e^{-ir} \sum_{j=0}^N \alpha_{m,j} r^{-(j+1/2)} + e^{ir} \sum_{j=0}^N \beta_{m,j} r^{-(j+1/2)} \\
&\quad + O(r^{-(N+3/2)}), \quad \text{as } r \rightarrow \infty, \quad (3.3)
\end{aligned}$$

for each $N \in \mathbb{Z}^+$. □

We continue the proof of Lemma 6 now. Denote

$$\tilde{I}(t, x) \equiv \left| \int_{\mathbb{R}^n} e^{-it(\varepsilon|\xi|^2 + |\xi|^4)} e^{ix\xi} \psi_k(|\xi|) d\xi \right|$$

$$= \frac{1}{r^{(n-2)/2}} \left| \int_0^\infty e^{-it(\varepsilon s^2 + s^4)} J_{\frac{n-2}{2}}(rs) \psi_k(s) s^{\frac{n}{2}} ds \right| \equiv \tilde{I}(t, r),$$

where $r = |x|$.

For $r \in (0, 1)$, we have

$$\tilde{I}(t, r) \leq \frac{1}{r^{\frac{n-2}{2}}} \int_0^\infty (rs)^{\frac{n-2}{2}} \psi_k(s) s^{\frac{n}{2}} ds \leq C \int_{I_k} s^{n-1} ds \leq C 2^{nk}.$$

For $r > 1$, we first consider the remainder term in (3.3) to obtain the bound

$$C \frac{1}{r^{\frac{n-2}{2}}} \int_{I_k} \frac{1}{(rs)^{N+3/2}} s^{\frac{n}{2}} ds \leq C \frac{2^{kn/2}}{r^n \cdot 2^{kn/2}} \leq \frac{C}{r^n},$$

where we have taken N such that $N > \frac{n-1}{2}$.

Next, we deal with the j -term in (3.3) for $0 \leq j \leq N$

$$\begin{aligned} & \frac{1}{r^{\frac{n-2}{2}}} \left| \int_0^\infty e^{-it(\varepsilon s^2 + s^4)} e^{isr} \frac{1}{(sr)^{j+1/2}} s^{n/2} \psi_k(s) ds \right| \\ & \leq \frac{1}{r^{\frac{n-2}{2}}} \frac{1}{r^{j+1/2}} 2^{-(k+1)(j+1/2)} 2^{nk/2} \left| \int_0^\infty e^{-it(\varepsilon s^2 + s^4)} e^{isr} \psi_k(s) ds \right| \\ & \leq C \frac{2^{\frac{kn}{2} - (j+\frac{1}{2})k}}{r^{\frac{n}{2} + j - \frac{1}{2}}} \cdot \begin{cases} \left(\frac{2^{3k}}{r}\right)^{\frac{1}{4}}, & r \in [1, C2^{3k}], \\ C_M r^{-M}, & r > C2^{3k}. \end{cases} \end{aligned}$$

Now, we consider the L^1 -norm of $\tilde{I}(t, x)$ with respect to x variable. For $r \in [0, 1]$, it is clear that $\int_{|x| \leq 1} |\tilde{I}(t, x)| dx \leq C \int_0^1 2^{nk} r^{n-1} dr \leq C 2^{nk}$. In the case $r \in [1, C2^{3k}]$, we have

$$\begin{aligned} \int_{1 \leq |x| \leq C2^{3k}} |\tilde{I}(t, x)| dx & \leq C \int_1^{C2^{3k}} \frac{2^{\frac{nk}{2} - (j+\frac{1}{2})k}}{r^{\frac{n}{2} + j - \frac{1}{2}}} \left(\frac{2^{3k}}{r}\right)^{\frac{1}{4}} r^{n-1} dr \\ & \leq C 2^{\frac{nk}{2} - (j-\frac{1}{4})k} \int_1^{C2^{3k}} r^{\frac{n}{2} - j - \frac{3}{4}} dr \leq C 2^{k(2n+1)}. \end{aligned}$$

For $r > C2^{3k}$, it follows that $\int_{|x| \geq C2^{3k}} |\tilde{I}(t, x)| dx \leq C 2^{kn}$. Thus, we obtain

$$\int_{\mathbb{R}^n} |\tilde{I}(t, x)| dx \leq C 2^{k(2n+1)}. \quad (3.4)$$

We decompose the operator $S(t)$ as follows according to the partition of unity $\{\psi_k\}$,

$$W_k(t)u_0 = \mathcal{F}^{-1} e^{-it(\varepsilon|\xi|^2 + |\xi|^4)} \psi_k(\xi) \mathcal{F} u_0.$$

To prove (3.1), it suffices to show that

$$\begin{aligned} & \left(\sum_{\gamma \in \mathbb{Z}^n} \sup_{|t| \leq 1} \sup_{x \in Q_\gamma} \left| \int_{-1}^1 W_k(t - \tau) g(\tau, \cdot) d\tau \right|^2 \right)^{1/2} \\ & \leq C 2^{k(2n+1)} \left(\sum_{\gamma \in \mathbb{Z}^n} \left(\int_{-1}^1 \int_{Q_\gamma} |g(t, x)| dx dt \right)^2 \right)^{1/2}. \end{aligned} \quad (3.5)$$

In fact, we have

$$\begin{aligned} & \left| \int_{-1}^1 W_k(t - \tau) g(\cdot, \tau) d\tau \right| \\ & = \left| \int_{-1}^1 \mathcal{F}^{-1} e^{-i(t-\tau)(\varepsilon|\xi|^2 + |\xi|^4)} \psi_k(\xi) \mathcal{F} g(\tau, \xi) d\tau \right| \\ & = \left| \int_{-1}^1 \int_{\mathbb{R}^n} \left[\mathcal{F}^{-1} e^{-i(t-\tau)(\varepsilon|\xi|^2 + |\xi|^4)} \psi_k(\xi) \right] (y) \cdot g(\tau, x - y) dy d\tau \right| \\ & = \left| \int_{\mathbb{R}^n} \int_{-1}^1 \left[\mathcal{F}^{-1} e^{-i(t-\tau)(\varepsilon|\xi|^2 + |\xi|^4)} \psi_k(\xi) \right] (y) \cdot g(\tau, x - y) d\tau dy \right| \\ & \leq \int_{\mathbb{R}^n} \int_{-1}^1 \left| \left[\mathcal{F}^{-1} e^{-i(t-\tau)(\varepsilon|\xi|^2 + |\xi|^4)} \psi_k(\xi) \right] (y) \right| |g(\tau, x - y)| d\tau dy \\ & \leq \int_{\mathbb{R}^n} \sup_{t \in [0, 2]} \tilde{I}(t, |y|) \int_{-1}^1 |g(\tau, x - y)| d\tau dy \\ & \leq \sum_{\alpha \in \mathbb{Z}^n} \left(\sup_{t \in [0, 2]} \sup_{y \in Q_\alpha} \tilde{I}(t, |y|) \right) \int_{-1}^1 \int_{Q_\alpha} |g(\tau, x - y)| dy d\tau. \end{aligned}$$

Thus, the left hand side of (3.5) is bounded by

$$\left(\sum_{\gamma \in \mathbb{Z}^n} \left[\sum_{\alpha \in \mathbb{Z}^n} \left(\sup_{t \in [0, 2]} \sup_{y \in Q_\alpha} \tilde{I}(t, |y|) \right) \sup_{x \in Q_\gamma} \int_{-1}^1 \int_{Q_\alpha} |g(\tau, x - y)| dy d\tau \right]^2 \right)^{1/2}. \quad (3.6)$$

Let $E_{\alpha, \gamma} := Q_\alpha - Q_\gamma = 2^n Q_\alpha - x_\gamma$, where x_γ denotes the center of Q_γ . Then, we can get

$$\sup_{x \in Q_\gamma} \int_{-1}^1 \int_{Q_\alpha} |g(\tau, x - y)| dy d\tau \leq \int_{-1}^1 \int_{E_{\alpha, \gamma}} |g(\tau, z)| dz d\tau,$$

and consequently (3.6) is bounded, with the help of the Minkowski inequality, by

$$\left(\sum_{\gamma \in \mathbb{Z}^n} \left[\sum_{\alpha \in \mathbb{Z}^n} \left(\sup_{t \in [0, 2]} \sup_{y \in Q_\alpha} \tilde{I}(t, |y|) \right) \int_{-1}^1 \int_{E_{\alpha, \gamma}} |g(\tau, z)| dz d\tau \right]^2 \right)^{1/2}$$

$$\begin{aligned}
&\leq \sum_{\alpha \in \mathbb{Z}^n} \left[\left(\sup_{t \in [0,2]} \sup_{y \in Q_\alpha} \tilde{I}(t, |y|) \right) \left(\sum_{\gamma \in \mathbb{Z}^n} \left(\int_{-1}^1 \int_{E_{\alpha,\gamma}} |g(\tau, z)| \, dz \, d\tau \right)^2 \right)^{1/2} \right] \\
&\leq C \sum_{\alpha \in \mathbb{Z}^n} \left[\left(\sup_{t \in [0,2]} \sup_{y \in Q_\alpha} \tilde{I}(t, |y|) \right) \left(\sum_{\gamma \in \mathbb{Z}^n} \left(\int_{-1}^1 \int_{Q_\gamma} |g(\tau, z)| \, dz \, d\tau \right)^2 \right)^{1/2} \right],
\end{aligned}$$

where we have used the fact that $E_{\alpha,\gamma}$ can be covered by a finite number (independent of α) of Q_γ in the last step.

From (3.4), we have the estimate

$$\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0,2]} \sup_{y \in Q_\alpha} \tilde{I}(t, |y|) = \sup_{t \in [0,2]} \int_{\mathbb{R}^n} \tilde{I}(t, |y|) \, dy \leq C 2^{(2n+1)k}.$$

Therefore, the proof is completed. \square

When we deal with the Cauchy problem (1.1)-(1.2) in the case $l = 2$, we shall use the l^1 -estimate of the maximal function $\sup_{[0,T]} |S(t)u_0(x)|$ as the following inequality.

Lemma 8 *We have the estimate*

$$\begin{aligned}
&\sum_{\alpha \in \mathbb{Z}^n} \sup_{[0,T]} \sup_{x \in Q_\alpha} |S(t)u_0(x)| \\
&\leq C(1+T)^{\lfloor \frac{n}{2} \rfloor + 2} \left(\|u_0\|_{n+3\lfloor \frac{n}{2} \rfloor + 7, 2} + \|u_0\|_{n+3\lfloor \frac{n}{2} \rfloor + 7, 2, 2\lfloor \frac{n}{2} \rfloor + 2} \right), \quad (3.7)
\end{aligned}$$

where $\|\cdot\|_{l,2,j}$ is defined as in (1.6).

Proof. Taking $t_0 \in [0, T]$ such that

$$|f(t_0)| \leq \frac{\int_0^T |f(t)| \, dt}{T},$$

we have for any t

$$f(t) = f(t_0) + f(t) - f(t_0) = f(t_0) + \int_{t_0}^t f'(s) \, ds.$$

Thus, we can get

$$\begin{aligned}
|f(t)| &\leq \frac{1}{T} \int_0^T |f(t)| \, dt + \int_{t_0}^t |f'(s)| \, ds \\
&\leq \frac{1}{T} \int_0^T |f(t)| \, dt + \int_0^T |\partial_t f(t)| \, dt,
\end{aligned}$$

namely,

$$\sup_{t \in [0, T]} |f(t)| \leq \frac{1}{T} \int_0^T |f(t)| dt + \int_0^T |\partial_t f(t)| dt. \quad (3.8)$$

Notice that

$$\|f\|_{L^1(\mathbb{R}^n)} \leq C \|f\|_{L^2(\mathbb{R}^n)} + C \left\| |x|^{\bar{n}} f \right\|_{L^2(\mathbb{R}^n)}, \quad (3.9)$$

where $\bar{n} > n/2$, and

$$\begin{aligned} x_j S(t) f &= S(t)(x_j f) + 4itS(t)(\partial_{x_j} \Delta f) - 2\varepsilon itS(t)(\partial_{x_j} f), \quad (3.10) \\ x_k x_j S(t) f &= ((-2\varepsilon it)^2 + 8it)S(t)(\partial_{x_k} \partial_{x_j} f) + 4itS(t)(\delta_{kj} \Delta f) - 2\varepsilon itS(t)(\delta_{kj} f) \\ &\quad + 16\varepsilon t^2 S(t)(\partial_{x_k} \partial_{x_j} \Delta f) + (4it)^2 S(t)(\partial_{x_k} \partial_{x_j} \Delta^2 f) \\ &\quad - 2\varepsilon itS(t)(x_k \partial_{x_j} f + x_j \partial_{x_k} f) + 4itS(t)(x_k \partial_{x_j} \Delta f + x_j \partial_{x_k} \Delta f) \\ &\quad + S(t)(x_k x_j f), \quad (3.11) \end{aligned}$$

where $\delta_{kj} = 1$ if $k = j$, $\delta_{kj} = 0$ if $k \neq j$.

We have, from (3.8)-(3.11), the Sobolev embedding theorem and the Fubini theorem, that

$$\begin{aligned} &\sum_{\alpha \in \mathbb{Z}^n} \sup_{[0, T]} \sup_{x \in Q_\alpha} |S(t)u_0(x)| \\ &\leq \sum_{\alpha \in \mathbb{Z}^n} \sup_{[0, T]} \left(\int_{Q_\alpha} |S(t)u_0(x)| dx + \sum_{|\beta| \leq n} \int_{Q_\alpha} |\partial_x^\beta S(t)u_0(x)| dx \right) \\ &\leq \sum_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \sup_{[0, T]} |S(t)u_0(x)| dx + \sum_{|\beta| \leq n} \int_{Q_\alpha} \sup_{[0, T]} |\partial_x^\beta S(t)u_0(x)| dx \right) \\ &\leq \sum_{\alpha \in \mathbb{Z}^n} \left\{ \frac{C}{T} \int_{Q_\alpha} \int_0^T |S(t)u_0(x)| dt dx + \int_{Q_\alpha} \int_0^T |\partial_t S(t)u_0(x)| dt dx \right. \\ &\quad \left. + \sum_{|\beta| \leq n} \int_{Q_\alpha} \left(\frac{C}{T} \int_0^T |S(t)\partial_x^\beta u_0(x)| dt + \int_0^T |\partial_t S(t)\partial_x^\beta u_0(x)| dt \right) dx \right\} \\ &\leq \frac{C}{T} \int_{\mathbb{R}^n} \int_0^T |S(t)u_0(x)| dt dx + \int_{\mathbb{R}^n} \int_0^T |S(t)\Delta u_0(x)| dt dx \\ &\quad + \int_{\mathbb{R}^n} \int_0^T |S(t)\Delta^2 u_0(x)| dt dx + \sum_{|\beta| \leq n} \frac{C}{T} \int_{\mathbb{R}^n} \int_0^T |S(t)\partial_x^\beta u_0(x)| dt dx \\ &\quad + \sum_{|\beta| \leq n} \int_{\mathbb{R}^n} \int_0^T |S(t)\partial_x^\beta \Delta u_0(x)| dt dx + \sum_{|\beta| \leq n} \int_{\mathbb{R}^n} \int_0^T |S(t)\partial_x^\beta \Delta^2 u_0(x)| dt dx \\ &\leq \frac{C}{T} \int_0^T \int_{\mathbb{R}^n} |S(t)u_0(x)| dx dt + \frac{C}{T} \sum_{|\beta| \leq n} \int_0^T \int_{\mathbb{R}^n} |S(t)\partial_x^\beta u_0(x)| dx dt \\ &\quad + C \sum_{|\beta| \leq n+4} \int_0^T \int_{\mathbb{R}^n} |S(t)\partial_x^\beta u_0(x)| dx dt \end{aligned}$$

$$\begin{aligned}
&\leq \frac{C}{T} \int_0^T [\|S(t)u_0\|_2 + \||x|^{\bar{n}} S(t)u_0\|_2] dt \\
&\quad + \frac{C}{T} \sum_{|\beta| \leq n} \int_0^T [\|S(t)\partial_x^\beta u_0\|_2 + \||x|^{\bar{n}} S(t)\partial_x^\beta u_0\|_2] dt \\
&\quad + C \sum_{|\beta| \leq n+4} \int_0^T [\|S(t)\partial_x^\beta u_0\|_2 + \||x|^{\bar{n}} S(t)\partial_x^\beta u_0\|_2] dt \\
&\leq C(1+T)^{\bar{n}+1} (\|u_0\|_{n+3\bar{n}+4, 2} + \|u_0\|_{n+3\bar{n}+4, 2, 2\bar{n}}),
\end{aligned}$$

which implies the desired result if we choose $\bar{n} = \left\lceil \frac{n}{2} \right\rceil + 1$. \square

4 The Wellposedness

We will give the proofs of the main Theorems in this section.

Proof of Theorem 1. Similar to the proof in [17, Theorem 4.1], we shall only consider the most interesting case $s = s_0$. The general case follows by combining this result with the fact that the highest derivatives involved in that proof always appear linearly and with some commutator estimates (see [14]) for the cases where $s \neq k + 1/2$, $k \in \mathbb{Z}^+$. For simplicity of the exposition, we shall assume that

$$P((\partial_x^\alpha u)_{|\alpha| \leq 2}, (\partial_x^\alpha \bar{u})_{|\alpha| \leq 2}) = \partial_{x_j}^2 u \partial_{x_k}^2 u \partial_{x_m}^2 u$$

For $u_0 \in H^{s_0}(\mathbb{R}^n)$, we denote by $u = \mathcal{F}(v) = \mathcal{F}_{u_0}(v)$ the solution of the linear inhomogeneous Cauchy problem

$$i\partial_t u = -\varepsilon \Delta u + \Delta^2 u + \partial_{x_j}^2 v \partial_{x_k}^2 v \partial_{x_m}^2 v, \quad t \in \mathbb{R}, x \in \mathbb{R}^n, \quad (4.1)$$

$$u(0, x) = u_0(x), \quad (4.2)$$

In order to construct \mathcal{F} being a contraction mapping in some space, we use the integral equation

$$u(t) = \mathcal{F}(v)(t) = S(t)u_0 - i \int_0^t S(t-\tau) \partial_{x_j}^2 v(\tau) \partial_{x_k}^2 v(\tau) \partial_{x_m}^2 v(\tau) d\tau. \quad (4.3)$$

We introduce the following work space

$$\begin{aligned}
Z_T^E = \left\{ w : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{C} : \sup_{t \in [0, T]} \|w(t)\|_{s_0, 2} \leq E; \right. \\
\left. \sum_{|\beta| = s_0 + 1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta w(t, x)|^2 dx dt \right)^{1/2} \leq T^\delta; \right.
\end{aligned}$$

$$\left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} |D_x^2 w(t, x)|^2 \right)^{1/2} \leq E \},$$

where $\delta < 1/3$ is a constant.

We notice, for any $\beta \in \mathbb{Z}^n$ with $|\beta| = s_0 - 3/2$, that

$$\begin{aligned} \partial_x^\beta (\partial_{x_j}^2 v \partial_{x_k}^2 v \partial_{x_m}^2 v) &= \partial_x^\beta \partial_{x_j}^2 v \partial_{x_k}^2 v \partial_{x_m}^2 v + \partial_{x_j}^2 v \partial_x^\beta \partial_{x_k}^2 v \partial_{x_m}^2 v + \partial_{x_j}^2 v \partial_{x_k}^2 v \partial_x^\beta \partial_{x_m}^2 v \\ &\quad + R((\partial_x^\gamma v)_{2 \leq |\gamma| \leq s_0 - 1/2}), \end{aligned}$$

where $|\beta'| = |\beta| - 1$ and $|\beta''| = 1$ with $\beta', \beta'' \in \mathbb{Z}^n$.

Thus, from the integral equation (4.3), (2.6), (2.1), the Sobolev embedding theorem and the commutator estimates [14], we can get, for $v \in Z_T^D$, that

$$\begin{aligned} & \sum_{|\beta|=s_0+1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta u(t, x)|^2 dx dt \right)^{1/2} \\ & \leq \sum_{|\beta|=s_0+1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |S(t) \partial_x^\beta u_0(x)|^2 dx dt \right)^{1/2} \\ & \quad + \sum_{|\beta|=s_0+1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} \left| \int_0^t S(t-\tau) \partial_x^\beta (\partial_{x_j}^2 v \partial_{x_k}^2 v \partial_{x_m}^2 v) \right|^2 dx dt \right)^{1/2} \\ & \leq CT^{\frac{1}{3}} \|u_0\|_{s_0, 2} + CT^{\frac{1}{4}} \sum_{|\beta|=s_0-3/2} \sum_{j, k, m=1}^n \sum_{\alpha \in \mathbb{Z}^n} \left\| \partial_x^\beta \partial_{x_j}^2 v \partial_{x_k}^2 v \partial_{x_m}^2 v \right\|_{L_x^2(Q_\alpha; L_t^2([0, T]))} \\ & \quad + C \int_0^T \left\| D_x^{\frac{1}{2}} R((\partial_x^\gamma v)_{2 \leq |\gamma| \leq s_0 - 1/2}) \right\|_2 dt \\ & \leq CT^{\frac{1}{3}} \|u_0\|_{s_0, 2} + CT^{\frac{1}{4}} \sum_{|\beta|=s_0+1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta v(t, x)|^2 dx dt \right)^{1/2} \\ & \quad \cdot \left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} |D_x^2 v(t, x)|^2 \right) + CT \sup_{t \in [0, T]} \|v\|_{s_0, 2}^3 \\ & \leq CT^{\frac{1}{3}} \|u_0\|_{s_0, 2} + CT^{\frac{1}{4}} T^\delta E^2 + CTE^3 \\ & \leq T^\delta, \end{aligned} \tag{4.4}$$

where we have taken T so small that

$$CT^{\frac{1}{3}-\delta} \|u_0\|_{s_0, 2} + CT^{\frac{1}{4}} E^2 + CT^{1-\delta} E^3 \leq 1, \tag{4.5}$$

in the last step.

By the Sobolev embedding theorem, (2.2) and the Hölder inequality, we

have

$$\begin{aligned}
\sup_{t \in [0, T]} \|u(t)\|_{s_0, 2} &\leq \|u_0\|_{s_0, 2} + \sup_{t \in [0, T]} \int_0^t \left\| S(t-\tau) \partial_{x_j}^2 v(\tau) \partial_{x_k}^2 v(\tau) \partial_{x_m}^2 v(\tau) \right\|_2 d\tau \\
&\quad + \sup_{t \in [0, T]} \left\| D_x^{\frac{3}{2}} \int_0^t S(t-\tau) D_x^{s_0 - \frac{3}{2}} \partial_{x_j}^2 v(\tau) \partial_{x_k}^2 v(\tau) \partial_{x_m}^2 v(\tau) d\tau \right\|_2 \\
&\leq \|u_0\|_{s_0, 2} + T \sup_{t \in [0, T]} \left\| \partial_{x_j}^2 v(t) \partial_{x_k}^2 v(t) \partial_{x_m}^2 v(t) \right\|_2 \\
&\quad + \sum_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_0^T \left| D_x^{s_0 - \frac{3}{2}} (\partial_{x_j}^2 v(t) \partial_{x_k}^2 v(t) \partial_{x_m}^2 v(t)) \right|^2 dt dx \right)^{1/2} \\
&\leq \|u_0\|_{s_0, 2} + T \sup_{t \in [0, T]} \|v\|_{\frac{3}{3}+2, 2}^3 \\
&\quad + \sum_{j, k, m=1}^n \sum_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_0^T \left| D_x^{s_0 - \frac{3}{2}} \partial_{x_j}^2 v(t) \partial_{x_k}^2 v(t) \partial_{x_m}^2 v(t) \right|^2 dt dx \right)^{1/2} \\
&\quad + \sum_{\alpha \in \mathbb{Z}^n} \left(\int_{Q_\alpha} \int_0^T \left| (D_x^\gamma v)_{2 \leq |\gamma| \leq s_0 - \frac{1}{2}} \right|^2 dt dx \right)^{1/2} \\
&\leq \|u_0\|_{s_0, 2} + T \sup_{t \in [0, T]} \|v\|_{\frac{3}{3}+2, 2}^3 \\
&\quad + \sum_{|\beta|=s_0+\frac{1}{2}} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta v|^2 dx dt \right)^{1/2} \cdot \sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} |D_x^2 v|^2 \\
&\quad + T^{\frac{1}{2}} \sup_{t \in [0, T]} \|v\|_{s_0, 2}^3 \\
&\leq \|u_0\|_{s_0, 2} + (T + T^{\frac{1}{2}}) E^3 + T^\delta E^2 \\
&\leq E, \tag{4.6}
\end{aligned}$$

where in the last step, we have chosen T small enough such that

$$(T + T^{\frac{1}{2}}) E^2 + T^\delta E \leq \frac{1}{2}. \tag{4.7}$$

Similar to the derivation of (4.4) and (4.6), we obtain, by inserting (3.2) in (4.3), that

$$\begin{aligned}
&\left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} |D_x^2 u(t, x)|^2 \right)^{1/2} \\
&\leq C \|u_0\|_{s_0, 2} + C \left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} \left| \int_0^t S(t-\tau) D_x^2 \partial_{x_j}^2 v(\tau) \partial_{x_k}^2 v(\tau) \partial_{x_m}^2 v(\tau) \right|^2 \right)^{1/2} \\
&\leq C \|u_0\|_{s_0, 2} + CT \sup_{t \in [0, T]} \|v\|_{s_0, 2}^3 \\
&\leq \|u_0\|_{s_0, 2} + CTE^3 \\
&\leq E, \tag{4.8}
\end{aligned}$$

where we have taken

$$E = 2C \|u_0\|_{s_0,2}, \quad (4.9)$$

and T sufficiently small such that

$$CTE^2 \leq 1. \quad (4.10)$$

Therefore, choosing an E as in (4.9) and then taking a T sufficiently small such that (4.5), (4.7) and (4.10) hold, we obtain that the mapping

$$\mathcal{F} = \mathcal{F}_{u_0} : Z_T^E \rightarrow Z_T^E$$

is well defined.

For convenience, we denote

$$\Lambda_T(w) = \max \left\{ ET^{-\delta} \sum_{|\beta|=s_0+1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta w(t,x)|^2 dx dt \right)^{1/2}; \right. \\ \left. \sup_{t \in [0,T]} \|w(t)\|_{s_0,2}; \left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0,T]} \sup_{x \in Q_\alpha} |D_x^2 w(t,x)|^2 \right)^{1/2} \right\}.$$

To show that \mathcal{F} is a contraction mapping, we apply the estimates obtained in (4.4), (4.6) and (4.8) to the following integral equation

$$\mathcal{F}v(t) - \mathcal{F}w(t) = \int_0^t S(t-\tau) \left[\partial_{x_j}^2 v(\tau) \partial_{x_k}^2 v(\tau) \partial_{x_m}^2 v(\tau) \right. \\ \left. - \partial_{x_j}^2 w(\tau) \partial_{x_k}^2 w(\tau) \partial_{x_m}^2 w(\tau) \right](\tau) d\tau,$$

and obtain, for $v, w \in Z_T^E$, that

$$\Lambda_T(\mathcal{F}v - \mathcal{F}w) \leq CT^\delta \Lambda_T(v-w) \cdot [\Lambda_T^2(v) + \Lambda_T^2(w)] \\ \leq 2CT^\delta E^2 \Lambda_T(v-w).$$

where the constant C depends only on the form of $P(\cdot)$ and the linear estimates (2.1), (2.3), (2.9) and (3.1).

Thus, we can choose $0 < T \ll 1$ satisfying (4.5), (4.7), (4.10) and

$$2CT^\delta E^2 \leq 1/2. \quad (4.11)$$

Therefore, for those T , satisfying (4.5), (4.7), (4.10) and (4.11), the mapping \mathcal{F}_{u_0} is a contraction mapping in Z_T^E . Consequently, by the Banach contraction mapping principle, there exists a unique function $u \in Z_T^E$ such that $\mathcal{F}_{u_0}u = u$ which solves the Cauchy problem.

By the method given in [17, Theorem 4.1], we can prove the persistence property of $u(t)$ in H^{s_0} , *i.e.*

$$u(t, x) \in C([0, T]; H^{s_0}(\mathbb{R}^n)),$$

the uniqueness and the continuous dependence on the initial data of solution. For simplicity, we omit the rest of the proof. \square

Proof of Theorem 2. For simplicity, we assume

$$P((\partial_x^\alpha u)_{|\alpha| \leq 2}, (\partial_x^\alpha \bar{u})_{|\alpha| \leq 2}) = |\Delta u|^2.$$

It will be clear, from the argument presented below, that this does not represent any loss of generality. And as in the proof of Theorem 1, we consider the case $s = s_0 = 2n + 3 \lfloor \frac{n}{2} \rfloor + 15 + \frac{1}{2}$.

We introduce the following work space

$$\begin{aligned} X_T^E = \left\{ w : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{C} : \sup_t \|w(t)\|_{s_0, 2} \leq E; \sup_t \|w(t)\|_{n + \lfloor \frac{n}{2} \rfloor + 8, 2, 2 \lfloor \frac{n}{2} \rfloor + 2} \leq E; \right. \\ \left. \sum_{|\beta| = s_0 + 1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta w(t, x)|^2 dx dt \right)^{1/2} \leq T^\delta; \right. \\ \left. (1 + T)^{-\lfloor \frac{n}{2} \rfloor - 2} \left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} |D_x^2 w(t, x)|^2 \right)^{1/2} \leq E \right\}, \end{aligned}$$

where $0 < \delta < 1/4$ is a constant and the norm is defined as

$$\begin{aligned} \|v(t)\|_{X_T^E} = \max \left\{ \sup_t \|v(t)\|_{\frac{23}{2}}; \sup_t \|w(t)\|_{n + \lfloor \frac{n}{2} \rfloor + 8, 2, 2 \lfloor \frac{n}{2} \rfloor + 2}; \right. \\ \left. ET^{-\delta} \sum_{|\beta| = s_0 + 1/2} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta w(t, x)|^2 dx dt \right)^{1/2}; \right. \\ \left. (1 + T)^{-\lfloor \frac{n}{2} \rfloor - 2} \left(\sum_{\alpha \in \mathbb{Z}^n} \sup_{t \in [0, T]} \sup_{x \in Q_\alpha} |D_x^2 w(t, x)|^2 \right)^{1/2} \right\}. \end{aligned}$$

For $u_0 \in H^{s_0}(\mathbb{R}^n) \cap H^{n+4 \lfloor \frac{n}{2} \rfloor + 8}(\mathbb{R}^n; |x|^{2 \lfloor \frac{n}{2} \rfloor + 2} dx)$ and $v \in X_T^E$, we denote by $u = \mathcal{T}(v) = \mathcal{T}_{u_0}(v)$ the solution of the linear inhomogeneous Cauchy problem

$$i\partial_t u = -\varepsilon \Delta u + \Delta^2 u + |\Delta v|^2, \quad (4.12)$$

$$u(0, x) = u_0(x), \quad (4.13)$$

and consider the corresponding integral equation

$$u(t) = \mathcal{T}(v)(t) = S(t)u_0 - i \int_0^t S(t - \tau) |\Delta v|^2 d\tau. \quad (4.14)$$

We notice that

$$\partial_x^\beta (\Delta v \Delta \bar{v}) = \partial_x^\beta \Delta v \Delta \bar{v} + \Delta v \partial_x^\beta \Delta \bar{v} + R_0 \left((\partial_x^{\gamma_1} v \partial_x^{\gamma_2} v)_{|\gamma_1|, |\gamma_2| \leq s_0 - \frac{1}{2}} \right),$$

where $\beta, \gamma_1, \gamma_2 \in \mathbb{Z}^n$.

From the integral equation (4.14), (2.6), the Sobolev embedding theorem and (2.9), we can get, as in (4.4), that

$$\begin{aligned} & \sum_{|\beta|=s_0+\frac{1}{2}} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta u(t, x)|^2 dx dt \right)^{1/2} \\ & \leq \sum_{|\beta|=s_0+\frac{1}{2}} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta S(t) u_0(x)|^2 dx dt \right)^{1/2} \\ & \quad + \sum_{|\beta|=s_0+\frac{1}{2}} \sup_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} \left| \int_0^t S(t-\tau) \partial_x^\beta |\Delta v(\tau)|^2 d\tau \right|^2 dx dt \right)^{1/2} \\ & \leq CT^{\frac{1}{3}} \|u_0\|_{s_0,2} + CT^{\frac{1}{4}} \sum_{|\beta|=s_0-\frac{3}{2}} \sum_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta \Delta v \Delta \bar{v}|^2 dx dt \right)^{1/2} \\ & \quad + C \int_0^T \left\| D_x^{\frac{1}{2}} R_0 \right\|_2 dt \\ & \leq CT^{\frac{1}{3}} \|u_0\|_{s_0,2} + CT^{\frac{1}{4}} \sum_{|\beta|=s_0+\frac{1}{2}} \sum_{\alpha \in \mathbb{Z}^n} \left(\int_0^T \int_{Q_\alpha} |\partial_x^\beta v|^2 dx dt \right)^{1/2} \\ & \quad \cdot \left(\sum_{\alpha \in \mathbb{Z}^n} \sup_t \sup_{x \in Q_\alpha} |D_x^2 v| \right) + CT \sup_t \|v\|_{s_0,2}^2 \\ & \leq CT^{\frac{1}{3}} \|u_0\|_{s_0,2} + CT^{\frac{1}{4}} T^\delta E + CTE^2 \\ & \leq T^\delta, \end{aligned} \tag{4.15}$$

if we take T sufficiently small such that

$$CT^{\frac{1}{3}-\delta} \|u_0\|_{s_0,2} + CT^{\frac{1}{4}} E + CT^{1-\delta} E^2 \leq 1. \tag{4.16}$$

We can rewrite (4.14) as

$$u(t) = S(t) \left(u_0 - i \int_0^t S(-\tau) |\Delta v(\tau)|^2 d\tau \right). \tag{4.17}$$

From (3.7), we have

$$\begin{aligned} & (1+T)^{-[\frac{n}{2}]-2} \sum_{\alpha \in \mathbb{Z}^n} \sup_t \sup_{x \in Q_\alpha} |D_x^2 u(t, x)| \\ & \leq C (\|u_0\|_{n+3[\frac{n}{2}]+9,2} + \|u_0\|_{n+3[\frac{n}{2}]+9,2,2[\frac{n}{2}]+2}) \end{aligned}$$

$$+ \sum_{\alpha \in \mathbb{Z}^n} \sup_t \sup_{x \in Q_\alpha} \left| D_x^2 S(t) \int_0^t S(-\tau) |\Delta v(\tau)|^2 d\tau \right|,$$

where the second term in the right hand side can be bounded by

$$\begin{aligned} & C \sup_t \left\| \int_0^t S(-\tau) |\Delta v|^2 d\tau \right\|_{n+3[\frac{n}{2}]+9,2} + \sup_t \left\| \int_0^t S(-\tau) |\Delta v|^2 d\tau \right\|_{n+3[\frac{n}{2}]+9,2,2[\frac{n}{2}]+2} \\ & \leq C \int_0^T \left\| |\Delta v|^2 \right\|_{n+3[\frac{n}{2}]+9,2} d\tau + C \int_0^T \left\| S(-\tau) |\Delta v|^2 \right\|_{n+3[\frac{n}{2}]+9,2,2[\frac{n}{2}]+2} d\tau \\ & \leq CT \sup_t \|v\|_{n+3[\frac{n}{2}]+11,2}^2 + CT \sup_t \sum_{|\gamma| \leq n+3[\frac{n}{2}]+9} \left\| |x|^{[\frac{n}{2}]+1} S(-\tau) \partial_x^\gamma |\Delta v|^2 \right\|_2 \\ & \leq CT \sup_t \|v\|_{n+3[\frac{n}{2}]+11,2}^2 + CT \left\{ \sup_t \sum_{|\gamma| \leq n+3[\frac{n}{2}]+9} \left\| |x|^{[\frac{n}{2}]+1} \partial_x^\gamma |\Delta v|^2 \right\|_2 \right. \\ & \quad \left. + \sum_{|\gamma| \leq n+6[\frac{n}{2}]+12} t^{[\frac{n}{2}]+1} \left\| \partial_x^\gamma |\Delta v|^2 \right\|_2 + C \sum_{\substack{|\beta|+|\gamma| \leq n+6[\frac{n}{2}]+10 \\ 1 \leq |\beta| \leq [\frac{n}{2}]} t^{|\beta|} \left\| x^\beta \partial_x^\gamma |\Delta v|^2 \right\|_2 \right\} \\ & \leq CT \sup_t \|v\|_{n+3[\frac{n}{2}]+11,2}^2 + CT \sup_t \sum_{|\gamma| \leq n+3[\frac{n}{2}]+9} \sum_{\substack{|\gamma_1|+|\gamma_2|=|\gamma|+4 \\ |\gamma_1| \leq |\gamma_2|}} \left\| \partial_x^{\gamma_2} v \right\|_\infty \left\| |x|^{[\frac{n}{2}]+1} \partial_x^{\gamma_1} v \right\|_2 \\ & \quad + CT^{[\frac{n}{2}]+2} \sup_t \|v\|_{n+6[\frac{n}{2}]+14,2} \\ & \quad + \sum_{\substack{|\beta|+|\gamma| \leq n+4[\frac{n}{2}]+12 \\ 1 \leq |\beta| \leq [\frac{n}{2}]} T^{|\beta|+1} \sup_t \sum_{\substack{|\gamma_1|+|\gamma_2|=|\gamma|+4 \\ |\gamma_1| \leq |\gamma_2|}} \left\| \partial_x^{\gamma_2} v \right\|_\infty \left\| x^\beta \partial_x^{\gamma_1} v \right\|_2 \\ & \leq CT \sup_t \|v\|_{n+3[\frac{n}{2}]+11,2}^2 + CT \sup_t \|v\|_{n+4[\frac{n}{2}]+14,2} \sup_t \|v\|_{n+[\frac{n}{4}]+7,2,2[\frac{n}{2}]+2} \\ & \quad + CT^{[\frac{n}{2}]+2} \sup_t \|v\|_{n+6[\frac{n}{2}]+14,2} \\ & \quad + C \sum_{1 \leq |\beta| \leq [\frac{n}{2}]} T^{|\beta|+1} \sup_t \|v\|_{\frac{3n}{2}+4[\frac{n}{2}]+15+\frac{1}{2},2} \sup_t \|v\|_{n+[\frac{n}{2}]+8,2,2[\frac{n}{2}]+2} \\ & \leq CT \sup_t \|v\|_{s_0,2}^2 + CT \sup_t \|v\|_{s_0,2} \sup_t \|v\|_{n+[\frac{n}{2}]+8,2,2[\frac{n}{2}]+2} + CT^{[\frac{n}{2}]+2} \sup_t \|v\|_{s_0,2}^2 \\ & \quad + CT(T + T^{[\frac{n}{2}]}) \sup_t \|v\|_{s_0,2} \sup_t \|v\|_{n+[\frac{n}{2}]+8,2,2[\frac{n}{2}]+2} \\ & \leq CT(1 + T^{[\frac{n}{2}]+1})E^2 \\ & \leq \frac{1}{2}E, \end{aligned}$$

if we take $E \geq 2C(\|u_0\|_{n+3[\frac{n}{2}]+9,2} + \|u_0\|_{n+4[\frac{n}{2}]+8,2,2[\frac{n}{2}]+2})$ and T so small that

$$CT(1 + T^{[\frac{n}{2}]+1})E \leq \frac{1}{2}. \quad (4.18)$$

Thus, we obtain

$$(1 + T)^{-[\frac{n}{2}] - 2} \sum_{\alpha \in \mathbb{Z}^n} \sup_t \sup_{x \in Q_\alpha} |D_x^2 u(t, x)| \leq E. \quad (4.19)$$

Similar to (4.19), we have

$$\begin{aligned} & \sup_t \|u(t)\|_{n+[\frac{n}{2}]+8, 2, 2[\frac{n}{2}]+2} = \sup_t \sum_{|\gamma| \leq n+[\frac{n}{2}]+8} \left\| |x|^{[\frac{n}{2}]+1} \partial_x^\gamma u(t) \right\|_2 \\ & \leq \sum_{|\gamma| \leq n+[\frac{n}{2}]+8} \left[\sup_t \left\| |x|^{[\frac{n}{2}]+1} S(t) \partial_x^\gamma u_0 \right\|_2 + \sup_t \left\| \int_0^t |x|^{[\frac{n}{2}]+1} S(t-\tau) \partial_x^\gamma |\Delta v|^2 d\tau \right\|_2 \right] \\ & \leq \sup_t \sum_{|\gamma| \leq n+[\frac{n}{2}]+8} \left\| |x|^{[\frac{n}{2}]+1} \partial_x^\gamma u_0 \right\|_2 + C \sup_t \sum_{|\gamma| \leq n+4[\frac{n}{2}]+11} t^{[\frac{n}{2}]+1} \|\partial_x^\gamma u_0\|_2 \\ & \quad + C \sup_t \sum_{\substack{|\beta|+|\gamma| \leq n+4[\frac{n}{2}]+9 \\ 1 \leq |\beta| \leq [\frac{n}{2}]} } t^{|\beta|} \|x^\beta \partial_x^\gamma u_0\|_2 + \sup_t \int_0^t \sum_{|\gamma| \leq n+[\frac{n}{2}]+8} \left\| |x|^{[\frac{n}{2}]+1} \partial_x^\gamma |\Delta v|^2 \right\|_2 \\ & \quad + C \sup_t \sum_{|\gamma| \leq n+4[\frac{n}{2}]+11} t^{[\frac{n}{2}]+1} \|\partial_x^\gamma |\Delta v|^2\|_2 + C \sup_t \sum_{\substack{|\beta|+|\gamma| \leq n+4[\frac{n}{2}]+9 \\ 1 \leq |\beta| \leq [\frac{n}{2}]} } t^{|\beta|} \|x^\beta \partial_x^\gamma |\Delta v|^2\|_2 \\ & \leq C(1 + T)^{[\frac{n}{2}]+1} (\|u_0\|_{n+4[\frac{n}{2}]+8, 2, 2[\frac{n}{2}]+2} + \|u_0\|_{s_0, 2}) \\ & \quad + T \sup_t \|v\|_{s_0, 2} \|v\|_{n+[\frac{n}{2}]+8, 2, 2[\frac{n}{2}]+2} + CT^{[\frac{n}{2}]+1} \|v\|_{s_0, 2}^2 \\ & \leq E. \end{aligned} \quad (4.20)$$

As in the derivation of (4.6), we can get

$$\sup_t \|u(t)\|_{s_0, 2} \leq E. \quad (4.21)$$

The estimates (4.15), (4.19), (4.20) and (4.21) yield $u \in X_T^E$. Thus, we can fix E and T as above such that \mathcal{S} is a contraction mapping from X_T^E to itself. By the standard argument used in the proof of Theorem 1, we can complete the proof for which we omit the details. \square

Acknowledgment

The authors would like to thank the referees for valuable comments and suggestions for the original manuscript.

References

- [1] M. Ben-Artzi, H. Koch and J. C. Saut, Dispersion estimates for fourth order Schrödinger equations, *C. R. Acad. Sci. Paris Sér. I Math.*, 330 (2) (2000) 87–92.
- [2] J. Bergh and J. Löfström, *Interpolation Spaces, An Introduction*, Springer-Verlag, 1976.
- [3] T. Cazenave, An introduction to nonlinear Schrödinger equations, *Textos de Métodos Matemáticos*, 26, Universidade Federal do Rio de Janeiro.
- [4] H. Chihara, Local existence for semilinear Schrödinger equations, *Math. Japonica*, 42 (1995) 35–52.
- [5] P. Constantin and J. C. Saut, Local smoothing properties of dispersive equations, *J. Amer. Math. Soc.*, 1 (1988) 413–446.
- [6] K. B. Dysthe, Note on a modification to the nonlinear Schrödinger equation for application to deep water waves, *Proc. R. Soc. Lond. A*369 (1979) 105–114.
- [7] J. Ginibre and Y. Tsutsumi, Uniqueness for the generalized Korteweg-de Vries equations, *SIAM J. Math. Anal.*, 20 (1989) 1388–1425.
- [8] B. L. Guo and B. X. Wang, The global Cauchy problem and scattering of solutions for nonlinear Schrödinger equations in H^s . *Diff. Integral Eqs.*, 15 (9) (2002) 1073–1083.
- [9] C. C. Hao, L. Hsiao and B. X. Wang, Wellposedness for the fourth order nonlinear Schrödinger equations, *J. Math. Anal. Appl.*, 320(1) (2005) 246–265.
- [10] N. Hayashi and T. Ozawa, Remarks on nonlinear Schrödinger equations in one space dimension, *Diff. Integral Eqs.*, 2 (1994) 453–461.
- [11] Z. H. Huo and Y. L. Jia, The Cauchy problem for the fourth-order nonlinear Schrödinger equation related to the vortex filament, *J. Differential Equations* 214 (2005) 1–35.
- [12] V. I. Karpman, Stabilization of soliton instabilities by higher-order dispersion: fourth order nonlinear Schrödinger-type equations, *Phys. Rev. E*, 53 (2) (1996) R1336–R1339.
- [13] T. Kato, On the Cauchy problem for the (generalized) Korteweg-de Vries equation, *Advances in Math. Supp. Studies, Studies in Applied Math.*, 8 (1983) 93–128.
- [14] T. Kato and G. Ponce, Commutator estimates and the Euler and Navier-Stokes equations, *Comm. Pure Appl. Math.*, 41 (1988) 891–907.
- [15] C. E. Kenig, G. Ponce and L. Vega, Oscillatory integrals and regularity of dispersive equations, *Indiana University Math. J.*, 40 (1991) 33–69.
- [16] C. E. Kenig, G. Ponce and L. Vega, Well-posedness of the initial value problem for the Korteweg-de Vries equation, *J. Amer. Math. Soc.*, 4 (1991) 323–347.

- [17] C. E. Kenig, G. Ponce and L. Vega, Small solutions to nonlinear Schrödinger equations, *Ann. Inst. Henri Poincaré, Nonlinear Anal.*, 10 (1993) 255–288.
- [18] C. E. Kenig, G. Ponce and L. Vega, Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations, *Invent. Math.* 134 (1998) 489–545.
- [19] C. E. Kenig, G. Ponce and L. Vega, The Cauchy problem for quasi-linear Schrödinger equations. *Invent. Math.* 158 (2004) 343–388.
- [20] S. N. Kruzhkov, A. V. Faminskiĭ, Generalized solutions of the Cauchy problem for the Korteweg-de Vries equation, *Math. USSR Sbornik* 48 (1984), 93–138.
- [21] H. Pecher and W. von Wahl, Time dependent nonlinear Schrödinger equations. *Manuscripta Math.*, 27(2) (1979) 125–157.
- [22] J. Segata, Well-posedness for the fourth order nonlinear Schrödinger type equation related to the vortex filament, *Diff. Integral Eqs.*, 16 (7) (2003) 841–864.
- [23] J. Segata, Remark on well-posedness for the fourth order nonlinear Schrödinger type equation, *Proc. Amer. Math. Soc.*, 132 (2004) 3559–3568.
- [24] P. Sjölin, Regularity of solutions to the Schrödinger equation, *Duke Math. J.*, 55 (1987) 699–715.
- [25] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis in Eucliden Spaces*, Princeton University Press, 1971.
- [26] L. Vega, Schrödinger equations: pointwise convergence to the initial data. *Proc. Amer. Math. Soc.* 102 (1988) 874–878.