

THEORETICAL AND EXPERIMENTAL EVIDENCES OF MATERIAL PHASE CAUSALITY. DYNAMICS-STATISTICAL INTERPRETATION OF THE WAVE FUNCTION.

I. G. Koprinkov

Department of Applied Physics, Technical University of Sofia, 1756 Sofia, Bulgaria

e-mail: igk@tu-sofia.bg

Abstract

Theoretical and experimental evidences of causal relation of the phase of the wave function with the dynamics of the quantum system are presented. Dynamics-statistical interpretation of the quantum mechanics is proposed. A particle-wave duality picture incorporated in the wave function through its phase and amplitude is considered.

PACS: 03.65.-w ; 03.65.Ta ; 42.50.Ct

Keywords: Quantum mechanics; State vector; Material phase; Physical meaning; Interpretation

I. Introduction

The physical state in the quantum mechanics is described not by a definite number of dynamical variables, *e.g.*, coordinates, linear momentum, etc., as in the classical mechanics, but by an abstract quantity – the state vector, or the wave function in a given representation, which obeys the Schrödinger equation [1]. Although the wave function describes the state of localized physical objects, *i.e.*, particle or system of particles, and the Hamiltonian of the system is written usually tacitly assuming point-like particles, the wave function itself, and, thus, the quantum state, is not localized, in general, but is distributed in space as for the continuous objects, *i.e.*, the waves. The wave-like nature of such point-like particle dynamics is further enforced by the validity of the quantum superposition principle and the quantum mechanical interference due to the linearity of the quantum evolution equation. Two ways of evolution of the wave function are known. If no measurement takes place, the wave function is subject to continuous, local, unitary (assuming Hermitian Hamiltonian), causal evolution ruled by the Schrödinger equation. The quantum state may evolve into a linear/coherent superposition of eigenstates of given dynamical observable and may reveal various wave-like features. During the measurement, the wave function undergoes a sudden, discontinuous, non-local, probabilistic collapse, von Neumann's *process 1* [2], onto some of the eigenstates of the observable being measured selected by the measuring apparatus (or the environment), a process called decoherence. Such a collapse, in fact, is a prerequisite to consider given process as a measurement. Although the collapse “shrinks” the state, it still remains distributed in space in view of the non-locality of the eigenstates of the quantum system. Thus, the particle-wave duality is built in the very nature of the quantum mechanics. The particle-wave duality and the nonlocality of the quantum phenomena are clearly demonstrated by the Feynman's “which-way” interference type experiments [3], Einstein-Podolsky-Rosen (EPR) type correlations [4], etc. The inherent duality of the quantum mechanical description predetermines the consequent problems with the interpretation of some results, sometimes considered as paradoxes. The description of the state of a localized physical object by a substantially non-local quantity, the wave function, does not have an analog within the classical physics. That is why, the wave function does not possess an obvious physical meaning that can be determined, *e.g.*, applying the correspondence principle between the classical and the quantum physics, and its relation with the physical reality is subject to additional considerations. Thus, the question of the relation between the wave function and the physical reality, and, on that ground – its physical meaning, becomes a fundamental problem in the quantum mechanics.

Since Einstein, in his famous debate with Bohr, has not succeed to present convincing arguments that “*God does not play dice*”, the so called standard, or Copenhagen interpretation of quantum mechanics has been widely accepted. It defeats the early proposed ontological point of view, according to which the wave function represents an objectively existing wave, *e.g.*, the de Broglie's pilot-wave associated to the particle. The ontological meaning was also supported by Schrödinger when he introduced the wave function. The Copenhagen interpretation encompasses several principles and conventions. Among these is the convention of epistemological meaning of the wave function, Heisenberg's uncertainty principle, Bohr's complementary principle, etc. Within the Copenhagen interpretation, the wave function is only a tool in the determination of the probability to find a quantum system in given space position at given instant of time. In some aspect, the wave function is considered as the observer's knowledge (epistemology) about the state of the physical object.

Beside of the Copenhagen interpretation, a number of other interpretations have been proposed. Some of them involve substantial additional concepts and should be considered as new formulation rather than interpretation of the original quantum mechanics. Among the most famous non-Copenhagen interpretations/formulations of the quantum mechanics are: de Broglie-Bohm ontological interpretation [5], Feynman's path integral formulation [6], consistent histories interpretation [7], many-worlds interpretation [8], environment-induced superselection [9], etc. In the ontological de Broglie-Bohm interpretation, one assumes that not only the particle (the primary object of the quantum mechanics) but also its state, the wave-function, represents a real physical field. The evolution of that field is described by the Schrödinger equation, while the evolution of the particle position (considered as a *hidden variable* in Bohm's theory) and its trajectory is ruled by the physical action. An alternative approach to the quantum trajectories is developed based on the Feynman's path integral. In the many-worlds interpretation, the postulate of the instantaneous collapse of

the wave function at the process of measurement, that takes place in the Copenhagen interpretation, is replaced by the decoherence of the quantum state at its interaction with the environment. At the same time, an assumption of existence of many parallel worlds is introduced so that every possible outcome of the experiment is realized in some of these worlds. The consistent histories interpretation is based on the concept of finite ordered sequence of events taken at successive times, called history. The criterion of consistency allows selecting histories that have physical meaning. The consistent history is used in combination with the quantum decoherence. A complementary view to the quantum decoherence is the environment-induced superselection. It originates from the continuous decoherence between the states of the quantum system caused by the environment. This selects a small set of substates, pointer states, from the Hilbert space, which remain stable under further interaction with the environment. The above interpretations introduce some important concepts in quantum mechanics but their application still remains limited.

The Copenhagen interpretation confers physical, probabilistic [10], meaning to the amplitude of the wave function, only. In contrast to the amplitude, the role of the phase of the wave function has been strongly neglected. Thus, for example, the time dependent phase, which appears in a factor of unit modulus, $\exp(-i\lambda t/\hbar)$, is considered, in principle, as unobservable [2], or, else, any true constant phase factor $\exp(i\alpha)$ is considered as irrelevant to any physical results [11]. Such an irrelevance of the phase of the wave function to the physical reality is subject to convention and, to the best of our knowledge, it has never been investigated and proved in an especially designed experiment prior the acceptance of the Copenhagen interpretation. In fact, the Copenhagen interpretation has been formulated based on limited experimental evidences and knowledge about the quantum phenomena, obtained by rather simple apparatuses. A number of *gedanken* experiments [3, 4] have been proposed and exploited to analyze the behavior of the quantum objects that has not been directly accessible to experimental verification at that time. Nowadays, very refined laboratory equipment and experimental methods exist that offer an unprecedented precision in the performance of the experiments. This allows testing experimentally some of the foundations of the quantum mechanics, as well as to realize some of the *gedanken* experiments [12-15]. Analyzing some experiments, one may clearly distinguish physically observable appearances of the material phase (MP) [16-19]. Some theoretical considerations and critical inspection of experiments with material wave-packets reveal that the relation of the wave function with the physical reality is not restricted to the amplitude only, but its phase is also strongly involved [20-22]. Here we present systematic theoretical and experimental evidences showing that the MP is causally related with the dynamics of the quantum system that may lead to observable physical consequences. All considerations are done within the frame of the quantum mechanics. The conclusions found here may serve as a base of more consistent, to our opinion, interpretation of the quantum mechanics, and, in particular, the elements of the wave function - phase and amplitude.

II. Preliminaries and definitions

In the beginning, we will specify some terminological questions and definitions.

First, considering the MP, we have in mind *dynamical* part of the MP, geometric, or Berry phase [23] is not considered here.

Second, the initial (constant) and the time dependent parts of the MP are considered together at an equivalent ground because of a number of arguments [20]: *i*) they have equivalent appearance in the initial and subsequent mathematical expressions, *ii*) they both behave as constants within the Hilbert space, whose vectors are functions on the configuration space, only, *iii*) changing the phase even by a constant value has a dramatic effect on the interference phenomena with material wave packets and it is accessible to experimental observations [16-19], *iv*) the initial phase, if retained in the wave function, shows sensible physical behavior, see Sec. III.1, *v*) the initial phase represents actually the time-dependent phase acquired by the quantum system during its preceding history up to the moment of creation of the state whose dynamical behavior is studied. The time of existence of the created state should be counted namely from this moments, considered as initial moment of the state. The exact determination of that moment is subject of definition because a transient time exists between the preceding and the subsequently created states. The preceding history and the time of creation of given state are not under concern in the usual treatment of the quantum mechanical problems, partially because, until recently, such questions were beyond the reach of the experimental method. The present achievements in the ultrafast laser technology allow tracing the internal dynamics of the quantum systems. These include the nuclear motion inside the molecules in the femtosecond time scale [17-19], and the electronic motion inside the atoms, in the pico/femtosecond time scale (for some highly excited Rydberg atomic states) [16]. Since recently, tracing the attosecond time dynamics seems feasible [24-26]. Hence, the ability to trace the atomic and molecular transients gives sense to the question of the time of formation of given quantum state, and, on that ground, the initial phase of the state.

Third, not only the relative (phase difference) but the absolute value, too, of the MP is considered to be important. They can be done equal depending on the choice of the reference phase. Knowledge of the absolute value of the MP refers the problem to the absolute internal clocking of the state of the quantum system. Since this problem seems intractable using ordinary laboratory equipment, while not meaningless for the advanced experimental methods, it is usually considered the relative phase, or the phase difference, if, by exception, the MP is discussed at all [2, 27, 28]. Here, we have in mind the absolute value of the MP while switching to the relative phase is trivial.

The MP causality will be considered as a particular case of a more general understanding of causality that will be introduced here. To our understanding, the causality is related with the existence of *fundamental physical relations* between *the objects of the physical reality*, particles, fields, and hierarchic constructs of them. The relations between the objects of the physical reality impose definite relations between the *elements of the physical reality* and the corresponding to them *physical quantities*. According to Einstein's point of view, "If, without in any way disturbing a system, we can

predict with certainty (*i.e.*, with probability equal to unity) the value of the physical quantity, then there exists an element of physical reality corresponding to this physical quantity” [4]. Undoubtedly, this is the ideal for most of the physicists but its applicability is disputable from what is known from the present state of the theory and the experiment. In particular, it invokes the problem of the type of causality and determinism in the quantum mechanics [29]. In this work, the elements of the physical reality will be recognized based on the fundamental relations that exist between the objects of the physical reality. Such relations are undisputable and exist irrespectively on the way of description of the physical reality. The contemporary physics reveals the existence of two different kinds of relations between the physical objects. The first one is based on the usual fundamental interactions. The second one involves EPR type correlations, and, according to Einstein, can be considered as a “spooky interaction at a distance”. It may cause relation between the physical objects that are separated by a space-like interval, in sharp contrast with the requirement of local realism [4]. The EPR type correlations appear proved experimentally [12, 13] while their physical mechanism remains unclear. According to Einstein’s understanding of a complete theory [4], “every element of the physical reality must have a counterpart in the physical theory”. This again reflects the ideal of the orthodoxly thinking physicists, but it is hardly to believe that all elements of the physical reality sometime will be known. In fact, we do not know at present all elements of the physical reality. That is why, the accent will be put here on the physical quantities rather than on the related, sometimes abstract, elements of the physical reality. The physical quantities and their relationships are so important that they, in fact, constitute the essence of our knowledge about the physical objects and the physical phenomena. The meaningful physical behavior of given physical quantity is an indication of its relation with the physical reality. Sometimes, namely a particular behavior of given quantity (*e.g.*, conservation of momentum or energy) is a base of discovery of new physical objects. The following *definition of physical reality - physical quantity* relationship will be used here [20]: *If we find that a change (direct or indirect) of given quantity affects at least one other quantity, subject of direct or indirect observation/measurement, then the quantity under consideration is causally related with the physical reality. The quantity that is causally related with the physical reality will be called physical quantity, and (if necessary, Einstein’s extension [4] can be associated at this point) “...there exists an element of physical reality corresponding to this physical quantity”. On the same ground, the affected quantity is, of course, also a physical quantity.* Object/element of the physical reality (if one exists at all) that is not related to any other object/element of the physical reality is, in principle, unobservable and it seems meaningless to speak about it. Such kind of causality can be relatively easy and unambiguously established and this, to our opinion, is the most important and general problem that should be solved about given quantity. To such a quantity, after (if necessary) some additional studies and considerations, can be ascribed a definite *physical meaning*. The ultimate (at given stage of the science) qualitative and quantitative relationships between the physical quantities and their relevance to the respective elements of the physical reality can be completely clarified within given physical theory. It must be underlined that both understandings, the present one and that one of Ref. [4], do not contradict each other in the sense that the requirement of relationship between the physical objects and the related physical quantities does not exclude the possibility that such a relationship may predict with certainty the values of the physical quantities. The opposite, however, does not hold, *i.e.*, a relationship may exist even if the behavior of the physical quantities cannot be predicted with certainty. Consequently, the present approach is more general and it can be applied if any of absolute (Laplacian) or partial (quantum mechanical) [29] determinism takes place in the physical reality.

The problem with the MP causality will be considered based namely on the above understanding of causality: *if a change of the MP causes observable physical consequences, e.g., change of other quantity that is subject of direct or indirect observation/measurement, then we may consider that the MP is causally related with the physical reality.* That is why, looking for a sensible physical behavior of the MP and its causal relationship with other observable physical quantities will be the main goal in the forthcoming considerations.

III. Evidences of material phase causality

The basic arguments in support of the MP causality will be classified in the following groups: special theoretical arguments, general theoretical arguments, and experimental evidences.

III. 1. Special theoretical arguments.

In this section we present the results of an extended study on the dynamics of the internal state of a quantum system, paying spatial attention to the MP [20]. To reveal the MP dynamics, the quantum system must be involved in a definite physical process. Isolated quantum system in stationary bare states is not suitable for that purpose. That is why, the quantum system will be subject to interaction with an electromagnetic field and the environment. Nonadiabatic factors from the field (field time derivatives) and the environment (dumping) naturally appear in the rigorous solution [21] of the field-matter interaction problem.




Fig.1. Bare (zero external field) (a) and dressed (non-zero external field) states (b) and (c) of the quantum system and the respective transitions at ground (b) and excited (c) initial conditions. The bold arrows show the optical pumping, the hollow arrows show transitions due to the nonadiabatic factors, and the wavy arrows show the dumping. The black spots show the initial state (initial condition) from which the process of excitation begins.

The problem of the MP causality will be treated within an analytic solution $|\Psi(\vec{r}, t)\rangle$ of the time dependent Schrödinger equation

$$\hat{H}|\Psi(\vec{r}, t)\rangle = i\hbar\partial_t|\Psi(\vec{r}, t)\rangle \quad , \quad (1)$$

applying a generalized adiabatic condition [20]. The Hamiltonian of the (two-level) quantum system under consideration (in standard notations) is

$$\hat{H} = \sum_{j=1}^2 \hbar\omega_j |j\rangle \langle j| - \mu E (|1\rangle \langle 2| + h.c.) - i\hbar \sum_{j=1}^2 \frac{\gamma_j}{2} |j\rangle \langle j| \quad . \quad (2)$$

The original bare states $|1\rangle$ and $|2\rangle$ of the quantum system, are coupled by a nonzero electric dipole moment. If $|1\rangle \equiv |g\rangle$ and $|2\rangle \equiv |e\rangle$ are ground and excited state, respectively, the dumping rate γ_g of $|g\rangle$ can be taken zero. The states will be called “ground” $|1\rangle \equiv |g\rangle$ and “excited” $|2\rangle \equiv |e\rangle$, although the first one is not necessarily to be a real ground state.

The electromagnetic (optical) field is expressed in the form

$$E = (1/2) E_o(t) [\exp(i\Phi_F(t)) + c. c.] \quad , \quad (3)$$

where $E_o(t)$ and $\Phi_F(t) = \omega t + \varphi(t)$ are the amplitude/envelope and the total phase of the field, respectively, ω is the carrier frequency of the field, and $\varphi(t)$ is the envelope-carrier relative phase. The above representation of the electromagnetic field in terms of envelope and carrier wave is quite general and holds even for ultrashort pulses of envelope time duration as short as the carrier wave period [30], which is in the order of few femtoseconds for the electromagnetic fields in the optical range. The original bare states of the quantum system, from which the dressed states arise, are taken in the form [20]

$$\begin{aligned} |g(\vec{r}, t)\rangle &= |g(\vec{r})\rangle \exp[-i\Phi_g(t)] \\ |e(\vec{r}, t)\rangle &= |e(\vec{r})\rangle \exp[-i\Phi_e(t)] \quad , \end{aligned} \quad (4)$$

where $\Phi_g(t) = \varphi_g + \omega_g t$, $\Phi_e(t) = \varphi_e + \omega_e t$ are the total dynamical phases; ω_g , ω_e are eigenfrequencies/energies and φ_g , φ_e are the respective initial phases. The latter are considered because of the arguments given in Sec. II.

The total wave function $|\Psi(\vec{r}, t)\rangle$ is found as a solution of the time dependent Schrödinger equation (1) of Hamiltonian (2) and can be presented as

$$|\Psi(\vec{r}, t)\rangle = \frac{1}{\sqrt{\tilde{\Omega}'}} \left\{ \frac{\tilde{\Lambda}'_{10}}{\sqrt{\tilde{\Omega}'_0}} \text{COS}^{-1} \frac{\theta}{2} |G\rangle + \frac{\tilde{\Lambda}'_{20}}{\sqrt{\tilde{\Omega}'_0}} \text{SIN}^{-1} \frac{\theta}{2} |E\rangle \right\} \quad , \quad (5)$$

at ground state initial conditions, and

$$|\Psi(\vec{r}, t)\rangle = \frac{1}{\sqrt{\tilde{\Omega}'}} \left\{ \frac{\Omega_0}{2\sqrt{\tilde{\Omega}'_0}} \text{COS}^{-1} \frac{\theta}{2} |G\rangle + \frac{\Omega_0}{2\sqrt{\tilde{\Omega}'_0}} \text{SIN}^{-1} \frac{\theta}{2} |E\rangle \right\} \quad , \quad (6)$$

at *excited state initial conditions* [31].

New kind of states naturally arises from the initial bare states in the course of interaction of the quantum system with the electromagnetic field in presence of the nonadiabatic factors. Such states of the combined “atom”-field system will be called *phase sensitive nonadiabatic dressed states* (PSNADSs) [32]. Each of the PSNADSs, ground $|G\rangle$ and excited $|E\rangle$, consists of real, $|G\rangle_r$ and $|E\rangle_r$, and virtual, $|G\rangle_v$ and $|E\rangle_v$, components

$$|G\rangle = \text{COS} \frac{\theta}{2} |G\rangle_r + \text{SIN} \frac{\theta}{2} |G\rangle_v \quad (7)$$

$$|E\rangle = \text{COS} \frac{\theta}{2} |E\rangle_r - \text{SIN} \frac{\theta}{2} |E\rangle_v \quad (8)$$

The quantities $\text{COS} \theta/2 = \sqrt{\tilde{\Lambda}'_1/\tilde{\Omega}'}$ and $\text{SIN} \theta/2 = \sqrt{-\tilde{\Lambda}'_2/\tilde{\Omega}'}$ are complex functions that correspond to the usual $\cos \theta/2$ and $\sin \theta/2$ in the case of adiabatic dressed states [33], $\tilde{\Omega}'$ is generalized (nonadiabatic) Rabi frequency. The subscript "0" in the respective quantities means their initial value at the initial moment $t = 0$ and come from the respective initial conditions.

The real and the virtual components of the PSNADSs can be expressed in the following form:

$$\begin{aligned} |G\rangle_r &= |g\rangle \exp(-i\Phi_{G,r}) \\ |G\rangle_v &= |e\rangle \exp(-i\Phi_{G,v}) \\ |E\rangle_r &= |e\rangle \exp(-i\Phi_{E,r}) \\ |E\rangle_v &= |g\rangle \exp(-i\Phi_{E,v}) \end{aligned} \quad (9)$$

where $\Phi_{G,r}$, $\Phi_{G,v}$, $\Phi_{E,r}$, and $\Phi_{E,v}$ are the total phases of the respective real and virtual components of the dressed states.

At *ground state initial conditions*, the total phases of the dressed state components are:

$$\begin{aligned} \Phi_{G,r} &= \varphi_g + \int_0^t \tilde{\omega}'_G dt' \\ \Phi_{G,v} &= \Phi_{G,r} + \Phi_F = (\varphi_g + \varphi(t)) + \int_0^t (\tilde{\omega}'_G + \omega) dt' \\ \Phi_{E,r} &= \Phi_{G,r} + \Phi_F + \Phi_{NAD} = (\varphi_g + \varphi(t)) + \int_0^t \tilde{\omega}'_E dt' \\ \Phi_{E,v} &= \Phi_{E,r} - \Phi_F = \varphi_g + \int_0^t (\tilde{\omega}'_E - \omega) dt' \end{aligned} \quad (10)$$

At *excited state initial conditions*, the total phases of the dressed state components are:

$$\begin{aligned} \Phi_{E,r} &= \varphi_e + \varphi(t) - \varphi_0 + \int_0^t \tilde{\omega}'_E dt' \\ \Phi_{E,v} &= \Phi_{E,r} - \Phi_F = \varphi_e - \varphi_0 + \int_0^t (\tilde{\omega}'_E - \omega) dt' \\ \Phi_{G,r} &= \Phi_{E,r} - \Phi_F - \Phi_{NAD} = \varphi_e - \varphi_0 + \int_0^t \tilde{\omega}'_G dt' \\ \Phi_{G,v} &= \Phi_{G,r} + \Phi_F = \varphi_e + \varphi(t) - \varphi_0 + \int_0^t (\tilde{\omega}'_G + \omega) dt' \end{aligned} \quad (11)$$

Here, $\tilde{\omega}'_G$ and $\tilde{\omega}'_E$ are nonadiabatic frequencies/energies of the ground and the excited dressed states, respectively, $\Phi_{NAD} = \int_0^t \Delta \tilde{\omega}'_{NAD} dt' = \int_0^t (\tilde{\omega}'_E - \tilde{\omega}'_G - \omega) dt'$ is nonadiabatic phase, and $\Delta \tilde{\omega}'_{NAD} = \tilde{\omega}'_E - \tilde{\omega}'_G - \omega$ is nonadiabatic frequency detuning. The frequencies of the ground $\tilde{\omega}'_G$ and excited $\tilde{\omega}'_E$ dressed states are Stark shifted and modified by means of the nonadiabatic contributions from the field ($\partial_t \varphi$, $\Omega^{-1} \partial_t \Omega$) and the dumping (γ_g , γ_e). The mathematical details of the approach can be found in [21].

The behavior of the dynamical phase can be revealed analyzing Eqs. (9)-(11). Starting from $|G\rangle_r$, *ground state initial condition* or $|g\rangle \dashrightarrow |G\rangle_r \rightarrow |G\rangle_v \Rightarrow |E\rangle_r \rightarrow |E\rangle_v$ sequence (the dashed arrow shows continuous evolution of the initial bare state toward the respective real dressed state component, the continuous arrow shows radiative transition within given dressed state, and the hollow arrow shows nonradiative nonadiabatic transition between the different dressed states), Fig.1(b), and using Eqs. (9) and (10) one may find the following development of the states and their phases along with the relevant physical processes. The bare ground state $|g(\vec{r}, t)\rangle$ evolves into a new ground state, the dressed ground state $|G\rangle$, while the eigenfrequency/energy ω_g of $|g\rangle$ evolves continuously into the eigenfrequency/energy $\tilde{\omega}'_G$ of the

real component $|G\rangle_r$ of $|G\rangle$. At the same time, the electromagnetic field “peaks up” the initial phase φ_g of the bare ground state and modifying the instantaneous energy of the state forces the total phase Φ_g of $|g(\vec{r},t)\rangle$ to evolve into the total phase $\Phi_{G,r}$ of $|G\rangle_r$. Once $|G\rangle_r$ is created, the process of creation/population of the other dressed state components and their phases can be traced out starting from $|G\rangle_r$. The next step is the formation of $|G\rangle_v$ from $|G\rangle_r$. The phase $\Phi_{G,v}$ of the virtual component $|G\rangle_v$ results from the phase $\Phi_{G,r}$ of the real component $|G\rangle_r$ adding the field (optical) phase Φ_F . At the same time, physically, the virtual component of the ground state results from the real component of the ground state by temporal association (and reemission) of one photon from (to) the field, stimulated by the field itself. Next, the phase $\Phi_{E,r}$ of the real component $|E\rangle_r$ results from the phase $\Phi_{G,v}$ of the virtual component $|G\rangle_v$ adding the nonadiabatic phase Φ_{NAD} . As a consequence, the phase of $|G\rangle_v$ transfers to $|E\rangle_r$ acquiring in this process nonadiabatic contributions from the field and the dumping factors while the quantum system overcomes the nonadiabatic frequency detuning $\Delta\tilde{\omega}'_{NAD}$. At the same time, in accordance with the adiabatic theorem of quantum mechanics [34], the physical transition between different states, in this case $|G\rangle_v$ and $|E\rangle_r$ components of the ground and excited dressed states, results namely from the nonadiabatic factors acting on the quantum system. This is accompanied by irreversible absorption of one photon from the field thus leading to continuous (within the lifetime of the state) population of the real excited state $|E\rangle_r$. Finally, the phase $\Phi_{E,v}$ of the virtual component $|E\rangle_v$ results from the phase $\Phi_{E,r}$ of the real component $|E\rangle_r$ subtracting the optical phase Φ_F . At the same time, physically, the virtual component of the excited state results from the real component of the excited state by temporal emission (and reabsorption) of one photon to (from) the field, stimulated by the field itself. The above analysis shows exact correspondence between the MP behavior and the respective physical processes. In particular, it naturally explains why the initial MP φ_g of the ground state $|g\rangle$, from which the process of formation of the dressed states begins at ground state initial conditions, appears in all dressed state components, whereas the excited state initial MP φ_e totally disappears. Similar consequences can be revealed at *excited state initial condition*, or $|e\rangle \rightarrow |E\rangle_r \rightarrow |E\rangle_v \Rightarrow |G\rangle_r \rightarrow |G\rangle_v$ sequence, Fig.1(c), using Eqs. (9) and (11). The bare excited state $|e(\vec{r},t)\rangle$ evolves into a new excited state, the dressed excited state $|E\rangle$, while the eigenfrequency/energy ω_e of $|e\rangle$ evolves continuously into the eigenfrequency/energy $\tilde{\omega}'_E$ of the real component $|E\rangle_r$ of $|E\rangle$. At the same time, the electromagnetic field “peaks up” the initial phase φ_e of the bare excited state and modifying the instantaneous energy of the state forces the total phase Φ_e of $|e(\vec{r},t)\rangle$ to evolve into the total phase $\Phi_{E,r}$ of $|E\rangle_r$. The formation/population of the dressed state components can be traced in a similar way as in the case of the ground state initial conditions, starting in this case from $|E\rangle_r$. The next step is the formation of $|E\rangle_v$ from $|E\rangle_r$. The phase $\Phi_{E,v}$ of $|E\rangle_v$ results from the phase $\Phi_{E,r}$ of $|E\rangle_r$ subtracting the optical phase Φ_F . Physically, the virtual component $|E\rangle_v$ results from the real component $|E\rangle_r$ by a temporal release (and reabsorption) of one photon to (from) the field. The transition from the virtual state $|E\rangle_v$ to the real state $|G\rangle_r$ results again from the action of nonadiabatic factors, acquiring the respective nonadiabatic phase Φ_{NAD} . This is accompanied by irreversible emission of one photon to the field leading to continuous (within the lifetime of the state) population of the real ground state $|G\rangle_r$. Finally, the phase $\Phi_{G,v}$ of $|G\rangle_v$ results from the phase $\Phi_{G,r}$ of $|G\rangle_r$ adding the field phase Φ_F . The latter corresponds to the physical formation of $|G\rangle_v$ from $|G\rangle_r$ as a result of temporal association (and reemission) of one photon from (to) the field. Again, the initial phase φ_e of the state from which the process begins, in this case - the excited state $|e\rangle$, appears in all dressed states components, while the ground state initial phase φ_g totally disappears, Eq.(11). It is important to note that such behavior of the initial phase does not result from a kind of trivial elimination of the respective phases due to the initial conditions but represents a general behavior of the MP. It is a consequence of the specified way of formation of the respective quantum states and their phases.

Although the MP behavior is the main subject of this work, the behavior of the optical phase is also important for the phase behavior of the quantum system because both phases additively constitute the total phase of the PSNADS. The appearance of the optical phase at ground state initial conditions is rather simple and transparent and it promotes the understanding of the MP dynamics. It shows that, whereas the energy of the original ground state $|g\rangle$ continuously evolves into that one of $|G\rangle_r$, the initial phase φ_g of $|g\rangle$ transfers without change into initial phase of $|G\rangle_r$, Eq. (10). At excited state initial conditions, however, the total phase of the PSNADS is subject to more complicated and non-intuitive

way of formation. If the phase $\varphi(t)$ of the optical pulse remains constant, $\varphi(t) = \varphi_0 = \text{const}$, it cancels the term $\varphi(t) - \varphi_0$ in Eq.(11), and the phase behavior at excited state initial conditions coincides with that one at ground state initial conditions. In that case, the initial phase φ_e of the original bare excited state $|e\rangle$ again transfers without change into initial phase φ_e of $|E\rangle_r$. If the phase $\varphi(t)$ changes in time, a nonzero phase difference $\varphi(t) - \varphi_0 = \sum_{k=1} (k!)^{-1} \varphi^{(k)}(t=0) t^k$ appears in the total phase $\Phi_{E,r}$ of the initial state $|E\rangle_r$ in the case of excited dressed state formation sequence, Eq. (11). With exception of absolute envelope-carrier phase $\varphi_0 = \varphi(t=0)$, the MP of the initial dressed state component $|E\rangle_r$ “absorbs” (in the course of its formation from $|e\rangle$) all time dependent phase contributions from the forcing optical field coming from carrier frequency shift, $\varphi_1 = \partial_t^1 \varphi(t=0)$, linear chirp, $\varphi_2 = \partial_t^2 \varphi(t=0)$, quadratic chirp, $\varphi_3 = \partial_t^3 \varphi(t=0)$, etc. It can be considered as a kind of nonadiabatic penetration of the time dependent optical phase into the phases of the PSNADSS. In that case, the initial phase φ_e of $|e\rangle$ evolves into $\varphi_e + \varphi(t) - \varphi_0$ and including the above mentioned time dependent phase contribution from the field, leads to the formation of the total phase of the real excited state component $|E\rangle_r$, Eq.(11).

The dynamics of the quantum system within the PSNADSSs shows that *the MP behaves as an additive dynamical quantity that causally follows the initial conditions (the initial, constant part of the MP) and the physical process (the time dependent part of the MP) responsible for the formation/population of given quantum state*. Such a behavior of the MP has been called MP tracking [20]. It is important to note that the MP tracking takes place within both, the adiabatic [35] and the nonadiabatic [20] dressed states. This means that it is more fundamentally related to the dynamics of the quantum system than the particular way, adiabatic or nonadiabatic, of the field-matter interaction. The above analysis reveals not only a well expressed physical behavior of the MP but also that such a behavior cannot be well understood without taking into account the initial conditions and the relevant physical processes. To the best of our knowledge, such detail dynamical behavior of the material and the field phases is presented for the first time. The theoretically and experimentally observed manifestation of the sensitivity of the MP with respect to the physical processes and the initial conditions is actually what it is usually considered as a *sensible physical behavior* of given quantity. Elements of the discussed above behavior of the MP can be distinguished in a number of experiments with material wave packets, (for details, see section III.3). In the light of the above considerations, we may also speculate that it is the phase/action that is actually added in response to the physical phenomena whereas the addition of the frequency/energy (the time derivative of the phase/action) appears to be a secondary result.

III. 2. General theoretical arguments.

In this section we present a general theoretical argument showing that the MP is causally related with the physical reality. In the polar representation, the wave function $\Psi = R(\vec{r}, t) \exp[-i\Phi(\vec{r}, t)] = R(\vec{r}, t) \exp[iS(\vec{r}, t)/\hbar]$ consists of amplitude $R(\vec{r}, t)$ and phase $\Phi(\vec{r}, t) = -\hbar^{-1}S(\vec{r}, t)$, where S is the physical action. In other words, up to a constant factor (determined by a fundamental physical constant \hbar), the MP coincides with the physical action. It can be shown that the amplitude R and the action S , or the phase Φ of the wave function, are not independent but become mutually related. Such a relation can be revealed within the so called "hydrodynamic" representation of the quantum mechanics. The latter is known since the works of de Broglie [36] and Madelung [37]. Later on, it is extensively exploited by D. Bohm [5] in the ontological (causal) interpretation of the quantum mechanics based on hidden variables concept [4]. Substituting the wave function in the Schrödinger equation leads to the following coupled differential equations for the action/phase and the amplitude

$$\partial S / \partial t + (\nabla S)^2 / 2m + V(\vec{r}, t) - (\hbar^2 / 2m) (\Delta R / R) = 0 \quad (12)$$

$$\partial (R^2) / \partial t + \nabla \cdot (R^2 \nabla S / m) = 0 \quad (13)$$

The first equation (12) is the so called quantum mechanical equation of Hamilton-Jacobi for the action/phase, where $V(\vec{r}, t)$ is the usual interaction potential, and $U(\vec{r}, t) = -(\hbar^2 / 2m) (\Delta R / R)$ is called quantum potential. The second equation (13) is the continuity equation for the quantum probability density $\rho = R^2 = |\Psi|^2$ [38]. The quantum potential, according to the Bohm's interpretation of quantum mechanics, originates from a real interaction between the wave function Ψ , considered as “a objectively real field” [5], and the particle. This field exerts a force on the respective particle by means of the quantum potential and, together with the usual interaction potential, governs its propagation. The quantum potential is responsible for all the non-classical features of the physical object. The hydrodynamic representation is used in the Bohmian mechanics to introduce a quantum concept of particle moving along a trajectory. Within this concept, an ensemble of Bohmian trajectories arises from the integration of the velocity $d_t \vec{r} = \vec{v} = m^{-1} \nabla S(\vec{r}, t)$. In this work, the hydrodynamic representation is used not to exploit the concept of trajectory (although it is close to our understandings) but to prove the MP causality. We neither consider the wave function as an objectively existing field, nor the quantum potential as a really existing potential that is capable to exert a force on a particle, see Sec.IV. The phase/action Φ/S and the amplitude R obey coupled differential equations (12) and (13). Therefore, they are not independent but codetermine each other. Any change of the amplitude of the wave function influences the action/phase and vice versa. It is hardly to believe that the square of the amplitude $R^2 = |\Psi|^2$ of the wave function is related to some element of the physical reality,

while its action/phase S/Φ , is irrelevant to the same reality. Taking into account the $S/\Phi \leftrightarrow R$ relationship that follows from Eqs. (12), (13) and the widely accepted $R \leftrightarrow \text{physical reality}$ relationship, as, *e.g.*, in the Copenhagen interpretation, existence of a complete $S/\Phi \leftrightarrow R \leftrightarrow \text{physical reality}$ relationship can be established. *Such $S/\Phi \leftrightarrow R \leftrightarrow \text{physical reality}$ relationship is fundamental and exists independently on the particular interpretation of the quantum mechanics.* The latter, however, can be considered as a basic reference point in the interpretation of the quantum mechanics. What is also important, it is namely the action/phase S/Φ , but not the amplitude R , that is ruled by the real dynamical equation (12). The amplitude R obeys the continuity equation (13), which plays only an auxiliary role with respect to the quantum dynamics, *i.e.*, to rule the conservation of the probability density. In view of the above considerations, the phase/action appears to be much more deeply related to the dynamics of the quantum system than the amplitude. Therefore, one may expect that much more fundamental physics is related namely with the MP. The logical relation $S/\Phi \leftrightarrow R \leftrightarrow \text{physical reality}$ is the most important, to our opinion, theoretical argument that proves the MP causality. It is completely general because it results solely from the structure of the Schrödinger equation and the wave function. The only the assumption involved here is the widely accepted (not only in the Copenhagen interpretation) relation between the amplitude of the wave function and the physical reality. Whereas the hydrodynamic representation of the quantum mechanics is well established, it has not been used till now (to the best of our knowledge) to reveal the existence of a fundamental relation between the phase of the wave-function and the physical reality.

It is usually considered that the constant action/phase does not matter for the physical phenomena [39] because it participates only in derivatives in the classical and quantum Hamilton-Jacobi equation (12), as well as in the continuity equation (13). Such point of view is not supported here, especially if the interference phenomena are considered. The action/phase is an additive quantity and the action/phase acquired during the dynamical evolution is added to its initial, constant, part, which is determined by the initial conditions. The sensible physical behavior of the initial (constant) value of the phase/action has been already discussed in Sec. II and III.1. What is more important, however, any constant shift of the phase/action may lead to observable effects at suitably designed (interference) experiments.

While the above argument is enough to prove the MP causality at a fundamental level, another general consideration worth also to be mentioned. It arises from the quantum superposition principle. While not explicitly included within the postulates of quantum mechanics, *e.g.*, [28], the quantum superposition principle is one of the foundations of the quantum mechanics. It constitutes the essence of the quantum mechanical interference. The interference between the quantum states, as well as between the quantum transitions, represents the very nature of the quantum mechanical phenomena. Having in mind the quantum superposition principle [11]

$$\Psi(\vec{r}, t) = \sum_i C_i \Psi_i(\vec{r}, t) \quad , \quad (14)$$

it is a trivial mathematical result that both, the amplitudes $|\Psi_i|$ and the phases Φ_i of the superimposed states Ψ_i participate in the determination of the amplitude $|\Psi|$ and the phase Φ of the resultant state Ψ . Any change of the phases of the superimposed states may cause observable results on the entire resultant state, *i.e.*, its amplitude and phase. This is another evidence of the mutual relation between the phase and the amplitude of the wave function.

III. 3. Experimental evidences.

The most important point in the present considerations is that the above theoretical arguments have experimental confirmations. In fact, a large number of experimental evidences from various area in physics exist in support of the MP causality. Here we will only summarize the outcome and the main conclusions from some of the most convincing, to our opinion, experimental studies, without pretending for completeness of this survey. Phase sensitivity of observable quantities takes place with respect to, both, the field phase (the phase of the optical pulses that create the material wave-packets), which is a well recognized effect, and, what is particularly important here - the MP (the phase of the material wave-packet itself). Here, the accent will be put on experiments that clearly demonstrate the MP dependence of the physical processes, while some important aspects of optical phase dependence will also be considered.

The interference of the matter waves is a basic quantum mechanical phenomenon. A general outcome of the matter-wave experiments consists in the fact that, changing the MP, one may affect the interference picture with material wave-packets, *i.e.*, the population of given internal state of the quantum system, or the number density distribution of the particles in space, and all these are subject to experimental observation. The experiments under consideration have not been particularly designed to study the MP causality and critical inspection of the conditions and the experimental results are required to make correct conclusions.

The material wave packets can be classified into two well distinguishable pure types: (i) internal wave-packets (superposition of bound states in a quantum system - atom, molecule) [16-19] and (ii) external, de Broglie wave-packets (external motion state of free or quasi-free particle - electron, neutron, atom, molecule, etc.) [14, 15]. To illustrate the discussion, the following rather general form of MP will be considered: $\Phi(\vec{r}, t) = \hbar^{-1} \left(\int_{t_0}^t E dt' - \int_{\vec{r}_0}^{\vec{r}} \vec{p} \cdot d\vec{r}' \right)$, where E is the energy and \vec{p} is the linear momentum. The total MP can be ruled either by control of its initial, constant part, or by control of its variable part. The initial phase depends on the conditions at the initial moment t_0 and the initial position \vec{r}_0 of the creation of the wave packet. It can be manipulated, *e.g.*, by the phase of the electromagnetic field that creates the wave packet, see Eqs. (10) and (11) and the related discussion in sec.III.1. The variable phase depends on the particular dynamics in which the quantum system becomes involved. It can be controlled changing the energy E of the quantum system and/or the evolution time (from t_0 to t), or changing the linear momentum \vec{p} and/or the evolution path (from

\vec{r}_0 to \vec{r}) to the time/position of interference with another, reference wave packet. For the case of internal wave packets, the energy of the quantum system can be changed by a selective excitation of a particular superposition of electronic states in an atom [16], or a particular superposition of rovibration states of given electronic state in a molecule [17-19]. Change of the linear momentum or the whole phase/action path integral can be realized by a selective excitation of given electron trajectory [40]. For convenience, the phase sensitive experiments in the forthcoming discussion will be divided into the following types: (i) experiments with bound intraatomic/intramolecular wave-packets; (ii) experiments with quasi-free, tunnel electron wave-packets; and (iii) experiments with free wave-packets.

(i) experiments with bound (intraatomic/intramolecular) wave packets

MP dependence of the physical observables can be distinguished in an analog of Young's double slit type interferometer within an atom [16], or Michelson type interferometer within a molecule [17-19]. In these experiments, the light beams in the usual optical interferometers are replaced by electron (within the atom) or nuclear (within the molecule) wave packets. The wave packets are created by a sequence of usually two (pump) laser pulses of controllable phase. A third (probe) laser pulse is used to probe the evolution of the superposition of the wave packets. To achieve well expressed MP effects, the wave packets, *i.e.*, the pump pulses, must be substantially shorter than the electron orbiting time in the atoms or the nuclear vibration time in the molecules. Once created, the wave packets are ruled solely by the intraatomic/intramolecular Hamiltonian in the time between the pulses. After some propagation time, the wave-packets may overlap and interfere. The result of the interference can be detected by fluorescence interferogram from the excited state [17-19], by ion current due to ionization of the atom/molecule from the superposition state [16], or any other process capable to reveal the excited state population. Beside of other factors, the local population in the interference state depends on the MP acquired by the wave packets during their evolution. The MP difference between the wave packets inside the atoms/molecules can be controlled by a number of ways: (i) changing the phases of the pump pulses, and, in this way, the initial phase of the created wave packets [16], (ii) changing the delay time between the pump pulses, and, in this way, the relative MP between the created wave packets [16-19], (iii) changing the carrier frequency of the pump pulses, which changes the carrier vibration frequency (the mean energy) of the created intramolecular wave packets, and, thus, the time rate of the MP acquisition [17, 18], or (iv) any combination from above. The most important outcome from these experiments is that a change of the MP in some of the specified ways leads to observable physical effects [16-19]. In particular, even a constant phase shift of the phase-locked laser pulses that create the material wave packets leads to a dramatic change in the observed result [16-18]. In addition, the well defined behavior of the field-matter phenomena is a strong evidence that the material wave packet is able to "carry" in a deterministic way the optical phase ("absorbed" from the optical field at the process of creation of the wave packets), adding to it a definite value of the MP acquired during the evolution of the wave packets inside the quantum system (atom, molecule, etc.).

(ii) experiments with quasi-free tunnel electron wave packets

One of the most amazing advances in the ultrafast physics is the generation of single attosecond pulses by means of atomic level control of the emission of high-harmonics from tunnel electrons driven by high-intensity femtosecond laser pulses. The underlying physical processes of the high-harmonic generation are strongly sensitive to the particular trajectories of the tunnel electrons. Electron trajectory concept in the spirit of Feynman's path integral formulation of quantum mechanics [6] becomes a very successful approach in the description of some high-field phenomena: high-harmonic generation, above-threshold ionization [40]. Within this approach, the action/phase acquired by the tunnel electrons along definite trajectories, quantum paths, plays the key role in the development of above phenomena. The acquired action/phase differs in strength, energy region [40], as well as in spatial and temporal coherence [41]. In this way, the contributions of the different trajectories can be experimentally distinguished, leading even to macroscopically visible results [41]. In the same way as for the PSNADS, section III.1, the created electron wave packet "absorbs" the optical phase and adding the acquired pure MP forms the total phase of the tunnel electron wave packet. Influence of the phase modulation (chirp) of the exciting pulse on the spectra of the generated high-harmonics [42] is another indication that the field and the matter (atomic dipole) phases add together in a predictable and controllable way. Changing the phase of the laser field and/or MP by means of selective excitation of definite electron trajectories affects the total, optical plus material, phase of the wave packet, and it is accessible to experimental observation [40-42].

(iii) experiments with free wave-packets

Every experiment on interference between particles (matter waves) [13-15, 43-45] is, in fact, an example of phase sensitive phenomenon. Nevertheless, below we will point out only experiments which explicitly show that the interference picture is sensitive to *the change* of the MP.

Interference of wave packets of free electrons is observed experimentally ionizing argon atoms by extreme ultraviolet attosecond pulses and a suitable infrared laser field to induce a momentum transfer of the released electrons [46]. Dependence of the interference picture on the phase of the electron wave packets, *i.e.*, the MP, can be distinguished in these experiments. The phase of the wave packets is manipulated by momentum transfer from the infrared laser field, or changing the time delay between the infrared and attosecond pulses. The change of the phase of the electronic wave packets leads to observable change in the interference picture [46].

The atomic/molecular interference is basic phenomenon in the atomic/molecular matter wave interferometry. The latter forms a well established field of research with strong impact on the fundamental studies, high-precision metrology, etc. Change of the phases of the atomic partial waves, and thus, the created (Ramsey) fringes, by means of (i) change of the phase of the laser fields that are used to split and recombine the atomic beam in a Ramsey interferometer, (ii) ac-Stark shift, or (iii) rotation of the atomic interferometer (Sagnac effect) is well known observable result in the matter wave

interferometry [47, 48]. Thus, the matter-wave interferometry is another source of evidences of the dependence of the physical phenomena on the MP.

The above theoretical (III.1, III.2) and experimental (III.3) investigations conclusively establish the existence of a fundamental relation of the MP with the physical reality and we may summarize that *the MP is causally related with the dynamics of the quantum system*. The latter, together with the fact that a Hermitian operator can be associated to the phase [49, 50], allows considering the MP as a physical observable on the same formal basis as any other physical observable [Appendix A]. Hence, although the entire wave function Ψ does not have real physical meaning, each of its elements, amplitude R and phase $\Phi(S)$, have observable appearances in the physical phenomena. The MP causality brings the matter waves closer to the electromagnetic waves, for which the physical meaning of the phase is undisputable [Appendix B]. In the light of above arguments, it seems apparent that the Copenhagen interpretation unnecessarily restricts the physical meaning of the wave function to its amplitude, only. Investigations on the MP behavior may lead to much deeper understanding of the intimate nature of the quantum phenomena.

IV. Dynamics-statistical interpretation of quantum mechanics

In spite of the well expressed relation between the physical reality and the MP, the latter yet remains irrelevant to the interpretation of the quantum mechanics. Some basic features of the MP announced here are not consistent with the Copenhagen interpretation, or some of the most significant non-Copenhagen interpretations [5-9]. This invokes considering of another interpretation. It will be based solely on the basic principles of the quantum mechanics, without involving additional hypotheses. Also, we hope that it is not influenced by the observer's prejudice about the physical reality.

Let us first begin with the ontological point of view according to which the wave function is an objectively existing field Ψ that is capable to exert a force on the related particle by means of the quantum potential [5]. The quantum potential depends on the instantaneous positions $\vec{r}_1, \dots, \vec{r}_n$ of all particles in the system at given moment of time t , $U(\vec{r}_1, \dots, \vec{r}_n, t) = -(\hbar^2/2m)\Delta R(\vec{r}_1, \dots, \vec{r}_n, t)/R(\vec{r}_1, \dots, \vec{r}_n, t)$. It is considered as a non-local connection between the individual physical objects of the system. The extrapolation of this approach to the universe leads to the Bohm's idea of existence of an implicate order and an undivided wholeness of the entire universe through its quantum potential. Such a physical picture, however, should be introduced in the theory since the very beginning with the subsequent dramatic reconstruction of the foundations of the physics. In that case, instead of motion and interaction between individual particles and fields (considered as relatively independent entities), the physical phenomena should also reflect the additional contribution of all material objects through the quantum potential of the system. If this is so, solving of any particular problem would require knowledge of the state of the whole universe, more particularly, the instantaneous positions of all particles in it. While it could be a correct approach, there are not enough evidences that it is namely the quantum potential that may play role of such an instantaneous unification factor. A useful reference point when considering the problem with the interpretation of the quantum mechanics, in general, and the quantum potential, in particular, is to go back to the origins. The physical processes involved in given physical problem are specified in advance and it is, in fact, a precondition to construct the Hamiltonian in the Schrödinger equation and to solve the problem. Any material system consists of particles, each of which, at first, moves (finitely or infinitely) in space and, consequently, possesses kinetic energy described by kinetic energy operator $\hat{T} = \hat{p}^2/2m$ and, at second, interacts with the other particles or the environment by means of the well known ordinary physical fields and, consequently, possesses energy of the interaction described by interaction operator $\hat{U}(\vec{r}, t)$ (potential energy operator $\hat{U}(\vec{r})$, for conservative systems) in the Hamiltonian. Such a general understanding exhausts the widely accepted picture of the physical phenomena. If the quantum potential is considered as a real interaction, it should correspond to completely new physics that is not founded in the theory in advance. However, no any "unusual" physics, *e.g.*, interaction with unknown physical fields, the Ψ -field [5], is being considered for the quantum system and its relation with the physical reality at the formulation of the problem. Hence, the real existence of the quantum potential should be subject to a lucky coincidence. Although the latter cannot be excluded, no direct evidences (to the best of our knowledge) of existence of additional non-local interaction between given particle and the rest of the system (in fact, the rest of the universe) due to the quantum potential have been shown. The Bohm's idea of unified and undivided world is highly inspiring and beautiful but more convincing arguments are required so as to be accepted. That is why, the problem of the interpretation of the quantum mechanics will be considered within the usual understanding of the physical phenomena.

The appearance of the quantum potential is, in fact, a formal consequence from the particular way of description of the physical state accepted in the quantum mechanics, *i.e.*, by means of wave function of particular structure that obeys dynamical equation of particular structure. The quantum potential is responsible for all non-classical features in the quantum Hamilton-Jacobi equation. Without the quantum potential, it reduces to the classical Hamilton-Jacobi equation that determines the classical trajectory of the particle. A class of Bohmian trajectories arises from of the quantum potential after integration of the velocity $\partial_t \vec{r} = \vec{v} = m^{-1}\nabla S(\vec{r}, t)$. An important indication of the meaning of the quantum potential consists in the fact that it is determined only by the amplitude R of the wave function, $U(\vec{r}, t) = -(\hbar^2/2m)(\Delta R/R)$. The amplitude of the wave function has well recognized statistical meaning (probability density), not only in the Copenhagen interpretation, but also in the Bohmian mechanics [5]. Hence, the quantum potential introduces statistical properties in the quantum Hamilton-Jacobi equation and it can be referred to the probabilistic meaning of the amplitude of the wave function rather than to a real interaction. That is why, instead of quantum potential, it can be more reasonably called *statistical term*, as will be used hereafter. The statistical properties of the quantum phenomena arise from the fact that, in the quantum mechanics, the dynamics of an individual quantum system is described actually within an ensemble of similar systems. The latter, however, well corresponds to the typical

experimental approach in the study of physical objects of microscopic size. What is important, the results predicted by such an approach are consistent with the experimental observations [51].

Summarizing the theoretical and experimental arguments presented here, we cannot entirely accept either the Copenhagen or the ontological interpretation [5] of the quantum mechanics. In fact, the wave function is a complex construct from the amplitude and the phase that have different physical meaning. The action/phase is ruled by the real dynamical equation (12) (the quantum Hamilton-Jacobi equation), which, beside of pure dynamical terms, includes in addition the statistical term. That is why, the action/phase has a quantum-dynamical meaning, which, however, is influenced by the statistical properties coming from the statistical term. The amplitude of the wave function, which is ruled by the continuity equation of the probability density, Eq. (13), has a statistical meaning, which is influenced by the dynamical properties of the action/phase through the term including the velocity, $\vec{v} = m^{-1}\nabla S(\vec{r}, t)$. Also, we consider that the amplitude of the wave function is related with objectively existing probability that is only registered by the observer, but not simply its knowledge (epistemology) about the state of the quantum object [52]. Thus, the dynamical and the statistical properties are entangled [53] in the wave function by means of its phase and amplitude, respectively. Such relation between the elements of the wave function is clearly expressed within the hydrodynamic representation of the quantum mechanics but not apparent within the Schrödinger's representation. Nevertheless, it exists irrespectively on the particular representation and becomes a fundamental feature of the wave function. Accepting the MP causality does not necessarily require acceptance of the ontological interpretation, considering Ψ as a real physical field [5]. We consider that the wave function as a whole does not have a definite physical meaning although its elements, amplitude and phase, are causally related with the physical reality and physical meaning can be attributed to both of them. Such an understanding lies between the Copenhagen interpretation (according to which the amplitude, but not the phase, has a physical (epistemological) meaning), and the deBroglie-Bohm causal interpretation (according to which, the entire wave function, being a real physical object/field that may exert a force on the particle, has physical meaning). Our understandings can be put into a kind of a *dynamics-statistical interpretation* of the wave function. It is remarkable that the dynamical and the statistical nature of the state become so well combined in the wave function that it gives the best known description of the physical reality at a quantum level.

Two main approaches of extraction of the physical information from the wave function can be distinguished, *probabilistic* and *dynamical* ones. The probabilistic approach is based on a bilinear product of wave functions, the matrix element, as in the standard quantum mechanics. The MP dependence within this approach is not easy to be seen due to its appearance in the matrix elements in a product of exponential phase factor and its complex conjugated. The $S/\Phi \leftrightarrow \text{physical reality}$ relationship suggests a dynamical way of extraction of the physical information. The dynamical approach, applied in the Bohmian mechanics [5], is based on the action/phase, from which the velocity/linear momentum and trajectory of the particle can be determined. The dynamical approach clearly reveals the dependence of the physical processes on the MP. Due to the mutual dependence between the amplitude and phase, both ways of extraction of the physical information are not independent but become *complimentary* each other. On the other hand, according to Bohr complimentary principle, there two kinds of (complementary) pictures in the description of the physical reality, particle-like and wave-like. Both pictures can be distinguished here based on the elements of the wave function. The particle-like picture is primary related with the action/phase S/Φ , influenced by the amplitude R due to their mutual relation in Eq. (12). For example, a particle-like feature like the quantum trajectory can be introduced by the integration of the velocity $\partial_t \vec{r} = \vec{v} = m^{-1}\nabla S(\vec{r}, t)$ [5]. The wave-like picture is primary related with the amplitude R , influenced by the action/phase S/Φ due to their mutual relation in Eq. (13). The square of the amplitude determines the probability density distribution, which reveals the wave-like picture at suitably designed experiments. Each of these pictures can be stronger or lesser expressed depending on the particular conditions. Both pictures become entangled in the wave function through the phase/action and the amplitude. Such a point of view can be considered as a further extension of the Bohr complementary principle, this time, based on the elements of the wave function. As can be seen, the truth (as we believe) does not recognize the border between the different formulations/interpretations of the quantum mechanics and reasonable concepts can be found in most of them, [Appendix C].

V. Conclusion

Theoretical and experimental evidences of a fundamental relation between the phase of the wave function and the physical reality are found. Existence of material phase causality is conclusively established on that ground. The phase/action and the amplitude of the wave function are mutually related by coupled dynamical and continuity equations. The phase of the wave function is ruled by the real dynamical equation, the quantum Hamilton-Jacobi equation, modified statistically by the amplitude through a statistical term. Hence, the material phase is primary related with the dynamics of the quantum system. The amplitude of the wave function is ruled by the continuity equation of the probability density, influenced dynamically by the action/phase. Hence, the amplitude of the wave function is primary related with the statistical properties of the quantum system. The amplitude is related with the real physical probability instead of the observer's knowledge (epistemology) about the state of the physical object. The wave function, as a whole, does not have a definite physical meaning (as in the standard interpretation of the quantum mechanics) while each of its elements, the amplitude and the phase/action, are causally related with the physical reality and physical meaning can be attributed to each of them. In view of above, the material phase appears to be a *missed parameter* within the standard interpretation. A new dynamics-statistical interpretation of quantum mechanics is formulated. The wave function incorporates the particle-wave duality through its phase and amplitude. It can be considered as a further extension of the Bohr complementary principle.

References

- [1] P.A.M. Dirac, *The principles of Quantum Mechanics*, Oxford, Clarendon Press, 1958.
- [2] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton University Press, 1955.
- [3] R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics, Vol.3, Quantum mechanics*, Addison-Wesley, Reading, MA, 1965.
- [4] A. Einstein, B. Podolsky, and N. Rosen, *Phys. Rev.* 47 (1935) 777
- [5] D. Bohm, *Phys. Rev.* 85 (1952) 166.
- [6] R. Feynman, *Rev. Mod. Phys.* 20 (1948) 367.
- [7] R. B. Griffiths, *Journal of Statistical Physics.* 36 (1984) 219.
- [8] H. Everett, *Reviews of Modern Physics*, 29 (1957) 454.
- [9] W. H. Zurek, *Reviews of Modern Physics* 75 (2003) 715.
- [10] M. Born, *Z. Phys.* 37 (1926) 863.
- [11] L.D. Landau, E.M. Lifshitz, *Quantum Mechanics, Nonrelativistic Theory*, Nauka, Moscow, 1989 (in Russian).
- [12] A. Aspect, J. Dalibard, and G. Roger, *Phys. Rev. Lett.* 49 (1982)1804.
- [13] A. Zeilinger, *Rev. Mod. Phys.* 71 (1999) S288.
- [14] M.S. Chapman, T. D. Hammond. A. Lenef, J. Schmiedmayer, R. A. Rubenstein, E. Smith, and D. Pritchard, *Phys. Rev. Lett.* 75 (1995) 3783.
- [15] S. Durr, T. Nonn, G. Rempe, *Nature London* 395 (1998) 33
- [16] M. W. Noel and C. R. Stroud, Jr, *Phys. Rev. Lett.* 75 (1995) 1252.
- [17] N.F. Scherer, A.J. Ruggiero, M. Du, G.R. Fleming, *J. Chem. Phys.* 93 (1990) 856.
- [18] N. F. Scherer, et al., *J. Chem. Phys.* 95 (1991) 1487.
- [19] J. J. Gerdy, M. Dantus, R. M. Bowman, A. H. Zewail, *Chem. Phys. Lett.* 171 (1990) 1.
- [20] I. G. Koprnikov, *Phys. Lett. A* 272 (2000) 10.
- [21] I. G. Koprnikov, *J. Phys. B: At. Mol. and Opt. Phys.* 34 (2001) 3679.
- [22] I. G. Koprnikov, *ICPEAC 2003 (2003) Mo196*.
- [23] M.V. Berry, *Proc. R. Soc. London A392 (1984) 45*.
- [24] R. Kienberger, E. Goulielmakis, M. Uiberacker, A. Baltuska, V. Yakovlev, F. Bammer, A. Scrinzi, Th. Westerwalbesloh, U. Kleineberg, U. Heinzmann, M. Drescher, and F. Krausz, *Nature*, 427 (2004) 817.
- [25] G. Sansone, E. Benedetti, F. Calegari, C. Vozzi, L. Avaldi, R. Flammini, L. Poletto, P. Villoresi, C. Altucci, R. Velotta, S. Stagira, S. De Silvestri, M. Nisoli, *Science* 314 (2006) 443.
- [26] M. Schultze, E. Goulielmakis, M. Uiberacker, M. Hofstetter, J. Kim, D. Kim, F. Krausz and U. Kleineberg, *New J. of Phys.* 9 (2007) 243.
- [27] D. Bohm, *Quantum Theory, Chapters 6 and 22*, Prentice-Hall Inc., New York, 1952.
- [28] C.C-Tannoudji, B. Diu, F. Laloë, *Quantum mechanics*, J. Wiley & Sons, N.Y., 1991
- [29]. The quantum mechanics is deterministic theory because the quantum state $|\Psi\rangle$ evolves deterministically obeying the respective dynamical/Schrödinger equation, $\hat{H}|\Psi\rangle = i\hbar\partial_t|\Psi\rangle$. The abstract quantum state $|\Psi\rangle$, however, does not have a definite physical meaning as a whole and does not corresponds to some known classical quantity, although quantities that have physical meaning can be constructed from it. On the other hand, the quantities that have well understandable meaning from a macroscopic point of view, *i.e.*, position, momentum, etc., cannot be followed deterministically within the quantum mechanics but can be predicted only statistically. Due to such a reason, the quantum mechanics can be considered as a partially deterministic theory.
- [30] T. Brabec and F. Krausz, *Rev. Mod. Phys.* 72 (2000) 545
- [31] The quantity $\Omega_0/\sqrt{2\sqrt{\Omega'_0}}$ is not extracted from the brackets in Eq. (6) in order to keep the form of the wave function similar to that one in Eq. (5)
- [32] Some discussion will be useful to clarify the terminology used here. The dressed states arise from the simplest, bare states of an isolated quantum system (atom or molecule, *e.g.*, the atomic hydrogen states, Born–Oppenheimer type molecular states, etc.) when it interacts with an electromagnetic field. If the field changes adiabatically, *i.e.*, it satisfies given adiabatic condition, adiabatic dressed states arise because the nonadiabatic factors are neglected, see Ref. 33 of this work. The introduction of generalized adiabatic condition [20] allows constructing dressed states of more general form, in which the nonadiabatic factors (up to given order) from the field (time derivatives of the field amplitude and phase) and the dumping are retained. For complete treatment of the material phase effects, the initial material phases are also subject to consideration. Such states were called Phase Sensitive Adiabatic States or PSASs [20] because a (generalized) adiabatic condition is still required for their derivation. In contrast to other known adiabatic states, however, the PSASs explicitly contain nonadiabatic factors. As the latter is of crucial importance for many physical problems, hereafter, these states will be more correctly called Phase Sensitive Nonadiabatic Dressed States, or PSNADS.
- [33] D. C. Hanna, M. A. Yuratich, D. Cotter, *Nonlinear Optics of Free Atoms and Molecules* (Springer-Verlag), 1979
- [34] M. Born, V. A. Fock, *Zeitschrift für Physik A Hadrons and Nuclei* 51 (1928) 165
- [35] I. G. Koprnikov, *Proceedings of the Int. Conference on Lasers'97*, Ed. J. J. Carroll, T. A. Goldman, STS Press 1998, pp. 291.
- [36] L. de Broglie, *C. R. Acad. Sci. Paris* 183 (1926) 447
- [37] von E. Madelung, *Z. Phys.*, 40 (1926) 322
- [38] In the Bohmian mechanics, the square of the amplitude of the wave function also has meaning of probability density.
- [39] E. Schmutzer, *Grundprinzipien der classischen mechanik und und der calssischen feldtheorie (kanonisher apparat)* VEB Deutscher Verlag der Wissenschaften, Berlin 1973.

- [40] P. Salières, B. Carré, L. Le Déroff, F. Grasbon, G. G. Paulus, H. Walther, R. Kopold, W. Becker, D. B. Milošević, A. Sanpera, M. Lewenstein, *Science* 292 (2001) 902
- [41] M. Bellini, C. Lyngå, A. Tozzi, M. B. Gaarde, T. W. Hänsch, A. L’Huillier, C.-G. Wahlström, *Phys. Rev. Lett.* 81 (1998) 297.
- [42] Z. Chang, A. Rundquist, H. Wang, I. Christov, H. C. Kapteyn, M. M. Murnane, *Phys. Rev. A* 58 (1998) R30.
- [43] M. O. Scully, B.-G. Englert, H. Walther, *Nature*, 351 (1991) 111.
- [44] M. S. Chapman, C. R. Ekstrom, T. D. Hammond, R. A. Rubenstein, J. Schmiedmayer, S. Wehinger, D. E. Pritchard, *Phys. Rev. Lett.* 74 (1995) 4783.
- [45] F. Riehle, A. Witte, Th. Kisters, J. Helmeke, *Appl. Phys. B* 54 (1992) 333.
- [46] T. Remetter, P. Johnsson, J. Mauritsson, K. Varjú, Y. Ni, F. Lépine, E. Gustafsson, M. Kling, J. Khan, R. López-Martens, K. J. Schafer, M. J. J. Vrakking, A.L’Huillier, *Nature Physics*, 2 (2006) 323.
- [47] U. Sterr, K. Sengstock, J. H. Müller, D. Bettermann, W. Ertmer, *Appl. Phys. B* 54 (1992) 341.
- [48] F. Riehle, A. Witte, Th. Kisters, J. Helmcke, *Appl. Phys. B* 54 (1992) 333.
- [49] L. Susskind, J. Glogower, *Physics*, 1 (1964) 49.
- [50] D. T. Pegg, S. M. Barnett, *Phys. Rev. A* 39 (1989) 1665.
- [51] The quantum mechanics is the most successful theory for computation of physical phenomena at a quantum level and it is considered that no contradictions exist with the experimental results. While we definitely agree with the first statement, the second one will be accepted after some comment. The experimentally observed phenomena agree with the predictions of the quantum mechanics if the experiment is done “at once” with a large number of quantum systems, or with a large number of subsequent experiments with single quantum systems, superimposing the outcome of all individual observations/measurements. For example, in the famous two-slit experiment, a single particle reproduces only “a point” of the whole interference picture predicted by the quantum mechanics. The latter can predict only the probability of the particle to hit given point on the screen. To reproduce the whole interference picture, one requires to superimpose a large number of individual interference events of single particles sent through the slits [13]. The conventional quantum mechanics does not reveal, in fact, the dynamics of a single particle and its behavior can be predicted only statistically.
- [52] We consider that the wave function, respectively its amplitude, describes objectively existing reality but not the observer’s knowledge about it. The evidence of that can be found in particularly designed EPR type experiments with entangled photons whose state of polarization is measured by arbitrary switched polarizers that remain completely unknown for the observer until the experiment is completed and the results are processed and stored, *e.g.*, G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, A. Zeilinger, *Phys. Rev. Lett.* 81 (1988) 5039. The well known correlation between the polarization states of the entangled photons is found even at the above mentioned special precautions against the influence of the observer on the process of acquisition and processing the data during the measurement. This means that the EPR type correlations reflect properties that are directly related with the physical reality but not with the observer’s knowledge about it. This entails the understanding that the wave function is related with objectively existing physical but not epistemological meaning.
- [53] The term “entanglement” is associated here with the appearance of the amplitude and phase/action in the coupled equations (12) and (13), as well as with the fact that the wave function incorporates both, the dynamical and the statistical characteristics through its phase and amplitude, respectively. It differs from the notion of entanglement between the wave functions of different quantum systems, as in the case of EPR correlations, Schrödinger’s cat states, etc.
- [54] A. Zellinger, R. Gähler, W. Treimer and W. Hampe, *Rev. Mod. Phys.* 60 (1988) 1067
- [55] T. S. Rose, M. J. Rosker, A. H. Zewail, *J. Chem. Phys.* 88 (1988) 6672.

Appendix A.

The phase/action is not simply one of the physical observables, but one may find a more distinguished role of this quantity in physics. In the wave-like phenomena, the phase/action $\Phi(t) = \hbar^{-1} \int_0^t H dt'$ / $S(t) = -\int_0^t H dt'$ of the state (the electric and magnetic field vectors, the wave function, etc.) of a physical object plays a crucial role, together with the amplitude. In the particle-like phenomena, the main role play the dynamical quantities, which arise from the action/phase by derivatives: velocity $\vec{v} = m^{-1} \nabla S$, linear momentum $\vec{p} = \nabla S$, the Lagrangian $\partial_t S = L$, Hamiltonian (energy, for conservative systems) $H = -\partial_t S$ ($E = -\partial_t S$), etc. The widely accepted understanding is that the wave-like picture (*e.g.*, the interference picture) and the particle-like picture (*e.g.*, the trajectory of the particle) are complementary. In fact, the interference phenomena are capable to reveal not only the wave-like features, but also some aspects of the particle-like features, for example, the localized positions of the particles. In the case of very low intensity fields, the interference picture is formed “point” by “point” after each particle (quantum of the field, or more complex object - atom, molecule, etc.) hits the phosphorescent screen [54], or other kind of detector. In this case, the particle manifests itself as a localized object, nevertheless that its actual dimensions is subject of additional investigation. On the other side, particle-like features, *e.g.*, trajectory, loose their definiteness in an experiment particularly designated to reveal the wave-like features of the object [13]. Once the interference picture is completed, it reveals the wave-like feature of the same physical object (more correctly – a large number of identical physical objects). It is naturally to expect that this also holds in the case of high-intensity fields or flux of particles, but, in that case, the process of formation of the interference picture cannot be as easily resolved in time as in the case of low-intensity fields. Hence, the interference phenomena are capable to reveal elements of both sides of the entire physical nature of the quantum systems and the phase/action plays a crucial role in its description.

The action/phase is an integral quantity that acquires contributions from all motions and interactions of the quantum system in an additive way to its initial value. Any motion, inertial or accelerated, of the quantum system makes definite contribution to the action/phase and it can be made observable in a suitably designed interference experiment in which the wave packet under consideration interferes with other, reference wave packet. From such a point of view, even the simple motion of the quantum system across the empty space seems not to be a trivial matter. Even for an isolated quantum system at rest, which does not interact with any other quantum system, its phase/action makes “idle” running in time thus acquiring a definite phase. The time rate of the action/phase acquisition, *i.e.*, energy/frequency, is ruled by the Hamiltonian of the free quantum system at rest, H_0 . Such a rate can be considered as a kind of *internal time* “clock” of the quantum system. The Hamiltonian H_0 is non-zero because of the internal motions and interactions of the building parts of the quantum system. This results in a non-zero action/phase, $S_0(t) = -\int_0^t H_0 dt'$, $\Phi_0(t) = \hbar^{-1} \int_0^t H_0 dt'$. In principle, the latter can be made observable if compare interferometrically with the phase of given reference state.

The action/phase stores the information from all events that have happened to the quantum system. This, in fact, constitutes the *physical history* of the quantum system. Due to the integral nature of the phase, it is not uniquely related to the corresponding physical history. Different systems having different histories may acquire same phase due to the difference in their Hamiltonians and/or the duration of their histories. Hence, the phase/action does not give detail information about the local/instantaneous behavior of the quantum system, as, *e.g.*, the linear momentum, energy, etc. do, although the “local” quantities can be found considering the rate of change of the phase/action. They, however, lose the relevant physical information carried by true constants, or functions not depending on the variable in the respective derivative. In the interference phenomena, it is the integral (phase/action) but not the local (energy, momentum) quantities that directly influence the observable results, the interference picture. Consequently, the material phase/action seems to encompass most completely the details of both, the particle-like and the wave-like, pictures of the physical reality.

Appendix B

Although the electromagnetic waves can well be described in terms of classical waves (at least for high-intensity fields), they are closer in some aspects to the matter (de Broglie) waves and they both strongly differ from the typical classical waves in the matter as, *e.g.*, elastic/sound waves. The elastic waves represent collective oscillations of a large number of “classical” particles with variable phase delay joined by elastic forces. Such waves cannot be formed if the number (number density) of the involved particles is too low, and, consequently, the elastic forces between the particles are negligible. Such kind of waves will be generally called *oscillating waves* and will be considered as classical waves. On the other side, the electromagnetic and the matter waves do not represent collective oscillations of particles but they arise from the specific way of propagation of particles or quanta of the respective fields (photons, electrons, neutrons, atoms, etc.). Such propagating particles are considered as waves in some phenomena *e.g.*, interference and diffraction, because they manifest a behavior that is similar to that of the usual oscillating waves at the same phenomena and, from given point of view, can be explained in terms of waves. In contrast to the oscillating waves, the electromagnetic and the matter waves retain their wavelike features at very low density of the propagating particles, including even the case of single particles. The interference picture in the latter case is formed after large number of consecutive events involving single particles. To explain the interference in that case, it is consider that each particle interfere by itself [1]. Such kind of waves will be generally called *particle waves*. While the wavelike behavior of the particle waves is well proved, the wavelike “displacement” (if in reality such a picture takes place at all) of the individual quanta of such waves has never been “seen” directly, partially because of the microscopic nature of the optical and the de-Broglie waves usually studied. By implementation of the cutting edge ultrafast laser technology (using high-intensity phase-locked femtosecond laser pulses and extreme ultraviolet single attosecond pulses), it has been shown (at least for high-intensity/high-number density fields) that the electrons, ionized by the attosecond pulse, have oscillation-like momentum distribution due to the electromagnetic field of the femtosecond pulse [24]. This, however, is not sufficient to deduce a definite conclusion about the particular way of motion of the electromagnetic field quanta. One might suppose from such an experiment that electromagnetic field quanta exert an oscillation-like force on the ionized electrons that results in an oscillation-like momentum distribution.

Appendix C

The results in Sec. IV follow from strict theoretical and experimental arguments. The forthcoming discussion is an attempt to show (at an intuitive level) that the concept of particle moving along a trajectory [5, 6] could be reconciled with some basic concepts in the quantum mechanics. The concept of trajectory is not accepted within the standard quantum mechanics due to the limitations on the simultaneous determination of the coordinates and the respective components of the linear momentum coming from the Heisenberg’s uncertainty relations. On the other hand, the concept of trajectory is intuitively and logically clear, has a macroscopic correspondence and seems to be one of the preconditions to set the question about the physical picture and, on that ground, the physical mechanism of the quantum phenomena. Apart from the qualitative and the quantitative consistency, the pictorial presentation of the physical processes and the relevant physical mechanism (that seem not to be within the main priorities of the present quantum mechanics) should be, to our opinion, the final goal of any physical description and it is the most significant difference between the physics and mathematics.

The physical state in the quantum mechanics is substantially distributed in space with no apparent indications of trajectories. The ordinary experiments seem to support such an understanding. In some phenomena with high-intensity fields, however, the concept of trajectory leads to excellent quantitative agreement with the exact quantum mechanical results, and, in addition, gives a clear pictorial explanation of the underlying physical mechanism [40]. To our opinion,

the lack of indications of trajectories in most of the cases could be a subject to suitable interpretation rather than to their real absence. Although in theory the state of a single quantum system can be studied within the Schrödinger equation, the typical characteristics of the quantum systems, microscopic size and huge number, impose that the experimental studies are usually performed within an ensemble of similar systems. Alternatively, superposition of a large number of experimental results with single quantum systems may also be used. For ergodic systems, both types of experiments must lead to same results that, in addition, cover the quantum mechanical predictions [51]. To our opinion, the quantum mechanical results (space distributed states and the associated characteristics) can be understood in terms of trajectories if one consider a superposition of all possible *real* trajectories (instead of imaginary quantum paths [6]), associated with the external motion of the quantum system, and/or the internal motion of its building parts, all of them subject to external perturbations. For example, the electron moves in an internal field of higher (atoms) or lower (molecules) symmetry, and a weak external perturbation (collision with other members of the ensemble, influence of external electromagnetic fields, including the background of zero point vacuum fluctuations), only, may become sufficient to change, at least, *within the symmetry limits of the state*, the trajectory orientation, apart from the orientation of the atom as a whole. These have randomization effect on the electron motion and may completely wash out the effect of the individual trajectories. At lack of angular momentum, the state will simply reflect the internal and external symmetry properties and may result in highly symmetric state, as in $1s$ ground state of the hydrogen atom. The non-zero angular momentum breaks the spherical symmetry and leads to particular spatial orientation, as in p, d , etc. atomic states. The states formed by such strongly randomized trajectories entail statistical description, as in the usual quantum mechanics. From the other side, a strong external field of particular polarization (comparable or stronger than the intra-atomic field) may peak up from the background of the randomized trajectories a relatively low number of electron trajectories of strongly dominating contribution. It enforces the particle-like picture and entails dynamical description. This allows reconciling the idea of locality, *i.e.*, particle-trajectory, with a nonlocal aspect of the quantum mechanics, *i.e.*, the distribution of the quantum state in space. The idea of trajectory can be brought to a quantitative level if one considers that the particles of the ensemble move along perturbed trajectories in a way that depicts the probability distribution $|\psi|^2$ ($|\sum_i \psi_i|^2$, for a superposition state) and the associated characteristics of the state if observed for long enough time. Then, the probability distribution in given space “point” can be related with the relative time spent by the particle in that “point”. The electron orbiting time, say, for the ground state of the hydrogen atom is about 150 attoseconds [24]. Hence, even for some typical microscopic time scales, *e.g.*, the lifetime of an electronic state, which is in the order of nanoseconds or much longer, may become long enough to depict the quantum mechanical probability distribution by means of electron motion along trajectory. Another problem is related with the “instantaneous” collapse of the wave function during the measurement. The collapse can be considered as a fast, but continuous in time, rearrangement of the way of motion (and the associated characteristics of the state) corresponding to the superposition state into a way of motion (and the associated characteristics) corresponding to the measured state.

It is not clear (for now) how to reconcile the local concept of particle moving along a trajectory with the substantially non-local effects as the interference of single particle (Feynman’s “which-way” interference experiment) and the EPR type correlations, where seem to be hidden the main mysteries in the quantum mechanics [13]. In any case, recent advances in the attosecond metrology [24-26] set the question of the real time dynamics of atomic/molecular electrons, as has been done some time ago with the nuclear motion inside the molecules [55]. It is important to emphasize that the MP causality established here is not related with the validity of the concept of trajectory. The latter would only help to build a clear pictorial presentation of the quantum phenomena. To what extend, if ever, such a concept can be forwarded to encompass entirely the quantum phenomena is a question addressed to the future.