

# Invariants of stationary AF-algebras and torsion subgroup of elliptic curves with complex multiplication

Igor Nikolaev \*

## Abstract

Let  $G_A$  be an  $AF$ -algebra given by periodic Bratteli diagram with the incidence matrix  $A \in GL(n, \mathbb{Z})$ . For a given polynomial  $p(x) \in \mathbb{Z}[x]$  we assign to  $G_A$  a finite abelian group  $Ab_{p(x)}(G_A) = \mathbb{Z}^n / p(A)\mathbb{Z}^n$ . It is shown that if  $p(0) = \pm 1$ , then  $Ab_{p(x)}(G_A)$  is an invariant of the strong stable isomorphism class of  $G_A$ . For  $n = 2$  and  $p(x) = x - 1$  we conjecture a formula linking values of the invariant and torsion subgroup of elliptic curves with complex multiplication.

*Key words and phrases:*  $AF$ -algebras, elliptic curves

*MSC:* 11G15 (complex multiplication); 46L85 (noncommutative topology)

## 1 Introduction

Let  $A \in GL(n, \mathbb{Z})$  be a non-negative integer matrix, which (if necessary) becomes strictly positive after a proper power of  $A$  is taken. We shall consider the following two objects, naturally attached to  $A$ . The first one, which we denote by  $(G_A, \sigma_A)$ , is a pair consisting of an  $AF$ -algebra,  $G_A$ , given by an infinite periodic Bratteli diagram with the incidence matrix  $A$  and a shift automorphism,  $\sigma_A$ , canonically attached to  $G_A$ . (The definitions of an  $AF$ -algebra, a Bratteli diagram and a shift automorphism are given in Section 2.)

---

\*Partially supported by NSERC.

The second object is an abelian group, which can be introduced as follows. Let  $p(x) \in \mathbb{Z}[x]$  be a polynomial over  $\mathbb{Z}$ , such that  $p(0) = \pm 1$ . Consider the following (finite) abelian group:

$$\mathbb{Z}^n / p(A)\mathbb{Z}^n := Ab_{p(x)}(G_A), \quad (1)$$

which we shall call an *abelianized*  $G_A$  at the polynomial  $p(x)$ . Recall that the *AF*-algebras  $G_A$  and  $G_{A'}$  are said to be stably isomorphic, whenever  $G_A \otimes \mathcal{K} \cong G_{A'} \otimes \mathcal{K}$ , where  $\mathcal{K}$  is the  $C^*$ -algebra of compact operators on a Hilbert space  $\mathcal{H}$ . We shall say that the *AF*-algebras  $G_A$  and  $G_{A'}$  are *strongly* stably isomorphic if they are stably isomorphic and  $\sigma_A, \sigma_{A'}$  are the conjugate shift automorphisms. The main result of the present note is the following theorem.

**Theorem 1** *For each polynomial  $p(x) \in \mathbb{Z}[x]$ , such that  $p(0) = \pm 1$ , the abelian group  $Ab_{p(x)}(G_A)$  is an invariant of the strong stable isomorphism class of the *AF*-algebra  $G_A$ .*

Let  $E_{CM}$  be elliptic curve with complex multiplication by an order of conductor  $f \geq 1$  in imaginary quadratic field  $\mathbb{Q}(\sqrt{-d})$ , where  $d \neq 1$  [8], p. 96. Consider a periodic continued fraction  $f\omega = [a_0, \overline{a_1, \dots, a_n}]$ , where  $\omega = \frac{1+\sqrt{d}}{2}$  if  $d \equiv 1 \pmod{4}$  and  $\omega = \sqrt{d}$  if  $d \equiv 2, 3 \pmod{4}$ . We shall introduce an integer matrix  $A = \prod_{i=1}^n \begin{pmatrix} a_i & 1 \\ 1 & 0 \end{pmatrix}$ , see Section 4.1 for a motivation.

**Conjecture 1** *For each  $E_{CM}$  there exists an imaginary number field  $K$  such that  $E_{CM} \cong E(K)$  and a twist of  $E(K)$  such that  $E_{tors}(K) \cong Ab_{x-1}(G_A)$ , where  $E_{tors}(K)$  is the torsion subgroup of  $E(K)$ .*

The note is organized as follows. The preliminary facts are brought together in Section 2. Theorem 1 is proved in Section 3. In Section 4 conjecture 1 is explained and some examples are given.

## 2 Preliminaries

An *AF*-algebra (approximately finite-dimensional  $C^*$ -algebra) is defined to be the norm closure of an ascending sequence of the finite-dimensional  $C^*$ -algebras  $M_n$ 's, where  $M_n$  is the  $C^*$ -algebra of the  $n \times n$  matrices with the entries in  $\mathbb{C}$ . Here the index  $n = (n_1, \dots, n_k)$  represents a semi-simple matrix

algebra  $M_n = M_{n_1} \oplus \dots \oplus M_{n_k}$ . The ascending sequence mentioned above can be written as  $M_1 \xrightarrow{\varphi_1} M_2 \xrightarrow{\varphi_2} \dots$ , where  $M_i$  are the finite dimensional  $C^*$ -algebras and  $\varphi_i$  the homomorphisms between such algebras. The set-theoretic limit  $A = \lim M_n$  has a natural algebraic structure given by the formula  $a_m + b_k \rightarrow a + b$ ; here  $a_m \rightarrow a, b_k \rightarrow b$  for the sequences  $a_m \in M_m, b_k \in M_k$ . The homomorphisms  $\varphi_i$  can be arranged into a graph as follows. Let  $M_i = M_{i_1} \oplus \dots \oplus M_{i_k}$  and  $M_{i'} = M_{i'_1} \oplus \dots \oplus M_{i'_k}$  be the semi-simple  $C^*$ -algebras and  $\varphi_i : M_i \rightarrow M_{i'}$  the homomorphism. One has the two sets of vertices  $V_{i_1}, \dots, V_{i_k}$  and  $V_{i'_1}, \dots, V_{i'_k}$  joined by the  $a_{rs}$  edges, whenever the summand  $M_{i_r}$  contains  $a_{rs}$  copies of the summand  $M_{i'_s}$  under the embedding  $\varphi_i$ . As  $i$  varies, one obtains an infinite graph called a *Bratteli diagram* of the  $AF$ -algebra. The Bratteli diagram defines a unique  $AF$ -algebra.

If the homomorphisms  $\varphi_1 = \varphi_2 = \dots = \text{Const}$  in the definition of the  $AF$ -algebra  $A$ , the  $AF$ -algebra  $A$  is called *stationary*. The Bratteli diagram of a stationary  $AF$ -algebra looks like a periodic graph with the incidence matrix  $A = (a_{rs})$  repeated over and over again. Since matrix  $A$  is a non-negative integer matrix, one can take a power of  $A$  to obtain a strictly positive integer matrix – which we always assume to be the case. We shall denote the above  $AF$ -algebra by  $G_A$ . Recall that in the case of  $AF$ -algebras, the abelian monoid  $V_{\mathbb{C}}(\mathcal{A})$  defines the  $AF$ -algebra up to an isomorphism and is known as a *dimension group* of  $\mathcal{A}$ . We shall use a standard dictionary existing between the  $AF$ -algebras and their dimension groups [6], Section 7.3. Instead of dealing with the  $AF$ -algebra  $G_A$ , we shall work with its dimension group  $(G, G_A^+)$ , where  $G \cong \mathbb{Z}^n$  is the lattice and  $G_A^+$  is a positive cone inside the lattice given by a sequence of the simplicial dimension groups:

$$\mathbb{Z}^n \xrightarrow{A} \mathbb{Z}^n \xrightarrow{A} \mathbb{Z}^n \xrightarrow{A} \dots \quad (2)$$

(Equivalently, one can talk about the  $K_0$ -group of  $G_A$ ; in such terms,  $K_0(G_A) \cong (G_A, G_A^+)$ , see [6], p.122 for details.) There exists a natural automorphism,  $\sigma_A$ , of the dimension group  $(G_A, G_A^+)$ , see [1], p.37. It can be defined as follows. Let  $\lambda_A > 1$  be the Perron-Frobenius eigenvalue and  $v_A \in \mathbb{R}_+^n$  the corresponding eigenvector of the matrix  $A$ . It is known that  $G_A^+$  is defined by the inequality  $\mathbb{Z}v_A^{(1)} + \dots + \mathbb{Z}v_A^{(n)} \geq 0$ . One can multiply the basis  $\{v_A^{(1)}, \dots, v_A^{(n)}\}$  of the above  $\mathbb{Z}$ -module by  $\lambda_A$ . It is easy to see that such a multiplication defines an automorphism of the dimension group  $(G_A, G_A^+)$ . The automorphism is called a *shift automorphism* and denoted by  $\sigma_A$ . The shift automorphisms  $\sigma_A, \sigma_{A'}$  are said to be conjugate, if  $\sigma_A \circ \theta = \theta \circ \sigma_{A'}$  for any order-isomorphism

$\theta$  between the dimension groups  $(G_A, G_A^+)$  and  $(G_{A'}, G_{A'}^+)$ . We shall write this fact as  $(G_A, \sigma_A) \cong (G_{A'}, \sigma_{A'})$  (an isomorphism).

**Lemma 1** *The pairs  $(G_A, \sigma_A)$  and  $(G_{A'}, \sigma_{A'})$  are isomorphic if and only if the matrices  $A$  and  $A'$  are similar.*

*Proof.* By Theorem 6.4 of [1],  $(G_A, \sigma_A) \cong (G_{A'}, \sigma_{A'})$  if and only if the matrices  $A$  and  $A'$  are shift equivalent, see [9] for a definition of the shift equivalence. On the other hand, since the matrices  $A$  and  $A'$  are unimodular, the shift equivalence between  $A$  and  $A'$  coincides with a similarity of the matrices in the group  $GL(n, \mathbb{Z})$ , see Corollary 2.13 of [9].  $\square$

**Corollary 1** *The AF-algebras  $G_A$  and  $G_{A'}$  are strongly stably isomorphic if and only if the matrices  $A$  and  $A'$  are similar.*

*Proof.* By a dictionary between the dimension groups and AF-algebras, the order-isomorphic dimension groups correspond to the stably isomorphic AF-algebras, see Theorem 2.3. of [1]. Since  $\sigma_A$  and  $\sigma_{A'}$  are conjugate, one gets a strong stable isomorphism.  $\square$

### 3 Proof of theorem 1

Our proof is based on the following criterion ([1], Theorem 6.4): the dimension groups

$$\mathbb{Z}^n \xrightarrow{A} \mathbb{Z}^n \xrightarrow{A} \mathbb{Z}^n \xrightarrow{A} \dots \quad \text{and} \quad \mathbb{Z}^n \xrightarrow{A'} \mathbb{Z}^n \xrightarrow{A'} \mathbb{Z}^n \xrightarrow{A'} \dots \quad (3)$$

are order-isomorphic and  $\sigma_A, \sigma_{A'}$  are conjugate *iff* the matrices  $A$  and  $A'$  are similar in the group  $GL(n, \mathbb{Z})$ , i.e.  $A' = BAB^{-1}$  for a  $B \in GL(n, \mathbb{Z})$ . The rest of the proof follows from the structure theorem for the finitely generated modules given by the matrix  $A$  over a principal ideal domain, see e.g. [7], p. 43. The result says the normal form of the module (in our case – over  $\mathbb{Q}[x]$ ) is independent of the particular choice of a matrix in the similarity class of  $A$ . The normal form, evaluated at  $x = 0$ , coincides with the invariant  $Ab_{p(x)}(G_A)$ . Let us pass to a step by step argument.

Let  $\mathbb{Q}[x]$  be the ring of polynomials in one variable over the rational numbers. Since  $\mathbb{Q}$  is a field, the ring  $\mathbb{Q}[x]$  is a principal ideal domain, see e.g. [7], p.26. We shall consider the following  $\mathbb{Q}[x]$ -module. If  $A \in M_n(\mathbb{Z})$  is an

$n \times n$  integer matrix, one endows the abelian group  $\mathbb{Z}^n$  with a  $\mathbb{Q}[x]$ -module structure by defining:

$$p_n(x)v = (p_n(A) \pm I)v, \quad p_n(x) \in \mathbb{Q}[x], v \in \mathbb{Z}^n, \quad (4)$$

where  $I$  is the identity matrix. Note that the obtained module depends on the matrix  $A$  and we write it as  $(\mathbb{Z}^n)^A$ .

Fix a basis  $\{\varepsilon_1, \dots, \varepsilon_n\}$  in  $(\mathbb{Z}^n)^A$ . Let  $p(x) \in \mathbb{Q}[x]$  be a polynomial, such that  $p(0) = \pm 1$ . A set of generators  $\{p(x)\varepsilon_1, \dots, p(x)\varepsilon_n\}$  defines a submodule of  $(\mathbb{Z}^n)^A$ , which we identify with the kernel of a homomorphism  $\phi_{p(x)} : (\mathbb{Z}^n)^A \rightarrow \mathbb{Z}^n$ . A simple calculation gives us:

$$p(x)\varepsilon_j = (p(A) \pm I)\varepsilon_j = p(A)\varepsilon_j \pm \varepsilon_j. \quad (5)$$

Since  $p(0) = \pm 1$ , it follows from the last equation that  $p(A)\varepsilon_j = 0$ . Thus, one obtains a set of relations between the generators of the module  $(\mathbb{Z}^n)^A$ . The above relations depend on the polynomial  $p(x)$  and define completely the structure of a  $\mathbb{Q}[x]$ -module

$$(\mathbb{Z}^n)_{p(x)}^A := \langle \varepsilon_1, \dots, \varepsilon_n \mid p(A)\varepsilon_j = 0 \rangle = \mathbb{Z}^n / \text{Ker } \phi_{p(x)} \cong \mathbb{Z}^n / p(A)\mathbb{Z}^n. \quad (6)$$

Let  $A'$  be a matrix similar to  $A$ , i.e.  $A' = BAB^{-1}$  for a matrix  $B \in GL(n, \mathbb{Z})$ . By a basic property of the finitely generated modules over a principal ideal domain, the modules  $(\mathbb{Z}^n)_{p(x)}^A$  and  $(\mathbb{Z}^n)_{p(x)}^{A'}$  are isomorphic to the same direct sum of the cyclic modules [7], p.43. Namely, the relations  $p(A)\varepsilon_j = 0, j = 1, \dots, n$  can be written as:

$$\begin{pmatrix} g_{11} & \cdots & g_{1n} \\ \vdots & & \vdots \\ g_{n1} & \cdots & g_{nn} \end{pmatrix} \begin{pmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}, \quad (7)$$

where  $(g_{ij}) = p(A) = b_n A^n + \dots + b_1 A \pm I$ . It is well known that by the elementary transformations (the Euclidean algorithm) consisting of (i) an interchange of two rows, (ii) a multiplication of a row by  $-1$ , (iii) addition a multiple of one row to another and similar operations on columns, brings the matrix  $(g_{ij})$  to a diagonal form (the Smith form):

$$D = \begin{pmatrix} g_1 & & & & & \\ & \ddots & & & & \\ & & g_r & & & \\ & & & 0 & & \\ & & & & \ddots & \\ & & & & & 0 \end{pmatrix}, \quad (8)$$

where  $g_i$  are positive integers, such that  $g_i \mid g_{i+1}$ . The elementary transformations are equivalent to a matrix equation  $D = PGQ$ , where  $G = (g_{ij})$  and  $P, Q \in GL(n, \mathbb{Z})$ . Thus, the module  $(\mathbb{Z}^n)_{p(x)}^A$ , evaluated at the point  $x = 0$ , can be written as:

$$(\mathbb{Z}^n)_{p(0)}^A \cong \mathbb{Z}_{g_1} \oplus \dots \oplus \mathbb{Z}_{g_r} \oplus \mathbb{Z}^{n-r}, \quad (9)$$

where  $\mathbb{Z}_{g_i} = \mathbb{Z}/g_i\mathbb{Z}$ . The same set of integers  $g_i$  will appear in the diagonal form of the matrix  $A'$ . Thus,  $Ab_{p(x)}(G_A) \cong Ab_{p(x)}(G_{A'})$  for every choice of the polynomial  $p(x) \in \mathbb{Z}[x]$ . (In the practical considerations, we often have  $r = n$  so that our invariant is a finite abelian group. It is easy to see that this is always the case, provided  $G_A$  is a simple  $AF$ -algebra.) The theorem 1 follows now from the corollary 1.

The most important special case of the above invariant is the case when  $p(x) = x - 1$  (the Bowen-Franks invariant). The invariant takes the form:

$$Ab_{x-1}(G_A) = \mathbb{Z}^n / (A - I)\mathbb{Z}^n. \quad (10)$$

The Bowen-Franks invariant is covered extensively in the literature and has a geometric meaning of tracking an algebraic structure of the periodic points of an automorphism of the lattice  $\mathbb{Z}^n$  defined by the matrix  $A$ . In particular, the cardinality of the group  $Ab_{x-1}(G_A)$  is equal to the total number of the isolated fixed points of the automorphism  $A$ . It is easy to see, that such a number coincides with  $|\det(A - I)|$ .  $\square$

## 4 Torsion conjecture

An introduction to elliptic curves, complex multiplication, etc. can be found in [8]; torsion of rational elliptic curves with complex multiplication was studied in [4]. A link between complex multiplication and  $G_A$  was the subject of [3].

### 4.1 Teichmüller functor

Let  $\theta \in [0, 1)$  be an irrational number. The universal  $C^*$ -algebra  $\mathcal{A}_\theta$  generated by the unitaries  $u$  and  $v$  satisfying the commutation relation  $vu = e^{2\pi i\theta}uv$  is called a *noncommutative torus* [5]. A pair of such tori is said to be stably isomorphic (Morita equivalent) whenever  $\mathcal{A}_\theta \otimes \mathcal{K} \cong \mathcal{A}_{\theta'} \otimes \mathcal{K}$ , where  $\mathcal{K}$

is the  $C^*$ -algebra of compact operators. The  $\mathcal{A}_\theta$  is stably isomorphic to  $\mathcal{A}_{\theta'}$  if and only if  $\theta' = (a\theta + b)/(c\theta + d)$ , where  $a, b, c, d \in \mathbb{Z}$  and  $ad - bc = 1$ . The K-theory of  $\mathcal{A}_\theta$  is Bott periodic with  $K_0(\mathcal{A}_\theta) = K_1(\mathcal{A}_\theta) \cong \mathbb{Z}^2$ . The range of trace on projections of  $A_\theta \otimes \mathcal{K}$  is a subset  $\Lambda = \mathbb{Z} + \mathbb{Z}\theta$  of the real line; the set  $\Lambda \cong K_0(\mathcal{A}_\theta)$  is known as a pseudo-lattice [2]. The noncommutative torus  $\mathcal{A}_\theta$  is said to have *real multiplication*, if  $\theta$  is a quadratic irrationality; we denote such an algebra by  $\mathcal{A}_{RM}$ . Real multiplication implies non-trivial endomorphisms of the pseudo-lattice  $\Lambda_{RM}$  given as a multiplication by real numbers – hence the name. Such endomorphisms make a ring under addition and composition of the endomorphisms; the latter is isomorphic to an order of conductor  $f \geq 1$  in the ring of integers of quadratic field  $\mathbb{Q}(\theta)$ . Recall that each order of  $\mathbb{Q}(\sqrt{d})$  has the form  $\mathbb{Z} + (f\omega)\mathbb{Z}$ , where  $\omega = \frac{1+\sqrt{d}}{2}$  if  $d \equiv 1 \pmod{4}$  and  $\omega = \sqrt{d}$  if  $d \equiv 2, 3 \pmod{4}$ . It is known that continued fraction of  $\theta = f\omega$  is periodic and has the form  $[a_0, \overline{a_1, \dots, a_n}]$ ; we shall consider a matrix  $A = \prod_{i=1}^n \begin{pmatrix} a_i & 1 \\ 1 & 0 \end{pmatrix}$ .

**Lemma 2** *The  $C^*$ -algebras  $G_A$  and  $\mathcal{A}_{RM}$  are stably isomorphic.*

*Proof.* It follows easily from definition of  $A$ , that  $K_0(G_A) \cong \mathbb{Z} + \mathbb{Z}\theta'$ , where  $\theta' = \theta - a_0$ . Thus,  $K_0(G_A) \cong K_0(\mathcal{A}_{RM})$  and, therefore,  $G_A \otimes \mathcal{K} \cong \mathcal{A}_{RM} \otimes \mathcal{K}$ .  $\square$

Let  $\mathbb{H} = \{x + iy \in \mathbb{C} \mid y > 0\}$  be the upper half-plane and for  $\tau \in \mathbb{H}$  let  $\mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$  be a complex torus; we routinely identify the latter with a non-singular elliptic curve via the Weierstrass  $\wp$  function [8], pp. 6-7. Recall that two complex tori are isomorphic, whenever  $\tau' = (a\tau + b)/(c\tau + d)$ , where  $a, b, c, d \in \mathbb{Z}$  and  $ad - bc = 1$ . If  $\tau$  is an imaginary quadratic number, elliptic curve is said to have *complex multiplication*; we shall denote such curves by  $\mathcal{E}_{CM}$ . Complex multiplication means that lattice  $L = \mathbb{Z} + \mathbb{Z}\tau$  admits non-trivial endomorphisms given as multiplication of  $L$  by certain complex (quadratic) numbers. Again, such endomorphisms make a ring under addition and composition of the endomorphisms; the latter is isomorphic to an order of conductor  $f \geq 1$  in the ring of integers of imaginary quadratic field  $\mathbb{Q}(\tau)$ .

Our calculations of torsion are based on a covariant functor between elliptic curves and noncommutative tori; roughly speaking, the functor maps isomorphic curves to the stably isomorphic tori. To give an idea, let  $\phi$  be a closed form on a topological torus; the trajectories of  $\phi$  define a measured

foliation on the torus. By the Hubbard-Masur theorem, such a foliation corresponds to a point  $\tau \in \mathbb{H}$ . The map  $F : \mathbb{H} \rightarrow \partial\mathbb{H}$  is defined by the formula  $\tau \mapsto \theta = \int_{\gamma_2} \phi / \int_{\gamma_1} \phi$ , where  $\gamma_1$  and  $\gamma_2$  are generators of the first homology of the torus. The following is true: (i)  $\mathbb{H} = \partial\mathbb{H} \times (0, \infty)$  is a trivial fiber bundle, whose projection map coincides with  $F$ ; (ii)  $F$  is a functor, which maps isomorphic complex tori to the stably isomorphic noncommutative tori. We shall refer to  $F$  as the *Teichmüller functor*. Remarkably, functor  $F$  maps  $E_{CM}$  to  $\mathcal{A}_{RM}$  [3]; more specifically, complex multiplication by order of conductor  $f$  in imaginary field  $\mathbb{Q}(\sqrt{-d})$  goes to real multiplication by an order of conductor  $f$  in the real field  $\mathbb{Q}(\sqrt{d})$ <sup>1</sup>.

## 4.2 Numerical examples

We conclude by examples supporting conjecture 1; they cover all rational  $E_{CM}$  [4], except  $d = -1$  and  $d = -163$ .

---

<sup>1</sup>This fact follows easily from an explicit formula for  $F$  [3], p.524.

$-d$	$f$	$E_{tors}(\mathbb{Q})$ , see [4] p.196	continued fraction of $\sqrt{f^2 d}$	$A$	$Ab_{x-1}(G_A)$
-2	1	$\mathbb{Z}_2$	$[1, \overline{2}]$	$\begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}$	$\mathbb{Z}_2$
-3	1	$\mathbb{Z}_1$ or $\mathbb{Z}_2$	$[1, \overline{1, 2}]$	$\begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix}$	$\mathbb{Z}_2$
-7	1	$\mathbb{Z}_2$	$[2, \overline{1, 1, 1, 4}]$	$\begin{pmatrix} 14 & 3 \\ 9 & 2 \end{pmatrix}$	$\mathbb{Z}_{14}$
-11	1	$\mathbb{Z}_1$	$[3, \overline{3, 6}]$	$\begin{pmatrix} 19 & 3 \\ 6 & 1 \end{pmatrix}$	$\mathbb{Z}_3 \oplus \mathbb{Z}_6$
-19	1	$\mathbb{Z}_1$	$[4, \overline{2, 1, 3, 1, 2, 8}]$	$\begin{pmatrix} 326 & 39 \\ 117 & 14 \end{pmatrix}$	$\mathbb{Z}_{13} \oplus \mathbb{Z}_{26}$
-43	1	$\mathbb{Z}_1$	$[6, \overline{1, 1, 3, 1, 5, 1, 3, 1, 1, 12}]$	$\begin{pmatrix} 6668 & 531 \\ 3717 & 296 \end{pmatrix}$	$\mathbb{Z}_{59} \oplus \mathbb{Z}_{118}$
-67	1	$\mathbb{Z}_1$	$[8, \overline{5, 2, 1, 1, 7, 1, 1, 2, 5, 16}]$	$\begin{pmatrix} 96578 & 5967 \\ 17901 & 1106 \end{pmatrix}$	$\mathbb{Z}_{221} \oplus \mathbb{Z}_{442}$
-3	2	$\mathbb{Z}_2$ or $\mathbb{Z}_6$	$[3, \overline{2, 6}]$	$\begin{pmatrix} 13 & 2 \\ 6 & 1 \end{pmatrix}$	$\mathbb{Z}_2 \oplus \mathbb{Z}_6$
-7	2	$\mathbb{Z}_2$	$[5, \overline{3, 2, 3, 10}]$	$\begin{pmatrix} 247 & 24 \\ 72 & 7 \end{pmatrix}$	$\mathbb{Z}_6 \oplus \mathbb{Z}_{42}$
-3	3	$\mathbb{Z}_1$	$[5, \overline{5, 10}]$	$\begin{pmatrix} 51 & 5 \\ 10 & 1 \end{pmatrix}$	$\mathbb{Z}_5 \oplus \mathbb{Z}_{10}$

**Remark 1** Note that  $E_{tors}(\mathbb{Q}) \subseteq E_{tors}(K)$  since  $K$  is a non-trivial extension of  $\mathbb{Q}$ . The third column lists all twists of  $E(\mathbb{Q})$  satisfying conjecture 1.

## References

- [1] E. G. Effros, Dimensions and  $C^*$ -Algebras, in: Conf. Board of the Math. Sciences, Regional conference series in Math., No.46, AMS, 1981.
- [2] Yu. I. Manin, Real multiplication and noncommutative geometry, in "Legacy of Niels Hendrik Abel", 685-727, Springer, 2004; arXiv:math/0202109.
- [3] I. V. Nikolaev, Remark on the rank of elliptic curves, Osaka J. Math. 46 (2009), 515-527.

- [4] L. D. Olson, Points of finite order on elliptic curves with complex multiplication, *Manuscripta Math.* 14 (1974), 195-205.
- [5] M. A. Rieffel, Non-commutative tori – a case study of non-commutative differentiable manifolds, *Contemp. Math.* 105 (1990), 191-211. Available <http://math.berkeley.edu/~rieffel/>
- [6] M. Rørdam, F. Larsen and N. Laustsen, An introduction to  $K$ -theory for  $C^*$ -algebras. London Mathematical Society Student Texts, 49. Cambridge University Press, Cambridge, 2000.
- [7] I. R. Shafarevich, Basic Notions of Algebra, in: *Algebra I*, E.M.S, vol. 11, Springer, 1990.
- [8] J. H. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, GTM 151, Springer 1994.
- [9] J. B. Wagoner, Strong shift equivalence theory and the shift equivalence problem, *Bull. Amer. Math. Soc.* 36 (1999), 271-296.

THE FIELDS INSTITUTE FOR MATHEMATICAL SCIENCES, TORONTO,  
ON, CANADA, E-MAIL: [igor.v.nikolaev@gmail.com](mailto:igor.v.nikolaev@gmail.com)

*Current address: 616-315 Holmwood Ave., Ottawa, ON, Canada, K1S  
2R2*