Continuous Wavelets on Compact Manifolds

Daryl Geller

Department of Mathematics, Stony Brook University, Stony Brook, NY 11794-3651

daryl@math.sunysb.edu

*Azita Mayeli

Department of Mathematics, Stony Brook University, Stony Brook, NY 11794-3651

amayeli@math.sunysb.edu

June 5, 2018

Abstract

Let \mathbf{M} be a smooth compact oriented Riemannian manifold, and let $\Delta_{\mathbf{M}}$ be the Laplace-Beltrami operator on \mathbf{M} . Say $0 \neq f \in \mathcal{S}(\mathbb{R}^+)$, and that f(0) = 0. For t > 0, let $K_t(x, y)$ denote the kernel of $f(t^2\Delta_{\mathbf{M}})$. We show that K_t is well-localized near the diagonal, in the sense that it satisfies estimates akin to those satisfied by the kernel of the convolution operator $f(t^2\Delta)$ on \mathbb{R}^n . We define continuous \mathcal{S} -wavelets on \mathbf{M} , in such a manner that $K_t(x, y)$ satisfies this definition, because of its localization near the diagonal. Continuous \mathcal{S} -wavelets on \mathbf{M} are analogous to continuous wavelets on \mathbb{R}^n in $\mathcal{S}(\mathbb{R}^n)$. In particular, we are able to characterize the Hölder continuous functions on \mathbf{M} by the size of their continuous \mathcal{S} -wavelet transforms, for Hölder exponents strictly between 0 and 1. If \mathbf{M} is the torus \mathbb{T}^2 or the sphere S^2 , and $f(s) = se^{-s}$ (the "Mexican hat" situation), we obtain two explicit approximate formulas for K_t , one to be used when t is large, and one to be used when t is small.

Keywords and phrases: Frames, Wavelets, Continuous Wavelets, Spectral Theory, Schwartz Functions, Time-Frequency Analysis, Manifolds, Sphere, Torus, Pseudodifferential Operators, Höder Spaces.

AMS Classification; 42C40, 42B20, 58J40, 58J35, 35P05.

Table of Contents

- Section 1: Introduction and Historical Comments
- Section 2: Applying the Spectral Theorem
- Section 3: Preliminaries on Manifolds
- Section 4: Kernels
- Section 5: Continuous S-Wavelets on Manifolds
- Section 6: Homogeneous Manifolds
- Section 7: A Technical Lemma

^{*}This work was partially supported by the Marie Curie Excellence Team Grant MEXT-CT-2004-013477, Acronym MAMEBIA.

1 Introduction

Let $\mathcal{S}(\mathbb{R}^+)$ denote the space of restrictions to \mathbb{R}^+ of functions in $\mathcal{S}(\mathbb{R})$. Say $0 \neq f_0 \in \mathcal{S}(\mathbb{R}^+)$, and let

$$f(s) = sf_0(s).$$

One then has the *Calderón formula* for f: if $c \in (0, \infty)$ is defined by

$$c = \int_0^\infty |f(t)|^2 \frac{dt}{t} = \int_0^\infty t |f_0(t)|^2 dt$$

then for all s > 0,

$$\int_0^\infty |f(ts)|^2 \frac{dt}{t} = c < \infty.$$
(1)

The even function $h(\xi) = f(\xi^2)$ is then in $\mathcal{S}(\mathbb{R})$ and satisfies

$$\int_0^\infty |h(t\xi)|^2 \frac{dt}{t} = \frac{c}{2} < \infty.$$
⁽²⁾

(In fact, all even functions in $\mathcal{S}(\mathbb{R})$ satisfying (2) arise in this manner).

Its inverse Fourier transform $\psi = \check{h}$ is admissible (i.e. is a continuous wavelet). That is, for some c' > 0,

$$\int_{0}^{\infty} \|F * \psi_t\|_2^2 \frac{dt}{t} = c' \|F\|_2^2 \tag{3}$$

for all $F \in L^2(\mathbb{R}^n)$. Here, as usual, $\psi_t(x) = t^{-1}\psi(x/t)$.

We prefer to write, formally,

$$\check{h} = f(-d^2/dx^2)\delta;$$

the formal justfication being that

$$(f(-d^2/dx^2)\delta)^{\hat{}} = f(\xi^2) = h(\xi)$$

Thus $f(-d^2/dx^2)\delta$ is a continuous wavelet on \mathbb{R} .

Our program is to construct analogues of continuous wavelets, on much more general spaces, by replacing the positive number s in (1) by a positive self-adjoint operator T on a Hilbert space \mathcal{H} . If P is the projection onto the null space of T, by the spectral theorem we obtain the relation

$$\int_{0}^{\infty} |f|^{2} (tT) \frac{dt}{t} = c(I - P), \tag{4}$$

where the integral in (4) converges strongly. (4) will be justified in the next section. Taking T to be $-d^2/dx^2$ on \mathbb{R} leads to the continuous wavelet $f(-d^2/dx^2)\delta$ on \mathbb{R} . (Of course, on \mathbb{R} , P = 0.)

We began our program of looking at more general positive self-adjoint operators T, in order to construct continuous wavelets, in our article [17]. There we took T to be the sublaplacian L on $L^2(G)$, where G is a stratified Lie group, and thereby obtained continuous wavelets and frames on such G. In fact, in that context, $f(L)\delta$ was a continuous wavelet. (Again, in that context, P was zero.) (Our article [17] was motivated by the second author's thesis [32], where it was shown that if $f(s) = se^{-s}$, then the "Mexican hat" $f(L)\delta$ is a continuous wavelet on the Heisenberg group.)

In this article we will look at the (much more practical!) situation in which T is the Laplace-Beltrami operator on $L^2(\mathbf{M})$, where \mathbf{M} is a smooth compact oriented Riemannian manifold without boundary. We will construct analogues of continuous Schwartz wavelets in this context, and will obtain explicit formulas for them if \mathbf{M} is the sphere or the torus. In a sequel article ([18]) we shall use a discrete analogue of (4) to construct nearly tight frames in this situation, and show that they are appropriately well-localized (specifically, that they satisfy a space-frequency analysis). In both of these articles, P will be the projection onto the one-dimensional space of constant functions.

We now summarize our results and methods. We discuss prior work on wavelets on manifolds, especially the work of Narcowich, Petrushev and Ward on the sphere [35], [36], at the end of this introduction.

In the model situation (1) – (3) on \mathbb{R} , note that the kernel of the operator $F \to F * \psi_t$ is $K_t(x, y) := t^{-1}\psi((x-y)/t)$. Since h(0) = 0, we have $\int K_t(x, y)dx = 0$ for all y and $\int K_t(x, y)dy = 0$ for all x. Also $K_t(x, y)$ is smooth in t, x, y for t > 0.

Motivated by this model case, we define continuous wavelets on \mathbf{M} as follows. Suppose that the function $K_t(x, y)$ is smooth for t > 0, $x, y \in \mathbf{M}$. For t > 0, define $T_t : L^2(\mathbf{M}) \to C^{\infty}(\mathbf{M})$ to be the operator with kernel K_t , We define $K_t(x, y)$ to be a *continuous wavelet* on \mathbf{M} , provided that for some c > 0,

$$\int_0^\infty \|T_t F\|_2^2 \frac{dt}{t} = c \|(I - P)F\|_2^2,$$
(5)

for all $F \in L^2(\mathbf{M})$, and that for any t > 0, $\int_{\mathbf{M}} K_t(x, y) d\mu(x) = 0$ for all y and $\int_{\mathbf{M}} K_t(x, y) d\mu(y) = 0$ for all x. (Here μ is the measure on \mathbf{M} arising from integration with respect to the volume form.)

It is then easy to see that, if K_t is the kernel of $f(t^2\Delta)$ (f as before), where $\Delta = \Delta_{\mathbf{M}}$ is the Laplace-Beltrami operator on \mathbf{M} , then $K_t(x, y)$ is a continuous wavelet on \mathbf{M} . In particular, (5) follows easily from (4), simply by applying both sides of (4) to F, taking the inner product of both sides with F, and making the change of variables which replaces t by t^2 in the integral. Note that $K_t(x, y) = [f(t^2\Delta)\delta_y](x)$, which is analogous to $f(-d^2/dx^2)\delta$ being a continuous wavelet on \mathbb{R} .

This then gives an L^2 theory of continuous wavelets on manifolds. However, in practice, one wishes to go beyond this theory, looking at continuous wavelets on other function spaces, and discretizing them to obtain frames. For this, one needs the wavelets to have further properties. Fortunately, as we shall see, if K_t is the kernel of $f(t^2\Delta)$, these properties are present.

Specifically, let us return to our model situation; let us however work on \mathbb{R}^n , not just \mathbb{R}^1 . Then $K_t(x, y)$, the kernel of $f(t^2\Delta)$, would be of the form $t^{-n}\psi((x-y)/t)$ for some $\psi \in S$. (Here Δ is the usual Laplacian on \mathbb{R}^n , and $\hat{\psi} = G$, where $G(\xi) = f(|\xi|^2)$.) For any N, α, β , there would thus exist $C_{N,\alpha,\beta}$ such that

$$t^{n+|\alpha|+|\beta|} \left| \left(\frac{x-y}{t} \right)^N \partial_x^{\alpha} \partial_y^{\beta} K_t(x,y) \right| \le C_{N,\alpha,\beta}$$

for all t, x, y. Such estimates are essential in the theory of wavelets on \mathbb{R}^n . We therefore make the following definition:

Definition 1.1. Let $K_t(x, y)$ be a continuous wavelet on **M**. Then we say that $K_t(x, y)$ is a continuous S-wavelet on **M** provided that:

For every pair of C^{∞} differential operators X (in x) and Y (in y) on **M**, and for every integer $N \ge 0$, there exists $C_{N,X,Y}$ as follows. Suppose deg X = j and deg Y = k. Then

$$t^{n+j+k} \left| \left(\frac{d(x,y)}{t} \right)^N XYK_t(x,y) \right| \le C_{N,X,Y}$$
(6)

for all t > 0 and all $x, y \in \mathbf{M}$.

Here d is the geodesic distance on **M**. We use the terminology "continuous S-wavelet" to express the idea that these wavelets are analogous to Schwartz wavelets on \mathbb{R}^n .

In Lemma 4.1, we will show the following key result:

Lemma 4.1 If K_t is the kernel of $f(t^2\Delta)$ (f as before), then $K_t(x, y)$ satisfies (6) (and hence is a continuous *S*-wavelet on **M**).

Our proof of this lemma uses the theory of pseudodifferential operators (most crucially, a result of Strichartz [43]), and Huygens' principle.

In Theorem 5.5 we shall show that continuous S-wavelets are well adapted to the study of certain other function spaces. Specifically, we shall show the following generalization of a theorem of Holschneider and Tchamitchian ([25]), who worked on the real line, and characterized the wavelet transforms of the spaces of Hölder continuous functions (for Hölder exponent strictly between 0 and 1).

Theorem 5.5

Let $K_t(x, y)$ be a continuous S-wavelet on \mathbf{M} , and, for t > 0, let T_t be the operator on L^2 with kernel K_t . Suppose $F \in L^2(\mathbf{M})$. Then:

(a) If F is Hölder continuous, with Hölder exponent α ($0 < \alpha \leq 1$), then for some C > 0,

$$\|T_t F\| \le C t^{\alpha} \tag{7}$$

for all t > 0. (Here || || denotes sup norm.)

(b) Conversely, say $0 < \alpha < 1$, C > 0, and that F satisfies (7) for all t > 0. Then F is Hölder continuous, with Hölder exponent α .

The proof of this result will be a straightforward generalization of the argument of Holschneider and Tchamitchian ([25]).

Our construction of nearly tight frames in our sequel article [18] will also rely heavily on Lemma 4.1. In another sequel article (already available [19]), we will use Lemma 4.1 to show that one can determine whether F is in a Besov space, solely from a knowledge of the size of its frame coefficients. In a future article, we hope to study the same question for Triebel-Lizorkin spaces. (The analogous problems on \mathbb{R}^n were solved in [12] and [13].)

It should be noted that in (6) we assume nothing about the t derivatives of $K_t(x, y)$, and no such information is needed in proving Theorem 5.5. However, in fact, if K_t is the kernel of $f(t^2\Delta)$, one does have the following improvement on (6):

For every pair of C^{∞} differential operators X (in x) and Y (in y) on **M**, and for all integers $m, N \ge 0$, there exists $C_{N,m,X,Y}$ as follows. Suppose deg X = j and deg Y = k. Then

$$t^{m+n+j+k} \left| \left(\frac{d(x,y)}{t} \right)^N \left(\frac{\partial}{\partial t} \right)^m XYK_t(x,y) \right| \le C_{N,m,X,Y}$$
(8)

for all t > 0 and all $x, y \in \mathbf{M}$.

This is in fact an easy consequence of (6). For example, $\frac{\partial}{\partial t}K_t(x, y)$ is the kernel of $2t\Delta f(t^2\Delta)$, and hence equals $2t\Delta_x K_t(x, y)$, which we can estimate by using (6).

We should explain in what sense $K_t(x, y)$ (the kernel of $f(t^2\Delta)$) deserves to be called a *wavelet*. In our model situation on \mathbb{R}^n , $K_t(x, y) = t^{-n}\psi((x - y)/t)$ behaves in evident, well-known ways under dilation and translation:

$$K_{rt}(rx, ry) = r^{-n} K_t(x, y)$$
 for $r > 0$; and (9)

$$K_t(x+a,y+a) = K_t(x,y) \text{ for } a \in \mathbb{R}^n$$
(10)

Now (9) says that, in the model case, $K_t(x, y)$ is homogeneous of degree -n in (t, x, y), so that for any $m, \alpha, \beta, t^{n+m+|\alpha|+|\beta|} \partial_t^m \partial_x^\alpha \partial_y^\beta K_t(x, y)$ is homogeneous of degree 0, and is hence bounded as a function of (t, x, y). An analogous fact to this boundedness holds on **M**, by (8) with N = 0.

As for (10), a general manifold **M** has nothing akin to translations, but in Section 5 we shall discuss the situation in which **M** has a transitive group G of smooth metric isometries. In that case one can easily see that $K_t(Tx, Ty) = K_t(x, y)$ for all $T \in G$, which is analogous to (10). (Manifolds with such a group G are usually called *homogeneous*. Obvious examples are the sphere and the torus.)

Of course, to apply our algorithm in practice, one would need to compute the kernels K_t approximately. We give some examples where this can be done, in Section 5. In the special case $f(s) = se^{-s}$, K_t can be thought of, very naturally, as a "Mexican hat" continuous wavelet for **M**. (On \mathbb{R}^n , the Mexican hat wavelet is a multiple of $\Delta e^{-\Delta}\delta$, the second derivative of a Gaussian, the function whose Fourier transform is $|\xi|^2 e^{-|\xi|^2}$.) In Section 5 we compute these continuous wavelets in the special cases where **M** is the *n*-sphere S^n and the *n*-torus T^n . For the 2-torus, we show that K_t can be evaluated by use of either of two different sums, one (obtained from the eigenfunction expansion) which converges very quickly for t large, and the other which (obtained from the proof of the Poisson summation formula) converges very quickly for t small. (The method can be extended to general n.) For the sphere, the eigenfunction expansion again gives a sum which converges very quickly for t large. When n = 2, for t small, by use of heat trace methods, we obtain a formula which converges very quickly, and which appears (from numerical evidence) to be an excellent approximation to K_t . Specifically, in [37], I. Polterovich obtains a completely explicit formula for the heat trace asymptotics on the sphere. (Earlier, less explicit formulae were found earlier in ([4]) and ([5]).) It is not hard to see that, on the sphere, $K_t(x,y) := h_t(x \cdot y)$ is a function of $x \cdot y$. Using Polterovich's result, we show how one can compute the Maclaurin series for $4\pi h_t(\cos\theta)$. In this manner we obtain an approximation

$$4\pi h_t(\cos\theta) \sim \frac{e^{-\theta^2/4t^2}}{t^2} [(1 - \frac{\theta^2}{4t^2})p(t,\theta) - t^2q(t,\theta)],\tag{11}$$

where

$$p(t,\theta) = 1 + \frac{t^2}{3} + \frac{t^4}{15} + \frac{4t^6}{315} + \frac{t^8}{315} + \frac{\theta^2}{4} \left(\frac{1}{3} + \frac{2t^2}{15} + \frac{4t^4}{105} + \frac{4t^6}{315}\right)$$

$$q(t,\theta) = \frac{1}{3} + \frac{2t^2}{15} + \frac{4t^4}{105} + \frac{4t^6}{315} + \frac{\theta^2}{4}\left(\frac{2}{15} + \frac{8t^2}{105} + \frac{4t^4}{105}\right)$$

Maple says that when t = .1, the error in the approximation (11) is never more than 9.5×10^{-4} for any $\theta \in [-\pi, \pi]$, even though both sides have a maximum of about 100. (To obtain rigorous bounds on the error is research in progress, which we expect to complete soon.) We derive a similar approximation to the heat kernel itself, when n = 2. The method can be extended to general n. Note that, if in (11) we approximate $p \sim 1$ and $q \sim 0$, we would obtain the formula for the usual Mexican hat wavelet on the real line, as a function of θ .

1.1 Historical comments

A number of groups of researchers have been studying continuous wavelets and frames on manifolds, and some have obtained important real-world applications. While our method, based on the new formula (4) is original, certain of the ideas that we have presented in this introduction have arisen in other forms before. We now discuss the work of these other researchers.

Weaker forms of Lemma 4.1 have appeared before; here is the history, as best we can determine it. If f has compact support away from 0, in 1989, Seeger-Sogge [40] showed (6) modulo a remainder term that they must handle separately. (We would not be able to handle this remainder in the applications we seek.) Next, in 1996, Tao showed ([44]. Proposition 3.1 (ii)) the case j = k = 0 of Lemma 4.1, under the restriction that \hat{h} has compact support. (Recall that $h(\xi) = f(\xi^2)$. An assumption that \hat{h} has compact support would not be natural in our context.)

Most relevantly, in 2006 Narcowich, Petrushev and Ward ([35], [36]) showed a slight variant of Lemma 4.1 if $\mathbf{M} = S^n$, the sphere, provided f had compact support away from 0. (In their variant, $K_t(x, y)$ was the kernel not of $f(t^2\Delta)$ but rather of $f(t^2\mathcal{M})$, where \mathcal{M} is a particular first-order pseudodifferential operator which is similar to Δ . Specifically, Δ multiplies spherical harmonics of degree l by l(l + n - 1), while \mathcal{M} multiplies them by l^2 . This is a minor distinction, however.)

Narcowich, Petrushev and Ward do not discuss continuous wavelets, but use spectral theory arguments to construct tight frames on S^n . They then apply their variant of Lemma 4.1 for purposes similar to ours, including characterizations of Besov and Triebel-Lizorkin spaces through frame coefficients on the sphere. Their frames have been dubbed "needlets", and have been used by statisticians and astrophysicists to study cosmic microwave background radiation (CMB). (See, for instance, [2], [31], [22] and the references therein.) We present a detailed comparison of their approach to frames and ours in our sequel article [18].

Returning to our own approach, but still restricting to the sphere, the most important cases to consider are the case in which f has compact support away from 0 (the "needlet" case, essentially considered by Narcowich, Petrushev and Ward), and the case in which $f(s) = s^r e^{-s}$ for some integer $r \ge 1$. In our sequel article [18], we construct frames from the latter f; we call these frames *Mexican needlets*. Needlets and Mexican needlets each have their own advantages. Needlets are a tight frame, and frame elements at non-adjacent scales are orthogonal. Mexican needlets, though not tight, are nearly tight; they have the advantage that one can work with them directly on the sphere, because of the formula (11). (This formula is only for r = 1, but can be readily generalized to general r. As we said before, estimating the error in (11) is work in progress.) Assuming this formula, Mexican needlets have strong Gaussian decay at each scale, and do not oscillate (for small r), so they can be implemented directly on the sphere, which is desirable if there is missing data (such as the "sky cut" of the CMB).

The statistical properties of needlets were investigated in [2]. Also, the statistical properties of Mexican needlets are already being investigated, by the second author in [29], and by Lan and Marinucci in [28]. It would be worthwile to utilize both needlets and Mexican needlets in the analysis of CMB, and the results should be compared.

A number of other researchers have studied wavelets and frames on manifolds. In all of the works mentioned below, when orthonormal bases were constructed, they are not known to give rise to a space-frequency analysis; and when frames were constructed, they are not known to be tight or to give rise to a space-frequency analysis.

Let us begin by discussing earlier works on manifolds, which contain some ideas related to those in this article. In alphabetical order:

- Antoine, Vandergheynst, and collaborators ([1], [3]) have constructed smooth continuous wavelets on the sphere and related manifolds, by use of stereographic dilations (replacing the usual dilations), rotations (replacing the usual translations), and spherical convolution (replacing the usual convolution). They obtained frames by discretizing these continuous wavelets.
- Coifman, Maggioni, and collaborators ([6], [30]) used the heat equation on manifolds for the rather different purpose of constructing orthonormal wavelet bases through a diffusion process, leading to a multiresolution analysis. They exploit the idea (which they attribute to Stein) of $e^{-t\Delta}$ being a sort of dilate of $e^{-\Delta}$.
- Freeden and collaborators ([15], [16]) defined continuous wavelets on the sphere S^2 , and applied them to the geosciences. Their continuous wavelets were of the form $f(t^2\Delta)$ for various f (not all in the Schwartz space), although they did not formulate them in that manner. One of their many examples was our Mexican hat wavelet, which they called the Gauss-Weierstrass wavelet of order zero. They did not have Lemma 4.1, so they restricted to an (extensive) L^2 theory of continuous wavelets. In the context of S^2 , they had, in particular, results equivalent to our (4).
- Han ([23], [24]) constructed frames on general spaces of homogeneous type (including manifolds). His method is to discretize a discrete version of Calderón's formula in this general setting. He also used the T(1) theorem to estimate errors,

The following researchers have also worked on wavelets and frames on manifolds. They used methods which seem unrelated to those in the present article. In alphabetical order:

- Dahlke ([8]) constructed an analogue of Haar wavelets on Riemannian manifolds.
- Dahmen and Schneider ([9]) have used parametric liftings from standard bases on the unit cube to obtain wavelet bases on manifolds which are the disjoint union of smooth parametric images of the standard cube.
- Schröder and Sweldens ([39]) used a lifting scheme to build wavelets on manifolds. This lifting scheme uses no invariance properties, and regularity information is not easily obtained.

2 Applying the Spectral Theorem

In this section, we give the proof of (4), as well as the proof of a discrete analogue, which will be used in our sequel article [18] to construct nearly tight frames. (The proofs are quite elementary, and the reader who is willing to accept (4) can go on to the next section.) Specifically, we shall show:

Lemma 2.1. Let T be a positive self-adjoint operator on a Hilbert space \mathcal{H} , and let P be the projection onto the null space of T. Suppose $l \geq 1$ is an integer, $f_0 \in \mathcal{S}(\mathbb{R}^+)$, $f_0 \not\equiv 0$, and let $f(s) = s^l f_0(s)$. Set $c = \int_0^\infty |f(t)|^2 \frac{dt}{t}$.

(a) Then for any $F \in \mathcal{H}$,

$$\lim_{\varepsilon \to 0^+, N \to \infty} \left[\int_{\varepsilon}^N |f|^2 (tT) \frac{dt}{t} \right] F = c(I - P)F,$$
(12)

Thus

$$\int_0^\infty |f|^2 (tT) \frac{dt}{t} := \lim_{\varepsilon \to 0^+, N \to \infty} \left[\int_\varepsilon^N |f|^2 (tT) \frac{dt}{t} \right]$$

exists in the strong operator topology, and equals c(I - P).

(b) Suppose that a > 0 ($a \neq 1$) is such that the Daubechies condition holds: for any s > 0,

$$0 < A_a \le \sum_{j=-\infty}^{\infty} |f(a^{2j}s)|^2 \le B_a < \infty,$$

$$\tag{13}$$

Then $\lim_{M,N\to\infty} \left[\sum_{j=-M}^{N} |f|^2 (a^{2j}T)\right]$ exists in the strong operator topology on \mathcal{H} ; we denote this limit by $\sum_{j=-\infty}^{\infty} |f|^2 (a^{2j}T)$. Moreover

$$A_a(I-P) \le \sum_{j=-\infty}^{\infty} |f|^2 (a^{2j}T) \le B_a(I-P).$$
(14)

Remark (b) is a discrete analogue of (a), since the sum in (13) is a multiple of a Riemann sum for the integral $\int_0^\infty |f(st)|^2 \frac{dt}{t} = c$, while the spectral theorem will show that it is valid to replace s in (13) by T, to obtain (14).

Proof. We prove (a) and (b) together. Let $T = \int_0^\infty \lambda dP_\lambda$ be the spectral decomposition of T; thus, in particular, $P = P_{\{0\}}$.

Observe that, by the spectral theorem, if g is a bounded Borel function on \mathbb{R}^+ , then $||g(T)|| \leq \sup_{s\geq 0} |g(s)|$. It follows readily that the integrand in (12) is a family of operators in $\mathcal{B}(\mathcal{H})$ (indexed by t), which depends continuously on $t \in [\varepsilon, N]$ (in the norm topology on $\mathcal{B}(\mathcal{H})$). (Use the mean value theorem for s in a suitable compact interval and the rapid decay of f at ∞ .) Thus $\int_{\varepsilon}^{N} |f|^2 (tT) \frac{dt}{t}$ makes sense as a bounded operator on \mathcal{H} . (As usual, it is defined as the unique bounded operator S with $\langle S\varphi, \psi \rangle = \int_{\varepsilon}^{N} \langle |f|^2 (tT)\varphi, \psi \rangle \frac{dt}{t}$ for all $\varphi, \psi \in \mathcal{H}$.)

For $0 < \varepsilon < N < \infty$, define $g_{\varepsilon,N} : [0,\infty) \to [0,\infty)$ by

$$g_{\varepsilon,N}(\lambda) = \int_{\varepsilon}^{N} |f|^2 (t\lambda) \frac{dt}{t}.$$
(15)

Then

$$g_{\varepsilon,N}(T) = \int_{\varepsilon}^{N} |f|^2 (tT) \frac{dt}{t}$$

(This follows from an elementary application of Fubini's theorem to $\int_{\varepsilon}^{N} \int_{0}^{\infty} |f|^{2}(t\lambda) d\langle P_{\lambda}\varphi,\psi\rangle \frac{dt}{t}$.) For $M, N \geq 0$ we also define $h_{M,N} : [0,\infty) \to [0,\infty)$ by

$$h_{M,N}(\lambda) = \sum_{j=-M}^{N} |f|^2 (a^{2j}\lambda);$$

$$h(\lambda) = \sum_{j=-\infty}^{\infty} |f|^2 (a^{2j}\lambda).$$
 (16)

and we also set

To prove the lemma it is enough to show that $g_{\varepsilon,N}(T) \to c(I-P)$ (strongly, as $\varepsilon \to 0^+$ and $N \to \infty$) and that $h_{M,N}(T) \to h(T)$ (strongly, as $M, N \to \infty$). Indeed, (a) would then be immediate. (b) would then also follow at once from (13), which implies, by the spectral theorem, that $A_a(I-P) \leq h(T) \leq B_a(I-P)$.

To establish these conclusions, we first note that this strong convergence need only be proved on $(I-P)\mathcal{H}$, since all the operators vanish identically on $P\mathcal{H}$. Next we note that for any ε, M, N , $||g_{\varepsilon,N}||_{\infty} \leq c$, $||h_{M,N}||_{\infty} \leq B_a$, and $||h||_{\infty} \leq B_a$, whence $||g_{\epsilon,N}(T)|| \leq c$, $||h_{M,N}(T)||_{\infty} \leq B_a$, and $||h(T)|| \leq B_a$, Thus the needed strong convergence need only be proved on a dense subset of $(I-P)\mathcal{H}$. Set

$$V = \bigcup_{0 < \eta < L < \infty} P_{[\eta, L]} \mathcal{H};$$

observe that V is dense in $(I - P)\mathcal{H}$ (since $P = P_{\{0\}}$). Thus, it suffices to show the following: fix $0 < \eta < L < \infty$, and say $F \in P_{[\eta,L]}\mathcal{H}$. Then $g_{\varepsilon,N}(T)F \to cF$ and $h_{M,N}(T)F \to h(T)F$.

This, however, is immediate from the spectral theorem and the evident facts that $g_{\epsilon,N} \to c$ and $h_{M,N} \to h$, uniformly on $[\eta, L]$. Although this uniform convergence is easily shown, for later purposes, we carefully express it quantitatively in the next lemma. (Note that we may assume a > 1; otherwise replace it by 1/a.)

Lemma 2.2. Notation as in Lemma 2.1, and as in equations (15) through (16). Suppose $J \ge 1$ is an integer, and let $M_J = \max_{r>0} |r^J f(r)|$. Suppose $0 < \eta < L < \infty$, and let I be the closed interval $[\eta, L]$. Let $\| \|$ denote the sup norm on I.

(a) If $0 < \varepsilon < N$, then

$$||g_{\varepsilon,N} - c|| \le c_L \varepsilon^{2l} + \frac{C_\eta}{N^{2J}},$$

where we may take $c_L = L^{2l} ||f_0||_{\infty}^2/(2l)$, and $C_{\eta} = M_J^2/(2J\eta^{2J})$.

(b) If $M, N \ge 0$, and a > 1, then

$$||h_{M,N} - h|| \le \frac{c'_L}{a^{4Ml}} + \frac{C'_{\eta}}{a^{4NJ}}.$$

where we may take $c'_L = (L^{2l} \|f_0\|_{\infty}^2)/(a^{4l}-1)$, and $C'_{\eta} = M_J^2/[(a^{4J}-1)\eta^{2J}]$.

Proof. Say $s \in I$. Then

$$|g_{\varepsilon,N}(s) - c| = \int_0^\varepsilon |f|^2 (st) \frac{dt}{t} + \int_N^\infty |f|^2 (st) \frac{dt}{t}$$

(a) follows from noting $|f|^2(st) \leq (Lt)^{2l} ||f_0||_{\infty}^2$ in the first integral, and that $|f|^2(st) \leq M_J^2/(\eta t)^{2J}$ in the second. Similarly,

$$|h_{M,N}(s) - h(s)| = \sum_{j < -M} |f|^2 (a^{2j}s) + \sum_{j > N} |f|^2 (a^{2j}s).$$

(b) follows from noting $|f|^2(a^{2j}s) \leq (La^{2j})^{2l} ||f_0||_{\infty}^2$ in the first summation, and that $|f|^2(a^{2j}s) \leq M_J^2/(\eta a^{2j})^{2J}$ in the second. This completes the proof.

We can now express the strong convergence in Lemma 2.1 in the following very quantitative manner:

10

Lemma 2.3. Notation as in Lemmas 2.1 and 2.2, and again let $T = \int_0^\infty \lambda dP_\lambda$ be the spectral decomposition of T. Then for any $F \in \mathcal{H}$, we have:

$$\|[g_{\varepsilon,N}(T) - c]F\| \le \left[c_L \varepsilon^{2l} + \frac{C_{\eta}}{N^{2J}}\right] \|F\| + 2c \left\|(I - P_{[\eta,L]})F\right\|,$$
(17)

and, if a > 1,

$$\|[h_{M,N}(T) - h(T)]F\| \le \left[\frac{c'_L}{a^{4Ml}} + \frac{C'_{\eta}}{a^{4NJ}}\right]\|F\| + 2B_a \left\|(I - P_{[\eta,L]})F\right\|.$$
(18)

Proof. For (17), we need only substitute $F = P_{[\eta,L]}F + (I - P_{[\eta,L]})F$ in the left side. This gives

$$\begin{split} \| \left[g_{\varepsilon,N}(T) - c \right] F \| &\leq \| \left[g_{\varepsilon,N}(T) - c \right] P_{[\eta,L]} F \| + \| g_{\varepsilon,N}(T) \| \| (I - P_{[\eta,L]}) F \| + c \| (I - P_{[\eta,L]}) F \| \\ &\leq \| \left[g_{\varepsilon,N}(T) - c \right] \chi_{[\eta,L]} \|_{\infty} \| F \| + 2c \| (I - P_{[\eta,L]}) F \|, \end{split}$$

which, because of Lemma 2.2, establishes (17). The proof of (18) is entirely analogous.

(17) and (18) are of significance for numerical calculations. Say, for instance, that F = (I - P)F, and one wants to compute h(T)F to a certain precision. This involves summing an infinite series, so one instead seeks to compute $[h_{M,N}(T)]F$ for M, N large enough; how large must one take them to be? One first chooses η, L so that the second term on the right side of (18) is very small. Then one chooses M, N to make the first term on the right of (18) very small as well.

We will return to this point in our discussion of space-frequency analysis, in our sequel article [18].

3 Preliminaries on Manifolds

For the rest of the article, (\mathbf{M}, g) will denote a smooth, compact, connected, oriented Riemannian manifold of dimension n, and μ will denote the measure on \mathbf{M} arising from integration with respect to the volume form on \mathbf{M} . In this section we assemble several well-known facts about analysis on \mathbf{M} (preceded, below, by bullets), which we shall need in this article and in sequel articles.

For $x, y \in \mathbf{M}$, we let d(x, y) denote the infimum of the lengths of all piecewise C^1 curves joining x to y; then (\mathbf{M}, d) is evidently a metric space. It is well-known (see, e.g., [34]) that there is a geodesic joining x to y with length d(x, y), but this fact, though basic, is not so relevant for this article. Most of what we need to know about the metric d is contained in the simple proposition which follows.

For $x \in \mathbf{M}$, we let B(x, r) denote the ball $\{y : d(x, y) < r\}$.

Proposition 3.1. Cover **M** with a finite collection of open sets U_i $(1 \le i \le I)$, such that the following properties hold for each *i*:

- (i) there exists a chart (V_i, ϕ_i) with $\overline{U}_i \subseteq V_i$; and
- (*ii*) $\phi_i(U_i)$ is a ball in \mathbb{R}^n .

Choose $\delta > 0$ so that 3δ is a Lebesgue number for the covering $\{U_i\}$. Then, there exist $c_1, c_2 > 0$ as follows:

For any $x \in \mathbf{M}$, choose any $U_i \supseteq B(x, 3\delta)$. Then, in the coordinate system on U_i obtained from ϕ_i ,

$$d(y,z) \le c_2|y-z|$$

for all $y, z \in U_i$; and

$$|c_1|y-z| \le d(y,z)$$

for all $y, z \in B(x, \delta)$.

11

Proof. Say $y, z \in U_i$. We work in the coordinate system on U_i obtained from ϕ_i . Then d(y, z) is at most the length (in the Riemannian metric) of the straight line joining y to z, which is $\leq c_2|y-z|$ for some c_2 . (By assumption (ii), that straight line is contained in U_i .) On the other hand, if $y, z \in B(x, \delta) \subseteq U_i$, we may take a sequence of piecewise C^1 curves γ_k , joining y to z, whose lengths $l(\gamma_k)$ approach d(y, z)as $k \to \infty$. Surely $d(y, z) < 2\delta$. Thus, for some N > 0, if k > N, then $l(\gamma_k) < 2\delta$. Therefore each point on γ_k is at distance at most 2δ from y, hence at most 3δ from x. Thus $\gamma_k \subseteq U_i$. Letting $\|\gamma'_k(t)\|$ denote the length of the tangent vector $\gamma'_k(t)$ in the Riemannian metric, we see that

$$l(\gamma_k) = \int_0^1 \|\gamma'_k(t)\| dt \ge c_1 \int_0^1 |\gamma'_k(t)| dt \ge c_1 |z - y|,$$

since $z - y = \int_0^1 \gamma'_k(t) dt$. Letting $k \to \infty$, we see that $d(y, z) \ge c_1 |y - z|$ as well. This completes the proof.

We fix collections $\{U_i\}, \{V_i\}, \{\phi_i\}$ and also δ as in Proposition 3.1, once and for all.

• Notation as in Proposition 3.1, there exist $c_3, c_4 > 0$, such that

$$c_3 r^n \le \mu(B(x,r)) \le c_4 r^n \tag{19}$$

whenever $x \in \mathbf{M}$ and $0 < r \leq \delta$, and such that

$$c_3\delta^n \le \mu(B(x,r)) \le \mu(\mathbf{M}) \le c_4 r^n \tag{20}$$

whenever $x \in \mathbf{M}$ and $r > \delta$.

To see (19), note that, since the collection $\{U_i\}$ is finite, we may fix i and prove it for all x with $B(x, 3\delta) \subseteq U_i$.

We work in the coordinate system on U_i obtained from ϕ_i ; in that coordinate system, U_i is a Euclidean ball, say $\{y : |y - x_0| < R\}$. (See Proposition 3.1). By compactness and a simple contradiction argument, there is an $\eta > 0$ such that, for all x with $B(x, 3\delta) \subseteq U_i$, one has that $|x - x_0| < R - \eta$. Accordingly, for such an x, if $|y - x| < \eta$, then $y \in U_i$.

Thus, by Proposition 3.1, we have that

$$\{y : |y - x| < \min(r/c_2, \eta)\} \subseteq B(x, r) \subseteq \{y : |y - x| < r/c_1\},\$$

for all $r < \delta$. (19) now follows from the fact that the determinant of the metric tensor g is bounded above and below on U_i . For (20), one need only note that if $r > \delta$, then $\mu(B(x,r)) \ge \mu(B(x,\delta)) \ge c_3\delta^n$, while $\mu(B(x,r)) \le \mu(\mathbf{M}) \le [\mu(\mathbf{M})/\delta^n]\delta^n \le [\mu(\mathbf{M})/\delta^n]r^n$.

• (\mathbf{M}, d, μ) is a space of homogeneous type, in the sense of [7].

Indeed, d is a metric and μ is a positive Borel measure, so one only needs to check the doubling condition: $\mu(B(x,2r)) \leq C\mu(B(x,r))$ with C independent of x, r. But this is immediate from (19) and (20).

• For any N > n there exists C_N such that

$$\int_{\mathbf{M}} [1 + d(x, y)/t]^{-N} d\mu(y) \le C_N t^n$$
(21)

for all $x \in \mathbf{M}$ and t > 0.

(21) is proved by the "dyadic annulus" method. Fix x, t and let $A_j = B(x, 2^j t) \setminus B(x, 2^{j-1}t)$, so that, by (19) and (20), $\mu(A_j) \leq c_4 2^{nj} t^n$. (21) now follows at once, if one breaks up the integral in (21) into integrals over $B(x, t), A_1, A_2, \ldots$, and notes that $\sum_{j=0}^{\infty} 2^{(n-N)j} < \infty$.

• For any N > n there exists C'_N such that

$$\int_{d(x,y)\ge t} d(x,y)^{-N} d\mu(y) \le C'_N t^{n-N}$$
(22)

for all $x \in \mathbf{M}$ and t > 0.

(22) follows at once from (21), once we observe that, if $d(x,y) \ge t$, then $[d(x,y)/t]^{-N} \le C[1+d(x,y)/t]^{-N}$, if $C=2^N$.

• For any N > n there exists C''_N such that

$$\int_{\mathbf{M}} [1 + d(x, z)/t]^{-N} [1 + d(z, y)/t]^{-N} d\mu(z) \le C_N'' t^n [1 + d(x, y)/t]^{-N}$$
(23)

for all $x, y \in \mathbf{M}$ and t > 0.

To see this, break up the integral into integrals over $H_1 = \{z : d(x,z) \leq d(y,z)\}$ and $H_2 = \{z : d(x,z) > d(y,z)\}$ (which, by the way, are hemispheres if **M** is a round sphere and $x \neq y$). By symmetry we need only estimate the integral over H_1 . But if z is in H_1 , $d(x,y) \leq 2d(z,y)$, so $[1 + d(z,y)/t]^{-N} \leq C[1 + d(x,y)/t]^{-N}$ (where $C = 2^N$). Thus the integral over H_1 is no greater than $C[1 + d(x,y)/t]^{-N} \int_{H_1} [1 + d(x,z)/t]^{-N} d\mu(z)$). Estimating the latter integral through (21), we obtain (23).

• For all M, t > 0, and for all $E \subseteq \mathbf{M}$ with diameter less than Mt, if $x_0 \in E$, then one has that

$$\frac{1}{M+1}[1+d(x,y)/t] \le [1+d(x_0,y)/t] \le (M+1)[1+d(x,y)/t]$$
(24)

for all $x \in E$ and all $y \in \mathbf{M}$.

This is true simply because d is a metric.

4 Kernels

 Δ will now denote the Laplace-Beltrami operator on **M** (equal to $-d^*d$, where * is taken with respect to the given Riemannian metric). We apply Lemma 2.1 to $T = \Delta$.

In order to carry out the plan explained in the introduction to this article, we must study the kernel $K_{\sqrt{t}}(x, y)$ of the operator $f(t\Delta)$ for $f \in \mathcal{S}(\mathbb{R}^+)$, and we do so in this section. Before proving the crucial Lemma 4.1, we will present some well-known information about $K_{\sqrt{t}}(x, y)$ for large t and also, off the diagonal, for small t. (This information will be preceded, below, by \triangleright signs.) Concerning the Laplace-Beltrami operator Δ , we first recall:

• Δ , as an operator on $C^{\infty}(\mathbf{M})$, has an orthonormal basis of eigenfunctions $\{u_l : 0 \leq l < \infty\}$; all are in $C^{\infty}(\mathbf{M})$. We may, and do, choose the u_l to be real-valued. We order the u_l so that the corresponding eigenvalues λ_l form a non-decreasing sequence. Then u_0 is constant, $\lambda_0 = 0$ and all other $\lambda_l > 0$.

This easily implies:

• Δ , as an operator on $C^{\infty}(\mathbf{M})$, has a unique extension to a self-adjoint operator on $L^2(\mathbf{M})$ (which we also denote by Δ). Its domain is $\{F = \sum a_l u_l \in L^2(\mathbf{M}) : \sum |\lambda_l a_l|^2 < \infty\}$, and for such an F, $\Delta F = \sum \lambda_l a_l u_l$.

Of great importance is Weyl's Lemma, which says [26], in sharp form:

• For $\lambda > 0$, let $N(\lambda)$ denote the number of eigenvalues of Δ which are less than or equal to λ (counted with respect to multiplicity). Then for some c > 0, $N(\lambda) = c\lambda^{n/2} + O(\lambda^{(n-1)/2})$.

Since $N(\lambda_l) = l + 1$, we conclude:

• For some constants $c_1, c_2 > 0$, we have $c_1 l^{2/n} \le \lambda_l \le c_2 l^{2/n}$ for all l.

Since $\Delta^m u_l = \lambda_l^m u_l$, and Δ^m is an elliptic differential operator of degree 2m, Sobolev's lemma, combined with the last fact, implies:

• For any integer $k \ge 0$, there exists $C_k, \nu_k > 0$ such that $||u_l||_{C^k(\mathbf{M})} \le C_k(l+1)^{\nu_k}$.

(In fact, by Sobolev's lemma, we may, for any $\varepsilon > 0$, take $\nu_k = (2k + n + \varepsilon)/2n$. By [41], Lemma 4.2.4, with $\lambda = \lambda_l$ in that lemma, we may in fact take $\nu_0 = (n-1)/n$.)

From these facts one sees at once:

• The mapping $\sum a_l u_l \to (a_l)_{l\geq 0}$ gives a Fréchet space isomophism of $C^{\infty}(\mathbf{M})$ with the space of rapidly decaying sequences.

For the rest of this section, say $f \in \mathcal{S}(\mathbb{R}^+)$. We conclude:

 \triangleright For $t > 0, x, y \in \mathbf{M}$, let (for the rest of this section)

$$K_{\sqrt{t}}(x,y) = \sum_{l=0}^{\infty} f(t\lambda_l) u_l(x) u_l(y).$$
(25)

Then $K_{\sqrt{t}}$ is the kernel of the operator $f(t\Delta)$, in the sense that if $F \in L^2(\mathbf{M})$, then

$$[f(t\Delta)F](x) = \int_{\mathbf{M}} K_{\sqrt{t}}(x, y)F(y)d\mu(y),$$

and $K_{\sqrt{t}}(x,y)$ is smooth in (t, x, y) (for $t > 0, x, y \in \mathbf{M}$).

From (25), our estimates on the λ_l and the $||u_l||_{C^k}$, and the rapid decay of Schwartz functions, we conclude:

 \triangleright If f(0) = 0, then for any $M, N \ge 0$,

$$\lim_{t \to \infty} t^M \frac{\partial^N}{\partial t^N} K_{\sqrt{t}} = 0$$

in $C^{\infty}(\mathbf{M} \times \mathbf{M})$.

Note also, that whenever l > 0, $\int_{\mathbf{M}} u_l d\mu = C \int_{\mathbf{M}} u_l \overline{u_0} d\mu = 0$. Accordingly:

$$\triangleright \text{ If } f(0) = 0, \text{ then } \int_{\mathbf{M}} K_{\sqrt{t}}(x, y) d\mu(x) = 0 \text{ for all } y \in \mathbf{M}, \text{ and } \int_{\mathbf{M}} K_{\sqrt{t}}(x, y) d\mu(y) = 0 \text{ for all } x \in \mathbf{M}.$$

Next we discuss the behavior of $K_{\sqrt{t}}$, as $t \to 0^+$. For this, we utilize some of the theory of pseudodifferential operators.

We will need some facts about symbols on \mathbb{R} . For $m \in \mathbb{R}$, let $S_1^m(\mathbb{R})$ denote the space of standard symbols $p(\xi)$ of order m, which depend only on the "dual variable" ξ . Let $\{\| \|_{m,N}\}$ denote the natural nondecreasing family of seminorms defining the Fréchet space topology of $S_1^m(\mathbb{R})$; thus

$$\|p\|_{m,N} = \sum_{0 \le j \le N} \sup_{\xi} \left[(1+|\xi|)^{j-m} |p^{(j)}(\xi)| \right]$$

for $p \in S_1^m(\mathbb{R})$. If $G \in S_1^m(\mathbb{R})$, let $G_t(\xi) = G(t\xi)$. It is evident that $G_t \in S_1^m(\mathbb{R})$. In fact we have:

• Say that $G \in S_1^m(\mathbb{R})$. Let k be the least integer which is greater than or equal to m; if k > 0, suppose further that G vanishes to order at least k at 0. Then for any N, there exists C > 0 such that

$$\|G_t\|_{m,N} \le Ct^m \tag{26}$$

 $whenever \; 0 < t < 1.$

To see this, say $j \ge 0$ is an integer. One needs only note:

(i) if $0 \le j < m$ (so that k > 0) and $t|\xi| \le 1$, then

$$|(G_t)^{(j)}(\xi)| \le Ct^j(t^{k-j}|\xi|^{k-j}) \le Ct^j(t^{m-j}|\xi|^{m-j}) \le Ct^m(1+|\xi|)^{m-j};$$

(*ii*) if $0 \leq j < m$ and $t|\xi| > 1$, then

$$|(G_t)^{(j)}(\xi)| \le Ct^j (1+t|\xi|)^{m-j} \le Ct^j (t^{m-j}|\xi|^{m-j}) \le Ct^m (1+|\xi|)^{m-j};$$

(*iii*) if $j \ge m$, then

$$|(G_t)^{(j)}(\xi)| \le Ct^j (1+|t\xi|)^{-(j-m)} \le Ct^m (1+|\xi|)^{-(j-m)}$$

This proves the claim.

In particular, say $G \in S_1^1(\mathbb{R})$ is arbitrary. Then $G - G(0) \in S_1^1(\mathbb{R})$, and vanishes to order at least 1 at 0. Accordingly, for any N, $\|G_t - G(0)\|_{1,N} \leq Ct$ for 0 < t < 1, so that $G_t \to G(0)$ in $S_1^1(\mathbb{R})$ as $t \to 0^+$.

We return now to \mathbf{M} , where we need to look at the class $OPS_{1,0}^m(\mathbf{M})$ of pseudodifferential operators of order $m \in [-\infty, \infty)$. As is familiar, $T : C^{\infty}(\mathbf{M}) \to C^{\infty}(\mathbf{M})$ is in $OPS_{1,0}^m(\mathbf{M})$ provided that the following conditions hold for $\varphi, \psi \in C^{\infty}(\mathbf{M})$:

- 1. If $\operatorname{supp} \varphi \cap \operatorname{supp} \psi = \emptyset$, then the operator $\varphi T \psi$ has a smooth kernel; and
- 2. If $\operatorname{supp} \varphi \cup \operatorname{supp} \psi$ is contained in a chart (V, Φ) , then $\varphi T \psi$ is the pullback to **M** of a pseudodifferential operator $\Phi_*(\varphi T \psi) \in OPS_{1,0}^m(\mathbb{R}^n)$.

(Of course, here, $(\varphi T \psi)F = \varphi T(\psi F)$.) One places a Fréchet space structure on $OPS_{1,0}^m(\mathbf{M})$ in a natural manner. (A brief sketch: First note that $OPS^{-\infty}(\mathbf{M})$ is the space of operators with smooth kernels, so it has a natural Fréchet space structure, inherited from $C^{\infty}(\mathbf{M} \times \mathbf{M})$. For other m, one chooses a finite atlas $\{W_k\}$ on \mathbf{M} with the property that if two charts in the atlas intersect, their union is contained in a chart. One chooses a partition of unity $\{\varphi_k\}$ subordinate to this atlas. One notes that if $T \in OPS_{1,0}^m(\mathbb{R}^n)$, then $T = \sum_{i,j} \varphi_i T \varphi_j$. One notes that if $W_i \cap W_j = \emptyset$, then $\varphi_i T \varphi_j \in OPS^{-\infty}(\mathbf{M})$, a Fréchet space; while if $W_i \cap W_j \neq \emptyset$, then $W_i \cap W_j \subseteq V$ for some chart (V, Φ) , and $\Phi_*(\varphi T \psi) \in OPS_{1,0}^m(\mathbb{R}^n)$, also a Fréchet space. Finally one defines seminorms on $OPS_{1,0}^m(\mathbb{R}^n)$ of the form $\sum_{i,j} \|\varphi_i T \varphi_j\|$, where, in the summation, one uses appropriate seminorms coming from $OPS^{-\infty}(\mathbf{M})$ if $W_i \cap W_j = \emptyset$, or from $OPS_{1,0}^m(\mathbb{R}^n)$ if $W_i \cap W_j \neq \emptyset$. The Fréchet space topology thereby placed on $OPS_{1,0}^m(\mathbf{M})$ is independent of all choices made.)

One has the following theorem of Strichartz ([43], or Theorem 1.3, page 296, of [45]):

• If
$$p(\xi) \in S_1^m(\mathbb{R})$$
, then $p(\sqrt{\Delta}) \in OPS_{1,0}^m(\mathbf{M})$.

In fact, the map $p \to p(\sqrt{\Delta})$ is continuous from $S_1^m(\mathbb{R})$ to $OPS_{1,0}^m(\mathbf{M})$. Indeed, by the closed graph theorem for Fréchet spaces, it is enough to observe that if $u \in C^{\infty}(\mathbf{M})$, then the maps $p \to \langle p(\sqrt{\Delta})u, u_l \rangle$ are continuous from $S_1^m(\mathbb{R})$ to \mathbb{C} for every l, and this is clear.

As a consequence, if $\varphi, \psi \in C^{\infty}(\mathbf{M})$ are as in #1 above, then the map from $S_1^m(\mathbb{R})$ to $OPS^{-\infty}(\mathbf{M})$, which takes p to $\varphi p(\sqrt{\Delta})\psi$, is continuous. If $\varphi, \psi \in C^{\infty}(\mathbf{M})$ are as in #2 above, then the map from $S_1^m(\mathbb{R})$ to $OPS_{1,0}^m(\mathbb{R}^n)$, which takes p to $\Phi_*(\varphi p(\sqrt{\Delta})\psi)$ is continuous.

As usual, $f \in \mathcal{S}(\mathbb{R}^+)$; let $G(\xi) = f(\xi^2)$. Then $G \in \mathcal{S}(\mathbb{R})$. (In fact, if we allow f to vary, the map $f \to G$ is evidently a bijection between $\mathcal{S}(\mathbb{R}^+)$ and the space of even Schwartz functions on \mathbb{R} .) Now $G_{\sqrt{t}}(\xi) = f(t\xi^2)$, and $G_{\sqrt{t}}(\sqrt{\Delta}) = f(t\Delta)$. Since $G_{\sqrt{t}} \to G(0)$ in $S_1^1(\mathbb{R})$ as $t \to 0^+$, we infer:

• $f(t\Delta) \to f(0)I$ in $OPS_{1,0}^1(\mathbf{M})$ as $t \to 0^+$.

Let D denote the diagonal of $\mathbf{M} \times \mathbf{M}$. We can now show:

 \triangleright For any N > 0,

$$\lim_{t \to 0} \frac{\partial^N}{\partial t^N} K_{\sqrt{t}} = 0$$

in $C^{\infty}((\mathbf{M} \times \mathbf{M}) \setminus D)$.

To prove this, we adapt the arguments of [45], page 313. Say $\varphi, \psi \in C^{\infty}(\mathbf{M})$, have disjoint supports. Suppose further that $\varphi \equiv 1$ in an open set U and $\psi \equiv 1$ in an open set V.

It is enough to show that, for any C^{∞} differential operator Y on \mathcal{M} , acting in the y variable,

$$\lim_{t\to 0} Y \frac{\partial^N}{\partial t^N} K_{\sqrt{t}}(x,y) \text{ (regarded as a function of } x) = 0$$

in $C^{\infty}(U)$, uniformly for $y \in V$. But if $x \in U$ and $y \in V$, then

$$Y\frac{\partial^N}{\partial t^N}K_{\sqrt{t}}(x,y) = \varphi(x)\left[Yf^{(N)}(t\Delta)(\psi\Delta^N\delta_y)\right](x) = \left[S_t(w_y)\right](x), \text{ say },$$

where S_t is the pseudodifferential operator $\varphi Y f^{(N)}(t\Delta)\psi$, and $w_y = \Delta^N \delta_y$. For some s > 0, the set $\{w_y : y \in V\}$ is a bounded subset of $H^{-s}(\mathbf{M})$. Also, as $t \to 0^+$, $f^{(N)}(t\Delta) \to f^{(N)}(0)I$ in $OPS_{1,0}^{1}(\mathbf{M})$, so $Yf^{(N)}(t\Delta) \to f^{(N)}(0)Y$ in $OPS_{1,0}^{k+1}(\mathbf{M})$, if $k = \deg Y$. But φ, ψ have disjoint supports, so the map $R \to \varphi R \psi$ is continuous from $OPS_{1,0}^{k+1}(\mathbf{M})$ to $OPS_{1,0}^{-\infty}(\mathbf{M})$. Therefore $S_t \to \varphi \left[f^{(N)}(0)Y \right] \psi \equiv 0$ in $OPS_{1,0}^{-\infty}(\mathbf{M})$. Thus:

$$S_t w_y \to 0$$
 in $C^{\infty}(\mathbf{M})$, uniformly for $y \in V$,

as desired.

Applying the mean value theorem in the t variable repeatedly to the last fact about $K_{\sqrt{t}}$, we see:

 \triangleright Let *E* be any fixed compact subset of $(\mathbf{M} \times \mathbf{M}) \setminus D$, and let \mathcal{U} be the interior of *E*. Then for any k, N there exists $C_{k,N}$ such that

$$\|K_{\sqrt{t}}\|_{C^k(\mathcal{U})} \le C_{k,N} t^N$$

whenever 0 < t < 1.

So far, nearly everything in this section has been well-known, but now we must consider the behavior of K_t near the diagonal for small t. As we have explained and motivated in the introduction, this behavior is described by (6):

Lemma 4.1. Say f(0) = 0. Then for every pair of C^{∞} differential operators X (in x) and Y (in y) on **M**, and for every integer $N \ge 0$, there exists $C_{N,X,Y}$ as follows. Suppose deg X = j and deg Y = k. Then

$$t^{n+j+k} \left| \left(\frac{d(x,y)}{t} \right)^N XYK_t(x,y) \right| \le C_{N,X,Y}$$
(27)

for all t > 0 and all $x, y \in \mathbf{M}$.

Proof. Of course d is bounded on $\mathbf{M} \times \mathbf{M}$. Thus, by what we already know about K_t , it suffices to prove (27) for 0 < t < 1. In fact, with notation as in Proposition 3.1, it suffices to show that (27) holds whenever 0 < t < 1 and $d(x, y) < \delta/3$. Cover \mathbf{M} by a finite collection of balls $\{B(z_l, \delta/3) : 1 \le l \le J\}$. Then any pair of points (x, y) with $d(x, y) < \delta/3$ lie together in one of the balls $B(z_l, 2\delta/3)$. Thus, it suffices to show that (27) holds whenever 0 < t < 1 and $x, y \in B(z_l, 2\delta/3)$ for some l. Moreover, we may fix a positive integer M and prove that (27) holds for all $N \le M$.

Now K_t is the kernel of $f(t^2\Delta) = G(t\sqrt{\Delta})$, where $G(\xi) = f(\xi^2)$. We claim that it is enough to prove (27) in each of the following two cases:

- (i) $\operatorname{supp} \widehat{G} \subseteq (-1, 1)$; and
- (*ii*) G vanishes to order at least M at 0.

Indeed, any even G in $\mathcal{S}(\mathbb{R})$ can be written as the sum of two even functions G_1 and G_2 , where G_1 is of type (i) and G_2 is of type (ii). (To see this, say that, for $0 \leq l \leq M - 1$, $G^{(l)}(0) = a_l$. It is enough to show that there exists an even function G_1 with $\operatorname{supp}\widehat{G_1} \subseteq (-1,1)$, such that $G_1^{(l)}(0) = a_l$, for $0 \leq l \leq M - 1$, for then we can set $G_2 = G - G_1$. For this, see Lemma 7.1 in the Appendix (Section 7).)

In case (i), note that, by Huygens' principle, the support of K_t , the kernel of

$$G(t\sqrt{\Delta}) = c \int_{-\infty}^{\infty} \hat{G}(s) e^{-ist\sqrt{\Delta}} ds,$$

is contained in $\{(x, y) : d(x, y) \le t\}$. Thus, in this case, we may take M = 0. In either case (i) or case (ii), it is sufficient to show that, for every $\varphi, \psi \in C_c^{\infty}(B(z_l, \delta))$, we have that

$$t^{n+j+k}\left(\frac{d(x,y)}{t}\right)^N |XY[\varphi(x)K_t(x,y)\psi(y)]| \le C$$

whenever deg X = j and deg Y = k, for all 0 < t < 1, all $N \leq M$, and all $x, y \in B(z_l, \delta)$. (Indeed, we could then take $\varphi, \psi \equiv 1$ on $B(z_l, 2\delta/3)$.) Select U_i as in Proposition 3.1, with $B(z_l, 3\delta) \subseteq U_i$. Now, $\varphi(x)K_t(x, y)\psi(y)$ is the kernel of the pseudodifferential operator $\varphi G(t\sqrt{\Delta})\psi$. We can use the coordinate map ϕ_i to pull this kernel over to \mathbb{R}^n , thereby obtaining a smooth, compactly supported kernel L_t , with support in $\mathbb{R}^n \times \mathbb{R}^n$. Let us change our notation and now use x and y to denote points in \mathbb{R}^n . By Proposition 3.1, it is enough to show that:

$$t^{n+|\alpha|+|\beta|} \left(\frac{|x-y|}{t}\right)^N \left|\partial_x^{\alpha} \partial_y^{\beta} L_t(x,y)\right| \le C$$

for any multiindices α, β , for all 0 < t < 1, all $N \leq M$, and all $x, y \in \mathbb{R}^n$. Now let $p_t(x,\xi)$ denote the symbol of the operator with kernel L_t . Then

$$L_t(x,y) = \int e^{i(y-x)\cdot\xi} p_t(x,\xi) d\xi$$

Thus, $\partial_x^{\alpha} \partial_y^{\beta} L_t(x, y)$ is a finite linear combination of terms of the form

$$T = \int e^{i(y-x)\cdot\xi} \xi^{\gamma} \partial_x^{\delta} p_t(x,\xi) d\xi, \qquad (28)$$

where $|\gamma|, |\delta| \leq |\alpha| + |\beta|$.

In case (i) we may take M = 0, so we need only estimate |T|, the absolute value of the term T in (28). It will be enough to show that $|T| \leq Ct^{-n-|\gamma|}$ (for 0 < t < 1), since $Ct^{-n-|\gamma|} \leq Ct^{-n-|\alpha|-|\beta|}$. But

$$|T| \leq \int_{|\xi| \leq 1/t} |\xi^{\gamma} \partial_x^{\delta} p_t(x,\xi)| d\xi + \int_{|\xi| > 1/t} |\xi^{\gamma} \partial_x^{\delta} p_t(x,\xi)| d\xi$$
$$\leq C \left[A_t t^{-n-|\gamma|} + B_t \int_{1/t}^{\infty} r^{|\gamma|+n-1} r^{-|\gamma|-n-1} dr \right]$$
$$\leq C \left[A_t t^{-n-|\gamma|} + B_t t \right]$$

where $A_t = \sup_{x,\xi} |\partial_x^{\delta} p_t(x,\xi)|$, and $B_t = \sup_{x,\xi} |\xi|^{|\gamma|+n+1} |\partial_x^{\delta} p_t(x,\xi)|$. But, by (26) and the continuity of the map $p \to p(\sqrt{\Delta})$, from $S_1^m(\mathbb{R})$ to $OPS_{1,0}^m(\mathbb{M})$ in the cases m = 0 and $m = -(|\gamma| + n + 1)$, we find that $A_t \leq C$ (independent of 0 < t < 1) and $B_t \leq Ct^{-|\gamma|-n-1}$. Altogether $|T| \leq Ct^{-n-|\gamma|}$, as claimed. This completes the proof in case (i).

In case (ii), we need only show that for every *n*-tuple ν with $|\nu| \leq M$, we have that $|(x-y)^{\nu}T| \leq Ct^{-n-|\gamma|+|\nu|}$. Note that $(x-y)^{\nu}e^{i(y-x)\cdot\xi} = c\partial_{\xi}^{\nu}e^{i(y-x)\cdot\xi}$. Substituting this in the explicit expression for $(x-y)^{\nu}T$, and repeatedly integrating by parts in ξ , we see that $(x-y)^{\nu}T$ is a finite linear combination of terms of the form

$$T' = \int e^{i(y-x)\cdot\xi} \xi^{\kappa} \partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi) d\xi,$$

where $|\kappa| \leq |\gamma|, |\chi| \leq |\nu| \leq M$, and $|\gamma| - |\kappa| + |\chi| = |\nu|$. Just as in our estimate for T above, we see that

$$\begin{aligned} |T'| &\leq \int_{|\xi| \leq 1/t} \left| \xi^{\kappa} \partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi) \right| d\xi + \int_{|\xi| > 1/t} \left| \xi^{\kappa} \partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi) \right| d\xi \\ &\leq C \left[A_t t^{-n-|\kappa|} + B_t \int_{1/t}^{\infty} r^{|\kappa|+n-1} r^{-|\kappa|-n-1} dr \right] \\ &\leq C \left[A_t t^{-n-|\kappa|} + B_t t \right] \end{aligned}$$

where now

$$A_t = \sup_{x,\xi} \left| \partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi) \right| = \sup_{x,\xi} (1+|\xi|)^{-|\chi|+|\chi|} \left| \partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi) \right|,$$

and

$$B_t = \sup_{x,\xi} \left|\xi\right|^{|\kappa|+n+1} \left|\partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi)\right| \le \sup_{x,\xi} (1+|\xi|)^{|\kappa|+n+1-|\chi|+|\chi|} \left|\partial_x^{\delta} \partial_{\xi}^{\chi} p_t(x,\xi)\right|.$$

But, by (26) and the continuity of the map $p \to p(\sqrt{\Delta})$, from $S_1^m(\mathbb{R})$ to $OPS_{1,0}^m(\mathbf{M})$ in the cases $m = |\chi|$ and $m = |\chi| - (|\kappa| + n + 1)$, we find that $A_t \leq Ct^{|\chi|}$ and $B_t \leq Ct^{|\chi| - |\kappa| - n - 1}$. ((26) may be used here, since G vanishes to order at least M at 0, and both $|\chi|$ and $|\chi| - (|\kappa| + n + 1)$ are less than or equal to M.) Altogether $|T'| \leq Ct^{-n+|\chi|-|\kappa|} = Ct^{-n-|\gamma|+|\nu|}$, as claimed. This completes the proof.

Remark 4.2. Note that, in Lemma 4.1, the conclusion (27) holds even without the hypothesis f(0) = 0, provided t is restricted to lie in the interval (0, 1]. Indeed, after the second sentence of the proof of the lemma, we assumed 0 < t < 1 and never used the hypothesis that f(0) = 0. Of course (27) holds also for t = 1 by continuity.

5 Continuous S-Wavelets on Manifolds

We now turn to our definitions of continuous wavelets and continuous S-wavelets on \mathbf{M} , which we have motivated in the introduction.

Definition 5.1. Suppose that the function $K_t(x, y)$ is smooth for t > 0, $x, y \in \mathbf{M}$. For t > 0, define $T_t : L^2(\mathbf{M}) \to C^{\infty}(\mathbf{M})$ to be the operator with kernel K_t , so that for all $F \in L^2(\mathbf{M})$ and all $x \in \mathbf{M}$,

$$(T_tF)(x) = \int_{\mathbf{M}} K_t(x, y)F(y)d\mu(y).$$

As usual, let P denote the projection in $L^2(\mathbf{M})$ onto the space of constant functions. Then we define $K_t(x, y)$ to be a continuous wavelet on \mathbf{M} , provided the following three conditions hold, for some c > 0:

(i) For all $F \in L^2(\mathbf{M})$,

$$\int_0^\infty \|T_t F\|_2^2 \frac{dt}{t} = c \|(I - P)F\|_2^2;$$
(29)

- (ii) $\int_{\mathbf{M}} K_t(x,y) d\mu(y) = 0$ for all t > 0 and all $x \in \mathbf{M}$ (or, equivalently, $T_t(1) = 0$ for all t > 0);
- (iii) $\int_{\mathbf{M}} K_t(x,y) d\mu(x) = 0$ for all t > 0 and all $y \in \mathbf{M}$ (or, equivalently, $T_t^*(1) = 0$ for all t > 0).

Definition 5.2. Suppose $K_t(x, y)$ is a continuous wavelet on **M**. We then say that $K_t(x, y)$ is a continuous *S*-wavelet on **M**, if the following additional condition holds:

(iv) For every pair of C^{∞} differential operators X (in x) and Y (in y) on **M**, and for every integer $N \ge 0$, there exists $C_{N,X,Y}$ as follows. Suppose deg X = j and deg Y = k. Then

$$t^{n+j+k}|(\frac{d(x,y)}{t})^N XYK_t(x,y)| \le C_{N,X,Y}$$
(30)

for all t > 0 and all $x, y \in \mathbf{M}$.

We then have the following result:

Theorem 5.3. Say $f_0 \in \mathcal{S}(\mathbb{R}^+)$, $f_0 \neq 0$, and let $f(s) = sf_0(s)$. For t > 0, let K_t be the kernel of $f(t^2\Delta)$. Then $K_t(x, y)$ is a continuous S-wavelet on **M**.

Proof. Of course, condition (iv) is Lemma 4.1. As we have seen, conditions (ii) and (iii) of Definitio 5.1 are immediate consequences of (25), as for condition (i), say $F \in L^2(\mathbf{M})$. We need only take the inner product of both sides of (12) with F to see that, if $c = \int_0^\infty |f(t)|^2 \frac{dt}{t}$, then

$$\int_0^\infty \|f(t\Delta)F\|_2^2 \frac{dt}{t} = c\|(I-P)F\|_2^2.$$

Replacing t by t^2 in this equation, we find that

$$\int_0^\infty \|f(t^2\Delta)F\|_2^2 \frac{dt}{t} = \frac{c}{2} \|(I-P)F\|_2^2$$

which yields condition (i) at once. This completes the proof.

As for properties of continuous wavelets, we first remark that it is a standard, simple matter to show the following result, which generalizes (12) (with $T = \Delta$ there):

Proposition 5.4. Suppose $K_t(x, y)$ is a continuous wavelet on \mathbf{M} , and, for t > 0, let T_t be the operator on $L^2(\mathbf{M})$ with kernel K_t . Then for any $F \in (I - P)L^2(\mathbf{M})$, we may reconstruct F through the identity

$$\int_0^\infty T_t^* T_t F \frac{dt}{t} = cF.$$
(31)

Here the integral on the left side of (31) conveges unconditionally in L^2 .

Proof. Let \mathcal{H}_1 be the Hilbert space $(I - P)L^2(\mathbf{M})$, and let \mathcal{H}_2 be the Hilbert space $L^2(\mathbb{R}^+, \mathcal{H}_1, dt/t)$. By our definition of continuous wavelet, we may define a bounded operator $U : \mathcal{H}_1 \to \mathcal{H}_2$ by

$$UF = (T_t F)_{t>0}$$

19

Moreover, we may define a bounded operator $V : \mathcal{H}_2 \to \mathcal{H}_1$ by

$$V(G_t)_{t>0} = \int_0^\infty T_t^* G_t \frac{dt}{t}$$

where the integral converges unconditionally in \mathcal{H}_1 . Indeed, let $\| \|$ denote $\| \|_{\mathcal{H}_1}$, and let $S = \{F \in \mathcal{H}_1 : \|F\| = 1\}$. If $E \subseteq (0, \infty)$ is measurable and contained in a compact subset of $(0, \infty)$, we have

$$\begin{split} \| \int_{E} T_{t}^{*} G_{t} \frac{dt}{t} \| &= \sup_{F \in S} | \int_{E} \langle T_{t}^{*} G_{t}, F \rangle \frac{dt}{t} | = \sup_{F \in S} | \int_{E} \langle G_{t}, T_{t} F \rangle \frac{dt}{t} | \\ &\leq \left[\int_{E} \| G_{t} \|^{2} \frac{dt}{t} \right]^{1/2} \left[\sup_{F \in S} | \int_{E} \| T_{t} F \|^{2} \frac{dt}{t} \right]^{1/2}; \end{split}$$

but, by (29), this is less than or equal to $\left[\int_{E} \|G_t\|^2 \frac{dt}{t}\right]^{1/2}$, and the unconditional convergence follows. One now readily checks that $V = U^*$. By (29), $\langle U^*UF, F \rangle = c \|F\|^2$ for all $F \in \mathcal{H}_1$. Polarizing this identity, we find (31), as desired.

As an example of the usefulness of continuous S-wavelets, we now show the following direct analogue of a theorem of Holschneider and Tchamitchian ([25]):

Theorem 5.5. Let $K_t(x, y)$ be a continuous S-wavelet on \mathbf{M} , and, for t > 0, let T_t be the operator on L^2 with kernel K_t . Suppose $F \in L^2(\mathbf{M})$. Then:

(a) If F is Hölder continuous, with Hölder exponent α (0 < $\alpha \leq 1$), then for some C > 0,

$$\|T_t F\| \le C t^{\alpha} \tag{32}$$

for all t > 0. (Here || || denotes sup norm.)

(b) Conversely, say $0 < \alpha < 1$, C > 0, and that F satisfies (32) for all t > 0. Then F is Hölder continuous, with Hölder exponent α .

Proof. For (a), we just note:

$$\begin{aligned} |(T_t F)(x)| &= \left| \int F(y) K_t(x, y) d\mu(y) \right| \\ &= \left| \int (F(x) - F(y)) K_t(x, y) d\mu(y) \right| \\ &\leq C t^{-n} \int d(x, y)^{\alpha} \left[1 + d(x, y)/t \right]^{-n-1-\alpha} d\mu(y) \\ &\leq C t^{\alpha-n} \int \left[1 + d(x, y)/t \right]^{-n-1} d\mu(y) \\ &\leq C t^{\alpha} \end{aligned}$$

by (21), as desired.

For (b), of course PF, being constant, is Hölder continuous of any exponent. Since $T_t 1 = 0$, we may assume that F = (I - P)F. Set $g_t = T_t F$, so that

$$\|g_t\| \le Ct^{\alpha}.\tag{33}$$

For $x, y \in \mathbf{M}$, set

$$K_{t,x}(y) = K_t^y(x) = K_t(x,y).$$

Thus, for all $x, g_t(x) = \langle K_{t,x}, \overline{F} \rangle$. Since we are assuming $F \in L^2$, by Cauchy-Schwartz and (30), we obtain the additional estimate

$$\|g_t\| \le Ct^{-n}.\tag{34}$$

By (30), we have that $|K_t(x,y)| \le Ct^{-n}(1 + d(x,y)/t)^{-n-1}$, so by (21),

$$\|K_t^y\|_1 \le C \tag{35}$$

for any y. Thus, for any y,

$$|(T_t^*T_tF)(y)| = |\langle g_t, K_t^y \rangle| \le ||g_t|| ||K_t^y||_1 \le C||g_t||.$$

Accordingly, by (33) and (34), for any y,

$$\int_{0}^{\infty} |(T_{t}^{*}T_{t}F)(y)| \frac{dt}{t} \leq C\left(\int_{0}^{1} t^{\alpha-1}dt + \int_{1}^{\infty} t^{-n-1}dt\right) \leq C$$

By Proposition 5.4, we now see that for almost every y,

$$cF(y) = \int_0^\infty T_t^* T_t F(y) \frac{dt}{t},$$

and from this, that $F \in L^{\infty}$.

To complete the proof, we claim that it suffices to show that if $d(y, z) \leq \min(t, \delta)$, then

$$\int_{\mathbf{M}} |K_t(x,y) - K_t(x,z)| d\mu(x) \le C d(y,z)/t.$$
(36)

For then, by (33), (34), (35) and (36), we would have, if $d(y, z) < \delta$, then

$$\begin{aligned} c|F(y) - F(z)| &\leq \int_0^\infty \int_{\mathbf{M}} |K_t(x,y) - K_t(x,z)| d\mu(x) \|g_t\| \frac{dt}{t} \\ &\leq C \left[\int_0^{d(y,z)} t^{\alpha - 1} dt + d(y,z) \int_{d(y,z)}^1 t^{\alpha - 2} dt + d(y,z) \int_1^\infty t^{-n-2} dt \right] \\ &\leq C d(y,z)^\alpha, \end{aligned}$$

as needed.

To prove (36), choose $U_i \supseteq B(y, 3\delta)$, and let us work in the local coordinates on U_i obtained from ϕ_i . We use the mean value theorem. By (30), for any $x \in \mathbf{M}$, there is point w_x on the line segment joining y to z such that

$$|K_t(x,y) - K_t(x,z)| \le C|y - z|t^{-n-1}(1 + d(x,w_x)/t)^{-n-1}.$$

By Proposition 3.1, if w is any point on that line segment,

$$d(y,w) \le c_2 |y-w| \le c_2 |y-z| \le c_1 c_2 d(y,z) \le c_1 c_2 t.$$

Thus the diameter of the line segment is at most $2c_1c_2t$, and so, by (24), we have

$$|K_t(x,y) - K_t(x,z)| \le Cd(y,z)t^{-n-1}(1 + d(x,y)/t)^{-n-1}.$$

(36) now follows from (21), as desired.

6 Homogeneous Manifolds

In this section look at the situation in which **M** has a transitive group G of smooth metric isometries. (Such manifolds are usually called homogneous manifolds.) Obvious examples of such manifolds are the sphere S^n , where we take G to be the group SO(n+1) of rotations, and the torus $\mathbf{T}^n = (S^1)^n$, where we take G to be the group $[SO(2)]^n$.

If $T \in G$ and F is a function on \mathbf{M} , we define the function TF on \mathbf{M} by $(TF)(x) = F(T^{-1}x)$. Then $T: L^2(\mathbf{M}) \to L^2(\mathbf{M})$ is a unitary operator which commutes with the Laplace-Beltrami operator Δ . Consequently, as operators on $L^2(\mathbf{M})$, $f(t^2\Delta)$ commutes with elements of G for any bounded Borel function f on \mathbb{R} , and in particular, if $f \in \mathcal{S}(\mathbb{R})$, which we now assume.

Thus, if $T \in G$, $F \in L^2(\mathbf{M})$ and $x \in \mathbf{M}$, we have

$$\int_{\mathbf{M}} K_t(Tx, Ty)F(y)d\mu(y) = \int_{\mathbf{M}} K_t(Tx, y)F(T^{-1}y)d\mu(y)$$
$$= [f(t^2\Delta)(TF)](Tx)$$
$$= T([f(t^2\Delta)(F)])(Tx);$$

but this is just $[f(t^2\Delta)(F)](x) = \int_{\mathbf{M}} K_t(x,y)F(y)d\mu(y)$, so

$$K_t(Tx, Ty) = K_t(x, y) \tag{37}$$

for all $x, y \in \mathbf{M}$. Thus, since G is transitive, if x_0 is any fixed point in \mathbf{M} , once one knows $K_t(x_0, y)$ for all y, then one knows $K_t(x, y)$ for all x, y.

In the analogous situation on \mathbb{R}^n , $K_t(x, y)$ has the form $t^{-n}\psi((x-y)/t)$ for some $\psi \in \mathcal{S}$, and so $K_t(x, y) = K_t(Tx, Ty)$ for any translation T on \mathbb{R}^n . Equation (37) is a natural analogue of this fact for **M**. It is interesting to note that one has

$$K_t(x,x) = \operatorname{tr}(f(t^2\Delta))/\operatorname{vol}(\mathbf{M})$$
(38)

for all x and all $f \in \mathcal{S}(\mathbb{R}^+)$. Indeed, by (37), $K_t(x, x)$ is constant for $x \in \mathbf{M}$. Accordingly

$$vol(\mathbf{M})K_{t}(x,x) = \int_{\mathbf{M}} K_{t}(y,y)d\mu(y) = \sum_{l} f(t^{2}\lambda_{l}) \int_{\mathbf{M}} |u_{l}(y)|^{2}d\mu(y) = \sum_{l} f(t^{2}\lambda_{l}) = tr(f(t^{2}\Delta))$$

as claimed.

Say now c > 0, and let us look at the special case

$$f(s) = (s/c)e^{-(s/c)}.$$
(39)

We have $\operatorname{tr}(f(t^2\Delta)) = \operatorname{tr}((t^2/c)\Delta e^{-(t^2\Delta)/c})$. A well known fact, usually associated with the heat kernel approach to index theorems ([33], [20], [37] and [21], pages 58 and 316), is that as $s \to 0^+$,

$$\operatorname{tr}(e^{-s\Delta}) \sim \sum_{m=0}^{\infty} s^{m-n/2} a_m,\tag{40}$$

where

$$a_0 = (4\pi)^{-n/2} \operatorname{vol}(\mathbf{M}).$$
 (41)

Differentiating with respect to s one finds that

$$\operatorname{tr}(\Delta e^{-s\Delta}) \sim \sum_{m=0}^{\infty} (\frac{n}{2} - m) s^{m-1-n/2} a_m$$

To lowest order, then,

$$K_t(x,x) = \operatorname{tr}((t^2/c)\Delta e^{-(t^2\Delta)/c})/\operatorname{vol}(\mathbf{M}) \sim \frac{nct^{-n}}{2(4\pi)^{n/2}}.$$
(42)

Again let f be general, but now let us look at the special case $\mathbf{M} = \mathbf{T}^n$, the torus. We write $\mathbf{T}^n = \{(e^{2\pi i r_1}, \ldots, e^{2\pi i r_n}) : -1/2 < r_1, \ldots, r_n \leq 1/2\}$. Here an orthonormal basis of eigenfunctions is given simply by $\{e^{2\pi i m \cdot r} : m \in \mathbb{Z}^n\}$. The Laplace-Beltrami operator Δ is just $-\sum_{l=1}^n (\partial/\partial r_l)^2$, and the eigenvalues are given through $\Delta e^{2\pi i m \cdot r} = 4\pi^2 ||m||^2 e^{2\pi i m \cdot r}$. (Here $||m||^2 = m_1^2 + \ldots m_n^2$.) The kernel $K_t(r, s)$ of $f(t^2\Delta)$ is given by

$$K_t(r,s) = \sum_{m \in \mathbb{Z}^n} f(4\pi^2 t^2 ||m||^2) e^{2\pi i m \cdot (r-s)}.$$

Thus, if $F \in L^2(\mathbf{T}^n)$,

$$[f(t^{2}\Delta)F](r) = \int_{-1/2}^{1/2} \dots \int_{-1/2}^{1/2} F(s)K_{t}(r,s)ds_{1}\dots ds_{n} = [F * h_{t}(r)]$$

where

$$h_t(s) = \sum_{m \in (\mathbb{Z})^n} f(4\pi^2 t^2 ||m||^2) e^{2\pi i m \cdot s}$$

and \ast denotes the natural convolution on $\mathbf{T}^n.$

We specialize now to the case n = 2, $f(u) = ue^{-u/4\pi}/4\pi^2$. We free the letter n for other uses. For t > 0 define the functions $U_t, V_t : \mathbb{R} \to \mathbb{C}$ by

$$U_t(x) = \sum_{n = -\infty}^{\infty} e^{-\pi t^2 n^2} e^{2\pi i n x},$$
(43)

$$V_t(y) = \sum_{n=-\infty}^{\infty} (nt)^2 e^{-\pi t^2 n^2} e^{2\pi i n y}.$$
(44)

It is then easy to calculate that

$$h_t(s_1, s_2) = U_t(s_1)V_t(s_2) + U_t(s_2)V_t(s_1).$$

 h_t is the "Mexican hat" on the torus \mathbf{T}^2 . One can use these equations to draw its graph. One of course needs to approximate the series in (43) and (44) by finite sums. This is a simple matter if t is greater than 1, but if t is small the series do not converge very quickly. Fortunately, however, we can give alternative series expansions for U_t and V_t which do converge very quickly for 0 < t < 1. We do this by using the Poisson summation formula, or rather, its proof: if $g \in \mathcal{S}(\mathbb{R})$, then the periodic function $G(x) = \sum_{n=-\infty}^{\infty} g(x+n)$ has Fourier series $\sum_{n=-\infty} \check{g}(n)e^{2\pi inx}$, and hence, G(x) equals the



Figure 1: h_t on \mathbf{T}^2 for t = 2 (left), t = 1/2 (middle), t = 1/8 (right)

latter series. (Here we use the inverse Fourier transform $\check{g}(x) = \int_{-\infty}^{\infty} g(\xi) e^{-2\pi i x \xi} d\xi$.) Applying this with $\check{g}(y) = e^{-\pi t^2 y^2}$, we obtain the formula

$$U_t(x) = \frac{1}{t} \sum_{n = -\infty}^{\infty} e^{-\pi (n+x)^2/t^2}.$$
(45)

Taking instead $\check{g}(y) = (ty)^2 e^{-\pi t^2 y^2}$, we obtain the formula

$$V_t(x) = \frac{1}{t} \sum_{n=-\infty}^{\infty} \left(\frac{1}{2\pi} - \left(\frac{n+x}{t}\right)^2\right) e^{-\pi(n+x)^2/t^2}.$$
(46)

(45) and (46) converge very quickly for t small, making it practical to draw pictures of h_t for t small. We include pictures, obtained by using Maple, of the Mexican hat functions $\pi h_t(r_1, r_2) - 1/2 < r_1, r_2 \leq 1/2$, for t = 2 (Figure 1, left), t = 1/2 (Figure 1, middle) and t = 1/8 (Figure 1, right). Note that the characteristic Mexican hat shape is obtained. To keep the pictures uncluttered, we have omitted the axes. However, one can ask Maple to evaluate $t^2\pi h_t(0,0)$ for various values of t. When t = 2, it is .00070; when t = 1, it is .59017; when t = 1/2, it is .99984, and when t = 1/8, it is 1.00000. This is consistent with the predictions of (39) with $c = 4\pi$, and (42).

Now let us look at the special case $\mathbf{M} = S^n$, and let f be general for now. Let $\mathbf{N} = (1, 0, \dots, 0)$, the "north pole". We shall now calculate $K_t(\mathbf{N}, y)$ as an explicit infinite series. (As we have explained, we

24

will then know $K_t(x, y)$ for all x, y.) Recall ([42]) that we may write $L^2(S^n) = \bigoplus_{l \ge 0} \mathcal{H}_l$, where \mathcal{H}_l is the space of spherical harmonics of degree l. If $P \in \mathcal{H}_l$, then

$$\Delta P = l(l+n-1)P.$$

Within each space \mathcal{H}_l is a unique zonal harmonic Z_l , which has the property that for all $P \in \mathcal{H}_l$, $P(\mathbf{N}) = \langle P, Z_l \rangle$. In particular, P is orthogonal to Z_l if and only if $P(\mathbf{N}) = 0$.

We wish to use (25) to evaluate $K_t(\mathbf{N}, y)$. To do so we choose an orthonormal basis for each \mathcal{H}_l , one of whose elements is $Z_l/||Z_l||_2$. Then any other element of this orthonormal basis vanishes at \mathbf{N} , so we find

$$K_t(\mathbf{N}, y) = \sum_{l=0}^{\infty} f(t^2 l(l+n-1)) Z_l(\mathbf{N}) Z_l(y) / \|Z_l\|_2^2$$

But surely $Z_l(\mathbf{N}) = \langle Z_l, Z_l \rangle$, so we simply have

$$K_t(\mathbf{N}, y) = \sum_{l=0}^{\infty} f(t^2 l(l+n-1)) Z_l(y).$$
(47)

However, $Z_l(y)$ is known explicitly. In fact ([42]), if ω_n is the area of S^n , then for some constant c_l ,

$$Z_l(y) = c_l P_l^{\lambda}(y_1), \tag{48}$$

where $y = (y_1, \ldots, y_n)$, $\lambda = (n-1)/2$, and P_l^{λ} is the ultraspherical (or Gegenbauer) polynomial of degree l associated with λ . The P_l^{λ} may be defined in terms of the generating function

$$(1 - 2r\tau + r^2)^{-\lambda} = \sum_{l=0}^{\infty} P_l^{\lambda}(\tau) r^l.$$
(49)

In particular, if $\tau = 1$, we see that

$$(1-r)^{-(n-1)} = \sum_{l=0}^{\infty} P_l^{\lambda}(1)r^l$$

so that

$$P_l^{\lambda}(1) = \binom{n+l-2}{l} = b_l \tag{50}$$

On the other hand,

$$Z_l(\mathbf{N}) = [\omega_n]^{-1} \dim \mathcal{H}_l = [\omega_n]^{-1} \left[\binom{n+l}{n} - \binom{n+l-2}{l} \right] = d_l.$$
(51)

Comparing (48),(50) and (51), we see that

$$c_l = d_l / b_l = \frac{\binom{n+l}{n} - \binom{n+l-2}{n}}{\omega_n \binom{n+l-2}{l}} = \frac{n+2l-1}{\omega_n (n-1)}.$$
(52)

From (47), we find

$$K_t(\mathbf{N}, y) = \sum_{l=0}^{\infty} \frac{(n+2l-1)}{\omega_n(n-1)} f(t^2 l(l+n-1)) P_l^{\lambda}(y_1) := h_t(y_1).$$
(53)

If $x \in S^n$, we can choose a rotation T with $Tx = \mathbf{N}$. If also $y \in S^n$, then by (37),

$$K_t(x,y) = K_t(Tx,Ty) = K_t(\mathbf{N},Ty) = h_t((Ty)_1) = h_t(\mathbf{N}\cdot Ty) = h_t(Tx\cdot Ty) = h_t(x\cdot y).$$

Thus, if $F \in L^2(S^n)$,

$$[f(t^2\Delta)F](x) = \int_{S^n} F(y)h_t(x \cdot y)d\mu(y),$$

the spherical convolution of F and the axisymmetric function h_t . Let us now take $f(s) = se^{-s}$. From (53), we find

$$h_t(y_1) = K_t(\mathbf{N}, y) = \sum_{l=0}^{\infty} \frac{l(l+n-1)(n+2l-1)}{\omega_n(n-1)} t^2 e^{-t^2 l(l+n-1)} P_l^{\lambda}(y_1).$$
(54)

One can use (54) to draw the graph of h_t for any t > 0. We do so, in the most practical situation, n = 2. We can go all around a great circle by using spherical coordinates, in which $y_1 = \cos(\theta)$, with θ going from $-\pi$ to π . We draw these graphs when t = 1 (Figure 2, left), t = .1 (Figure 2, middle) and t = .05 (Figure 2, right). Actually, since (42) predicts

$$4\pi h_t(1) = 4\pi K_t(\mathbf{N}, \mathbf{N}) \sim 1/t^2,$$

we draw the graphs of $4\pi h_t(\cos(\theta))$ instead, with θ going from $-\pi$ to π on the horizontal axis. The pictures do bear out the relation $4\pi h_t(1) \sim 1/t^2$. We also see the characteristic "Mexican hat" shape, familiar from the analogous situation on \mathbb{R}^1 , where $K_t(x, y)$ is the Schwartz kernel of $f(-t^2d^2/dx^2)$. Of course, in that situation, $[f(-t^2d^2/dx^2)F]^{(\xi)} = f(t^2\xi^2)\hat{F}(\xi) = t^2\xi^2e^{-t^2\xi^2}\hat{F}(\xi)$ for $F \in \mathcal{S}(\mathbb{R})$, so $K_t(x, y) = t^{-1}\psi((x-y)/t)$, where ψ is the second derivative of a Gaussian.

The series (54) converges quickly for t large, but not if t is small. Therefore, as in the case of the torus, for computational purposes, it is important to have a quickly converging alternate expression for this function for small t. Since the pictures indicate that $4\pi h_t(\cos\theta)$ is negligible outside a small neighborhood of $\theta = 0$ if t is small, one would assume that it is only necessary to compute the Maclaurin series of h_t . It is very fortunate that any number of terms of this series can be computed explicitly, through use of the work of Polterovich [38], together with some additional insights.

In fact, what Polterovich found in [38] was the heat trace asymptotics on the sphere, i.e. explicit formulae for the a_m of (40) for the manifold S^n , as sums of explicit finite series, for $m \ge 1$. (Earlier, less explicit formulae were found earlier in ([4]) and ([5]).) Using Polterovich's formula to evaluate some of these a_m on S^2 , (it is easiest to use Maple), we find that

$$\operatorname{tr}(e^{-s\Delta}) \sim \frac{1}{s} + \frac{1}{3} + \frac{s}{15} + \frac{4s^2}{315} + \frac{s^3}{315} + O(s^4);$$
(55)

there would be no difficulty in evaluating more terms. (Recall that a_0 is given by (41), for general **M**.) Using this formula we are going to evaluate the first few terms of the Maclaurin series of $4\pi h_t(\cos\theta)$, and we will show how any number of terms could be obtained.

Let $J_t(x, y)$ be the kernel of $e^{-t^2\Delta}$ on S^2 , so that, in particular, by (38), $J_t(x, x) = \operatorname{tr}(e^{-t^2\Delta})/4\pi$. From (53), with $f(r) = e^{-r}$, we find that if $s = t^2$, then

$$J_t(\mathbf{N}, y) = \sum_{l=0}^{\infty} \frac{(2l+1)}{4\pi} e^{-sl(l+1)} P_l^{\lambda}(y_1) := g_t(y_1).$$
(56)



Figure 2: $4\pi h_t(\cos\theta)$ on S^2 for t = 1 (left), t = .1 (middle), t = 0.05 (right)

and similarly, from (54),

$$\frac{1}{s}K_t(\mathbf{N}, y) = \sum_{l=0}^{\infty} \frac{l(l+1)(2l+1)}{4\pi} e^{-sl(l+1)} P_l^{\lambda}(y_1) = \frac{1}{s}h_t(y_1).$$
(57)

We would like to understand the Maclaurin series of $4\pi g_t(\cos\theta)$ and $4\pi h_t(\cos\theta)$. To this end, we shall use the following lemma.

Lemma 6.1. Suppose $U \in C^2(S^n)$ is a function of x_1 only, $U(x) = u(x_1)$, say, and that u is in fact C^2 in a neighborhood of [-1, 1]. Then ΔU is also a function of x_1 , and

$$(\Delta U)(x) = nx_1 u'(x_1) - (1 - x_1^2) u''(x_1).$$
(58)

In particular, if n = 1, one has

$$-\frac{d^2}{d\theta^2}[u(\cos\theta)] = (\cos\theta)u'(\cos\theta) - (\sin^2\theta)u''(\cos\theta).$$
(59)

Proof. By the results of [14], $\Delta = -\sum_{j < k} W_{jk}^2$, where

$$W_{jk} = x_j \frac{\partial}{\partial x_k} - x_k \frac{\partial}{\partial x_j}$$

if $1 \leq j, k \leq n+1$. Since U is a function of x_1 only we compute

$$\Delta U = -\sum_{k=1}^{n+1} W_{1k}^2 U$$
$$= -\sum_{j=2}^{n+1} [-x_1 u' + x_j^2 u'']$$

which proves (58), since on the unit sphere, $\sum_{j=2}^{n+1} x_j^2 = 1 - x_1^2$. Of course (59) is elementary, but we note that it can also be viewed as a special case of (58), since, in polar coordinates, the spherical Laplacian on S^1 is just $-\frac{d^2}{d\theta^2}$. This completes the proof.

In particular, if we set $x_1 = 1$ in (58), and $\theta = 0$ in (59), we find that

$$-\frac{d^2}{d\theta^2}[u(\cos\theta)]|_{\theta=0} = \frac{1}{n}(\Delta U)(\mathbf{N}),\tag{60}$$

since both sides equal u'(1).

Remark 6.2. We shall show below, in Lemma 6.3, that if u is C^{2m} on an open interval containing [-1,1], one can similarly obtain $\frac{d^{2m}}{d\theta^{2m}}[u(\cos\theta)]|_{\theta=0}$ from a knowledge of $(\Delta^i U)(\mathbf{N})$ for $i = 1, \ldots, m$. Thus, if u is C^{∞} on an open interval containing [-1,1], the entire Maclaurin series of $u(\cos\theta)$ (regarded as a function of θ) is completely determined from a knowledge of $(\Delta^i U)(\mathbf{N})$ for $i \ge 0$. (Of course, $u(\cos\theta)$ is an even function of θ , so all of its odd-order derivatives vanish at 0.)

For now, let us apply this lemma to $u = g_t$ as in (56) or $\frac{1}{s}h_t$ as in (57). To do so, we first explain why g_t and h_t have C^{∞} extensions to an open neighborhood of [-1, 1]. For this, it is evidently sufficient to show that for every $n \ge 1$ and every $m \ge 0$, there exists $c_{n,m}$ with

$$\left|\frac{d^m}{d\tau^m}P_l^{(n-1)/2}(\tau)\right| \le c_{n,m}l^{n+2m-1}.$$

From the generating function formula (49), we see the classical formula that the derivative of P_l^{λ} is $2\lambda P_{l-1}^{\lambda+1}$. Thus we may assume m = 0. By (48) and (52), we need only show that, on S^n , $|Z_l(y)| \leq Cl^n$, and for this, by (51), we need only show that $|Z_l(y)| \leq Z_l(\mathbf{N})$ for all $y \in S^n$. But we may choose an orthogonal transformation T on \mathbb{R}^n with $T\mathbf{N} = y$, and then

$$Z_{l}(y) = Z_{l} \circ T(\mathbf{N}) = \langle Z_{l} \circ T, Z_{l} \rangle \le ||Z_{l} \circ T||_{2} ||Z_{l}||_{2} = ||Z_{l}||_{2}^{2} = Z_{l}(\mathbf{N}),$$

as claimed.

Let us return to S^2 . Suppose as usual that t > 0, and again put $s = t^2$. By (60) and (56) we have that

$$\begin{aligned} -\frac{d^2}{d\theta^2} 4\pi [g_t(\cos\theta)]|_{\theta=0} &= \frac{4\pi}{2} (\Delta_y J_{\sqrt{s}})(\mathbf{N},\mathbf{N}) \\ &= -\frac{4\pi}{2} \frac{d}{ds} J_{\sqrt{s}}(\mathbf{N},\mathbf{N}) \\ &= -\frac{1}{2} \frac{d}{ds} \operatorname{tr}(e^{-s\Delta}). \end{aligned}$$

Thus, the first few terms of the Maclaurin series of $4\pi [g_t(\cos\theta)]$ are

$$4\pi g_t(\cos\theta) \sim \operatorname{tr}(e^{-s\Delta}) + \frac{\theta^2}{4} \frac{d}{ds} \operatorname{tr}(e^{-s\Delta}).$$

Similarly

$$\begin{aligned} -\frac{d^2}{d\theta^2} 4\pi \left[\frac{1}{s}h_t(\cos\theta)\right]|_{\theta=0} &= \frac{4\pi}{2} \left(\Delta_y \frac{1}{s}K_{\sqrt{s}}\right)(\mathbf{N},\mathbf{N}) \\ &= -\frac{4\pi}{2} \frac{d}{ds} \frac{1}{s}K_{\sqrt{s}}(\mathbf{N},\mathbf{N}) \\ &= +\frac{1}{2} \frac{d^2}{ds^2} \mathrm{tr}(e^{-s\Delta}). \end{aligned}$$

Thus, the first few terms of the Maclaurin series of $4\pi [g_t(\cos\theta)]$ and $4\pi [h_t(\cos\theta)]$ are

$$4\pi g_t(\cos\theta) \sim \operatorname{tr}(e^{-s\Delta}) + \frac{\theta^2}{4} \frac{d}{ds} \operatorname{tr}(e^{-s\Delta}).$$
(61)

$$4\pi h_t(\cos\theta) \sim -s\frac{d}{ds}\operatorname{tr}(e^{-s\Delta}) - \frac{\theta^2}{4}s\frac{d^2}{ds^2}\operatorname{tr}(e^{-s\Delta}).$$
(62)

In order to put these into a useful form, we now invoke a result of Kannai [27].

Set $z(\theta) = (\cos \theta, \sin \theta, 0, \dots, 0) \in S^n$. Since $g_t(\cos \theta)$ equals $J_t(\mathbf{N}, z(\theta))$, where $J_t(x, y)$ is the kernel of the heat operator $e^{-s\Delta}$, Kannai's results imply that

$$4\pi g_t(\cos\theta) \sim \frac{1}{s} e^{-\theta^2/4s} \sum_{j=0}^{\infty} v_j(\theta) s^j$$
(63)

at least for $0 < |\theta| < \pi$, and where the v_j are smooth functions. (Kannai's result for general **M**, proved through use of the Hadamard parametrix for the wave equation, is that the kernel of $e^{-s\Delta}$ has the form

$$\frac{1}{(4\pi s)^{n/2}}e^{-d(x,y)^2/4s}\sum_{j=0}^{\infty}V_j(x,y)s^j$$

for x sufficiently close to y, where the V_j are smooth. In our case the geodesic distance from **N** to $z(\theta)$ is just $|\theta|$, and we have set $v_j(\theta) = V_j(\mathbf{N}, z(\theta))$.) From (63) and the fact that (by (56) and (57)), one has

$$\frac{\partial}{\partial s}g_t(\cos\theta) = -\frac{1}{s}h_t(\cos\theta),\tag{64}$$

we are now motivated to find the first few terms in the Maclaurin series (in θ) of $e^{\theta^2/4s} 4\pi g_t(\cos\theta)$ and $e^{\theta^2/4s} 4\pi h_t(\cos\theta)$. For this, we need only multiply the right sides of (61) and (62) by

$$e^{\theta^2/4s} \sim \frac{1}{s}(s+\theta^2/4+\cdots).$$

This yields

$$e^{\theta^2/4s} 4\pi g_t(\cos\theta) \sim \frac{1}{s} [\operatorname{str}(e^{-s\Delta}) + \frac{\theta^2}{4} [s\frac{d}{ds} \operatorname{tr}(e^{-s\Delta}) + \operatorname{tr}(e^{-s\Delta})]$$

and

$$e^{\theta^2/4s} 4\pi h_t(\cos\theta) \sim -\frac{1}{s} [s^2 \frac{d}{ds} \operatorname{tr}(e^{-s\Delta}) + \frac{\theta^2}{4} [s^2 \frac{d^2}{ds^2} \operatorname{tr}(e^{-s\Delta}) + s \frac{d}{ds} \operatorname{tr}(e^{-s\Delta})], \tag{65}$$

where the error, for any fixed s, is $O(\theta^4)$. Combining this with Polterovich's result (55), and putting $s = t^2$ again, we obtain the approximations

$$4\pi g_t(\cos\theta) \sim \frac{e^{-\theta^2/4s}}{s} \left[\left(1 + \frac{s}{3} + \frac{s^2}{15} + \frac{4s^3}{315} + \frac{s^4}{315}\right) + \frac{\theta^2}{4} \left(\frac{1}{3} + \frac{2s}{15} + \frac{4s^2}{105} + \frac{4s^3}{315}\right) \right]$$
(66)

It would take considerably more analysis to estimate the error here, but Maple says that when t = .1, the error is never more than 6×10^{-4} for any $\theta \in [-\pi, \pi]$, even though both sides have a maximum of about 100. Maple says that the greatest error occurs at around $\theta = .3$, where both sides are about 10.655.

Similarly one could use (65) to give an approximation to $4\pi h_t(\cos\theta)$. But Maple says that the errors are smaller if we differentiate formally with respect to s in (66) (and then multiply by -s); here we are recalling (64). If we do this and finally replace s by t^2 , this yields the approximation

$$4\pi h_t(\cos\theta) \sim \frac{e^{-\theta^2/4t^2}}{t^2} [(1 - \frac{\theta^2}{4t^2})p(t,\theta) - t^2q(t,\theta)],$$
(67)

where

$$p(t,\theta) = 1 + \frac{t^2}{3} + \frac{t^4}{15} + \frac{4t^6}{315} + \frac{t^8}{315} + \frac{\theta^2}{4}(\frac{1}{3} + \frac{2t^2}{15} + \frac{4t^4}{105} + \frac{4t^6}{315})$$

and

$$q(t,\theta) = \frac{1}{3} + \frac{2t^2}{15} + \frac{4t^4}{105} + \frac{4t^6}{315} + \frac{\theta^2}{4}(\frac{2}{15} + \frac{8t^2}{105} + \frac{4t^4}{105})$$

This approximation differs from the one obtained from (65) only in terms which are fourth order in θ . Maple says that when t = .1, the error in the approximation (67) is never more than 9.5×10^{-4} for any

31

 $\theta \in [-\pi, \pi]$, even though both sides have a maximum of about 100. Maple says that the greatest error occurs at around $\theta = .4$, where both sides are about -5.593. Of course, if in (67) we approximate $p \sim 1$ and $q \sim 0$, we would obtain the formula for the usual Mexican hat wavelet on the real line, as a function of θ .

Let us then explain how one can readily obtain any number of terms of the Maclaurin series of g_t and h_t . Lemma 6.3. For any positive integer m, there are constants a_1, \ldots, a_m as follows. If, in the situation of Lemma 6.1, u is C^{2m} on an open interval containing [-1, 1], then

$$\frac{d^{2m}}{d\theta^{2m}}[u(\cos\theta)]|_{\theta=0} = \sum_{i=1}^m a_i(\Delta^i U)(\mathbf{N}).$$

Moreover, $a_m \neq 0$.

Proof. It is enough to show that, generalizing (58), there are polynomials p_1, \ldots, p_{2m} in x_1 such that

$$(\Delta^m U)(x) = \sum_{i=1}^m p_i(x_1)u^{(i)}(x_1) + \sum_{i=m+1}^{2m} (1 - x_1^2)^{i-m} p_i(x_1)u^{(i)}(x_1)$$
(68)

where $p_m(1) \neq 0$. For then the mapping

$$(u'(1), \cdots, u^{(m)}(1)) \mapsto ((\Delta U)(\mathbf{N}), \cdots, (\Delta^m U)(\mathbf{N}))$$

will be given by an invertible upper triangular matrix. In particular this is true if n = 1, so that the mapping

$$\left(u'(1),\cdots,u^{(m)}(1)\right)\mapsto \left(\frac{d^2}{d\theta^2}[u(\cos\theta)]|_{\theta=0},\cdots,\frac{d^{2m}}{d\theta^{2m}}[u(\cos\theta)]|_{\theta=0}\right)$$

is also given by an invertible upper triangular matrix. Thus the map

$$((\Delta U)(\mathbf{N}), \cdots, (\Delta^m U)(\mathbf{N})) \mapsto \left(\frac{d^2}{d\theta^2} [u(\cos\theta)]|_{\theta=0}, \cdots, \frac{d^{2m}}{d\theta^{2m}} [u(\cos\theta)]|_{\theta=0}\right)$$

is also given by an invertible upper triangular matrix, which is the desired result.

To prove (68), we recall first that by (58), $(\Delta U)(x) = Du(x_1)$, where $D = nx_1 \frac{d}{dx_1} - (1 - x_1^2) \frac{d^2}{dx_1^2}$. From this, (68), save for the statement that $p_m(1) \neq 0$, follows at once by a simple induction on m. If $p_m(1)$ were zero, then the mapping

$$(u'(1), \cdots, u^{(m)}(1)) \mapsto ((\Delta U)(\mathbf{N}), \cdots, (\Delta^m U)(\mathbf{N}))$$

would be given by a singular upper triangular matrix, so its range would not be all of \mathbb{R}^m . But the range is all of \mathbb{R}^m , since it contains all the vectors

$$\left((\Delta Z_l)(\mathbf{N}), \dots, (\Delta^m Z_l)(\mathbf{N}) \right) = Z_l(\mathbf{N})l(l+n-1)(1, l(l+n-1), \dots, [l(l+n-1)]^{m-1}),$$

for l = 1, ..., m, and since the rows of a Vandermonde matrix are linearly independent. This contradiction completes the proof.

To obtain further terms of the Maclaurin series of g_t or h_t , one need only use Lemma 6.3 together with the fact that applying Δ^i to the series in (56) or (57) is the same as applying $(-\partial/\partial s)^i$. Then Polterovich's formula for the heat trace asymptotics may be used.

Remark 6.4. Throughout this discussion, we have been considering (53) for $f(s) = se^{-s}$. If one instead used $f_m(s) = s^m e^{-s}$, where m is a nonnegative integer, one could still obtain explicit formulas, similar to (67), giving "generalized" Mexican hat wavelets on the sphere.) Indeed, if h_t^m corresponds to f_m in the same way that h_t corresponds to $f = f_1$, we have in place of (64) that $\frac{\partial^m}{\partial s^m}g_t(\cos\theta) = \frac{(-1)^m}{s^m}h_t^m(\cos\theta)$. Using f_m in place of f has certain advantages. In particular, considering functions (of Δ) which vanish more quickly at 0 (such as the f_m) is very important in the characterization of Besov spaces in [19].

7 Appendix: A Technical Lemma

In the proof of Lemma 4.1, we used the following fact.

Lemma 7.1. Suppose $a_0, \ldots, a_L \in \mathbb{C}$. Then there exists an even function $u \in \mathcal{S}(\mathbb{R})$, such that $u^{(2l)}(0) = a_l$, whenever $0 \leq l \leq L$, and such that $supp \hat{u} \subseteq (-1, 1)$. (Of course, all odd order derivatives of u vanish at 0.)

Proof. We construct $h = \hat{u}$. We need to show that, given numbers b_0, \ldots, b_L , there exists $h \in C_c^{\infty}(\mathbb{R})$, h even, supp $h \subseteq (-1, 1)$, with $\int_{-\infty}^{\infty} \xi^{2l} h(\xi) d\xi = b_l$ for $0 \le l \le L$. Select an even function $\eta \in C_c^{\infty}(\mathbb{R})$, with $\sup \eta \subseteq (-1, 1)$, and $\int \eta = 1/2$. Select L+1 different numbers $r_1, \ldots, r_{L+1} \in (0, 1/2)$. For 0 < t < 1, consider the even function $\eta_{t,m}(\xi) = t^{-1}[\eta((x+r_m)/t) + \eta((x-r_m)/t)]$. We claim that if t is sufficiently small, we may construct our h by setting $h = \sum_{m=1}^{L+1} c_m \eta_{t,m}$ for suitable $c_1, \ldots c_{L+1}$. Indeed, as $t \to 0^+, \eta_{t,m}$ becomes increasingly concentrated near r_m and near $-r_m$, so that $\int \xi^{2l} [\sum_{m=1}^{L+1} c_m \eta_{t,m}] d\xi \to \sum_{m=1}^{L+1} c_m (r_m)^{2l}$. The existence of suitable $c_1, \ldots c_{L+1}$ now follows from the nonvanishing of small perturbations of the Vandermonde determinant.

References

- J.-P. Antoine and P. Vandergheynst, Wavelets on the n-sphere and related manifolds, J. Math. Phys. 39 (1998) 3987-4008.
- [2] P. Baldi, G. Kerkyacharian, D. Marinucci, D. Picard, Asymptotics for Spherical Needlets, Annals of Statistics, in press, arxiv:math/0606599.
- [3] I. Bogdanova, P. Vandergheynst, J.-P. Antoine, L. Jacques and M. Morvidone, Stereographic wavelet frames on the sphere, Appl. Comput. Harmon. Anal. 19 (2005), 223-252.
- [4] R. S. Cahn and J. A. Wolf, Zeta functions and their asymptotics expansions for compact symmetric spaces of rank one, Comment. Math. Helv. 51 (1976), 1-21.
- [5] R. Camporesi, Harmonic analysis and propagators on homogeneous spaces, Phys. Rep. 196 (1990), 1-134.
- [6] R. Coifman and M. Maggioni, Diffusion wavelets, Appl. Comput. Harmon. Anal. 21 (2006), 53-94.
- [7] R. R. Coifman and G. Weiss, Analyse harmonique non-commutative sur certains espaces homogènes, Lecture Notes in Math. #242, Springer-Verlag, Berlin, 1971.
- [8] S. Dahlke, Multiresolution analysis, Haar bases and wavelets on Riemannian manifolds, in Wavelets: Theory, Algorithms and Applications, Taormina (1993), 33-52.

- [9] W. Dahmen and R. Schneider, Wavelets on manifolds I: construction and domain decomposition, Siam J. Math. Anal. 31 (1999), 184-230.
- [10] I.Daubechies, Ten Lectures on Wavelets, Philadelphia, Pennsylvania, 1992.
- [11] A. G. Doroshkevich, P. D. Naselsky, O. V. Verkhodanov, D. I. Novikov, V. I. Turchaninov, I. D. Novikov, P. R. Christensen and L. -Y. Chiang., *Gauss-Legendre Sky Pixelization (GLESP) for CMB maps*, Int. J. Mod. Phys. D. 14, (2005), 275-290.
- [12] M. Frazier and B. Jawerth, Decomposition of Besov spaces, Indiana Univ. Math. J. 34 (1985), 777-799.
- [13] M. Frazier and B. Jawerth, A discrete transform and decompositions of distribution spaces, J. Func. Anal. 93 (1990), 34-170.
- [14] D. Geller, The Laplacian and the Kohn Laplacian for the sphere, J. Diff. Geom. 15 (1980), 417-455.
- [15] W. Freeden, T. Gervens and M. Schreiner, Constructive Approximation on the Sphere with Applications to Geomathematics, Clarendon Press, Oxford, 1998.
- [16] W. Freeden and M. Volker, Multiscale Potential Theory with Applications to Geoscience, Birkhauser, Boston, 2004.
- [17] D. Geller and A. Mayeli, Continuous wavelets and frames on stratified Lie groups I, Journal of Fourier Analysis and Applications, 12 (2006), 543-579.
- [18] D. Geller and A. Mayeli, Nearly Tight Frames and Space-Frequency Analysis on Compact Manifolds, Math. Z., (2008), to appear.
- [19] D. Geller and A. Mayeli, Besov Spaces and Frames on Compact Manifolds, (2008), available on arXiv.
- [20] P. Gilkey, The spectral geometry of a Riemannian manifold, J. Diff. Geom. 10 (1975), 601-618.
- [21] P. Gilkey, Invariance Theory, the Heat Equation and the Atiyah-Singer Index Theorem, Publish or Perish, Inc., Wilmington, Delaware, 1984.
- [22] F. Guilloux, G. Faÿ and J-F. Cardoso, Practical wavelet design on the sphere, arXiv:0706.2598, (June 18, 2007).
- [23] Y.S. Han, Calderón-type reproducing formula and the Tb theorem, Rev. Mat. Ibero. 10 (1994), 51-91.
- [24] Y. Han, Discrete Calderón-type reproducing formula, Acta Math. Sinica 16 (2000), 277-294.
- [25] M. Holschneider and Ph. Tchamitchian, Régularité locale de la fonction 'non-différentiable' de Riemann, in Les ondelettes en 1989, Lecture Notes in Math no. 1438, Springer-Verlag, Berlin, 102-124.
- [26] L.Hörmander, The spectral function of an elliptic operator, Acta Math. 121 (1968), 193-218.
- [27] Y. Kannai, Off diagonal short term asymptotics for fundamental solutions of diffusion equations, Comm. PDE 2 (1977), 781-830.
- [28] X. Lan and D. Marinucci, On the dependence structure of wavelet coefficiets for spherical random fields, preprint (2008).

- [29] A. Mayeli, Asymptotic Uncorrelation for Mexican Needlets, preprint (2008).
- [30] M. Maggioni, G. L. Davis, F. J. Warner, F. B. Geshwind, A. C. Coppi, R. A. DeVerse and R.R. Coifman, Spectral analysis of normal and malignant tissue sections using a novel microoptoelectricalmechanical system, Mod. Pathol (2004) 17 Suppl 1; 358A.
- [31] D. Marinucci, D. Pietrobon, A. Balbi, P. Baldi, P. Cabella, G. Kerkyacharian, P. Natoli, D. Picard and N. Vittorio, *Spherical needlets for CMB data analysis* arXiv:0707.0844 (July 5, 2007).
- [32] A.Mayeli, Discrete and continuous wavelet transformation on the Heisenberg group, Ph.D thesis, Technische Universität München, 2005.
- [33] S. Minakshisundaram and A. Plegel, Some properties of the eignenfunctions of the Laplace-operator on Riemannian manifolds, Can. J. Math 1 (1949), 242-256.
- [34] J. Milnor, Morse Theory, Ann. of Math. Studies, Vol. 51, Princeton University Press, Princeton, N.J. 1963.
- [35] F.J. Narcowich, P. Petrushev and J. Ward, Localized tight frames on spheres, SIAM J. Math. Anal. 38 (2006), 574-594.
- [36] F.J. Narcowich, P. Petrushev and J. Ward, Decomposition of Besov and Triebel-Lizorkin spaces on the sphere, J. Func. Anal. 238 (2006), 530-564.
- [37] I. Polterovich, Heat invariants of Riemannian manifolds, Israel J. Math 119 (2000), 239-252.
- [38] I. Polterovich, Combinatorics of the heat trace on spheres, Canad. J. Math 54 (2002), 1086-1099.
- [39] P.Schröder and W. Sweldens, Spherical wavelets: efficiently representing functions on a sphere, Computer Graphics Proceedings (SIGGRAPH 95) (1985), 161-172.
- [40] A. Seeger and C.D. Sogge, On the boundedness of functions of (pseudo-) differential operators on compact manifolds, Duke Math. J. 59 (1989), 709-736.
- [41] C. Sogge, Fourier Integrals in Classical Analysis, Cambridge University Press, 1993.
- [42] E.M. Stein and G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, Princeton University Press, 1971.
- [43] R. Strichartz, A functional calculus for elliptic pseudodifferential operators, Amer. J. Math 94 (1972), 711-722.
- [44] T. Tao, Weak-Type Endpoint Bounds for Riesz Means, Proc. AMS **124** (1996), 2797-2805.
- [45] M. Taylor, *Pseudodifferential Operators*, Princeton University Press, 1981.