

Superfluidity of a perfect quantum crystal

V.A. Golovko

Moscow State Evening Metallurgical Institute

Lefortovsky Val 26, Moscow 111250, Russia

In recent years, experimental data were published which point to the possibility of the existence of superfluidity in solid helium. To investigate this phenomenon theoretically we employ a hierarchy of equations for reduced density matrices which describes a quantum system that is in thermodynamic equilibrium below the Bose-Einstein condensation point, the hierarchy being obtained earlier by the author. It is shown that the hierarchy admits solutions relevant to a perfect crystal (immobile) in which there is a frictionless flow of atoms, which testifies to the possibility of superfluidity in solids. The solutions are studied with the help of the bifurcation method and some their peculiarities are found out. We discuss also various physical aspects of the problem, among them experimental ones.

1. Introduction

The question as to the possibility of the existence of superfluidity in a crystal solid has come under scrutiny long ago [1-3]. However, rather firm experimental evidence for this possibility in solid ^4He was obtained only in recent years (for a review see [4] where the experimental as well as theoretical aspects of the question are discussed).

In the present paper, the superfluid state of a crystalline solid is investigated with the help of the approach in equilibrium quantum statistical mechanics proposed in [5] (see also [6] where a systematic exposition of the approach is presented and various results achieved with its use are discussed). The approach is based upon a hierarchy of equations for reduced density matrices obtained in the same paper [5] which goes over, in the classical limit, into the well-known equilibrium Bogoliubov-Born-Green-Kirkwood-Yvon (BBGKY) hierarchy. A characteristic feature of the approach is construction of thermodynamics without use made of the Gibbs method. In Ref. [7] that serves as starting point for the present investigation, a modification of the hierarchy of [5] was obtained to take account of the Bose-Einstein condensation in fluids.

It should be emphasized that the superfluidity of a perfect crystal is implied in the present paper whereas it is common practice to ascribe superfluidity in a solid to various disruptions of the ideal crystalline order [4,8,9]. In the present paper, theoretical treatment of the problem will first be carried out. In the concluding section of the paper, various physical aspects of the problem, among them experimental ones, will be discussed as well.

2. Basic equations

We consider a system of N spinless bosons (^4He atoms, for example) enclosed in a volume V . The particles of mass m interact via a two-body potential $K(|\mathbf{r}_j - \mathbf{r}_k|)$. The approach developed in [5] leans upon use of s -particle reduced density matrices $R_s(\mathbf{x}_s, \mathbf{x}'_s)$ with $s = 1, 2, 3, \dots$ where \mathbf{x}_s denotes a set of coordinates $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_s$. For a state of thermodynamic equilibrium, it is possible to deduce a hierarchy of equations in which figure only diagonal elements of the density matrices

$$\rho_s(\mathbf{x}_s) = R_s(\mathbf{x}_s, \mathbf{x}_s). \quad (2.1)$$

The following normalization for the density matrices is implied

$$\int_V \rho_1(\mathbf{r}) d\mathbf{r} = N. \quad (2.2)$$

Let us present first those results of Ref. [7] that will be needed in the present paper and which are valid both for homogeneous media (liquids and gases) and for crystals (see also [6] and [10])

where the premises used are discussed in greater detail). Below the Bose-Einstein condensation point, the density matrices break up into two parts:

$$R_s(\mathbf{x}_s, \mathbf{x}'_s) = R_s^{(c)}(\mathbf{x}_s, \mathbf{x}'_s) + R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s), \quad (2.3)$$

where $R_s^{(c)}(\mathbf{x}_s, \mathbf{x}'_s)$ is relevant to the condensate while $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$ to the normal fraction. The diagonal elements $\rho_s(\mathbf{x}_s)$ of (2.1) are broken up analogously. We shall not write down the equations that characterise $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$ because they will not be required in what follows, referring the reader to [7] and [5].

As to the condensate parts of the reduced density they are given by

$$R_s^{(c)}(\mathbf{x}_s, \mathbf{x}'_s) = \varphi_s(\mathbf{x}_s) \varphi_s^*(\mathbf{x}'_s), \quad (2.4)$$

the functions $\varphi_s(\mathbf{x}_s)$ being found from the equations

$$\frac{\hbar^2}{2m} \sum_{j=1}^s \nabla_j^2 \varphi_s(\mathbf{x}_s) + [\varepsilon_{(s)} - U_s(\mathbf{x}_s)] \varphi_s(\mathbf{x}_s) = 0. \quad (2.5)$$

The occurring effective potentials $U_s(\mathbf{x}_s)$ are determined by the equation

$$\rho_s(\mathbf{x}_s) \nabla_1 U_s(\mathbf{x}_s) = \rho_s(\mathbf{x}_s) \nabla_1 \sum_{j=2}^s K(|\mathbf{r}_1 - \mathbf{r}_j|) + \int \rho_{s+1}(\mathbf{x}_{s+1}) \nabla_1 K(|\mathbf{r}_1 - \mathbf{r}_{s+1}|) d\mathbf{r}_{s+1}, \quad (2.6)$$

in which the full matrices $\rho_s(\mathbf{x}_s) = \rho^{(c)}(\mathbf{x}_s) + \rho^{(n)}(\mathbf{x}_s)$ figure. Eq. (2.6) links $\rho_s(\mathbf{x}_s)$ and $\rho_{s+1}(\mathbf{x}_s)$ as well, thus yielding a hierarchy of equations for $\rho_s(\mathbf{x}_s)$. The constants $\varepsilon_{(s)}$, the normalization of $\varphi_s(\mathbf{x}_s)$ and the boundary conditions for the differential equation of (2.5) are found from the interrelation that should be satisfied by the density matrices at $s = 2, 3, \dots$ in compliance with their definition, namely,

$$(N - s + 1) R_{s-1}(\mathbf{x}_{s-1}, \mathbf{x}'_{s-1}) = \int_V R_s(\mathbf{x}_{s-1}, \mathbf{r}_s, \mathbf{x}'_{s-1}, \mathbf{r}_s) d\mathbf{r}_s. \quad (2.7)$$

This interrelation is satisfied by $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$ [7], and therefore it should hold for $R_s^{(c)}(\mathbf{x}_s, \mathbf{x}'_s)$ as well. Substituting (2.4) into (2.7) yields

$$\varphi_{s-1}(\mathbf{x}_{s-1}) \varphi_{s-1}^*(\mathbf{x}'_{s-1}) = \frac{1}{N} \int_V \varphi_s(\mathbf{x}_{s-1}, \mathbf{r}_s) \varphi_s^*(\mathbf{x}'_{s-1}, \mathbf{r}_s) d\mathbf{r}_s, \quad (2.8)$$

where account has been taken of the fact that $s \ll N$ in the case under study.

We turn now to Eq. (2.5) with $s = 1$:

$$\frac{\hbar^2}{2m} \nabla^2 \varphi_1(\mathbf{r}) + [\varepsilon_{(1)} - U_1(\mathbf{r})] \varphi_1(\mathbf{r}) = 0. \quad (2.9)$$

Up to this point we have nowhere taken into explicit account the fact that we are interested in solutions relevant to a crystal. The uniform case where $U_1(\mathbf{r}) = \text{constant}$ was studied in [7]. In the case of the crystal, the potential $U_1(\mathbf{r})$ is to be periodic. Eq. (2.9) is of the form of a Schrödinger equation whereas the character of solutions of the Schrödinger equation for a periodic potential is well known [11]:

$$\varphi_1(\mathbf{r}) = \sqrt{\rho_c} e^{i\frac{\mathbf{p}_0 \cdot \mathbf{r}}{\hbar}} u_1(\mathbf{r}), \quad (2.10)$$

where $u_1(\mathbf{r})$ is a periodic function with the same periods as the potential $U_1(\mathbf{r})$. Usually, in the exponent one writes $i\mathbf{k}\mathbf{r}$; however it is more convenient for us to write \mathbf{p}_0/\hbar instead of \mathbf{k} . Eq. (2.10) contains a yet unknown normalizing factor $\sqrt{\rho_c}$ introduced with the condition that the function $u_1(\mathbf{r})$ obeys the normalization

$$\int_V |u_1(\mathbf{r})|^2 d\mathbf{r} = V. \quad (2.11)$$

It is worth remarking that $u_1 = 1$ in the uniform case [7], which conforms to (2.11). For use later, we also write down Eq. (2.5) at $s = 2$:

$$\frac{\hbar^2}{2m} \left[\nabla_1^2 \varphi_2(\mathbf{r}_1, \mathbf{r}_2) + \nabla_2^2 \varphi_2(\mathbf{r}_1, \mathbf{r}_2) \right] + \left[\varepsilon_{(2)} - U_2(\mathbf{r}_1, \mathbf{r}_2) \right] \varphi_2(\mathbf{r}_1, \mathbf{r}_2) = 0. \quad (2.12)$$

Let us try to satisfy the condition of (2.8) at the least possible $s = 2$:

$$\varphi_1(\mathbf{r}_1) \varphi_1^*(\mathbf{r}'_1) = \frac{1}{N} \int_V \varphi_2(\mathbf{r}_1, \mathbf{r}_2) \varphi_2^*(\mathbf{r}'_1, \mathbf{r}_2) d\mathbf{r}_2. \quad (2.13)$$

At given \mathbf{r}_1 and \mathbf{r}'_1 , the main contribution to the last integral in the thermodynamic limit as $V \rightarrow \infty$ results from the regions where $|\mathbf{r}_1 - \mathbf{r}_2|$ and $|\mathbf{r}'_1 - \mathbf{r}_2|$ are large. If $|\mathbf{r}_1 - \mathbf{r}_2| \rightarrow \infty$, according to the relation (4.7) of [5] and to the remark concerning (4.9) of [5] one has

$$U_2(\mathbf{r}_1, \mathbf{r}_2) \rightarrow U_1(\mathbf{r}_1) + U_1(\mathbf{r}_2) - \bar{U}_1, \quad (2.14)$$

where \bar{U}_1 is the constant part of $U_1(\mathbf{r})$ (the character of the dependence of $U_2(\mathbf{r}_1, \mathbf{r}_2)$ on \mathbf{r}_1 and \mathbf{r}_2 in this event is evident from physical considerations as well). If (2.14) is placed in (2.12), the variables in the resulting equation will be separated and the solution of the equation will be, in view of Eq. (2.9), of the form

$$\varphi_2^{(\infty)}(\mathbf{r}_1, \mathbf{r}_2) = B_2 \varphi_1(\mathbf{r}_1) \varphi_1(\mathbf{r}_2), \quad (2.15)$$

where B_2 is a constant and the superscript (∞) underlines that this is the limiting value of $\varphi_2(\mathbf{r}_1, \mathbf{r}_2)$. Simultaneously, we find that

$$\varepsilon_{(2)} = 2\varepsilon_{(1)} - \bar{U}_1. \quad (2.16)$$

According to the foregoing, it is Eq. (2.15) that can be inserted into (2.13) in case the limit $V \rightarrow \infty$ is implied. Then the integral is calculated at once on account of (2.10) and (2.11) to yield

$$B_2 = \sqrt{\frac{\rho_0}{\rho_c}} \quad (2.17)$$

(without loss of generality the quantity B_2 can be taken to be real). Hence we have found $\varepsilon_{(2)}$ by (2.16) and established that the solution of Eq. (2.16) should be subject to the limiting condition of (2.15) that, together with (2.17), determines the normalization of $\varphi_2(\mathbf{r}_1, \mathbf{r}_2)$.

In a like manner, one can treat arbitrary s . The main contribution to the integral in (2.8) in the thermodynamic limit gives the regions where $|\mathbf{r}_s| \rightarrow \infty$. In this limit, according to [5] we have, similarly to (2.14), that

$$U_s(\mathbf{x}_s) \rightarrow U_{s-1}(\mathbf{x}_{s-1}) + U_1(\mathbf{r}_s) - \bar{U}_1. \quad (2.18)$$

If this is inserted into (2.5) with account taken of (2.9) and of Eq. (2.5) written for $\varphi_{s-1}(\mathbf{x}_{s-1})$, by analogy with (2.15) and (2.16) we shall obtain

$$\varphi_s^{(\infty)}(\mathbf{x}_s) = B_s \varphi_{s-1}(\mathbf{x}_{s-1}) \varphi_1(\mathbf{r}_s), \quad \varepsilon_{(s)} = \varepsilon_{(s-1)} + \varepsilon_{(1)} - \bar{U}_1. \quad (2.19)$$

Putting this limiting function into (2.8) yields, analogously to (2.17),

$$B_s = \sqrt{\frac{\rho_0}{\rho_c}}. \quad (2.20)$$

Thus, we have completely constructed the hierarchy of equations for the reduced density matrices which describes a crystal below the Bose-Einstein condensation point that can consequently be observed in the crystalline state as well. The normal part of the reduced density matrices is described by the equations for $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$ of [7] and [5] whereas their condensate part is determined by Eqs. (2.4)-(2.5) whose solutions should be subject to the first condition of (2.19) with B_s of (2.20). The constants $\varepsilon_{(s)}$ are to be found from the second equation of (2.9) that enables one to express them via $\varepsilon_{(1)}$. The condensate and normal parts are connected by the effective potentials $U_s(\mathbf{x}_s)$ determined by Eq. (2.6). The hierarchy obtained contains the arbitrary constants ρ_c and \mathbf{p}_0 as well as two constants (A and τ) in the formulae for $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$ [7,5].

Let us make three remarks as to the hierarchy obtained. The presence of ρ_c in the denominator of (2.17) and (2.20) should not cause difficulties in case the limit as $\rho_c \rightarrow 0$ is considered. The fact is that Eqs. (2.15) and (2.19) contain $\varphi_1(\mathbf{r}_s)$ as a factor while the function

$\varphi_1(\mathbf{r}_s)$ itself has a factor $\sqrt{\rho_c}$ in view of (2.10). As a result, the factor $\sqrt{\rho_c}$ disappears in fact from Eq. (2.19) that relates $\varphi_s(\mathbf{x}_s)$ and $\varphi_{s-1}(\mathbf{x}_{s-1})$, so that all $\varphi_s(\mathbf{x}_s)$'s will have $\sqrt{\rho_c}$ as a factor just as $\varphi_1(\mathbf{r}_s)$.

The second remark consists in the following. Equations of the type (2.9) with a periodic potential $U_1(\mathbf{r})$ are used in studies of movement of particles, e. g. of an electron, in a periodic field [11], as was mentioned above. In this case there exist an infinite set of values of $\varepsilon_{(1)}$ that form energy bands if the vector \mathbf{p}_0 (in our notation) changes. In our case, one should take the band that corresponds to the minimum of an appropriate thermodynamic potential while all subsequent $\varepsilon_{(s)}$'s will be obtained uniquely by (2.19). The situation here is analogous with that which occurs in the theory of an ordinary crystal. At a given interaction potential between particles, a great diversity of crystalline lattices can exist. Realized is the lattice that corresponds to the minimum of the relevant thermodynamic potential (see, for example, [12]).

The third remarks concerns the equations for $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$ that were not written down in this paper. Inasmuch as the condensate and the normal fraction are linked by the common potentials $U_s(\mathbf{x}_s)$, those equations should also have solutions corresponding to a periodic density once $U_1(\mathbf{r})$ is periodic. Such solutions do exist and are considered in [13]. They have properties characteristic of an ordinary crystal.

It is of interest to find the singlet density matrix. To this end, we substitute (2.4) with $s = 1$ into (2.3) and take account of (2.10), which gives

$$R_1(\mathbf{r}, \mathbf{r}') = \rho_c e^{i\frac{\mathbf{p}_0(\mathbf{r}-\mathbf{r}')}{\hbar}} u_1(\mathbf{r}) u_1^*(\mathbf{r}') + R_1^{(n)}(\mathbf{r}, \mathbf{r}'), \quad (2.21)$$

where $R_1^{(n)}(\mathbf{r}, \mathbf{r}')$ describes the normal fraction. The first term in (2.21) exhibits off-diagonal long range order (ODLRO) since it does not vanish in the limit as $|\mathbf{r} - \mathbf{r}'| \rightarrow \infty$. It is worthy of remark that ODLRO is characteristic of the phenomenon of superfluidity [14]. The second term in (2.21) does not display ODLRO as in an ordinary crystal, which can be proven rigorously. In the uniform case, the fact that this term tends to zero as $|\mathbf{r} - \mathbf{r}'| \rightarrow \infty$ is seen from Eq. (2.32) of [10].

In order to elucidate the physical sense of the results obtained let us calculate the momentum of the system that is given by Eq. (2.13) of [7]:

$$\mathbf{P} = -i\hbar \int [\nabla R_1(\mathbf{r}, \mathbf{r}')]_{\mathbf{r}'=\mathbf{r}} d\mathbf{r}. \quad (2.22)$$

Upon substituting the first term of (2.21) herein (the second term should not contribute to the macroscopic momentum as in an ordinary crystal), we get

$$\mathbf{P} = \rho_c \mathbf{p}_0 V - i\hbar \rho_c \int_V u_1^*(\mathbf{r}) \nabla u_1(\mathbf{r}) d\mathbf{r}, \quad (2.23)$$

where (2.11) has been taken into account. Hence the immobile crystal (its density $\rho_1(\mathbf{r})$ is time-independent) has a momentum that can be relevant only to a movement of the condensate. As long as the number of particles in the condensate is $N_c = \rho_c V$, upon dividing \mathbf{P} by mN_c we obtain the mean velocity of the directional movement of the condensate particles although this does not signify that all condensate particles move simultaneously. It should be emphasized that the flow of particles in the crystal is not accompanied by any dissipation for the system is in thermodynamic equilibrium. This result is completely analogous to the one obtained for uniform media in Ref. [7] (in the case of the uniform media the last term in (2.23) is lacking). We shall return to the physical aspect of the result in the concluding section of the present paper.

It should be added also that it may turn out that $\mathbf{p}_0 = 0$. Then we shall have a condensate phase ($\rho_c \neq 0$) without superfluidity (if $\mathbf{p}_0 = 0$, the second term in (2.23) vanishes as well, see Sec. 4). If $\mathbf{p}_0 \neq 0$, the condensate phase will be superfluid. Thus, formation of a condensate phase does not necessarily leads to superfluidity just as in the case of uniform media [7].

The next step in the employed approach of [5] consists in construction of thermodynamics, which enables one simultaneously to obtain equations for determination of the quantities ρ_c and \mathbf{p}_0 as well as of the two constants that characterize $R_s^{(n)}(\mathbf{x}_s, \mathbf{x}'_s)$. In the uniform case, the thermodynamics is built up in [7]. As to a crystal, the construction of thermodynamics is essentially complicated by an involved form of the pair correlation function in the crystal [15] while the function enters into expressions for thermodynamic quantities. Up to the present, even in the case of an ordinary crystal it has been possible to construct the thermodynamics in the framework of the approach only with use made of simplifying assumptions as to the form of the pair correlation function, and the quantum case [13] is noticeably more complicated than the classical [16]. The presence of the condensate adds complexity to the construction of thermodynamics. Inasmuch as in the present paper we are interested first of all in the possibility in principle concerning the existence of superfluidity in a crystal, in what follows we shall restrict ourselves to the case of zero temperature ($T = 0$) where the thermodynamics is not required and the equilibrium state of the system is determined by the condition that its energy is a minimum.

3. Periodic solutions

Having in mind the case of zero temperature (see the end of the preceding section) and trying to simplify, wherever possible, the problem under study, we shall presume that all particles

pertain to the condensate, that is to say, $\rho_c = \rho_0 \equiv N/V$. It should be remarked that this occurs in the case of the Bose liquid considered in [7], although in an analogous case of the Fermi liquid this does not happen [10]. If $\rho_c = \rho_0$, the full reduced density matrices coincide with $R_s^{(c)}(\mathbf{x}_s, \mathbf{x}'_s)$ while the spatial number density of particles is (henceforth we shall omit the subscript 1 of ρ_1 , u_1 and U_1)

$$\rho(\mathbf{r}) = \rho_0 |u(\mathbf{r})|^2, \quad (3.1)$$

where (2.10) has been taken into account (note that (3.1) agrees with the normalization condition of (2.2) owing to (2.11)). If (2.10) is placed in (2.9), there results the equation for $u(\mathbf{r})$:

$$\frac{\hbar^2}{2m} \nabla^2 u(\mathbf{r}) + \frac{i\hbar \mathbf{p}_0}{m} \nabla u(\mathbf{r}) + \left[\varepsilon_{(1)} - \frac{\mathbf{p}_0^2}{2m} - U(\mathbf{r}) \right] u(\mathbf{r}) = 0. \quad (3.2)$$

The effective potential $U(\mathbf{r})$ is determined by Eq. (2.6) at $s = 1$ which is of the form

$$\rho(\mathbf{r}) \nabla U(\mathbf{r}) = \int \rho_2(\mathbf{r}, \mathbf{r}') \nabla K(|\mathbf{r} - \mathbf{r}'|) d\mathbf{r}'. \quad (3.3)$$

We introduce also the pair correlation function $g(\mathbf{r}, \mathbf{r}')$ in line with the customary relation

$$\rho_2(\mathbf{r}, \mathbf{r}') = \rho(\mathbf{r}) \rho(\mathbf{r}') g(\mathbf{r}, \mathbf{r}'). \quad (3.4)$$

The function $\rho_2(\mathbf{r}, \mathbf{r}')$ and thereby $g(\mathbf{r}, \mathbf{r}')$ should be found from the subsequent hierarchy equations at $s = 2$ that contain $\rho_3(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3)$ as well. Putting aside the discussion of the problem as to how to close the hierarchy, the problem well known from the theory of the classical BBGKY hierarchy, for the moment we shall assume the function $g(\mathbf{r}, \mathbf{r}')$ to be given. Almost in all studies on statistical theory of crystals, one introduces a simplifying assumption that $g(\mathbf{r}, \mathbf{r}')$ depends only on $|\mathbf{r} - \mathbf{r}'|$ as in fluids. This assumption should not essentially affect results because in the case of a crystal the leading role for $\rho_2(\mathbf{r}, \mathbf{r}')$ of (3.4) is played by the periodic density $\rho(\mathbf{r})$ while $g(\mathbf{r}, \mathbf{r}')$ plays a secondary role [12]. If $g(\mathbf{r}, \mathbf{r}') = g(|\mathbf{r} - \mathbf{r}'|)$, Eq.(3.3) is readily integrated to give

$$U(\mathbf{r}) = \int K_g(|\mathbf{r} - \mathbf{r}'|) \rho(\mathbf{r}') d\mathbf{r}', \quad (3.5)$$

where

$$K_g(r) = \int_{-\infty}^r \frac{dK(r')}{dr'} g(r') dr'. \quad (3.6)$$

Here $r = |\mathbf{r}|$ and the limits of integration in (3.6) are chosen such that $K_g(\infty) = 0$ (only in this case the integral in (3.5) converges). It will be remarked that an arbitrary constant can be added to the right side of (3.5), which defines another choice for the origin of the energy scale. In case $g(r)$ is

taken to be given, one has two equations (3.2) and (3.5) that allow one to find the density in the crystal by (3.1).

We look for periodic solutions of the equations in terms of a Fourier series:

$$u(\mathbf{r}) = \sum_{l,m,n=-\infty}^{\infty} c_{lmn} e^{i\mathbf{A}\mathbf{r}}, \quad (3.7)$$

where $\mathbf{A} = l\mathbf{a}_1 + m\mathbf{a}_2 + n\mathbf{a}_3$ with the basic reciprocal-lattice vectors \mathbf{a}_1 , \mathbf{a}_2 and \mathbf{a}_3 [16]. Eq. (2.11) entails the following condition on the coefficients c_{lmn} :

$$\sum_{l,m,n} |c_{lmn}|^2 = 1. \quad (3.8)$$

The crystal density (3.1) is given by the series

$$\rho(\mathbf{r}) = \sum_{l,m,n} a_{lmn} e^{i\mathbf{A}\mathbf{r}}, \quad a_{lmn} = \rho_0 \sum_{l',m',n'} c_{l'm'n'} c_{l'-l,m'-m,n'-n}^*. \quad (3.9)$$

The noteworthy fact is that $a_{000} = \rho_0$ in view of (3.8). Substituting (3.9) into (3.5) yields

$$U(\mathbf{r}) = \sum_{l,m,n} a_{lmn} \sigma(\mathbf{A}) e^{i\mathbf{A}\mathbf{r}}, \quad (3.10)$$

where

$$\sigma(k) = \int K_g(|\mathbf{r}|) e^{i\mathbf{k}\mathbf{r}} d\mathbf{r} = \frac{4\pi}{k} \int_0^{\infty} r K_g(r) \sin kr dr = \frac{4\pi}{k^3} \int_0^{\infty} \frac{dK}{dr} g(r) (kr \cos kr - \sin kr) dr. \quad (3.11)$$

The last expression was obtained upon inserting $K_g(r)$ of (3.6) and carrying out one of the integrations.

If all these series are put into (3.2), we shall arrive at the set of equations that contain the coefficients c_{lmn} alone:

$$\left[\frac{(\hbar\mathbf{A} + \mathbf{p}_0)^2}{2m} + \rho_0 \sigma_0 - \varepsilon_{(1)} \right] c_{lmn} + \rho_0 \sum_{l',m',n'}' \sigma(\mathbf{A}') c_{l'-l,m'-m,n'-n} \sum_{l''m''n''} c_{l''m''n''} c_{l''-l',m''-m',n''-n'}^* = 0. \quad (3.12)$$

Here the primed summation denotes the omission of the term with $l' = m' = n' = 0$ because this term is separated out and gives the summand $\rho_0 \sigma_0$ in the square brackets with use made of (3.8) and with the notation $\sigma_0 = \sigma(0)$.

Let us find the expression for the momentum of the crystal in the present case. Substituting (3.7) into (2.23) and retaining only terms that increase with the volume V yields

$$\mathbf{P} = N \left(\mathbf{p}_0 + \hbar \sum_{l,m,n} \mathbf{A} |c_{lmn}|^2 \right). \quad (3.13)$$

It follows herefrom that in general the momentum \mathbf{P} of the crystal is not aligned with the vector \mathbf{p}_0 .

We turn now to the energy of the crystal. In the approach used, the energy of the system is specified by the formula [5]

$$E = -\frac{\hbar^2}{2m} \int \left[\nabla^2 R_1(\mathbf{r}, \mathbf{r}') \right]_{\mathbf{r}'=\mathbf{r}} d\mathbf{r} + \frac{1}{2} \int K(|\mathbf{r}_1 - \mathbf{r}_2|) \rho_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2. \quad (3.14)$$

We substitute the first term of (2.21) here, transform the resulting expression with the help of (3.2), and place the above Fourier series in the expression obtained. As a result, we have (cf. the derivation of Eq. (4.3) in [16])

$$E = \varepsilon_{(1)} N - V \sum_{l,m,n} |a_{lmn}|^2 \sigma_\varepsilon(A), \quad \sigma_\varepsilon(k) = \int \left[K_g(|\mathbf{r}|) - \frac{1}{2} K(|\mathbf{r}|) g(|\mathbf{r}|) \right] e^{i\mathbf{k}\mathbf{r}} d\mathbf{r}. \quad (3.15)$$

In an analogous way, one can calculate the stress tensor in the present case starting from the general formula (3.5) of [7].

It is desirable to estimate the pair correlation function $g(r)$ that figures in the above formulae. Beginning with the pioneering work on the statistical theory of a (classical) crystal [17] and in theories of crystals based upon the density functional (see the reviews [18,19]), for $g(r)$ is usually taken the pair correlation function of the corresponding liquid. We shall proceed along these lines. If $\rho_c = \rho_0$, according to [7] the condensate part of the pair correlation function of the liquid is $g(\mathbf{r}) = |u_2(\mathbf{r})|^2$ where the function $u_2(\mathbf{r})$ is to be found from Eq. (2.11) of [7]. This last equation contains the potential $U_2(\mathbf{r})$ that depends upon the triplet correlation function by virtue of Eq. (5.33) of [5], and therefore we have to resort to an approximation. We shall start from the quantum extension of the hypernetted-chain approximation proposed in [5]. The relevant equation (5.36) of [5] contains the quantity τ whose role goes over to $\tilde{\tau} = (1 - \rho_c/\rho_0)\tau$ below the Bose-Einstein condensation point [7,10]. Because of this, in our case where $\rho_c = \rho_0$ and thereby $\tilde{\tau} = 0$, one should set $\tau = 0$ in Eq. (5.36) of [5], which gives immediately that $U_2(r) = K(r)$ (it may be noted in passing that one obtains the same result from the classical hypernetted-chain equation (5.35) of [5] at $T = 0$). It should be added that the relation $U_2(r) = K(r)$ results also when one neglects triplet correlations [5]. After the replacement $U_2(r) = K(r)$ in the above-mentioned equation (2.11) of [7], in the spherically symmetric case we arrive at

$$\frac{d^2 u_2}{dr^2} + \frac{2}{r} \frac{du_2}{dr} - \frac{m}{\hbar^2} K(r) u_2(r) = 0. \quad (3.16)$$

This equation must be solved with the condition that $u_2(r) \rightarrow 1$ as $r \rightarrow \infty$.

For the atomic interaction potential we take the Lennard-Jones potential

$$K(r) = \varepsilon \left[\left(\frac{r_m}{r} \right)^{12} - 2 \left(\frac{r_m}{r} \right)^6 \right], \quad (3.17)$$

in which r_m corresponds to the minimum of $K(r)$ and $K(r_m) = -\varepsilon$, and which is equal to zero at $r = r_0 = r_m / 2^{1/6}$. The potential has a strong singularity as $r \rightarrow 0$, which considerably complicates numerical solution of Eq. (3.16). For this reason we shall proceed as follows. We take some $r_1 < r_0$ and set $K(r) = \infty$ if $r < r_1$, that is to say, we shall use a hard-sphere potential for $r < r_1$. When $r > r_1$, we shall solve Eq. (3.16) with the potential (3.17) and with the condition $u_2(r_1) = 0$. In numerical calculation, we took $r_1 = 0.85r_0$ where the potential (3.17) is sufficiently large: $K(r_1) \approx 18\varepsilon$. At lesser values of r_1 , the above-mentioned singularity of the potential (3.17) begins to manifest itself markedly.

If (3.17) is put into Eq. (3.16) and the equation is reformulated in terms of dimensionless quantities, in front of the dimensionless potential there appears the factor

$$\zeta = \frac{m\varepsilon r_m^2}{\hbar^2}. \quad (3.18)$$

For helium, $\varepsilon = 10.2 \text{ K}$, $r_m = 2.86 \text{ \AA}$ [20] and thereby $\zeta = 6.9$. Eq. (3.16) with this ζ was solved numerically with the help of the well-known Runge-Kutta method. We took $u_2(r_1) = 0$ and by the trial-and-error method we looked for a value of the derivative $u'_2(r_1)$ such that $u_2(r) \rightarrow 1$ as $r \rightarrow \infty$. The results of the calculation are presented in figure 1 where $\tilde{r} = r/r_m$.

Now we can compute the function $\sigma(k)$ of (3.11) that is the sole prescribed function in the set of equations of (3.12). The form of the function is shown in figure 2 where $\tilde{\sigma} = \sigma / (\varepsilon r_m^3)$ and $\tilde{k} = kr_m$. The noteworthy fact is that this form of $\sigma(k)$ is fully analogous with the one depicted in figure 1 of Ref. [15] in the classical case for zero temperature (that figure contains an error: in actual fact it is the quantity $\sigma / (\varepsilon r_m^3) = \tilde{\sigma}$ that is laid off on the vertical axis). The values characteristic of a crystal are $\tilde{k} \sim 2\pi$ [15]. They lie between points *A* and *B* in figure 2 where $\sigma(k) < 0$, which is necessary for the crystal to exist as will be shown in the next section.

4. Bifurcation method

One of the method employed for solving nonlinear equations describing a crystal is the bifurcation method. Although the bifurcation point is of little physical significance, the method enables one to find out the conditions for periodic solutions of the equations to exist and to investigate some their peculiarities [16]. The method consists in searching for the periodic

solutions characterizing the crystal that bifurcate off the uniform one relevant to the corresponding fluid.

In our case, the set of (3.12) has a solution $c_{000} = 1$ with the remaining $c_{lmn} = 0$, which corresponds to a liquid (this can, more simply, be seen from Eqs. (3.2) and (3.5) that admit a solution $u(\mathbf{r}) = 1$, $\rho(\mathbf{r}) = \text{const}$, $U(\mathbf{r}) = \text{const}$). We now look for a solution where some c_{lmn} connected by the symmetry are small upon assuming the remaining c_{lmn} to be of higher order of magnitude except for $c_{000} \approx 1$ by (3.8). If we take Eq. (3.12) at $l = m = n = 0$ ($\mathbf{A} = 0$) and discard all small terms (such will be all terms under the summation sign), in a zeroth approximation we shall get

$$\varepsilon_{(1)} = \frac{\mathbf{p}_0^2}{2m} + \rho_0 \sigma_0. \quad (4.1)$$

To obviate any confusion we note that this $\varepsilon_{(1)}$ differs by the last term from $\varepsilon_{(1)}$ written for the uniform case in Eq. (2.8) of [7], which is due to a different choice for the origin of the energy scale (see the remark concerning (3.5)).

In the remaining equations of (3.12), we now retain only terms of the first order of magnitude (recall that $c_{000} \approx 1$, Eq. (4.1) is also used) with the result that

$$\left(\frac{\hbar^2}{2m} A^2 + \frac{\hbar}{m} \mathbf{p}_0 \mathbf{A} \right) c_{lmn} + \rho_0 \sigma(A) (c_{-l,-m,-n}^* + c_{lmn}) = 0. \quad (4.2)$$

Upon changing the sign of l, m, n we take the complex conjugate:

$$\left(\frac{\hbar^2}{2m} A^2 - \frac{\hbar}{m} \mathbf{p}_0 \mathbf{A} \right) c_{-l,-m,-n}^* + \rho_0 \sigma(A) (c_{lmn} + c_{-l,-m,-n}^*) = 0. \quad (4.3)$$

The set of these two homogeneous equations for c_{lmn} and $c_{-l,-m,-n}^*$ has a nontrivial solution only if the determinant of the set is equal to zero. Before calculating the determinant it is worthwhile to establish what c_{lmn} 's are connected by the symmetry. For the sake of simplicity, we shall imply cubic lattices (SC, FCC or BCC of the simplest symmetry) although solid ^4He has a hexagonal close-packed (HCP) lattice at low pressures. However the difference in energy between the FCC and HCP structures are small; besides, ^4He can have a BCC or FCC lattice at some pressures and temperatures [20]. In addition, the Lennard-Jones potential used of (3.17) describes the interaction between helium atoms rather roughly in order to lead to the HCP lattice unequivocally. The aforementioned cubic lattices have a different number of basic coefficients c_{lmn} that correspond to the vectors \mathbf{a}_i obtainable from the basic reciprocal-lattice vectors \mathbf{a}_1 , \mathbf{a}_2 and \mathbf{a}_3 by the relevant symmetry transformations, all these \mathbf{a}_i 's having the same magnitude $|\mathbf{a}_i| =$

a (see, e.g., [16,13]). We substitute this a into the set of Eqs. (4.2)-(4.3) instead of A and equate the determinant of the set to zero, so that

$$\frac{\hbar^2}{2m}a^2 - \frac{2p_0^2 \cos^2 \xi_i}{m} + 2\rho_0 \sigma(a) = 0, \quad (4.4)$$

where ξ_i is the angle between the vectors \mathbf{p}_0 and \mathbf{a}_i . This equation is the condition of appearance of nonzero c_{lmn} 's, that is to say, the condition under which a periodic solution branches off from the uniform one (the bifurcation condition).

Inasmuch as the lattice period is inversely proportional to a , the average density of the crystal is proportional to a^3 , which amounts to saying that $\rho_0 = a^3/\eta$ where $\eta = \eta'\pi^3$ with $\eta' = 8, 6\sqrt{3}, 8\sqrt{2}$ for the SC, FCC, BCC lattices respectively. We now rewrite Eq. (4.4) as

$$\sigma(a) = -\frac{\hbar^2 \eta}{2ma} + \frac{p_0^2 \eta \cos^2 \xi_i}{ma^3}. \quad (4.5)$$

This equation is conveniently solved graphically. We consider first the case $p_0 = 0$. In this case, from (4.5) it follows immediately that for the crystal to exist it is necessary that $\sigma(a) < 0$. We are interested in solutions between points A and B in figure 2 (see the end of the preceding section). The right-hand side of (4.5) at $p_0 = 0$ is plotted by the broken curve in this figure when the solutions required occur. It should not be surprising that two solutions rather than one necessarily exist in this instance. To clarify the situation, let us resort to the classical case where the bifurcation condition has a form similar to (4.4) at $p_0 = 0$ if the quantity $\hbar^2 a^2/m$ is replaced by a quantity proportional to the temperature (see Eq. (6.3) of [16]), and thereby has two solutions as well. Account must be taken of the fact that the different values of a correspond to different pressures; besides, only stable solutions are of importance (the solutions that provide a minimum for the relevant thermodynamic potential). The solutions of equations for the classical crystal obtained in [12] not only in the vicinity of the bifurcation point but also in the entire region where the solutions exist show that at a given pressure and given temperature there is a unique stable solution of a specified symmetry.

It remains now to be seen whether there are relevant solutions in the helium case as long as the required solutions will be lacking in case the broken curve of figure 2 passes below the minimum of $\sigma(k)$ between points A and B . To this end, we rewrite (4.5) at $p_0 = 0$ in the dimensionless form:

$$\tilde{\sigma}(a) = -\frac{\eta}{4\zeta a r_m}, \quad (4.6)$$

where the dimensionless parameter ζ of (3.18) is used. According to the foregoing, $\eta \sim 10\pi^3$ for the cubic lattices, $ar_m \sim 2\pi$ between points A and B in figure 2, and $\zeta = 6.9$ for helium. Therefore the right side of Eq. (4.6) between these points is of the order of -2 whereas the minimum value of $\sigma(k)$ is equal to -4.5 there. Consequently, the situation in helium corresponds to that presented in figure 2, and the required periodic solutions do exist. At the same time, it is to be remarked that the broken curve of figure 2 for helium passes not too far from the minimum of $\sigma(k)$. If a more realistic interatomic potential is used instead of (3.17) and if the function $u_2(r)$ that should depend on the pressure is calculated with used made of more sophisticated approximations, it may turn out that at low pressures the broken curve of figure 2 will pass below the curve for $\tilde{\sigma}(\tilde{k})$. This will explain the fact that at low pressures helium remains liquid down to zero temperature. However, to explain the fact convincingly it needs to investigate Eq. (3.12) in case c_{lmn} 's are not small. As to other rare gases, the parameter ζ of (3.18) for them is substantially greater because their values of m , ε and r_m far exceed those of helium. As a result, according to (4.6) the broken curve of figure 2 for them will pass much closer to the abscissa and far from the minimum of $\sigma(k)$, so that the influence of the pressure should not be so crucial. For this reason, the rare gases solidify prior to attaining zero temperature at any pressure.

We turn now to the case $p_0 \neq 0$. If p_0 is small in (4.5), the broken curve of figure 2 will rise but slightly, and the periodic solutions will exist as before. However, the magnitude of a will now depend upon the angle ξ_i , and the lattice will cease to be cubic. Such deformation of the lattice at a nonzero p_0 is quite natural. When p_0 is sufficiently large, so that the broken curve will rise drastically, the required solutions with $\cos\xi_i \neq 0$ may disappear at all. However, for some directions of the vector \mathbf{p}_0 with respect to the vectors \mathbf{a}_i , when $\cos\xi_i = 0$, the periodic solutions will remain, although the periodicity will not be three-dimensional now. Insofar as the last term in (4.5) at $p_0 \cos\xi_i \neq 0$ tends to $+\infty$ as $a \rightarrow 0$ ($\propto 1/a^3$), there always exist solutions with $p_0 \neq 0$ and small a , that is, long-period structures, even if the solutions in the vicinity of the interval AB of figure 2 are lacking. On the other hand, if the solutions in the vicinity of the interval AB are nonexistent at $p_0 = 0$, they may appear at $p_0 \neq 0$ when the broken curve will rise sufficiently in conformity with (4.5). Thus, there is a theoretical possibility of the existence of periodic structures only with the simultaneous presence of a superflow.

In the expression for the momentum of the crystal of (3.13), in the present approximation it needs to sum over the above-specified vectors \mathbf{a}_i alone. We express c_{lmn}^* in terms of $c_{-l,-m,-n}$ with the help of (4.2) upon changing the sign of l , m , n and substitute into the sum of (3.13). With regard to the fact that the vectors \mathbf{a}_i and $-\mathbf{a}_i$, to which the coefficients c_{lmn} and $c_{-l,-m,-n}$

correspond, are of the same magnitude because (4.4) contains $\cos^2 \xi_i$, some terms in the sum obtained will cancel out. As a result, we shall have

$$\mathbf{P} = N\mathbf{p}_0 + \frac{2\hbar^2 V}{m} \sum_i' \frac{\mathbf{a}_i}{\sigma(a_i)} \mathbf{p}_0 \mathbf{a}_i c_{lmn} c_{-l,-m,-n}, \quad (4.7)$$

where the primed summation signifies that only one of the two vectors \mathbf{a}_i or $-\mathbf{a}_i$ should be taken into account, no matter which. Eq. (4.7) shows explicitly that the vector \mathbf{P} need not be directed along \mathbf{p}_0 although its magnitude is proportional to p_0 .

Subsequent calculation in the bifurcation method should be carried out separately for each type of the lattice. The calculations can be performed by analogy with an ordinary crystal, the classical [16] or quantum [13]. We shall not carry out the calculations as long as their results are not required for the aims of the present paper.

Of importance is the question as to whether the solutions with $p_0 \neq 0$ can correspond to a minimum of energy, that is to say, whether they can represent the ground state of the system. If (4.1) is substituted into (3.15), we shall see that the energy increases with increasing p_0 at small $|a_{lmn}|$. This, however, does not necessarily mean that the same will occur at $|a_{lmn}|$ relevant to a real crystal. By way of example we can point out the classical crystal where the crystalline state becomes energetically advantageous only at sufficiently large $|a_{lmn}|$ [12]. It is quite possible that advantageous will be superflows only with particular orientations of the vector \mathbf{p}_0 with respect to the crystal axes. To answer these questions, other approaches to solving the set of equations of (3.12) should be searched for besides the bifurcation method.

5. Concluding remarks

The present paper shows that a perfect quantum crystal can possess superfluidity. Let us discuss how this phenomenon can be understood from the physical point of view. The superfluidity of a crystal may be conceived as a peculiar kind of collective tunnelling of the crystal particles when the points corresponding to the density maximums remain immobile to form a regular crystalline lattice though deformed by the flow. At low temperatures, macroscopic bodies tend to have a crystalline order; on the other hand, helium tends to become superfluid. The results of the present paper demonstrate that these two tendencies do not contradict each other.

The approach used in the present paper enables one to treat both the superfluidity of a liquid and the one of a crystal in perfect analogy. It all depends on whether we look for uniform or periodic solutions of Eq. (2.9). From the viewpoint of the approach, the superflow can exist

irrespective of whether the atoms of the substance are located at random (the liquid) or in a perfect order (the crystal). Because of this, one can, for the crystal, utilize the analogies and arguments from the concluding section of Ref. [7] that will not be repeated here.

Let us take a brief look at the following question alone. We considered an unbounded crystal in the present paper. In finite crystalline specimens, the superflows must close upon themselves somehow as in liquids [7]. It is quite possible that the specimen will break up into cells with closed superflows. Perhaps, this explains the negative result obtained when trying to detect mass transport in solid helium [21]. At the same time, it should be emphasized once again that the superfluid solid does not flow like a fluid. It remains immobile and solid while the superflow exists in the interior of the solid (analogously with the flow of electrons that move in an immobile conductor that carries a current). When trying to detect the mass transport in a superfluid crystal experimentally, the challenge is to bring out the superflow through the surface of the specimen. This may be done, for example, if the supersolid is in contact with a superfluid on two sides. A superflow from the superfluid can penetrate the supersolid in which some of the closed superflows can open to escape through the second side. In this connection let us cite Ref. [22] in which experimental observation of mass transport in solid ^4He that is in contact with superfluid ^4He was reported (see, however, [23,24]).

It is not excluded also that the state with the least possible energy, that is, the ground state will be the one with no superflow ($p_0 = 0$) whereas the state with a superflow ($p_0 \neq 0$) will be excited (the same may occur in a liquid as well [7]). Under certain external conditions, this excited state may be metastable and may exist for an appreciable length of time. The state may even become ground. In rotatory movement, for example, the kinetic energy of a body is $E_k = I\omega^2/2$ where I is the moment of inertia of the body and ω is its angular velocity. If a superflow develops, the moment of inertia of the “normal” part of the body rotating with the angular velocity ω decreases, and the velocity of the “superfluid” part decreases as well if the superflow is opposite in direction to the rotation. As a result, the total energy of the rotating body may become less than at $p_0 = 0$. The situation here is analogous to that with a ferromagnet in which formation of magnetic domains is energetically unfavourable from the structural point of view, nevertheless the domains do form because this leads to a decrease in the magnetic energy of the specimen. It is worth remarking that superfluidity in solid ^4He was observed for the first time with confidence in torsional oscillator experiments [4] for which just the above considerations are valid. It should be added that the abovementioned observation of mass transport in solid ^4He that is in contact with superfluid ^4He [22] may be explained also in the case where the ground state of the crystal is not superfluid. The superflow injected from the superfluid into the crystal can cause the crystal to pass into an excited state that is superfluid.

Of interest is to discuss the possibility of superfluidity in solid ^3He whose atoms are fermions as distinct from the bosons considered in this paper. The superfluidity of a fermionic liquid in the framework of the approach proposed in [5] was considered in [10]. Although the treatment of the problem proves to be more involved than in the case of spinless bosons, by and large the ideas of Ref. [7] that were exploited in the present paper remain in force. In particular, Eq. (2.21) of [10] will coincide completely with Eq. (2.9) of the present paper if account is taken of the potential $U_1(\mathbf{r})$ that can be set equal to zero in the case of the liquid. Other equations will also coincide if $U_1(\mathbf{r})$ is allowed for. Therefore, there exists the possibility in principle as to the existence of superfluidity in solid ^3He as well.

References

- [1] Andreev A F and Lifshitz I M 1969 *Zh. Eksp. Teor. Fiz.* **56** 2057 [*Sov. Phys. JETP* **29** 1107]
- [2] Chester G V 1970 *Phys. Rev. A* **2** 256
- [3] Leggett A J 1970 *Phys. Rev. Lett.* **25** 1543
- [4] Prokof'ev N 2007 *Adv. Phys.* **56** 381
- [5] Golovko V A 1996 *Physica A* **230** 658
- [6] Golovko V A 2008 *Progress in Statistical Mechanics Research*, ed J. S. Moreno (Hauppauge, NY: Nova Science Publ.) ch 2
- [7] Golovko V A 1997 *Physica A* **246** 275
- [8] Balibar S and Caupin F 2008 *J. Phys.: Condens. Matter* **20** 173201
- [9] Pollet L, Boninsegni M, Kuklov A B, Prokof'ev N V, Svistunov B V and Troyer M 2008 *Phys. Rev. Lett.* **101** 097202
- [10] Golovko V A 2004 *J. Math. Phys.* **45** 1571
- [11] Lifshitz E M and Pitaevskii L P 1980 *Statistical Physics. Part 2* (Oxford: Pergamon) sec 55
- [12] Golovko V A 2004 *Physica A* **341** 340
- [13] Golovko V A 2002 *Physica A* **310** 39
- [14] Sonin E B 1982 *Usp. Fiz. Nauk* **137** 267 [*Sov. Phys. Usp.* **25** 409]
- [15] Golovko V A 2007 *Physica A* **374** 15
- [16] Golovko V A 2001 *Physica A* **300** 195
- [17] Kirkwood J G and Monroe E 1941 *J. Chem. Phys.* **9** 514
- [18] Baus M 1990 *J. Phys.: Condens. Matter* **2** 2111
- [19] Singh Y 1991 *Phys. Rep.* **207** 351
- [20] Guyer R A 1969 *Sol. Stat. Phys.* **23** 413
- [21] Day J and Beamish J 2006 *Phys. Rev. Lett.* **96** 105304
- [22] Ray M W and Hallock R B 2008 *Phys. Rev. Lett.* **100** 235301
- [23] Balibar S and Caupin F 2008 *Phys. Rev. Lett.* **101** 189601
- [24] Ray M W and Hallock R B 2008 *Phys. Rev. Lett.* **101** 189602




Figure 1. Function $u_2(r)$




Figure 2. Function $\sigma(k)$; the broken curve is explained in Sec. 4