### On a generalization of $P_3(n)$

#### AGUSTIN MORENO CAÑADAS

Departamento de Matemáticas, Universidad Nacional de Colombia Bogotá-Colombia amorenoca@unal.edu.co

We present a formula for  $P_3^s(n)$  the number of partitions of a positive number n into 3 s-gonal numbers, by using representations of posets over  $\mathbb{N}$ .

Keywords: partition, polygonal number, poset, representation, square number.

Mathematics Subject Classification 2000:05A17;11D25;11D45;11D85;11E25;11P83.

#### 1. Introduction

In [10] Lehmer denoted  $P_k(n)$  the number of partitions of a natural number n into k integral squares  $\geq 0$ , and solved almost completely the equation  $P_k(n) = 1$ . Lehmer claimed that the general problem of finding a formula for  $P_k(n)$  was a problem of great complexity. The case k = 3 was studied by Grosswald, A. Calloway, and J. Calloway in [4], and Grosswald solved (essentially) the problem, giving the number of partitions of an arbitrary integer n into k squares (taking into account that, he didn't distinguish between partitions that contains zeros and those that do not) [5].

The main goal of this paper is to give a formula for  $P_3^s(n)$ , where the s-th polygonal number of order or rank r,  $p_r^s$  is given by the formula (often, 0 is included as a polygonal number [3,7]),

$$p_r^s = \frac{1}{2}[(s-2)r^2 - (s-4)r].$$

We must note that, for the particular case s=3 Hirschhorn and Sellers proved the identity  $P_3^3(27n+12)=3P_3^3(3n+1)$ , via generating functions manipulations and some combination arguments [9]. Instead of generating functions, we shall give the formula for  $P_3^s(n)$  (note that  $P_3(n)=P_3^4(n)$ ) by using representations of posets over the set of natural numbers  $\mathbb{N}$ , which have been used in [11] to give criteria for natural numbers which are expressible as sums of three polygonal numbers of positive rank.

#### 2. Preliminaries

#### 2.1. Posets


An ordered set (or partially ordered set or poset) is an ordered pair of the form  $(\mathcal{P}, \leq)$  of a set  $\mathcal{P}$  and a binary relation  $\leq$  contained in  $\mathcal{P} \times \mathcal{P}$ , called the order (or the partial order) on  $\mathcal{P}$ , such that  $\leq$  is reflexive, antisymmetric and transitive [1]. The elements of  $\mathcal{P}$  are called the points of the ordered set.

Let  $\mathcal P$  be an ordered set and let  $x,y\in \mathcal P$  we say x is covered by y if x< y and  $x\leq z< y$  implies z=x.

Let  $\mathcal{P}$  be a finite ordered set. We can represent  $\mathcal{P}$  by a configuration of circles (representing the elements of  $\mathcal{P}$ ) and interconnecting lines (indicating the covering relation). The construction goes as follows.

- (1) To each point  $x \in \mathcal{P}$ , associate a point p(x) of the Euclidean plane  $\mathbb{R}^2$ , depicted by a small circle with center at p(x).
- (2) For each covering pair x < y in  $\mathcal{P}$ , take a line segment l(x,y) joining the circle at p(x) to the circle at p(y).
- (3) Carry out (1) and (2) in such a way that
  - (a) if x < y, then p(x) is lower than p(y),
  - (b) the circle at p(z) does not intersect the line segment l(x,y) if  $z \neq x$  and  $z \neq y$ .

A configuration satisfying (1)-(3) is called a Hasse diagram or diagram of  $\mathcal{P}$ . In the other direction, a diagram may be used to define a finite ordered set; an example is given below, for the ordered set  $\mathcal{P} = \{a, b, c, d, e, f\}$ , in which a < b < c < d < e, and f < c.



We have only defined diagrams for finite ordered sets. It is not possible to represent the whole of an infinite ordered set by a diagram, but if its structure is sufficiently regular it can often be suggested diagrammatically.

An ordered set C is called a *chain* (or a *totally ordered set* or a *linearly ordered set*) if and only if for all  $p, q \in C$  we have  $p \leq q$  or  $q \leq p$  (i.e., p and q are comparable).

Let  $\mathcal P$  be a poset and  $S \subset \mathcal P$ . Then  $a \in S$  is a maximal element of S if  $a \leq x$  and  $x \in S$  imply a = x. We denote the set of maximal elements of S by Max S. If S (with the order inherited from  $\mathcal P$ ) has a top element,  $\top_S$  (i.e.,  $s \leq \top_S$  for all  $s \in S$ ), then Max  $S = \{\top_S\}$ ; in this case  $\top_S$  is called the *greatest* (or *maximum*) element of S, and we write  $\top_S = \max S$ .

Suppose that  $\mathcal{P}_1$  and  $\mathcal{P}_2$  are (disjoint) ordered sets. The disjoint union  $\mathcal{P}_1 + \mathcal{P}_2$  of  $\mathcal{P}_1$  and  $\mathcal{P}_2$  is the ordered set formed by defining  $x \leq y$  in  $\mathcal{P}_1 + \mathcal{P}_2$  if and only if either

 $x, y \in \mathcal{P}_1$  and  $x \leq y$  in  $\mathcal{P}_1$  or  $x, y \in \mathcal{P}_2$  and  $x \leq y$  in  $\mathcal{P}_2$ . A diagram for  $\mathcal{P}_1 + \mathcal{P}_2$  is formed by placing side by side diagrams for  $\mathcal{P}_1$  and  $\mathcal{P}_2$ .

#### 2.2. Partitions and Representations of posets over $\mathbb{N}$

As usual in this paper  $\mathbb N$  denotes the set of natural numbers, while  $\mathbb N\setminus\{0\}$  is the set of positive integers.

We denote  $t_k=p_k^3=\frac{k(k-1)}{2}$  the k-th triangular number  $k\in\mathbb{Z}$ , and  $s_k=p_k^4=k^2$  is the k-th square number.

A partition of a positive integer n is a finite nonincreasing sequence of positive integers  $\lambda_1, \lambda_2, \ldots, \lambda_r$  such that  $\sum_{i=1}^r \lambda_i = n$ . The  $\lambda_i$  are called the parts of the partition [6]. A composition is a partition in which the order of the summands is considered.

Often the partition  $\lambda_1, \lambda_2, \dots, \lambda_r$  will be denoted by  $\lambda$  and we sometimes write  $\lambda = (1^{f_1} 2^{f_2} 3^{f_3} \dots)$  where exactly  $f_i$  of the  $\lambda_j$  are equal to i. Note that  $\sum_{i>1} f_i i = n$ .

The partition function p(n) is the number of partitions of n. Clearly p(n) = 0 when n is negative and p(0) = 1, where the empty sequence forms the only partition of zero.

Let  $(\mathbb{N}, \leq)$  be the set of natural numbers endowed with its natural order and  $(\mathcal{P}, \leq')$  a poset. A representation of  $\mathcal{P}$  over  $\mathbb{N}$  [11] is a system of the form

$$\Lambda = (\Lambda_0 \; ; \; (n_x, \lambda_x) \mid x \in \mathcal{P}), \tag{1}$$

where  $\Lambda_0 \subset \mathbb{N}$ ,  $\Lambda_0 \neq \emptyset$ ,  $n_x \in \mathbb{N}$ ,  $\lambda_x$  is a partition with parts in the set  $\Lambda_0$ , and  $|\lambda_x|$  is the size of the partition  $\lambda_x$ , in particular if  $n_x = 0$  then we consider  $\lambda_x = 0$ . Further

$$x \le y \Rightarrow n_x \le n_y, \quad |\lambda_x| \le |\lambda_y|, \quad \text{and} \quad \max\{\lambda_x\} \le \max\{\lambda_y\}.$$
 (2)

#### 2.3. The associated graph

A Graph is a pair G=(V,E) of sets satisfying  $E\subseteq V^2$ , thus the elements of E are 2-elements subsets of V, such that  $V\cap E=\varnothing$ . The elements of V are the vertices of the graph G, the elements of E are its edges. A graph with vertex set V is said to be a graph on V. The vertex set of a graph is referred to as V(G), its edge set as E(G). We write  $v\in G$  to a vertex  $v\in V(G)$  and  $e\in G$ , for an edge  $e\in E(G)$ , an edge  $\{x,y\}$  is usually written as xy or yx.

A vertex v is *incident* with an edge e; if  $v \in e$ , then e is an edge at v. The two vertices incident with an edge are its *endvertices* or *ends*, and an edge *joins* its ends. A *path* is a non-empty graph P = (V, E) of the form


$$V = \{x_0, x_1, \dots, x_k\}, \quad E = \{x_0x_1, x_1x_2, \dots, x_{k-1}x_k\},$$

where the  $x_i$  are all distinct. The vertices  $x_0$  and  $x_k$  are linked by P and are called its ends, the vertices  $x_1, \ldots, x_{k-1}$  are the inner vertices of P. The number of edges of a path is its length, and the path of length k is denoted by  $P^k$ . We often refer to a path

by the natural sequence of its vertices writing  $P = x_0 x_1 \dots x_k = x_0 || x_k$ , and calling P a path from  $x_0$  to  $x_k$  [2].

Given a representation  $\Lambda$  for an poset  $(\mathcal{P}, \leq)$  in [11,12] it was defined its associated graph,  $\Gamma_{\Lambda}$  which has as set of vertices the points of  $\mathcal{P}$ , and containing all information about partitions of the numbers  $n_x$ . That is  $\Gamma_{\Lambda}$  is represented in such a way that to each vertex of the graph it is attached, either a number  $n_x$  given by the representation or one part of a partition of some  $n_y$  representing some  $y \in \mathcal{P}$  such that  $x \leq y$ .

As an example, we consider [11] an infinite sum of infinite chains pairwise incomparable  $\mathbb{R}$  in such a way that  $\mathbb{R} = \sum_{i=0}^{\infty} C_i$ , where  $C_j$  is a chain such that  $C_j = v_{0j} < v_{1j} < v_{2j} < \ldots$ . It is defined a representation over  $\mathbb{N}$  for  $\mathbb{R}$ , by fixing a number  $n \geq 3$  and assigning to each  $v_{ij}$  the pair  $(n_{ij}, \lambda_{ij}) = (3 + (n-2)i + (n-1)j, (3 + (n-2)i + (n-1)j)^1)$ , we note  $\mathbb{R}_n$  this representation, and write  $v_{ij} \in \mathbb{R}_n$  whenever it is assigned the number  $n_{ij} = 3 + (n-2)i + (n-1)j$  to the point  $v_{ij} \in \mathbb{R}$  in this representation. Fig. 2 below suggests the Hasse diagram for this poset with its associated graph  $\Gamma_{p_k^n}$  which attaches to each vertex  $v_{ij}$  the number 3 + (n-2)i + (n-1)j,  $i, j \geq 0$ .



 $v_{t_{(i-1)i}} \in l.b.p$ , (left boundary path),  $t_{-1} = 0$ .

Fig. 2

Note that the vertices of the form  $v_{(2+7i+3k)(2(i+k)+3)}$ ,  $i, k \geq 0$ , do not lie on any non-trivial path of  $\Gamma_{p_k^n}$ .

The representations  $\mathcal{R}_n$  defined above induce an equivalence relation  $\sim$  on  $\mathcal{R}$  in such a way that  $v_{ij} \sim v_{kl}$  if and only if  $n_{ij} = n_{kl}$ . We denote  $[v_{ij}]$  the class of the point  $v_{ij} \in \mathcal{R}$ . Hence the points of

$$\mathcal{H} = \bigcup_{k \geq 1, m \geq 0} [v_{(2^{k-1}(72^k-3) + m(2^{2(k+1)}-1))(32^k+2m)}] \cup \bigcup_{i \geq 0, s \geq 0} [v_{(2+7i+3s)(2(i+s)+3)}] \subset \mathcal{R}$$

do not lie on any non-trivial path of  $\Gamma_{p_k^n}$ , if  $\mathcal{R}$  is represented in such a way that  $(n_{ij}, \lambda_{ij}) = (2i + j, (2i + j)^1), i, j \geq 0.$ 

We say that a path  $P \in \Gamma_{p_k^n}$  is admissible if and only if either  $P = P_{i0}$ , or  $P = P_{i0}P_{ij}$ , where  $P_{i0}$ ,  $P_{ij}$  are paths such that

$$P_{i0} = v_{00} \mid\mid v_{t_i(i+1)}, \ i > -1, \qquad P_{ij} = v_{t_i(i+1)} \mid\mid v_{(t_i + t_j)(i+j+2)},$$

and the inner vertices of  $P_{i0}$  have the form  $v_{t_h(h+1)}$ , -1 < h < i, while the inner vertices for  $P_{ij}$  have the form  $v_{(t_i+t_l)(i+l+2)}$ , -1 < l < j. Or  $P = P_{i0}P_{ij}P_{ijk}$ , where  $P_{ijk}$  is a path such that

$$P_{ijk} = v_{(t_i+t_i)(i+j+2)} \mid v_{(t_i+t_i+t_k)(i+j+k+3)}, -1 < k \le j.$$

If for  $s\geq 3$  fixed, we consider the representation  $\mathcal{R}_s$  then the numbers  $n_{t_i(i+1)}$  representing the vertices  $v_{t_i(i+1)}\in\mathcal{R}_s$  of the left boundary path, l.b.p are expressible in the form  $n_{t_i(i+1)}=p_1^s+p_1^s+p_{i+2}^s$ , s is an fixed index,  $i\geq -1$ . Furthermore if  $i_0\geq 0$  is a fixed number, and  $v_{t_{i_0}(i_0+1)}\in l.b.p$  then  $n_{(t_{i_0}+t_l)(i_0+l+2)}$  representing the vertex  $v_{(t_{i_0}+t_l)(i_0+l+2)}\in P_{i_0}P_{ij}$ , can be expressible in the form  $n_{(t_{i_0}+t_l)(i_0+l+2)}=p_{i_0+2}^s+p_{i+2}^s+p_1^s$ , and a number  $n_{(t_{i_0}+t_j+t_k)(i_0+j+k+3)}=p_{i_0+2}^s+p_{i+2}^s+p_{i+2}^s$ . In [11] has been used this facts to state the following theorem ;

**Theorem 1.** A number  $m \in \mathbb{N} \setminus \{0\}$  is the sum of three *n*-gonal numbers of positive rank if and only if *m* represents a vertex  $v_{ij} \in \mathcal{R}_n$  in a non-trivial component of  $\Gamma_{p_i^n}$ .

For example, if  $\mathcal{R}_3'$  is a representation such that  $(n_{ij}, \lambda_{ij}) = (i+j, (i+j)^1), i, j \geq 0$ , and  $\mathcal{G}_i$  is the family of subsets of  $\mathcal{R}$  such that for  $i \geq 0$ ,  $\mathcal{G}_i = \{[v_{(t_i+j)(i+2)}] \mid 0 \leq j \leq t_{i+1}\}$  then  $V(\Gamma_{p_k^n}) = \bigcup_i (\mathcal{G}_i \cup [v_{00}] \cup [v_{01}])$ , because every natural number is expressible as a sum of three or fewer triangular numbers.

## 3. The main result

If  $v_{i_0j} \in \Gamma_{p_k^n}$  belongs to an non-trivial component, and  $v_{i_0j} \in \mathcal{R}_s$  then there exists an admissible path  $P = v_{00} \mid\mid v_{i_0j}$ , which has associated a family of compositions (see 2.2) in such a way that if l > 0 then

$$\begin{split} n_{t_{i_0}(i_0+1)} &= p^s_{i_0+2} + p^s_1 + p^s_1, \quad i_0 \geq -1 \quad \text{if } P = P_{i_00}, \\ n_{(t_{i_0}+t_l)(i_0+l+2)} &= p^s_{l+2} + p^s_{i_0+2} + p^s_1 = p^s_{i_0+2} + p^s_{l+2} + p^s_1, \quad \text{if } P = P_{i_00}P_{i_0j}, \quad \text{(3)} \\ n_{(t_{i_0}+t_j+t_k)(i_0+j+k+3)} &= p^s_{i_0+2} + p^s_{j+2} + p^s_{k+2} = p^s_{k+2} + p^s_{j+2} + p^s_{i_0+2}, \quad \text{and} \\ n_{(t_{i_0}+t_j+t_k)(i_0+j+k+3)} &= p^s_{j+2} + p^s_{k+2} + p^s_{i_0+2}, \quad -1 \leq i_0 \leq k \leq j, \quad \text{if } P = P_{i_00}P_{i_0j}P_{i_0jk}. \end{split}$$

Thus we say that two admissible paths P, Q are equivalent if and only if they have associated the same partitions. If  $P = v_{00} \mid \mid v_{ij}$  is an admissible path then we note [P] the class of P. Therefore [P] can be determined by fixing the end  $v_{ij}$ , we note  $v_{ij}(P)$  whenever a vertex  $v_{ij}$  has been fixed due to this condition (i.e,  $v_{ij}(P) = [P]$ ,  $v_{ij}(Q) = \emptyset$  for  $Q \in [P]$ ,  $v_{ij} \in [v_{ij}]$ , and  $v_{ij} \neq v_{ij}$ ). For example  $v_{t_i(i+1)}(P_{t_i0}) = [P_{t_i0}]$  only contains the admissible path  $v_{t_i0}$ .

If  $v_{ij}(P)$  is the set of admissible paths of the form  $v_{00}||u_{rs}|$  where  $u_{rs} \in [v_{ij}]$  then we note

$$\mathcal{A}(v_{ij}) = \{w_{ij} \in [v_{ij}] \mid \overline{w_{ij}(P)} = [P], P \in v_{ij}(P)\}, \text{ therefore } v_{ij}(P) = \bigcup_{w_{ij} \in \mathcal{A}(v_{ij})} \overline{w_{ij}(P)},$$

If  $\mathcal{A}(v_{ij}) \neq \emptyset$  and  $w_{ij} \in \mathcal{A}(v_{ij})$  then  $\delta_g(w_{ij})$  denotes the number of classes of admissible paths at  $w_{ij}$ . If  $\mathcal{A}(v_{ij}) = \emptyset$  then  $\delta_g(v_{ij}) = 0$ .

The next theorem is a consequence of theorem 1 and the equations (3).

**Theorem 2.** Let  $P_3^s(n)$  denote the number of partitions of  $n \in \mathbb{N}$  into three s-gonal numbers,  $s \geq 3$ . then

$$P_3^s(n) = \begin{cases} \sum_{w_{ij} \in \mathcal{A}(v_{ij})} \delta_g(w_{ij}), & \text{if } \mathcal{A}(v_{ij}) \neq \varnothing, \\ 0, & \text{otherwise.} \end{cases}$$

where  $v_{ij} \in \mathcal{R}_s$  and  $n_{ij} = n$ .

For example in  $\Re_4$ ,  $[v_{(10,5)}] \cap \mathcal{A}(v_{(10,5)}) = \{v_{(10,5)}, v_{(7,7)}\}, \delta_g(v_{(10,5)}) = 1, \delta_g(v_{(7,7)}) = 1,$ 

If 
$$P = v_{(0,0)} ||v_{(0,1)}||v_{(6,5)}||v_{(7,7)}, Q = v_{(0,0)}||v_{(10,5)}||v_{(10,5)}||$$

 $\overline{v_{(7,7)}(P)} = \{v_{(0,0)}||v_{(0,1)}||v_{(6,5)}||v_{(7,7)},v_{(0,0)}||v_{(6,4)}||v_{(7,6)}||v_{(7,7)},v_{(0,0)}||v_{(2,1)}||v_{(7,6)}||v_{(7,7)}\}.$ 

$$\overline{v_{(10.5)}(Q)} = \{v_{(0.0)} || v_{(10.5)} \}.$$
 Therefore  $P_3^4(38) = 2$ .

# References

- B.A. Davey and H.A. Priestley, Introduction to Lattices and Order, Cambridge University Press, 2002.
- [2] R. Diestel, Graph Theory, Springer-Verlag, 1997.
- [3] Duke. W, Some old and new results about quadratic forms, Notices. Amer. Math. Soc 44 (1997), 190–196.
- [4] E. Grosswald, A. Calloway, and J. Calloway, The representations of integers by three positive squares, Proc. Amer. Math. Soc 10 (1959), 451–455.
- [5] E. Grosswald, Partitions into squares, L'Enseignement mathématique 30 (1984), 223–245.
- [6] \_\_\_\_\_\_, Representations of Integers as Sums of Squares, Springer-Verlag, New York,
- [7] R. Guy, Every number is expressible as a sum of how many polygonal numbers, A.M.M 101 (1994), 169–172.

- [8] \_\_\_\_\_\_, Unsolved Problems in Number Theory, 3rd ed., Springer-Verlag, New York, 2004.
- [9] M. D. Hirschhorn and J. A. Sellers, *Partitions into three triangular numbers*, Australasian Journal of Combinatorics **30** (2004), 307–318.
- [10] D. H. Lehmer, On the partition of numbers into squares, A.M.M 55 (1948), no. 8, 476–481.
- [11] A. Moreno, On sums of figurate numbers by using techniques of poset representation theory, Preprint (2008).