

Preferred basis without decoherence

Roberto Laura

*Facultad de Ciencias Exactas, Ingeniería y Agrimensura
(UNR) and Instituto de Física Rosario (CONICET - UNR),
Av. Pellegrini 250, 2000 Rosario, Argentina**

Leonardo Vanni

*Instituto de Astronomía y Física del Espacio (UBA - CONICET). Casilla de Correos 67,
Sucursal 28, 1428 Buenos Aires, Argentina.†*

(Dated: diciembre 2008)

Abstract

The aim of this paper is to argue that the “preferred basis problem” is not a real problem in measurement. We will show that, given an apparatus, among the infinite correlations that can be established in the final state by means of a change of basis, one and only one makes physical sense. It is the apparatus, through its interaction Hamiltonian, what selects a single basis and determines the observable to be measured, even without decoherence.

*Electronic address: rlaura@fceia.unr.edu.ar

†Electronic address: lv@iafe.uba.ar

I. INTRODUCTION

Let us consider the measurement of the observable $A = \sum a_i |a_i\rangle\langle a_i|$ of a system S , by means of an apparatus M with a pointer observable $Z = \sum z_i |z_i\rangle\langle z_i|$. Let us suppose that $\{|a_i\rangle\}$ is a basis of the Hilbert space of S , and $\{|z_i\rangle\}$ is a basis of the Hilbert space of M . Before the interaction between S and M , S is in the state $|\varphi\rangle = \sum c_i |a_i\rangle$, and M is prepared in a ready-to-measure state $|z_0\rangle$. According to the von Neumann model of measurement [1], the interaction introduces a correlation between the eigenstates of A and the eigenstates of Z :

$$|\varphi\rangle = \sum_i c_i |a_i\rangle |z_0\rangle \quad \longrightarrow \quad |\psi\rangle = U_{\Delta t} |\varphi\rangle = \sum_i c_i |a_i\rangle |z_i\rangle \quad (1)$$

where $U_{\Delta t} = e^{-iH_{int}\Delta t}$ is the evolution operator corresponding to the interaction Hamiltonian H_{int} . After the time Δt , the interaction ends and the correlation is established: since then it is assumed that, if the pointer Z acquires a value z_i , then the value of A measured in the process is a_i (see [2], [3]).

Eq. (1) expresses the biorthonormal decomposition of the state $|\psi\rangle$ of the composite system $S + M$, (see [4]): it establishes the correlation that defines the measurement. Nevertheless, such a decomposition is not always unique. In particular, if the coefficients c_i are all equal in absolute value, it is possible to introduce a change of basis $\{|a_i\rangle\} \rightarrow \{|a'_i\rangle\}$ and $\{|z_i\rangle\} \rightarrow \{|z'_i\rangle\}$ (see Appendix A) such that

$$|\psi\rangle = \sum_i c_i |a_i\rangle |z_i\rangle = \sum_i c'_i |a'_i\rangle |z'_i\rangle \quad (2)$$

This result is what leads to the so-called “preferred basis problem” [5]: since the two different sets of outcomes, $\{z_i\}$ and $\{z'_i\}$, are both admissible for same measurement, there is an intrinsic ambiguity about what observable is measured on the system S . In other words, if Z acquires a value z_i , A is measured according to the first expansion of $|\psi\rangle$; but if Z acquires a value z'_i , then A' is measured according to the second expansion. Therefore, the observer could not know whether the measured observable is A or A' , even in the case that $[A, A'] \neq 0$. From the viewpoint of the decoherence program, if the composite system $S + M$ remains isolated, this problem cannot be solved. For this reason, a further interaction with the environment is necessary in order to determine the preferred –pointer– basis: it is such an interaction what, via decoherence, selects the observable to be measured. This argument

has been widely accepted in the literature (see, for instance [5]).

The aim of this paper is conceptual: our purpose is to argue that the “preferred basis problem” is not a real problem in measurement. Instead of trying to solve the supposed difficulty by adding an interaction after the correlation given in eq. (2), we will analyze the process leading to that correlation. On this basis we will show that, given an apparatus, among the infinite correlations that can be established in the final state by means of a change of basis, one and only one makes physical sense. It is the apparatus, through its interaction Hamiltonian, what selects a single basis and determines the observable to be measured, even when the composite system $S + M$ is a closed system.

II. ONE SET OF OUTCOMES

As we have pointed out, if the coefficients c_i that span the initial state of the system are all equal, the change of basis is possible:

$$|\varphi\rangle = \sum_i c_i |a_i\rangle |z_0\rangle \quad \rightarrow \quad |\psi\rangle = U|\varphi\rangle = \sum_i c_i |a_i\rangle |z_i\rangle \quad (3)$$

$$= \sum_i c'_i |a'_i\rangle |z'_i\rangle \quad (4)$$

The equality between eqs. (3) and (4) leads to the preferred basis problem. However, this would really be a problem if there existed an apparatus where the two different sets $\{z_i\}$ and $\{z'_i\}$ could both be the outcomes of the measurement, with no modification at all in the experimental arrangement [13]. Only in this situation the decompositions (3) and (4) would be both physically meaningful, and their equality would actually imply the ambiguity introduced by the problem.

Let us suppose that such an apparatus exists and, as a consequence, eqs. (3) and (4) really imply different sets of outcomes of the same measuring apparatus. For simplicity, we consider an ideal measurement, where the sets of states $\{|z_i\rangle\}$ and $\{|z'_i\rangle\}$ share the ready-to-measure state, $|z_0\rangle = |z'_0\rangle$. Then, from the calibration of the measurement [2] in eq. (3), when we prepare the system S in the state $|a_i\rangle$, the pointer Z will acquire the value z_i corresponding to its eigenvector $|z_i\rangle$:

$$|a_i\rangle |z_0\rangle \quad \rightarrow \quad |a_i\rangle |z_i\rangle. \quad (5)$$

Nevertheless, from the calibration in eq. (4), when we prepare S in the state $|a'_i\rangle$, Z will acquire the value z'_i corresponding to $|z'_i\rangle$:

$$|a'_i\rangle|z_0\rangle \quad \rightarrow \quad |a'_i\rangle|z'_i\rangle. \quad (6)$$

Now, since $\{|a_i\rangle\}$ and $\{|a'_i\rangle\}$ are two bases of the Hilbert space of the system S , we can introduce the change of basis $|a'_i\rangle = \sum_k \alpha_k |a_k\rangle$. Then, if we introduce this change of basis in eq. (5), and given the linearity of the evolution induced by the apparatus, we also obtain

$$|a'_i\rangle|z_0\rangle = \left(\sum_k \alpha_k |a_k\rangle \right) |z_0\rangle \quad \rightarrow \quad \sum_k \alpha_k |a_k\rangle |z_k\rangle \quad (7)$$

Let us compare eqs. (6) and (7): eq. (6) corresponds to the outcome z'_i but, on the other hand, eq. (7) says that we will obtain one of the outcomes $\{z_i\}$. Independently of any macroscopicity condition that allows a human being to distinguish between the set $\{z_i\}$ and the set $\{z'_i\}$ [6], if these sets are different, they represent different physical outcomes [14]. In other words, eqs. (6) and (7) represent different physical processes with different physical results, one where the pointer Z acquires one of the values $\{z_i\}$, and the other where Z acquires one of the values $\{z'_i\}$. However, both processes begin in the same state $|a'_i\rangle$ of the system S : the only difference between them is that the state $|a'_i\rangle$ is decomposed in different bases. As a consequence, if there existed an apparatus where the two different sets $\{z_i\}$ and $\{z'_i\}$ could both be the outcomes of the same measurement, such an apparatus would distinguish between the two mathematical representations of the same initial state of S : the outcomes $\{z_i\}$ would indicate the representation of the state in the basis $\{|a_i\rangle\}$, and the outcomes $\{z'_i\}$ would indicate the representation of the state in the basis $\{|a'_i\rangle\}$. This means that the apparatus would distinguish mathematical entities, and this is physically absurd. Therefore, this kind of apparatus does not exist: any physical apparatus can only supply a single set of outcomes.

Summing up, independently of the mathematical equality between eqs. (6) and (7), one and only one of these correlations makes physical sense. Although the change of basis is mathematically correct, the previous argument shows that, with no modification in the apparatus, either the outcomes $\{z_i\}$ or the outcomes $\{z'_i\}$ can be obtained, but not both. Although the argument dissolves the supposed ambiguity about the preferred basis, it does not tell us yet which set is effectively obtained and, then, which the observable measured

on the system S is. As we will see, this question can be answered when measurement is considered as a process.

III. WHAT SET OF THE OUTCOMES?

The previous section proves that any measuring apparatus, if it is a quantum apparatus, can only activate a single set of outcomes, associated with a certain basis. In this section we will formulate an argument for specifying that basis.

As we have seen, only one of the correlations (3) or (4) makes physical sense, although the equality between them holds. This equality only expresses a change of basis on the final state, once it has been obtained. But endowing the mathematical change of basis with a physical content amounts to ignoring the fact that a measurement is a *quantum process* and, as such, it is governed by the dynamical equation of the theory. In fact, the dynamics of the process is given by the evolution operator U through the corresponding Hamiltonian: by beginning with an initial state, it is U what produces the correlation when applied to that initial state. Therefore, the correlation with physical meaning is that resulting from the application of U to the initial state. And since the evolution operator U is unitary, given the initial state, the correlation corresponding to this U is unique, in spite of the possibility of performing a mathematical change of basis on the final state.

In order to rephrase the argument in a formal way, let us consider the two following processes:

$$|\varphi_A\rangle = \sum_i c_i |a_i\rangle |z_0\rangle \quad \longrightarrow \quad |\psi_A\rangle = U|\varphi_A\rangle = \sum_i c_i |a_i\rangle |z_i\rangle \quad (8)$$

$$|\varphi_{A'}\rangle = \sum_i c'_i |a'_i\rangle |z'_0\rangle \quad \longrightarrow \quad |\psi_{A'}\rangle = U'|\varphi_{A'}\rangle = \sum_i c'_i |a'_i\rangle |z'_i\rangle, \quad (9)$$

According to the argument of the basis ambiguity, $|\psi_A\rangle = |\psi_{A'}\rangle$, but represented in different bases, which correspond to different correlations. But since the correlation is established by the evolution operator, different correlations require different evolution operators $U \neq U'$ and, then, different interaction Hamiltonians $H \neq H'$ which, in turn, correspond to different apparatuses. This means that the arrows of eqs. (8) and (9) represent different measurement processes: with the apparatus that induces the evolution given by U , we obtain an outcome belonging to the set $\{z_i\}$ and never an outcome of the set $\{z'_i\}$, and analogously for the apparatus corresponding to U' .

By contrast with the argument of the basis ambiguity, which only focuses on the final state, our argument traces the process back in order to see what dynamical evolution leads to that final state. It is represented in the next figure:

Measurement process before the correlation

$$\begin{array}{c}
 |\varphi_A\rangle \text{ --- } |\psi_A\rangle = U|\varphi_A\rangle \\
 \text{--- } U \neq U' \\
 |\varphi_{A'}\rangle \text{ --- } |\psi_{A'}\rangle = U'|\varphi_{A'}\rangle
 \end{array}
 \quad \Rightarrow \quad
 |\psi_A\rangle = |\psi_{A'}\rangle = \sum_i c_i |a_i\rangle |z_i\rangle = \sum_i c'_i |a'_i\rangle |z'_i\rangle$$

With this strategy, it is easy to show that different sets of outcomes finally imply different measuring apparatuses. As we can see, if one of the outcomes z_i is obtained, the preferred basis is the eigenbasis of A , and the correlation corresponding to $|\psi_A\rangle$ is introduced by U , that is, by the apparatus with Hamiltonian H : this apparatus measures the observable A of S . Analogously, if one of the outcomes z'_i is obtained, the preferred basis is the eigenbasis of A' , but now the correlation corresponding to $|\psi_{A'}\rangle$ is introduced by U' , that is, by a different apparatus with Hamiltonian H' : this second apparatus is what measures the observable A' of S .

Summing up, the supposed basis ambiguity vanishes once one acknowledges that a quantum measurement is not fully specified by the correlation expressed in the final state of the system $S + M$, but is defined by the process that, starting from the initial state, introduces the correlation in that final state. And decoherence has played no role in this argument.

IV. MEASUREMENT OF THE SPIN

A. Stern-Gerlach measurement

Let us consider a Stern-Gerlach device with a magnetic field in direction z , which correlates the projection S_z of the spin of the particle with its momentum p_z . The interaction Hamiltonian of this apparatus, after a Δt , is given by $H_{int} = -cz \otimes S_z$ [6], where c is

a constant related to the magnetic momentum of the particle. The evolution operator is, then, $U = e^{icz \otimes S_z}$. Let us suppose that the initial state of the composite system before the interaction is $|\varphi_A\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle)|p_0\rangle$, where $|p_0\rangle$ is the momentum state with $p_z = 0$. In this case, the supposed basis ambiguity is expressed as

$$|\varphi_A\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle)|p_0\rangle \quad \longrightarrow \quad |\psi_A\rangle = U|\varphi_A\rangle = \frac{1}{\sqrt{2}}|+\rangle|p_+\rangle + \frac{1}{\sqrt{2}}|-\rangle|p_-\rangle \quad (10)$$

$$= \frac{1}{\sqrt{2}}|\odot\rangle|p_\odot\rangle + \frac{1}{\sqrt{2}}|\otimes\rangle|p_\otimes\rangle \quad (11)$$

where

$$|\odot\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle), \quad |\otimes\rangle = \frac{1}{\sqrt{2}}(|+\rangle - |-\rangle), \quad (12)$$

are the eigenstates of S_x , and $|p_\odot\rangle, |p_\otimes\rangle$ are given by the combinations

$$|p_\odot\rangle = \frac{1}{\sqrt{2}}(|p_+\rangle + |p_-\rangle), \quad |p_\otimes\rangle = \frac{1}{\sqrt{2}}(|p_+\rangle - |p_-\rangle) \quad (13)$$

The evolution operator $U = e^{icz \otimes S_z}$ applied to the initial state $|\varphi_A\rangle$ produces the correlation given by eq. (10), and not that given by eq. (11). Then, the outcomes that may be obtained as a result of the measurement are p_+ or p_- , and not p_\odot or p_\otimes . In spite of the equality between the eqs. (10) and (11), only the correlation (10) has physical meaning, because it is produced by U : there is no ambiguity about whether the apparatus measures S_z or S_x . In other words, although the equality is valid, eq. (11) does not imply that S_x has been measured. If we want to measure S_x , we have to change the interaction introduced by the apparatus by rotating the magnetic field from direction z to direction x . But this amounts to change the apparatus itself, with its evolution operator: in this new case, $U' = e^{icx \otimes S_x}$.

This example has an additional particular feature that allows us to discard p_\odot and p_\otimes as possible outcomes of the measurement. Even under the ambiguity of the basis change, the states $|p_\odot\rangle$ and $|p_\otimes\rangle$, resulting from the combinations of the states $|p_+\rangle$ and $|p_-\rangle$, are not possible states of the apparatus that measures S_x , as eq. (11) might suggest. In fact, the values p_+ and p_- , corresponding to the states $|p_+\rangle$ and $|p_-\rangle$, are associated to wavepackets leading to “spot up” and “spot down” on the screen; they physically mean that S_z was measured with the value $1/2$ or $-1/2$, respectively. But, in this case, the values p_\odot and p_\otimes , corresponding to the states $|p_\odot\rangle$ and $|p_\otimes\rangle$, cannot be associated to “spot right”, “spot left”, that is, to the result that the measurement of S_x would produce: $|p_\odot\rangle$ and $|p_\otimes\rangle$ are linear

combinations of $|p_+\rangle$ and $|p_-\rangle$, and none combination of wavepackets “up” and “down” can produce “right” and “left” as a result.

B. Bit-by-bit measurement.

Let us now consider a system of spin $1/2$, for instance, a particle (the measured system S), measured by another system of spin $1/2$, for instance, an atom with two levels (the measuring apparatus M). The basis of the Hilbert space of the particle is $\{|+\rangle, |-\rangle\}$, and the basis of the Hilbert space of the atom is $\{|\uparrow\rangle, |\downarrow\rangle\}$. The interaction Hamiltonian after a Δt is $H_{int} = -\pi|-\rangle\langle-| \otimes |\leftarrow\rangle\langle\leftarrow|$ (see [7], [8]). The evolution operator is, then, $U = e^{i\pi|-\rangle\langle-| \otimes |\leftarrow\rangle\langle\leftarrow|}$. Let us suppose that the initial state of the composite system before the interaction is $|\varphi_A\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle)|\uparrow\rangle$, where $|\uparrow\rangle$ is the initial state of the atom. In this case, the supposed basis ambiguity is expressed as

$$|\varphi_A\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle)|\uparrow\rangle \quad \longrightarrow \quad |\psi_A\rangle = U|\varphi_A\rangle = \frac{1}{\sqrt{2}}|+\rangle|\uparrow\rangle + \frac{1}{\sqrt{2}}|-\rangle|\downarrow\rangle \quad (14)$$

$$= \frac{1}{\sqrt{2}}|\odot\rangle|\rightarrow\rangle + \frac{1}{\sqrt{2}}|\otimes\rangle|\leftarrow\rangle \quad (15)$$

where

$$|\odot\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle), \quad |\otimes\rangle = \frac{1}{\sqrt{2}}(|+\rangle - |-\rangle) \quad (16)$$

are the eigenstates of S_x of the particle, and

$$|\rightarrow\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle), \quad |\leftarrow\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle). \quad (17)$$

are the eigenstates of S_x of the atom. By contrast with the Stern-Gerlach measurement, in this case the states $|\rightarrow\rangle$ and $|\leftarrow\rangle$, resulting from the combinations of the states $|\uparrow\rangle$ and $|\downarrow\rangle$, are possible states of the apparatus that measures S_x in the correlation (15). Nevertheless, as we have already pointed out, only one of the correlations makes physical sense, and only one set of outcomes is activated. In fact, the evolution operator $U = e^{i\pi|-\rangle\langle-| \otimes |\leftarrow\rangle\langle\leftarrow|}$ applied to the initial state $|\varphi_A\rangle$ produces the correlation given by eq. (14), and not that given by (15). Therefore, the outcomes that may be obtained as a result of the measurement are $|\uparrow\rangle$ and $|\downarrow\rangle$, and not $|\rightarrow\rangle$ and $|\leftarrow\rangle$: this fact guarantees that the observable S_z of the particle is measured.

Again, if we want to measure S_x , we have to change the apparatus: the new apparatus has to produce the correlation given by eq. (15), which can be achieved if it induce an evolution described by the operator $U' = e^{i\pi|\otimes\rangle\langle\otimes|\otimes|\downarrow\rangle\langle\downarrow|}$.

V. CONCLUSIONS

In his 1981 seminal work, Zurek speaks for the first time about the “*ambiguity in the choice of the preferred apparatus basis*” in measurement (see [9], p. 1516). According to him, it is the decoherence resulting from the interaction between the apparatus and the environment what determines “*in what mixture the wave function appears to have collapsed*” ([9], p. 1517). This argument has been repeatedly appealed to through the years (see [10], [11]), and reappears in Zurek’s more recent works ([8], [12]). Moreover, the “problem of the preferred basis” has been considered (with the “definite outcomes problem”) as one of the two central problems of quantum measurement. In his review paper on decoherence, Schlosshauer considers that, although in the literature the definite outcomes difficulty is typically referred to as ‘the measurement problem’, “*the preferred-basis problem is at least equally important, since it does not make sense even to inquire about specific outcomes if the set of possible outcomes is not clearly defined*” (see [5], p. 1270). On the basis of this assumption, the solution to preferred basis problem supplied by the decoherence program is usually viewed as one of its main theoretical advantages.

In this paper we have argued that this uncritically accepted problem is not a legitimate difficulty, but a pseudo-problem. The supposed basis ambiguity is the result of considering only a formal property of state vectors, and of forgetting the physical process of measurement. In fact, when the measurement is correctly understood as a physical process obeying the dynamical law of quantum mechanics, the supposed ambiguity vanishes since the preferred basis is determined by the measuring apparatus. Although the argumentation leading to this conclusion was simple and straightforward, it deserves to be seriously taken into account: its conceptual relevance derives from the widespread acceptance of the basis ambiguity as a legitimate problem in measurement.

APPENDIX A: NON-UNIQUENESS OF THE BIORTHONORMAL DECOMPOSITION

Let us consider the final correlated state of the measurement:

$$|\psi_A\rangle = \sum_i c_i |a_i\rangle |z_i\rangle \quad (\text{A1})$$

If we introduce the change of basis $|a_i\rangle = \sum_j M_{ji} |a'_j\rangle$, then $|\psi_A\rangle$ reads

$$\begin{aligned} |\psi_A\rangle &= \sum_i c_i \sum_j M_{ji} |a'_j\rangle |z_i\rangle \\ &= \sum_j |a'_j\rangle \sum_i c_i M_{ji} |z_i\rangle \\ &= \sum_j c'_j |a'_j\rangle |z'_j\rangle \end{aligned}$$

The last equality holds only when

$$c'_j |z'_j\rangle = \sum_i c_i M_{ji} |z_i\rangle \quad (\text{A2})$$

If we multiply eq. (A2) by $c_l^* \langle z'_l| = \sum_m c_m^* M_{lm}^* \langle z_m|$, we obtain

$$\begin{aligned} c_l^* c'_j \langle z'_l | z'_j \rangle &= \sum_m c_m^* M_{lm}^* \langle z_m | \sum_i c_i M_{ji} |z_i\rangle \\ c_l^* c'_j \delta_{lj} &= \sum_i |c_i|^2 M_{li}^* M_{ji} \\ |c_l|^2 \delta_{jl} &= \sum_i |c_i|^2 M_{ji} M_{il}^\dagger, \end{aligned}$$

This shows that decomposition is not unique if, $\forall i$, $|c_i| = C = |c'_i|$. In this case, $|z_i\rangle = \sum_j M_{ij}^\dagger |z'_j\rangle$.

-
- [1] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton University Press, Princeton (1955).
 - [2] P. Mittelstaedt, *The Interpretation of Quantum Mechanics and the Measurement Process*, Cambridge University Press, Cambridge (1998).
 - [3] P. Busch, P. Lathi and P. Mittelstaedt, *The Quantum Theory of Measurement*, Springer-Verlag, Berlin, (1996).

- [4] E. Schmidt, *Math. Ann.* 63 (1907), 433.
- [5] M. Schlosshauer, *Rev. Mod. Phys.* 76 (2004), 1267.
- [6] L. Ballentine, *Quantum Mechanics. A Modern Development*, World Scientific, Singapore (1998).
- [7] A. Peres, *Am. Jour., Phys.* 42 (1974), 886.
- [8] J. P. Paz and W. H. Zurek, “Environment-induced decoherence and the transition from quantum to classical.” In D. Heiss (ed.), *Fundamentals of Quantum Information, Lecture Notes in Physics*, Vol. 587, Springer, Heidelberg-Berlin (2002) (also in arXiv:quant-ph/0010011).
- [9] W. H. Zurek, *Phys. Rev. D* 24 (1981), 1516.
- [10] W. H. Zurek, *Phys. Rev. D* 26 (1982), 1862.
- [11] W. H. Zurek, *Physics Today* 44 (1991), 36.
- [12] W. H. Zurek, *Rev. Mod. Phys.* 75 (2003), 715.
- [13] This requirement relies on the fact that any difference in the experimental arrangement would allow us to distinguish between the two bases and, as a consequence, to know what observable is measured. For instance, in the Stern-Gerlach measurement, the rotation of the magnetic field from direction z to direction x would allow us to know that S_z was measured in the first case, and S_x was measured in the second case.
- [14] Otherwise, we would not distinguish between the kets $\{|z_i\rangle\}$ and the kets $\{|z'_i\rangle\}$ in quantum mechanics. Under the assumption of the preferred basis problem, they are distinguished by a non-identity change of basis.