

End-to-end entanglement in Bose-Hubbard chains

Jose Reslen* and Sougato Bose†

*Dept. of Physics and Astronomy, University College London,
Gower Street, WC1E 6BT London, United Kingdom*

(Dated: May 9, 2019)

We study the ground state as well as the dynamics of chains of bosons with local interaction and nearest-neighbour exchange using numerical techniques based on renormalization group methods. We explore the development of entanglement between the sites at the ends of such chains as mechanisms are invoked to concentrate population in these sites. We find that long-range entanglement arises in chains with strong repulsion constants in the intermediate sites, both in ground state and through non-equilibrium dynamics. Additionally, it is shown that entanglement between the end sites can be made to increase above their ground state values by adding a perturbation.

PACS numbers:

As constant developments in experimental physics increasingly expand our possibilities of materializing a new generation of computing engineering, the need for a better understanding of quantum phenomena becomes greater and more important. Outstanding emphasis has been put in the concept of entanglement as, for example, a resource for quantum information processing and in general as a tool to study the contrast between classical and quantum physics and the ways of utilizing novel aspects of the latter to boost the state-of-art technology. One fundamental problem that is now the subject of intensive research is the emergence of quantum correlations between distant sites of a quantum system. This can be achieved both through statics [1] (for example, in the ground state of a system) or through non-equilibrium dynamics [2, 3, 4, 5, 6, 7, 8, 9]. Most of the work done so far in this field has been carried out in light-matter systems and spin chains, but more recently, the development on new numerical techniques and algorithms [10] has opened the possibility of addressing more complex systems such as chains of interacting bosons [5, 11, 12]. Usually, where bosonic chains have been studied, though true quantum correlations have indeed been found to emerge between distant individual sites, they are generically not “substantial” in magnitude [2]. Thus substantial entanglement between distant sites of a Bose-Hubbard model, even if with some reasonable engineering of the Hamiltonian, should be of great interest.

Chains of bosons are more often than not well described by the Bose-Hubbard model [13]. In this model bosons can hop between neighbour sites while undergoing local repulsion when several bosons stand in the same place. The transition from a superfluid to a Mott insulator has been verified in experiments of cold atoms in optical lattices [14, 15], and also the observation of second-order effects in the strong interacting regime has been reported in similar set-ups [16].




FIG. 1: Potential realization of a Bose-Hubbard chain in an optical lattice. The above finite lattice can be realized in one cell of a superlattice. The end sites of the lattice are depicted differently in the figure as we will require them to have vanishing repulsion in our study.

Motivated by these advancements in experimental physics, in this letter we analyse, from a fundamental point of view, the emergence of long-range quantum correlations in Bose-Hubbard chains (fig. 1). There has been one previous investigation on long range entanglement in a Bose-Hubbard chain where the dynamics is effectively reducible to a single particle propagation in a lattice [5]. Our study is, however, irreducibly a many-particle situation and can even result in more entanglement than what a single particle hopping in a lattice can ever do. It is also pertinent to point out that there has been much interesting work on the various forms of entanglement contained in Bose gases [17, 18], but here we concentrate on a specific type of entanglement, namely that between two individual sites at the ends. We aim to identify suitable conditions within the Bose-Hubbard Hamiltonian for the enhancement of entanglement as well as potential implications of such entanglement on both the ground state physics and the dynamics.

We assume a chain of size N with M bosons, nearest-neighbour hopping and open ended boundary conditions governed by a Bose-Hubbard Hamiltonian with variable coefficients,

$$\hat{H} = \sum_{k=1}^N V_k \hat{a}_k^\dagger \hat{a}_k (\hat{a}_k^\dagger \hat{a}_k - 1) - \sum_{k=1}^{N-1} J_k (\hat{a}_{k+1}^\dagger \hat{a}_k + \hat{a}_k^\dagger \hat{a}_{k+1}). \quad (1)$$

*Electronic address: reslen@theory.phys.ucl.ac.uk

†Electronic address: sougato@theory.phys.ucl.ac.uk

Constants V_k and J_k account for the on-site repulsion and hopping respectively while the annihilation and creation operators \hat{a}_k^\dagger and \hat{a}_k obey the usual commuting rules $[\hat{a}_k^\dagger, \hat{a}_l] = \delta_k^l$ and $[\hat{a}_k^\dagger, \hat{a}_l^\dagger] = [\hat{a}_k, \hat{a}_l] = 0$. Physical realizations of this Hamiltonian include experiments in optical lattices and magnetic traps where remarkable control of the ratio V/J can be achieved [14, 15, 16].

From the Heisenberg equations of motion we obtain

$$\frac{d\hat{\alpha}_k}{dt} = ie^{-it\hat{H}}[\hat{a}_k^\dagger, \hat{H}]e^{it\hat{H}}, \quad (2)$$

where $\hat{\alpha}_k = e^{-it\hat{H}}\hat{a}_k^\dagger e^{it\hat{H}}$. In the general case nonlinearities keep us from getting reliable expressions for the $\hat{\alpha}_k$ in terms of time. One fortunate instance is the repulsionless case where all V_k go to zero and we are left with

$$\frac{d}{dt} \begin{pmatrix} \hat{\alpha}_1 \\ \hat{\alpha}_2 \\ \hat{\alpha}_3 \\ \vdots \\ \hat{\alpha}_N \end{pmatrix} = -i \begin{pmatrix} 0 & J_1 & 0 & \cdots & 0 \\ J_1 & 0 & J_2 & \cdots & 0 \\ 0 & J_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & J_{N-1} \\ 0 & 0 & 0 & J_{N-1} & 0 \end{pmatrix} \begin{pmatrix} \hat{\alpha}_1 \\ \hat{\alpha}_2 \\ \hat{\alpha}_3 \\ \vdots \\ \hat{\alpha}_N \end{pmatrix}, \quad (3)$$

which resembles a Schroedinger equation with state vectors swapped for vectors of operators. Choosing a non-uniform hopping distribution (the reasons for which will be pointed out later) that matches an angular momentum representation of length $j = \frac{N-1}{2}$, namely $J_k = \frac{\lambda}{2}\sqrt{k(N-k)}$. The above couplings, when present in spin chains, are known to facilitate a perfect quantum state transfer [19], which has been developed in context of the idea of using spin chains to convey quantum states [20]. In a Hubbard model, the above coupling profile will transfer particles perfectly from one end of the chain to the other. In this case every \hat{a}_k can be mapped into an element of a basis that is being rotated with period $2\pi/\lambda$ by the action of a transverse angular momentum in such a way that any state is reflected with respect to the centre of the chain every π/λ units of time. The ground state of the chain can be obtained from the ground eigenvector of matrix (3) following an imaginary-time evolution analysis. Moreover, such eigenvector can be found by rotating an analogous state, namely, $|j, -j\rangle$. Thus, taking $M = N$ and up to a normalization coefficient the general ground state is in this way found to be

$$\left(\sum_{m=-j}^j \frac{(-1)^{j+m}}{2^j} \sqrt{\frac{(2j)!}{(j-m)!(j+m)!}} \hat{a}_{m+j+1}^\dagger \right)^N \prod_{l=1}^N |0l\rangle, \quad (4)$$

and similarly the ground state energy $E_G = -\lambda jN$ can be picked up out of the most general expression that leads to the ground state.

For analysing the more realistic case where repulsive terms are included, we will make use of renormalization-

based techniques as presented in [10], also known as *TEBD*, combined with conservation properties and symmetry considerations associated with the problem. Briefly, *TEBD* consists in writing the state in terms of canonical coefficients that are closely related to Schmidt vectors. These coefficients are updated through successive applications of semi-local unitary operations that correspondingly only modify coefficients associated with the sub-spaces on which they are applied. This fact is then skilfully exploited to work out an efficient way of simulating both real and complex evolution. We empower the method by introducing an extra set of integer coefficients, $\zeta_\alpha^{[n]}$, dimensionally identical to the set of $\lambda_\alpha^{[n]}$ in [10], which register the number of particles associated to each Schmidt vector. Following Vidal's notation the state of the system with explicit reference to two arbitrary neighbour sites can be written as

$$|\psi\rangle = \sum_{\alpha=1}^{\chi_A} \sum_{\gamma=1}^{\chi_F} \sum_{i=0}^M \sum_{l=0}^M \delta_{\zeta_\gamma+j}^{M-\zeta_\alpha-i} \Theta_{\alpha\gamma}^{ij} |\alpha i j \gamma\rangle, \quad (5)$$

where the inclusion of the delta function expressly accounts for the association of vectors with complementary number of particles. Although this association is implicit in $\Theta_{\alpha\gamma}^{ij}$, numerical errors cause small deviations that persistently corrupt canonical coefficients of chains with somehow high or moderate number of bosons compared to the number of sites. We tackle this problem by grouping states $|\alpha i\rangle$ and $|j \gamma\rangle$ bearing the same $\zeta_\alpha + i$ and $\zeta_\gamma + j$ respectively. Vidal methodology is then applied systematically to couples of mutually complementary families of vectors rather than to the whole spectrum of states [12]. Storage space is dynamically supplied according to the actual number of canonical coefficients developed in the quantum state at run time, making possible to obtain results without truncating many-boson states. Here we limit ourselves to Hamiltonians with symmetric distributions of parameters with respect to the centre of the chain and with an even number of sites and bosons. Correspondingly, the system can be genuinely simulated by focusing on the canonical coefficients of half of the chain plus the interaction between complementary half-chain blocks. The speed of the simulation is determined by χ , the dimension of the biggest $\lambda_\alpha^{[n]}$ and also a rough estimation of the amount of entanglement in the system. Hence, simulations in highly-entangled chains can be performed on systems of moderate size. The algorithm is applied recursively to simulate

$$|\psi_G\rangle = \lim_{\tau \rightarrow \infty} \frac{e^{-\tau\hat{H}}|\psi_0\rangle}{\|e^{-\tau\hat{H}}|\psi_0\rangle\|}, \quad (6)$$

until convergence to ground state $|\psi_G\rangle$ is accomplished. For simplicity we assumed $\lambda = 2$. Ground state simulations in repulsionless chains agree well with the exact solution given by Eq. (4) (In fact, this is one of the




FIG. 2: Ground state entanglement between the ends of a chain of eight places and eight bosons against repulsion. PTH and repulsionless ends are assumed everywhere. Fitting $y = 0.91 - 0.34e^{-0.07x}$, where y stands for the logarithmic negativity (entanglement), while x stands for repulsion. Inset: Particle distribution profile and fluctuations. (—) No repulsion at all. (---) $V_{2-7} = 50$.




FIG. 3: Dynamic entanglement between the ends of a boson chain with uniform unity hopping. Bosons are initially arranged in a Mott insulating state with one boson per site in a chain of eight places.

reasons that we chose PTH and analytically derived Eq. (4), so that we can verify the correctness of our numerical methodology through this explicit case).

In order to calculate the entanglement between the ends of the chain, we need to get the corresponding reduced density matrix. In terms of computational work, the number of basic operations involved in calculating such density matrix roughly scale as $\mathcal{O}(\chi^4)$ compared to the $\mathcal{O}(\chi^3)$ operations needed to update the coefficients. This is a considerable effort here even for 8 sites and 8 bosons as we need to retain a high χ for preserving high end to end entanglement. Once this density matrix is obtained, we have to quantify the entanglement using a

measure which is appropriate for mixed states of two arbitrary dimensional quantum systems. So we choose the *logarithmic negativity* as defined in Ref.[21].

From our initial simulations we observe that ground state entanglement is highly sensitive to the particle distribution profile along the chain. This is due to the effect of tracing out intermediate sites in order to get reduced density matrices. When population is accumulated in the middle of the chain we must trace out states with a considerable number of particles when obtaining the mixed state of two distant sites at the borders. Lower population at the ends implies a lower number of possibilities for distinct particle number states available at the ends and consequently a lower ability to correlate particle numbers between these sites and entangle them. Thus for repulsionless case and PTH, where population accumulates primarily in the middle of the chain (which is shown in the inset of fig. 2), the entanglement is vanishingly small. The choice of PTH, as opposed to a uniform J_k , thus clearly illustrates the negative effect of population concentration in the middle of the chain. To get a higher entanglement, we then proceed to deliberately frustrate boson accumulation in the central part of the chain by evenly increasing the repulsion in every site but the ends which in turn are left repulsionless (fig. 1). This approach allows us to look closely at the physical mechanisms underlying the interplay between hopping and repulsion and its effects on quantum correlations at long distances. In the inset of fig. 2 we can see how bosons in intermediate positions condensate in the lowest levels of occupation creating a fine arrangement of superfluid particles that connect the ends, which in turn become highly populated. Fig. 2 also depicts the behaviour of log negativity as a function of repulsion. It can be seen that entanglement grows monotonously, undergoing saturation for large repulsion, suggesting that maximum entanglement is achieved in the case of hardcore bosons. In this limit, bosons find difficult to stay at the intermediate positions as the hopping of particles from neighbour sites would give rise to multi-occupied states with strong repulsion. Instead, bosons organize in superposition states that on average contain only half a boson per site for the intermediate sites, creating a globally delocalized profile that favours long-range quantum correlations. Similarly, analogous plots corresponding to averages of number of particles also display saturation for high repulsion, establishing a clear signature of rescaling behaviour, that is, bosons reorganize under the influence of larger repulsion constants in such a way that the phenomenology of the system is altered, roughly, just by a proportionality factor. Similar conclusions can be drawn for chains with constant coefficients J_k and repulsionless ends, although PTH chains are more efficient at enhancing entanglement. This is, in fact, our principal reason for assuming PTH. For instance, in a chain of ten sites, ten bosons, PTH and high intermediate repulsion $V_{2-9} = 50$, log negativity was found to be 0.83 against 0.60 of the same chain with uniform unity hopping.

We also applied the routines developed for the ground state to find the dynamical behaviour of entanglement in a chain initially prepared in a Mott insulating state. This is the most natural scenario one can expect in a Bose-Hubbard chain where one first prepares a Mott insulating state with a boson per site and then suddenly turns off all the interactions globally. Here, we assume uniform hopping (as in this scenario we found no particular advantage of PTH) and evolve the system setting the repulsion to zero. End-to-end entanglement shows very little development in the course of evolution as can be seen in fig. 3. As in the case of ground state entanglement, dynamic entanglement does not build up unless a significant accumulation of particles assists the enhancement of fluctuations at the ends. Consequently, chains with repulsive terms in intermediate sites display a better register of entanglement than chains with no repulsion. Dynamics in repulsionless chains is, in fact, known to lead to a progressive thermalization of reduced density matrices of each site [22]. Two site entanglement is thus very low in this repulsionless scenario. An evolution with high repulsion in the intermediate sites, on the other hand, shows a less regular profile, product of interference among many unsynchronized phases accumulated locally. In the best cases, the amount of entanglement contained in the system after reasonable times is important (and this has been shown in Fig. 3), yet smaller than what can be obtained from the ground state in optimal configurations.

We would now ideally like to generate such an entanglement between the ends where the logarithmic negativity exceeds unity (the maximum value in a two qubit state) so that the possibilities of multiple occupation numbers of the ending sites is utilized highly beneficially. Looking at potential alternatives to the schemes discussed before, we now assume that a chain initially prepared in the ground state of a Hamiltonian with high intermediate repulsion is set to evolve under the action of a small perturbation potential corresponding to a physical mechanism that internally pumps bosons in and out the ends. The quantum state at any time t can be written as

$$|\psi(t)\rangle = e^{-it(\hat{H} + \Delta \sum_{j=1}^N \hat{h}_j)} |\psi_g\rangle, \quad (7)$$

where $|\psi_g\rangle$ is the ground state of \hat{H} , Δ is a small number and \hat{h}_j represents a local operator acting on site j . For consistency, we require $[\hat{H}_g, \sum_{i=1}^N \hat{h}_i] \neq 0$. Here we consider local operators of the form $\hat{h}_j = h_j \hat{a}_j^\dagger \hat{a}_j$ for every site but the ends. It is also practical to take $h_j = k$, where k is the integer distance between site j and the closest end. In order to determine the form of \hat{h} on the ends we require, first, that \hat{h} itself can be written as a function of the number of particles, and second, that to first order in Δ matter can be exchanged between the ends and the rest of the chain without unbalancing the perturbation energy. From the latter we get




FIG. 4: Dynamical entanglement between the ends of a chain of eight places and eight bosons for different values of Δ . Equivalent graphs corresponding to population averages deviate less than 2% from their initial values.

$$h(n_0 + \frac{1}{2}(\frac{N}{2} - k) + \frac{1}{2}) - h(n_0 + \frac{1}{2}(\frac{N}{2} - k)) = \frac{\Delta k}{2}, \quad (8)$$

where we have supposed an initial occupation of n_0 on each end and half a boson in the rest of the chain. For simplicity N is considered to be even. Inserting $h(x) = c_2 x^2 + c_1 x$ in eq. (8) and solving for c_1 and c_2 we find $c_1 = (\frac{N-1}{2} + 2n_0)\Delta$ and $c_2 = -\Delta$. Fig. 4 shows log-negativity as a function of time for a Hamiltonian with PTH, $V_{2-7} = 50$ and different perturbation intensities. Roundoff errors accumulated in the ground state during the imaginary time evolution cause slow oscillations in the dynamics at $\Delta = 0$, but these are tiny compared with the more complex evolution seen at finite Δ . In the initial stages of evolution the dynamics is characterized by a sharp increase of quantum fluctuations on the ends of the chain accompanied by little change in the average number of bosons. Hence, entanglement development is enhanced while avoiding massive migration of bosons towards the centre of the chain. This effect is maximal at around $\Delta = 0.1$ and becomes less important for higher values of Δ , where perturbative terms induce more irregular dynamics. Significantly, entanglement generation is improved not by adding interaction terms to the Hamiltonian but by dynamically unfolding quantum correlations already contained in the ground state.

In summary, we have studied the ground state entanglement as well as the dynamical entanglement displayed by Bose-Hubbard chains for several configurations of parameters. Our results indicate that chains with strong repulsion coefficients in intermediate places are suitable scenarios for the emergence of long-range quantum correlations. In addition, we showed that ground state entanglement can be improved through perturbative dynamics.

JR acknowledges an EPSRC-DHPA scholarship. We thank T.S. Monteiro and C. Hadley for very helpful com-

ments.

-
- [1] L. Campos Venuti, *et. al.*, Phys. Rev. Lett. **96**, 247206 (2006).
- [2] J. Eisert, M. B. Plenio, S. Bose and J. Hartley **93**, 190402 (2004).
- [3] M.-H. Yung and S. Bose, Phys. Rev. A **71**, 032310 (2005).
- [4] D.I. Tsomokos, M.J. Hartmann, S.F. Huelga and M.B. Plenio New Journal of Physics **9**, 79 (2007).
- [5] O. Romero-Isart, K. Eckert, C. Rodo and A. Sanpera, J. Phys. A: Math. Theor. **40** 8019 (2007).
- [6] I. D'Amico, B. W. Lovett and T. P. Spiller, Phys. Rev. A **76**, 030302(R) (2007).
- [7] H. Wichterich, S. Bose, arXiv: 0806.4568.
- [8] F. Galve, *et. al.*, arXiv:0809.3712.
- [9] P. Sodano, A. Bayat and S. Bose, arXiv:0811.2677
- [10] G. Vidal, Phys. Rev. Lett. **91**, 147902 (2003); G. Vidal, Phys. Rev. Lett. **93**, 040502 (2004); G. Vidal, Phys. Rev. Lett. **98**, 070201 (2007).
- [11] M. Moeckel and S. Kehrein, Phys. Rev. Lett. **100**, 175702 (2008); S. Gu *et al*, Phys. Rev. Lett. **93**, 086402 (2004); L. Tian *et al*, quant-ph/07052023.
- [12] A.J. Daley *et al*, Phys. Rev. A **72**, 043618 (2005); I. Danshita and P. Naidon, cond-mat/08091486.
- [13] M.P.A. Fisher *et al*, Phys. Rev. B **40**, 546 (1989).
- [14] D. Jaksch *et al*, Phys. Rev. Lett. **81**, 3108 (1998).
- [15] M. Greiner *et al*, Nature **415**, 39 (2002).
- [16] S. Fölling *et al*, Nature **448**, 1029 (2007).
- [17] C. Simon, Phys. Rev. A **66**, 052323 (2002).
- [18] L. Heaney, J. Anders, D. Kaszlikowski and V. Vedral, Phys. Rev. A, **76** 053605 (2007); L. Heaney and J. Anders, arXiv:0810.2882.
- [19] M. Christandl *et al*, Phys. Rev. Lett. **92**, 187902 (2004).
- [20] S. Bose, Phys. Rev. Lett. **91**, 207901 (2003).
- [21] G. Vidal and R. Werner, Phys. Rev. A **65**, 032314 (2002).
- [22] M. Cramer, C.M. Dawson, J. Eisert, T.J. Osborne, Phys. Rev. Lett. **100**, 030602 (2008); A. Flesch *et al*, cond-mat/08083779.