

**EXTENSIONS OF RAYNAUD SCHEMES
WITH TRIVIAL GENERIC FIBERS**

NAOKI IMAI

ABSTRACT. We calculate the number of extensions of Raynaud schemes over the ring of integers of an absolutely ramified p -adic field with the trivial generic fibers by counting the rational points of the moduli space of finite flat models.

INTRODUCTION

Let K be a totally ramified extension of degree e over \mathbb{Q}_p for $p > 2$, and \mathbb{F} be a finite field of characteristic p . We consider the constant group scheme $C_{\mathbb{F}}$ over $\text{Spec } K$ of the two-dimensional vector space over \mathbb{F} . Let $M(C_{\mathbb{F}}, K)$ be the isomorphism class of the pairs $(A, C_{\mathbb{F}} \xrightarrow{\sim} A_K)$ such that A is a finite flat group scheme over \mathcal{O}_K with a structure of an \mathbb{F} -vector space. Here A_K is the generic fiber of A . A finite flat group scheme over \mathcal{O}_K with a structure of one-dimensional vector space over a finite field is called a Raynaud scheme. So A is an extension of Raynaud schemes by [Tat, Proposition 4.2.1(a)]. If $e < p - 1$, then $M(C_{\mathbb{F}}, K)$ is one-point set by [Ray, Theorem 3.3.3]. However, if the ramification is big, there are surprisingly many finite flat models. In this paper, we calculate $|M(C_{\mathbb{F}}, K)|$, that is, the number of finite flat models of $C_{\mathbb{F}}$. The main theorem is the following.

Theorem. *Let q be the cardinality of \mathbb{F} . Then we have*

$$|M(C_{\mathbb{F}}, K)| = \sum_{n \geq 0} (a_n + a'_n) q^n.$$

Here a_n and a'_n are defined as in the following.

We express e and n by

$$e = (p-1)e_0 + e_1, \quad n = (p-1)n_0 + n_1 = (p-1)n'_0 + n'_1 + e_1$$

such that $e_0, n_0, n'_0 \in \mathbb{Z}$ and $0 \leq e_1, n_1, n'_1 \leq p-2$. Then

$$\begin{aligned} a_n &= \max\{e_0 - (p+1)n_0 - n_1 - 1, 0\} && \text{if } n_1 \neq 0, 1, \\ a_n &= \max\{e_0 - (p+1)n_0 - n_1 - 1, 0\} \\ &\quad + \max\{e_0 - (p+1)n_0 - n_1 + 1, 0\} && \text{if } n_1 = 0, 1, \end{aligned}$$

and

$$\begin{aligned} a'_n &= \max\{e_0 - e_1 - (p+1)n'_0 - n'_1 - 2, 0\} && \text{if } n'_1 \neq 0, 1, \\ a'_n &= \max\{e_0 - e_1 - (p+1)n'_0 - n'_1 - 2, 0\} \\ &\quad + \max\{e_0 - e_1 - (p+1)n'_0 - n'_1, 0\} && \text{if } n'_1 = 0, 1 \end{aligned}$$

except in the case where $n = 0$ and $e_1 = p - 2$, in which case we put $a'_0 = e_0$.

In the above theorem, we can easily check that $|M(C_{\mathbb{F}}, K)| = 1$ if $e < p - 1$.

Acknowledgment. The author is grateful to his advisor Takeshi Saito for his careful reading of an earlier version of this paper and for his helpful comments.

Notation. Throughout this paper, we use the following notation. Let $p > 2$ be a prime number, and K be a totally ramified extension of \mathbb{Q}_p of degree e . The ring of integers of K is denoted by \mathcal{O}_K , and the absolute Galois group of K is denoted by G_K . Let \mathbb{F} be a finite field of characteristic p . The formal power series ring of u over \mathbb{F} is denoted by $\mathbb{F}[[u]]$, and its quotient field is denoted by $\mathbb{F}((u))$. Let v_u be the valuation of $\mathbb{F}((u))$ normalized by $v_u(u) = 1$, and we put $v_u(0) = \infty$. For $x \in \mathbb{R}$, the greatest integer less than or equal to x is denoted by $[x]$.

1. PRELIMINARIES

To calculate the number of finite flat models of $C_{\mathbb{F}}$, we use the moduli spaces of finite flat models constructed by Kisin in [Kis].

Let $V_{\mathbb{F}}$ be the two-dimensional trivial representation of G_K over \mathbb{F} . The moduli space of finite flat models of $V_{\mathbb{F}}$, which is denoted by $\mathcal{GR}_{V_{\mathbb{F}},0}$, is a projective scheme over \mathbb{F} . An important property of $\mathcal{GR}_{V_{\mathbb{F}},0}$ is the following Proposition.

Proposition 1.1. *For any finite extension \mathbb{F}' of \mathbb{F} , there is a natural bijection between the set of isomorphism classes of finite flat models of $V_{\mathbb{F}'} = V_{\mathbb{F}} \otimes_{\mathbb{F}} \mathbb{F}'$ and $\mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}')$.*

Proof. This is [Kis, Corollary 2.1.13]. \square

By Proposition 1.1, to calculate the number of finite flat models, it suffices to count the number of the \mathbb{F} -rational points of $\mathcal{GR}_{V_{\mathbb{F}},0}$.

Let $\mathfrak{S} = \mathbb{Z}_p[[u]]$, and $\mathcal{O}_{\mathcal{E}}$ be the p -adic completion of $\mathfrak{S}[1/u]$. There is an action of ϕ on $\mathcal{O}_{\mathcal{E}}$ determined by identity on \mathbb{Z}_p and $u \mapsto u^p$. We choose elements $\pi_m \in \overline{K}$ such that $\pi_0 = \pi$ and $\pi_{m+1}^p = \pi_m$ for $m \geq 0$, and put $K_{\infty} = \bigcup_{m \geq 0} K(\pi_m)$. Let $\Phi M_{\mathcal{O}_{\mathcal{E}},\mathbb{F}}$ be the category of finite $\mathcal{O}_{\mathcal{E}} \otimes_{\mathbb{Z}_p} \mathbb{F}$ -modules M equipped with ϕ -semi-linear map $M \rightarrow M$ such that the induced $\mathcal{O}_{\mathcal{E}} \otimes_{\mathbb{Z}_p} \mathbb{F}$ -linear map $\phi^*(M) \rightarrow M$ is an isomorphism. We take the ϕ -module $M_{\mathbb{F}} \in \Phi M_{\mathcal{O}_{\mathcal{E}},\mathbb{F}}$ that corresponds to the $G_{K_{\infty}}$ -representation $V_{\mathbb{F}}(-1)$. Here (-1) denotes the inverse of Tate twist.

The moduli space $\mathcal{GR}_{V_{\mathbb{F}},0}$ is described via the Kisin modules as in the following.

Proposition 1.2. *For any finite extension \mathbb{F}' of \mathbb{F} , the elements of $\mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}')$ naturally correspond to free $\mathbb{F}'[[u]]$ -submodules $\mathfrak{M}_{\mathbb{F}'} \subset M_{\mathbb{F}} \otimes_{\mathbb{F}} \mathbb{F}'$ of rank 2 that satisfy $u^e \mathfrak{M}_{\mathbb{F}'} \subset (1 \otimes \phi)(\phi^*(\mathfrak{M}_{\mathbb{F}'})) \subset \mathfrak{M}_{\mathbb{F}'}$.*

Proof. This follows from the construction of $\mathcal{GR}_{V_{\mathbb{F}},0}$ in [Kis, Corollary 2.1.13]. \square

By Proposition 1.2, we often identify a point of $\mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}')$ with the corresponding finite free $\mathbb{F}'[[u]]$ -module.

For $A \in GL_2(\mathbb{F}((u)))$, we write $M_{\mathbb{F}} \sim A$ if there is a basis $\{e_1, e_2\}$ of $M_{\mathbb{F}}$ over $\mathbb{F}((u))$ such that $\phi \begin{pmatrix} e_1 \\ e_2 \end{pmatrix} = A \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}$. We use the same notation for any sublattice $\mathfrak{M}_{\mathbb{F}} \subset M_{\mathbb{F}}$ similarly.

Finally, for any sublattice $\mathfrak{M}_{\mathbb{F}} \subset M_{\mathbb{F}}$ with a chosen basis $\{e_1, e_2\}$ and $B \in GL_2(\mathbb{F}((u)))$, the module generated by the entries of $\left\langle B \begin{pmatrix} e_1 \\ e_2 \end{pmatrix} \right\rangle$ with the basis given by these entries is denoted by $B \cdot \mathfrak{M}_{\mathbb{F}}$. Note that $B \cdot \mathfrak{M}_{\mathbb{F}}$ depends on the

choice of the basis of $\mathfrak{M}_{\mathbb{F}}$. We can see that if $\mathfrak{M}_{\mathbb{F}} \sim A$ for $A \in GL_2(\mathbb{F}((u)))$ with respect to a given basis, then we have

$$B \cdot \mathfrak{M}_{\mathbb{F}} \sim \phi(B)AB^{-1}$$

with respect to the induced basis.

Lemma 1.3. *Suppose \mathbb{F}' is a finite extension of \mathbb{F} , and $x \in \mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}')$ corresponds to $\mathfrak{M}_{\mathbb{F}'}$. Put $\mathfrak{M}_{\mathbb{F}',i} = \begin{pmatrix} u^{s_i} & v_i \\ 0 & u^{t_i} \end{pmatrix} \cdot \mathfrak{M}_{\mathbb{F}'}$ for $1 \leq i \leq 2$, $s_i, t_i \in \mathbb{Z}$ and $v_i \in \mathbb{F}((u))$. Assume $\mathfrak{M}_{\mathbb{F}',1}$ and $\mathfrak{M}_{\mathbb{F}',2}$ correspond to $x_1, x_2 \in \mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}')$ respectively. Then $x_1 = x_2$ if and only if*

$$s_1 = s_2, \quad t_1 = t_2 \quad \text{and} \quad v_1 - v_2 \in u^{t_1}\mathbb{F}'[[u]].$$

Proof. The equality $x_1 = x_2$ is equivalent to the existence of $B \in GL_2(\mathbb{F}'[[u]])$ such that

$$B \begin{pmatrix} u^{s_1} & v_1 \\ 0 & u^{t_1} \end{pmatrix} = \begin{pmatrix} u^{s_2} & v_2 \\ 0 & u^{t_2} \end{pmatrix}.$$

It is further equivalent to the condition that

$$\begin{pmatrix} u^{s_2-s_1} & v_2 u^{-t_1} - u^{s_2-s_1-t_1} v_1 \\ 0 & u^{t_2-t_1} \end{pmatrix} \in GL_2(\mathbb{F}'[[u]]).$$

The last condition is equivalent to the desired condition. \square

2. MAIN THEOREM

Theorem 2.1. *Let q be the cardinality of \mathbb{F} . Then we have*

$$|M(C_{\mathbb{F}}, K)| = \sum_{n \geq 0} (a_n + a'_n) q^n.$$

Here a_n and a'_n are defined as in the introduction.

Proof. Since $V_{\mathbb{F}}$ is the trivial representation, $M_{\mathbb{F}} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for some basis. Let $\mathfrak{M}_{\mathbb{F},0}$ be the lattice of $M_{\mathbb{F}}$ generated by the basis giving $M_{\mathbb{F}} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. By the Iwasawa decomposition, any sublattice of $M_{\mathbb{F}}$ can be written as $\begin{pmatrix} u^s & v \\ 0 & u^t \end{pmatrix} \cdot \mathfrak{M}_{\mathbb{F},0}$ for $s, t \in \mathbb{Z}$ and $v \in \mathbb{F}((u))$. We put

$$\mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F}) = \left\{ \begin{pmatrix} u^s & v \\ 0 & u^t \end{pmatrix} \cdot \mathfrak{M}_{\mathbb{F},0} \in \mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}) \mid v \in \mathbb{F}((u)) \right\}.$$

Then

$$\mathcal{GR}_{V_{\mathbb{F}},0}(\mathbb{F}) = \bigcup_{s,t \in \mathbb{Z}} \mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F})$$

and this is a disjoint union by Lemma 1.3.

We put

$$\mathfrak{M}_{\mathbb{F},s,t} = \begin{pmatrix} u^s & 0 \\ 0 & u^t \end{pmatrix} \cdot \mathfrak{M}_{\mathbb{F},0}.$$

Then we have $\mathfrak{M}_{\mathbb{F},s,t} \sim \begin{pmatrix} u^{(p-1)s} & 0 \\ 0 & u^{(p-1)t} \end{pmatrix}$ with respect to the basis induced from $\mathfrak{M}_{\mathbb{F},0}$. The lattice $\mathfrak{M}_{\mathbb{F}}$ corresponding to a point of $\mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F})$ is written as $\begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \cdot \mathfrak{M}_{\mathbb{F},s,t}$ for some v in $\mathbb{F}((u))$. Then we have

$$\mathfrak{M}_{\mathbb{F}} \sim \begin{pmatrix} u^{(p-1)s} & -vu^{(p-1)s} + \phi(v)u^{(p-1)t} \\ 0 & u^{(p-1)t} \end{pmatrix}$$

with respect to the induced basis. The condition $u^e \mathfrak{M}_{\mathbb{F}} \subset (1 \otimes \phi)(\phi^*(\mathfrak{M}_{\mathbb{F}})) \subset \mathfrak{M}_{\mathbb{F}}$ is equivalent to the following:

$$0 \leq (p-1)s \leq e, \quad 0 \leq (p-1)t \leq e, \\ v_u(vu^{(p-1)s} - \phi(v)u^{(p-1)t}) \geq \max\{0, (p-1)(s+t) - e\}.$$

Conversely, $s, t \in \mathbb{Z}$ and $v \in \mathbb{F}((u))$ satisfying this condition gives a point of $\mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F})$ as $\begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \cdot \mathfrak{M}_{\mathbb{F},s,t}$. We put $r = -v_u(v)$.

We fix $s, t \in \mathbb{Z}$ such that $0 \leq s, t \leq e_0$. The lowest degree term of $vu^{(p-1)s}$ is equal to that of $\phi(v)u^{(p-1)t}$ if and only if $v_u(v) = s-t$, in which case $v_u(vu^{(p-1)s}) = ps-t$.

In the case where $ps-t \geq \max\{0, (p-1)(s+t) - e\}$, the condition $v_u(vu^{(p-1)s} - \phi(v)u^{(p-1)t}) \geq \max\{0, (p-1)(s+t) - e\}$ is equivalent to

$$\min\{v_u(vu^{(p-1)s}), v_u(\phi(v)u^{(p-1)t})\} \geq \max\{0, (p-1)(s+t) - e\},$$

and further equivalent to

$$r \leq \min\left\{(p-1)s, \frac{e - (p-1)s}{p}, e - (p-1)t, \frac{(p-1)t}{p}\right\}.$$

We put

$$r_{s,t} = \min\left\{(p-1)s, \left\lceil \frac{e - (p-1)s}{p} \right\rceil, e - (p-1)t, \left\lceil \frac{(p-1)t}{p} \right\rceil\right\}.$$

In this case, the number of the points of $\mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F})$ is equal to $q^{r_{s,t}}$ by Lemma 1.3.

In the case where $ps-t < \max\{0, (p-1)(s+t) - e\}$, the condition $v_u(vu^{(p-1)s} - \phi(v)u^{(p-1)t}) \geq \max\{0, (p-1)(s+t) - e\}$ is satisfied if and only if

$$v = \alpha u^{s-t} + v_+ \text{ for } \alpha \in \mathbb{F} \text{ and } v_+ \in \mathbb{F}((u)) \text{ such that } -v_u(v_+) \leq r_{s,t}.$$

In this case, the number of the points of $\mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F})$ is equal to $q^{r_{s,t}+1}$ by Lemma 1.3.

We put $h_{s,t} = \log_q |\mathcal{GR}_{V_{\mathbb{F}},0,s,t}(\mathbb{F})|$. Collecting the above results, we get the followings:

- If $s+t \leq e_0$ and $ps-t \geq 0$, then $h_{s,t} = [(p-1)t/p]$.
- If $s+t \leq e_0$ and $ps-t < 0$, then $h_{s,t} = (p-1)s + 1$.
- If $s+t > e_0$ and $ps-t \geq (p-1)(s+t) - e$, then $h_{s,t} = [(e - (p-1)s)/p]$.
- If $s+t > e_0$ and $ps-t < (p-1)(s+t) - e$, then $h_{s,t} = e - (p-1)t + 1$.

Now we have

$$|M(C_{\mathbb{F}}, K)| = \sum_{0 \leq s, t \leq e_0} q^{h_{s,t}}.$$

We put

$$S_n = \{(s, t) \in \mathbb{Z}^2 \mid 0 \leq s, t \leq e_0, h_{s,t} = n\},$$

and

$$\begin{aligned} S_{n,1} &= \{(s, t) \in S_n \mid s + t \leq e_0, ps - t \geq 0\}, \\ S_{n,2} &= \{(s, t) \in S_n \mid s + t \leq e_0, ps - t < 0\}, \\ S'_{n,1} &= \{(s, t) \in S_n \mid s + t > e_0, ps - t \geq (p-1)(s+t) - e\}, \\ S'_{n,2} &= \{(s, t) \in S_n \mid s + t > e_0, ps - t < (p-1)(s+t) - e\}. \end{aligned}$$

It suffices to show that $|S_{n,1}| + |S_{n,2}| = a_n$ and $|S'_{n,1}| + |S'_{n,2}| = a'_n$.

Firstly, we calculate $|S_{n,1}|$. We assume $(s, t) \in S_{n,1}$. In the case $n_1 \neq 0$, we have $t = pn_0 + n_1 + 1$ by $[(p-1)t/p] = (p-1)n_0 + n_1$. Then $ps \geq t = pn_0 + n_1 + 1$ implies $s \geq n_0 + 1$, and we have

$$n_0 + 1 \leq s \leq e_0 - pn_0 - n_1 - 1.$$

We note that if $t > e_0$, we have

$$(e_0 - pn_0 - n_1 - 1) - (n_0 + 1) + 1 = e_0 - (p+1)n_0 - n_1 - 1 < 0.$$

So we get

$$|S_{n,1}| = \max\{e_0 - (p+1)n_0 - n_1 - 1, 0\}.$$

In the case $n_1 = 0$, we have $t = pn_0$ or $t = pn_0 + 1$ by $[(p-1)t/p] = (p-1)n_0$. If $t = pn_0$, we have $n_0 \leq s \leq e_0 - pn_0$. If $t = pn_0 + 1$, we have $n_0 + 1 \leq s \leq e_0 - pn_0 - 1$. So we get

$$|S_{n,1}| = \max\{e_0 - (p+1)n_0 + 1, 0\} + \max\{e_0 - (p+1)n_0 - 1, 0\}.$$

Secondly, we calculate $|S_{n,2}|$. In the case $n_1 \neq 1$, we have $S_{n,2} = \emptyset$. In the case $n_1 = 1$, we assume $(s, t) \in S_{n,2}$. Then $s = n_0$, and we have $pn_0 + 1 \leq t \leq e_0 - n_0$. So we get

$$|S_{n,2}| = \max\{e_0 - (p+1)n_0, 0\}.$$

Collecting these results, we have $|S_{n,1}| + |S_{n,2}| = a_n$.

Next, we calculate $|S'_{n,1}|$. We assume $(s, t) \in S'_{n,1}$. In the case $n'_1 \neq 0$, we have $s = e_0 - e_1 - pn'_0 - n'_1 - 1$ by $[(e - (p-1)s)/p] = (p-1)n'_0 + n'_1 + e_1$. We note that $n = (p-1)n'_0 + n'_1 + e_1 \geq 0$ shows $s \leq e_0$, because $n'_0 \geq -1$. Then $ps - t \geq (p-1)(s+t) - e$ implies $pt \leq pe_0 - pn'_0 - n'_1 - 1$, and further implies $t \leq e_0 - n'_0 - 1$. So we have

$$e_1 + pn'_0 + n'_1 + 2 \leq t \leq e_0 - n'_0 - 1.$$

We note that $e_1 + pn'_0 + n'_1 + 2 = n + n'_0 + 2 \geq 1$ and that if $s < 0$, we have

$$(e_0 - n'_0 - 1) - (e_1 + pn'_0 + n'_1 + 2) + 1 = e_0 - e_1 - (p+1)n'_0 - n'_1 - 2 < 0.$$

So we get

$$|S'_{n,1}| = \max\{e_0 - e_1 - (p+1)n'_0 - n'_1 - 2, 0\}.$$

In the case $n'_1 = 0$, we have $s = e_0 - e_1 - pn'_0 - 1$ or $s = e_0 - e_1 - pn'_0$ by $[(e - (p-1)s)/p] = (p-1)n'_0 + e_1$. If $s = e_0 - e_1 - pn'_0 - 1$, we have $e_1 + pn'_0 + 2 \leq t \leq e_0 - n'_0 - 1$. If $s = e_0 - e_1 - pn'_0$, we have $e_1 + pn'_0 + 1 \leq t \leq e_0 - n'_0$. We note that $n'_0 \geq 0$, because $n'_1 = 0$. So we get

$$|S'_{n,1}| = \max\{e_0 - e_1 - (p+1)n'_0 - 2, 0\} + \max\{e_0 - e_1 - (p+1)n'_0, 0\}.$$

At last, we calculate $|S'_{n,2}|$. In the case $n'_1 \neq 1$, we have $S'_{n,2} = \emptyset$. In the case $n'_1 = 1$, we assume $(s, t) \in S'_{n,2}$. Then $t = e_0 - n'_0$, and we have $n'_0 + 1 \leq s \leq e_0 - e_1 - pn'_0 - 1$. Here we need some care, because there is the case $n'_0 = -1$, in which case $t > e_0$. Now $n'_0 = -1$ is equivalent to $n = 0$ and $e_1 = p - 2$. So we get

$$|S'_{n,2}| = \max\{e_0 - e_1 - (p+1)n'_0 - 1, 0\}$$

except in the case where $n = 0$ and $e_1 = p - 2$, in which case $S'_{n,2} = \emptyset$. Collecting these results, we have $|S'_{n,1}| + |S'_{n,2}| = a'_n$. This completes the proof. \square

Example 2.2. *If $K = \mathbb{Q}_p(\zeta_p)$ and $\mathbb{F} = \mathbb{F}_p$, we have $|M(C_{\mathbb{F}_p}, \mathbb{Q}_p(\zeta_p))| = p + 3$ by Theorem 2.1. We know that $\mathbb{Z}/p\mathbb{Z} \oplus \mathbb{Z}/p\mathbb{Z}$, $\mathbb{Z}/p\mathbb{Z} \oplus \mu_p$ and $\mu_p \oplus \mu_p$ over $\mathcal{O}_{\mathbb{Q}_p(\zeta_p)}$ have the generic fibers which is isomorphic to $C_{\mathbb{F}_p}$. There are $(p+1)$ -choices of an isomorphism $C_{\mathbb{F}_p} \xrightarrow{\sim} (\mathbb{Z}/p\mathbb{Z} \oplus \mu_p)_{\mathbb{Q}_p(\zeta_p)}$ which give the different elements of $M(C_{\mathbb{F}_p}, \mathbb{Q}_p(\zeta_p))$. So $|M(C_{\mathbb{F}_p}, \mathbb{Q}_p(\zeta_p))| = 1 + (p+1) + 1$ shows that there does not exist any other extension of Raynaud schemes whose generic fiber is isomorphic to $C_{\mathbb{F}_p}$.*

Remark 2.3. *Theorem 2.1 is equivalent to an explicit calculation of the zeta function of $\mathcal{GR}_{V_{\bar{z}},0}$, and we see that $\dim \mathcal{GR}_{V_{\bar{z}},0} = \max\{n \geq 0 \mid a_n + a'_n \neq 0\}$.*

REFERENCES

- [Kis] M. Kisin, *Moduli of finite flat group schemes, and modularity*, to appear in Ann. of Math.
- [Ray] M. Raynaud, *Schémas en groupes de type (p, \dots, p)* , Bull. Soc. Math. France **102** (1974), 241–280.
- [Tat] J. Tate, *Finite Flat Group Schemes*, in Modular Forms and Fermat's Last Theorem, G. Cornell, J. Silverman and G. Stevens (eds.), Springer-Verlag, 1997.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8914, JAPAN.

E-mail address: naoki@ms.u-tokyo.ac.jp