Theoretical study of the magnetism in the incommensurate phase of TiOCl

D.M. Mastrogiuseppe *, M.E. Torio, C.J. Gazza, A.O. Dobry

Facultad de Ciencias Exactas Ingeniería y Agrimensura, Universidad Nacional de Rosario and Instituto de Física Rosario, Bv. 27 de Febrero 210 bis., 2000 Rosario, Argentina

Abstract

Going beyond a recently proposed microscopic model [1] for the incommensurate transition in the spin-Peierls TiOX (X=Cl, Br) compounds, in the present work we start by studying the thermodynamics of the model with XY spins and adiabatic phonons. We find that the system enters in an incommensurate phase by a first order transition at a low temperature T_{c1} . At a higher temperature T_{c2} a continuous transition to a uniform phase is found. Furthermore, we study the magnetism in the incommensurate phase by Density Matrix Renormalization Group (DMRG) calculations on a 1D Heisenberg model where the exchange is modulated by the incommensurate atomic position pattern. When the wave vector q of the modulation is near π , we find local magnetized zones (LMZ) in which spins abandon their singlets as a result of the domain walls induced by the modulated distortion. When q moves far away enough from π , the LMZ disappear and the system develops incommensurate magnetic correlations induced by the structure. We discuss the relevance of this result regarding previous and future experiments in TiOCl.

 $\begin{tabular}{ll} Key\ words: & Spin-Peierls; TiOCl; DMRG \\ PACS: & 75.10.Pq; & 75.40.Mg; & 64.70.Rh \end{tabular}$

1. Introduction

The interest on spin-Peierls systems [2] has been renewed with the recent discovery of a dimerization in the TiOX (X = Cl, Br) compounds family. In them, the magnetically active Ti ions connected by O ones are arranged in a bilayer structure. Ti ion positions on a layer are shifted with respect to the other neighboring layer, forming something like an anisotropic triangular structure. Initially postulated for a resonating valence bond state [3], TiOCl turned out to be mainly a one-dimensional magnetic system [4] with the Ti d_{xy} orbitals pointing towards each other in the crystallographic b direction [5]. The high temperature magnetic

E-mail address: mastrogiuseppe@ifir-conicet.gov.ar

susceptibility is well described by the Bonner-Fisher curve, indicating a nearest neighbor magnetic exchange $J \sim 660 K$ [5]. The peculiar ingredient in the TiOCl phase diagram is the appearance of an incommensurate intermediate phase between the high temperature uniform phase and the low T dimerized phase [5,6] commonly found in a regular spin-Peierls system. The transition temperatures were experimentally found to be $T_{c1} \sim 66 K$ and $T_{c2} \sim 92 K$. In Ref. [6] a very large energy gap of about 430 K in the low temperature phase and a pseudo spin gap below 135 K have been also reported.

The origin of the intermediate phase remains controversial and not well understood yet. The shifted positions of a Ti ion in a chain between the two neighboring chains, make it plausible to entail that some type of competition between the in-chain and out-of-chain interactions could be the origin of the incommensurate

^{*} Corresponding author. Tel.: +54-341-4853200 ext. 127; fax: +54-341-480-8584.

phase.

Some of us recently proposed a microscopic mechanism for the incommensurate phase [1], where the antiferromagnetic Ti chains are immersed in the phonon bath of the bilayer structure. By using the Cross-Fisher theory [7], it was shown that the geometrically frustrated character of the lattice is responsible for the structural instability, leading the chains to an incommensurate phase without an applied magnetic field. For TiOCl, our results showed consistency with the temperature dependence of the phonon frequencies and the value of the incommensuration vector at the transition temperature. Moreover, we found that the dynamical structure factor shows a progressive softening of an incommensurate phonon near the zone boundary as the temperature decreases, along with a broadening of the peak. These features are in agreement with the experimental inelastic X-ray measurements [8].

The purpose of this work is double. As a first step we present the phase diagram as a function of the temperature of the model proposed in Ref. [1]. On the other hand, we theoretically analyze the evolution of the magnetism inside the incommensurate phase by solving a one-dimensional Heisenberg model with a distortion-dependent exchange, using an incommensurate wave as the modulation of the Ti positions. This study is performed using Density Matrix Renormalization Group (DMRG) calculations.

2. The model and its phase diagram

The model proposed in Ref. [1] contains the relevant magnetic interaction, the phonons and the spin-phonon coupling. It includes the Ti atoms on the bilayer structure which interact by harmonic forces. For simplicity, we consider ionic movements only in the direction of the magnetic chains. As the measured magnetic susceptibility of TiOCl is well reproduced by a 1D Heisenberg model, the assumption that the Ti atoms are magnetically coupled only along the b direction was taken into account. Moreover, as a direct exchange seems to be the dominant Ti-Ti interaction, it is reasonable to consider that J will be modulated by the movement of nearest neighbor Ti ions in the chain direction. Our spin-phonon Hamiltonian reads:

$$H = H_{ph} + H_{sph},$$

$$H_{ph} = \sum_{i,j} \frac{P_{i,j}^2}{2m} + \sum_{i,j} \left\{ \frac{K_{in}}{2} (u_{i,j} - u_{i+1,j})^2 + \frac{K_{inter}}{2} \left[(u_{i,j} - u_{i,j+1})^2 + (u_{i,j} - u_{i+1,j-1})^2 \right] \right\},$$

$$H_{sph} = \sum_{i,j} J_{i,j} \mathbf{S}_{i,j} \cdot \mathbf{S}_{i+1,j},$$
(2)

where $u_{i,j}$ is the displacement of the atom i of the chain j along the direction b of the magnetic chains and the $P_{i,j}$ are their respective momenta. $\mathbf{S}_{i,j}$ are spin- $\frac{1}{2}$ operators with exchange constant $J_{i,j} = J[1 + \alpha(u_{i,j} - u_{i+1,j})]$ along the b-axis of a non-deformed underlying lattice. K_{in} and K_{inter} are the harmonic force constants

The Hamiltonian (1) contains variables of different dimensionality: whereas the spins are one-dimensional, phonons are two-dimensional. An exact solution to this model at finite temperature is a formidable task, therefore we make two approximations to obtain the phase diagram. We consider only the XY coupling in the spin-spin interaction and phonons are treated in the adiabatic approximation. Spin variables are converted to spinless fermions by means of the Jordan-Wigner transformation. The resulting pseudofermion-phonon model is solved in a finite system by diagonalizing the one-electron problem in the background of a deformed lattice (let us call each eigenvalue λ^{γ} , $\gamma = 1, ..., N_s$, where N_s is the the number of sites in each chain).

Now we consider deformations of the form:

$$u_{i,j} = u_0 \cos(\mathbf{q} \cdot \mathbf{R}_{i,j} - \pi/4), \tag{3}$$

where the wave vector $\mathbf{q} = (0, q)$ is constrained to the chain direction and $\mathbf{R}_{i,j}$ is the position vector in the two-dimensional Bravais lattice. We have chosen the $\pi/4$ phase to have a continuous energy at $q = \pi$. The total free energy per site is:

$$f = \frac{u_0^2 M \omega^2(\mathbf{q})}{2} - \frac{T}{N_s} \sum_{\gamma=1}^{N_s} \ln(1 + e^{-\lambda^{\gamma}/T}), \tag{4}$$

where T is the temperature in units of J and $\omega^2(\mathbf{q}) = \frac{4}{M} \left[K_{in} \sin^2(\frac{q_y}{2}) + K_{inter} (1 - \cos(\frac{q_x}{2}) \cos(\frac{q_y}{2})) \right]$ is the dispersion relation of the two-dimensional phonons.

We have minimized f with respect to u_0 and q at different temperatures. The results are shown in Fig. 1. At low temperatures a dimerized phase is found. This is the usual result for an isolated spin-Peierls chain. At the first transition temperature T_{c1} , u_0 jumps discontinuously and q moves away from π . This is a structural incommensurate phase which arises as a competition between the tendency to the dimerization of each chain and the frustration in the elastic interchain coupling. The value of q remains constant up to a second temperature T_{c2} where u_0 vanishes and the system becomes uniform. The sequence of the two transitions and the first order character of the first transition [8,9] experimentally found in TiOCl can therefore be accounted by our model. However the precise fitting of the parameters is a quite complex task which would need the inclusion of the z-component of the spin interaction and the non adiabatic effects of the phonons [10].




Fig. 1. Phase diagram in the plane u_0, T of the model 1 as discussed in the text. Parameters are $\alpha=3$ and $\frac{K_{inter}}{K_{in}}=1.8$. We identify each phase with the value of the wave vector of the pattern that minimize the free energy.

3. Magnetic structure in the incommensurate phase

The feature that makes the compound under study different from an usual spin-Peierls system is the appearance of an incommensurate phase without an applied magnetic field. It is therefore quite important to characterize the magnetism in this phase. As no magnetic interchain interaction was included in the model (1), we will consider a one-dimensional Heisenberg model as given by (2) with deformation fixed by (3). The model will be solved by DMRG. We will focus on the local magnetization and spin-spin correlation functions.

As a reference state to discuss the results, we take the dimerized phase because the structural deformation (3) with q near π can be thought as a smooth modulation of the dimer pattern. In a dimerized phase the spins are paired in singlets and the local magnetization is zero. There is a gap in the spectrum and the magnetic correlation declines exponentially. When the pattern (3) is included with $q_l = \pi(1 - \frac{2l}{N_s})$ to have an integer number of wavelengths in the chain, the dimerization phase changes its sign 2l times as we travel along the chain from one extreme to the other (see the gray line of Fig. 2). When l is small enough, the situation has some similarities with the domain walls (or solitons) appearing in the presence of a high magnetic field in regular spin-Peierls systems [11]. We expect that for each zero of (3) free LMZ should appear. This is in fact the situation we found in Fig. 2a, b and c where the mean value $\langle S_i^z \rangle$ is shown. We see the appearance of LMZ in all the zeros of the deformation except in the first, which is greatly affected by the proximity to the edge. These LMZ are seen as clouds of polarized spins emerging from the singlet background. Moreover, they are uncorrelated among them because their distances are longer than the correlation length of the dimerized




Fig. 2. DMRG results for a lattice of $N_s=200$. We show in black the atomic positions for different $q_l=\pi(1-\frac{2l}{N_S})$, with l=1,2,3,4 from top to bottom, and $u_0=0.053$. Superimposed, we plot the $< S_i^z >$ to show the appearance of the LMZ.

state. They are randomly orientated. We expect that the magnetic susceptibility will be affected by these LMZ. It should increase compared to the dimerized phase. This feature has been experimentally observed in TiOCl [8].

When q_l moves further away from π , the zeros of (3) fall within a distance of the order of the correlation length and the LMZ disappear as seen in Fig. 2d. The effect of the structural incommensuration on the magnetism is more subtle and a careful analysis of the magnetic correlation function is necessary.

In Fig. 3 we show the spin-spin correlation function $C(|i-j|) \equiv |< \mathbf{S}_i \cdot \mathbf{S}_j > - < \mathbf{S}_i > < \mathbf{S}_j > |$ and its Fourier transform (the static magnetic structure factor S(q)). For the three q where the LMZ appear, the correlation decreases exponentially to zero. Moreover, it falls a bit faster the farther q is from π . This seems to be a finite size effect because the modulation interferes with the decrease of the spin correlation of the dimerized zones when a zero appears. So we expect that in the thermodynamic limit the dimerized region will be large enough for the correlation to decrease as in the dimerized phase.

A more dramatic change in the behavior of C(|i-j|) is found for q_4 where the LMZ do not appear. In this case the correlation decreases more slowly than in the previous cases. For comparison we show the result of a uniform Heisenberg model (UHM) where the quasi-long range antiferromagnetic order is expected. Note that C(|i-j|) for q_4 approaches the one of the UHM, a quite different behavior than that of q_1, q_2 and q_3 . We will undertake a careful analysis of this long distance behavior for different modulations in a forthcoming paper.

More importantly, there is a modulation of the long distance behavior with half the wavelength of that which has the imposed structure. The magnetic structure can no longer be interpreted in terms of a weak




Fig. 3. (a) DMRG results for the spin-spin correlation functions $C(\mid i-j\mid)$ for deformations with the q_l as given in Fig. 2. We also show the correlation of a UHM for comparison. In the inset we show a log-log plot of $C(\mid i-j\mid)$. (b) Fourier transforms of the correlation functions of (a).

perturbation of the dimerized state. Instead, the magnetism becomes incommensurate as a response to the modulated exchange. In some sense, the situation is the dual to the one recently studied in some multiferroic materials [12]. There, an helical magnetic structure induces a lattice distortion which finally produces the ferroelectricity, whereas in our case the lattice distortion is the driving force which produces an incommensuration of the magnetic state.

These features are even more apparent in the structure factor S(q) shown in Fig. 3b. For q_1 , q_2 and q_3 there is a single maximum at $q_{max} = \pi$ whose intensity decreases as q moves away from π . This maximum is rounded as in a pure dimerized chain [13]. It can be fitted near $q=\pi$ by the law $\frac{1}{(q-\pi)^2+\xi^{-2}}$ (corresponding to a free massive boson propagator [14]) with $\xi \sim 3.3$. This asserts our previous statement that imposing distortions with wavector q_l near enough to π , the long range magnetic correlation behaves as in a dimerized chain, decreasing exponentially with a correlation length of the order of ξ . For $q_l = q_4$ the behavior changes qualitatively. Two additional maxima appear at one side and the other of the highest peak at π , corresponding to the long distance magnetic modulation found in C(|i-j|). Moreover, the peak at the center is not longer rounded but it behaves similarly to the UHM. This signals a tendency towards a quasi-long range incommensurate magnetic correlation of the system. Let us make a brief discussion about the application of these results to TiOCl. As measured from X-ray experiments [8,9], the modulation wave vector is $q \sim 3.05$ at T_{c2} and it moves slightly to π when decreasing the temperature towards T_{c1} where it jumps discontinuously to π . So we expect a change of the magnetic behavior inside the incommensurate phase from a structure with LMZ to another without LMZ but with an incommensurate correlation function. This should be checked in future magnetic experiments.

In summary we have found that a system with one-dimensional antiferromagnetic chains coupled to two-dimensional phonons in an anisotropic triangular lattice accounts for the appearance of the incommensurate phase in TiOCl. The magnetism in this intermediate state includes uncorrelated LMZ near the dimerized phase, and incommensurate correlations with no LMZ towards the uniform phase. In the last case, the modulation found in the correlation function is related to the modulation of the imposed distortion by a factor of two in their wavelengths.

This work was supported by ANPCyT (PICT 1647), Argentina.

References

- [1] D. Mastrogiuseppe and A. Dobry, arXiv:0810.3018v1.
- [2] J. W. Bray et al., Extended linear chain compounds, edited by J. S. Miller, Plenun, New York 1983, Vol 3, pp 353-415.
- [3] R. Beynon and J. Wilson, J. Phys.: Condens. Matter 5, 1983 (1993).
- [4] M. Shaz et al., Phys. Rev. B. **71**, 100405(R) (2005).
- [5] A. Seidel, C. A. Marianetti, F. C. Chou, G. Ceder, P. A. Lee, Phys. Rev. B 67, 020405(R) (2003).
- [6] T. Imai and F. C. Chou, arXiv:cond-mat/0301425v1.
- [7] M.C. Cross and D.S. Fisher, Phys. Rev. B 19, 402 (1979).
- [8] E. T. Abel et al., Phys. Rev. B. **76**, 214304 (2007).
- [9] A. Krimmel et~al., Phys. Rev. B. **73**, 172413 (2006).
- [10] A. O. Dobry, D. C. Cabra, and G. L. Rossini, Phys. Rev. B 75, 045122 (2007).
- [11] A. E. Feiguin, J. A. Riera, A. Dobry, and H. A. Ceccatto, Phys. Rev. B 56, 14607 (1997). Note that in the case studied in this reference, states with finite magnetization where considered whereas in the present case no magnetic field is applied and the LMZ add up to give a zero total S_z state.
- [12] I. A. Sergienko and E. Dagotto, Phys. Rev. B 73, 094434 (2006).
- [13] R. Chitra et al., Phys. Rev B 52, 6581 (1995).
- [14] S. R. White and I. Affleck, Phys. Rev. B 54, 9862 (1996).