

Quantization of the Jackiw-Teitelboim model

Clisthenis P. Constantinidis^{*,†} and Olivier Piguet^{*,†}
Universidade Federal do Espírito Santo, Vitória, Brazil

Alejandro Perez
Centre de Physique Théorique[‡], Campus de Luminy, 13288 Marseille, France.

(Dated: May 1, 2019)

We study the phase space structure of the Jackiw-Teitelboim model in its connection variables formulation where the gauge group of the field theory is given by local $SL(2, \mathbb{R})$ (or $SU(2)$ for the Euclidean model), i.e. the de Sitter group in two dimensions. In order to make the connection with two dimensional gravity explicit, a partial gauge fixing of the de Sitter symmetry can be introduced that reduces it to spacetime diffeomorphisms. This can be done in different ways. Having no local physical degrees of freedom, the reduced phase space of the model is finite dimensional. The simplicity of this gauge field theory allows for studying different avenues for quantization, which may use various (partial) gauge fixings. We show that reduction and quantization are noncommuting operations: the representation of basic variables as operators in a Hilbert space depend on the order chosen for the latter. Moreover, a representation that is natural in one case may not even be available in the other leading to inequivalent quantum theories.

I. INTRODUCTION

The Jackiw-Teitelboim model [1, 2] is one of the simplest but nontrivial formulations of General Relativity (GR) in two-dimensional space-time with cosmological constant k . Its action is given by

$$S_{JT} = \frac{1}{2} \int d^2x \sqrt{-g} \psi (R - 2k). \quad (\text{I.1})$$

It is invariant under space-time diffeomorphisms and leads to the Liouville equation

$$R - 2k = 0, \quad (\text{I.2})$$

It contains only a finite number of degrees of freedom, namely one (here we assume the space time topology $M = S^1 \times \mathbb{R}$). The model may be quantized in the original variables of Jackiw and Teitelboim in a canonical framework including the two first class constraints corresponding to space-time diffeomorphism invariance [1, 3].

On the other hand, one may take profit of its equivalence with a BF theory [4, 5, 6], which has a structure similar to the first order formulation of four-dimensional GR in Ashtekar's variables. Here, instead of being the four-dimensional Lorentz group, the gauge group is that of two-dimensional de Sitter or anti-de Sitter symmetry $SO(1, 2)$, or $SO(3)$ in the Euclidean de Sitter case—or better their covering groups, $SL(2, \mathbb{R}) \approx SU(1, 1)$ or $SU(2)$, respectively. The fields are a gauge connection 1-form ω and a scalar ϕ in the adjoint representation. A quantization in the Euclidean case was presented in [7], using spin network and spinfoam techniques [8].

The canonical formulation of the BF theory gives rise to three first class constraints whose Poisson Bracket algebra reproduces the three-dimensional Lie algebra of the gauge group. The quantization may follow various roads, using some complete or partial gauge fixing [5, 6, 9, 10, 11, 12], or no gauge fixing at all [7, 13]. In [11], a time gauge has been used, which consists in the vanishing of the connection component ω^0_x —which is interpreted as the space component of the zweibein (2-bein) form e^0 —with the purpose of simulating the time gauge fixing of four-dimensional gravity leading to the Ashtekar variables formulation [8]. This partial gauge fixing leads to a reduction of the number of first class constraints to

* Work supported in part by the Conselho Nacional de Desenvolvimento Científico e Tecnológico—CNPq (Brazil).

† Work supported in part by the PRONEX project No. 35885149/2006 from FAPES—CNPq (Brazil).

‡ Unité Mixte de Recherche (UMR 6207) du CNRS et des Universités Aix-Marseille I, Aix-Marseille II, et du Sud Toulon-Var; laboratoire affilié à la FRUMAM (FR 2291)

two, corresponding to the space-time diffeomorphism invariance—namely one constraint generating the space diffeomorphisms and the other one playing the role of the Hamiltonian constraint.

Despite of the extensive literature studying the JT model, there is, to our knowledge, no complete treatment of the quantization of the Lorentzian sector in its first order formulation. The main difficulty is technical: the fact that the gauge group in that case is noncompact precludes the possibility of using the standard quantization techniques that are applicable in the Riemannian case.

The purpose of the present paper is twofold:

On the one hand we study in detail the quantization of the model in the Lorentzian sector. This is achieved through a minimalistic application of the techniques developed in [14] which can also be introduced from the point of view of [15]. It is by now known [16] that the general case of a gauge theory with noncompact internal gauge symmetries presents important difficulties that are not completely resolved by the methods proposed in [14]. Interestingly, those difficulties vanish in the 2-dimensional case and the quantization presented here is well defined.

On the other hand, we propose a new partial gauge fixing defined by the vanishing of one component of the scalar field ϕ , to compare the corresponding quantum theory with the theories already constructed [7] or under construction [12]. The question makes sense since it is well-known [17, 18] that, even in theories with a finite number of degrees of freedom, it may exist inequivalent quantization of the same classical theory if some assumption of the von Neumann theorem is invalid, such as for instance the existence of pairs of self-adjoint operators “ p , q ” obeying Heisenberg commutation relations. In a gauge theory, inequivalent quantizations can also arise from the possible inequivalence of the two customary procedures of quantization consisting in either reducing the unconstrained phase space and then quantizing or quantizing first and then imposing the constraints at the quantum level (Dirac procedure). This is particularly important in our case where, even though the number of physical degrees of freedom is finite, the unreduced phase space of the system is infinite dimensional. Consequently, in the second quantization procedure—the Dirac procedure—the von Newman theorem has no bearing and infinitely many inequivalent quantizations exist in principle.

II. THE JACKIW-TEITELBOIM MODEL IN THE BF FORMULATION

The model can be written as a BF theory [13] in 2 dimensions. The gauge group G is de Sitter or anti-de Sitter in Riemannian or Lorentzian space-time. The infinitesimal generators are

$$J_0 := P_0, \quad J_1 := P_1, \quad J_2 := \Lambda,$$

with commutation relations

$$[J_0, J_1] = kJ_2, \quad [J_0, J_2] = -J_2, \quad [J_1, J_2] = \sigma J_1,$$

where k is the cosmological constant and σ is the metric signature, equal to 1 in the Riemannian case and to -1 in the Lorentzian case. A redefinition of the Lie algebra basis allows to reduce its commutation rules to:

$$[J_i, J_j] = f_{ij}{}^k J_k, \quad \text{with} \quad f_{01}{}^2 = 1, \quad f_{12}{}^0 = \sigma, \quad f_{20}{}^1 = 1, \quad (i, j, \dots = 0, 1, 2), \quad (\text{II.1})$$

which, for $\sigma = -1$ or 1 , is the Lie algebra of $\text{SO}(3)$ or $\text{SO}(1,2)$. We shall consider in the following their covering groups $\text{SU}(2)$ or $\text{SL}(2, \mathbb{R})$. The invariant Killing form η_{ij} has the form of a Euclidian or Minkowskian 3-dimensional metric:

$$\eta_{ij} := -\frac{\sigma}{2} f_{ik}{}^l f_{jl}{}^k = \text{diag}(\sigma, 1, 1). \quad (\text{II.2})$$

The fields are an $\text{SU}(2)$ or $\text{SL}(2, \mathbb{R})$ connection 1-form ω^i and a scalar field ϕ^i in the adjoint representation of the gauge group. For this point on we shall denote the internal gauge group with G , and we will use \mathfrak{g} to designate its Lie algebra. We will use the explicit $\text{SL}(2, \mathbb{R})$ and $\text{SU}(2)$ when we specialize to the Lorentzian or Riemannian models respectively. The components ω^0 and ω^1 are interpreted as the zweibein components and ω^2 as the rotation, respectively Lorentz connection.

The theory is the 2d version of BF theory, and can be seen as the $g \mapsto 0$ limit (g being the coupling constant) of 2d Yang Mills theory. Its action takes the form

$$S = \int_{\mathcal{M}} \phi^i F^j(\omega) \eta_{ij}, \quad (\text{II.3})$$

and the field equations are

$$\begin{aligned} d_\omega \phi^i &:= d\phi^i + f_{jk}^i \omega^j \phi^k = 0, \\ F^i &:= d\omega^i + \frac{1}{2} f_{jk}^i \omega^j \omega^k = 0, \end{aligned} \quad (\text{II.4})$$

where d_ω is the ω -covariant exterior differential and F^i the curvature 2-form of the connection. At first sight the action is invariant under two kinds of gauge transformations: the conventional Yang-Mills-like local G transformations generated by a Lie algebra valued scalar field λ

$$\begin{aligned} \delta_\lambda \omega &= d_\omega \lambda, \\ \delta_\lambda \phi &= [\phi, \lambda], \end{aligned} \quad (\text{II.5})$$

whose exponentiated version gives

$$\begin{aligned} \omega' &= a\omega a^{-1} + da a^{-1}, \\ \phi' &= a\phi a^{-1}, \end{aligned} \quad (\text{II.6})$$

for $a = \exp(\lambda) \in G$. The active diffeomorphisms are generated by a vector field v

$$\begin{aligned} \delta_v \omega &= \mathcal{L}_v \omega, \\ \delta_v \phi &= \mathcal{L}_v \phi, \end{aligned} \quad (\text{II.7})$$

\mathcal{L}_v being the Lie derivative. However, on shell a diffeomorphism generated by v is the same transformation as a local G -transformation generated by the field

$$\lambda^i = v^a \omega_a^i, \quad (\text{II.8})$$

as can be easily checked by writing these equations in components and using the equations of motion. Therefore, in this theory the diffeomorphisms (acting on the space of solutions) can be considered as a subgroup of the local G gauge transformations.

Solutions are given by flat connections $\omega = gdg^{-1}$ for $g(x) \in G$, and (covariantly) constant ϕ fields. Locally, one can choose a gauge so that the connection $\omega = 0$. In this gauge the equation $d_\omega \phi = 0$ implies that $\phi = \text{constant}$. This is particularly important for the Lorentzian case since it implies that the causal type of ϕ (thought as a vector in three dimensional Minkowsky internal geometry) cannot change in the classical solutions. This conclusion is a global one since no gauge transformation $\phi \rightarrow g\phi g^{-1}$ can send a time-like ϕ into a space-like one or viceversa. In fact, this property is taken over to the quantum theory where we will show that super-selection sectors associated to ϕ being space-like or time-like appear.

III. THE HAMILTONIAN FORMULATION

When $\mathcal{M} = S^1 \times R$ one can quantize the theory in the canonical framework. General topologies can in principle be considered in the path integral approach. The Hamiltonian formulation is obtained through the standard 1 + 1 spacetime decomposition. More precisely, one introduces an arbitrary foliation of \mathcal{M} by choosing a time function. In terms of this foliation the action becomes

$$S = \int dt \int_{S^1} dx (\phi_i \dot{\omega}^i + \omega_i^i D\phi_i). \quad (\text{III.1})$$

We use the notations $D\phi^i := \partial\phi^i + f_{jk}^i \omega^j \phi^k$, $\omega^i := \omega_x^i$, $\partial := \partial_x$, x being the space coordinate. The Poisson bracket among the phase space variables is

$$\{\omega^j(x), \phi_i(y)\} = \delta_i^j \delta(x-y) \quad \text{or} \quad \{\omega^j(x), \phi^i(y)\} = \eta^{ij} \delta(x-y).$$

We have three first class constraints [19, 20, 21] corresponding to the three components of the Gauss law $g_i := D\phi^i = 0$. Explicitly these components are:

$$\begin{aligned} g_0 &= \partial\phi^0 + \sigma(\omega^1\phi^2 - \omega^2\phi^1) \approx 0, \\ g_1 &= \partial\phi^1 + \omega^2\phi^0 - \omega^0\phi^2 \approx 0, \\ g_2 &= \partial\phi^2 + \omega^0\phi^1 - \omega^1\phi^0 \approx 0. \end{aligned} \quad (\text{III.2})$$

The smeared Gauss constraint $g(\alpha) \equiv \int_{S^1} \text{Tr}[\alpha D\phi]$ for $\alpha \in \mathfrak{g}$ are first class—they satisfy the Poisson bracket identity $\{g(\alpha), g(\beta)\} = g([\alpha, \beta])$ —and generate infinitesimal G -gauge transformations:

$$\{g(\alpha), \omega^i\} = \partial\alpha^i + f_{jk}^i \omega^j \alpha^k, \quad \{g(\alpha), \phi^i\} = f_{jk}^i \phi^j \alpha^k. \quad (\text{III.3})$$

There are therefore three local first class constraints for the three configuration variables ω_a^i . Thus the naive counting of degrees of freedom gives zero physical degrees of freedom. However, the naive counting is only sensitive to local excitations. The theory has indeed global degrees of freedom. In particular, if $M = S^1 \times \mathbb{R}$, an algebraic basis for the gauge invariant (Dirac) observables is given by

$$O_1 = \phi_i \phi^i, \quad O_2 = \text{Tr}[P \exp(-\int_{S^1} \omega)]. \quad (\text{III.4})$$

The physical phase space being therefore 2-dimensional, the theory has a single (global) degree of freedom. We note for further use that the quantity

$$Q := \frac{\omega^i \phi_i}{\phi^i \phi_i} \quad (\text{III.5})$$

transforms as an Abelian connection under the special gauge transformations which leave ϕ invariant:

$$\{g(\alpha_{\text{Abel}}), Q\} \approx \partial a, \quad \text{with} \quad \alpha_{\text{Abel}}^i = a \phi^i. \quad (\text{III.6})$$

This holds up to the constraint $\partial(\phi_i \phi^i) \approx 0$ which follows from (III.2).

A. Partial gauge fixings

Spacetime diffeomorphisms are hidden inside the larger gauge group of BF theory that in two dimensions corresponds to the local G transformations given in (II.5). Notice that the former are generated by the two components of a vector field in \mathcal{M} while the latter are generated by the three components of $\lambda \in \mathfrak{g}$. In this section we partially gauge fix the symmetries of BF theory in order to establish a more direct relationship with diffeomorphisms, and hence emphasize the relationship of the model with two dimensional gravity.

We will partially gauge fix the G gauge symmetry by requiring the fourth constraint

$$g_3 = \phi_i n^i \approx 0, \quad (\text{III.7})$$

where n^i is a fixed normalized vector in the internal space. Before going into the technical details of the constraint algebra let us discuss the geometric interpretation of the partial gauge fixing introduced by the above equation. Due to the fact that $\phi \cdot \phi$ is a Dirac observable (a constant of motion) one can separate the analysis into three distinct (dynamically independent) cases in the Lorentzian case: $\phi \cdot \phi > 0$, $\phi \cdot \phi = 0$ and $\phi \cdot \phi < 0$ (in the Riemannian case there is only the first sector). The following discussion is restricted to the Lorentzian sector.

1. The “space-like” sector: $\phi \cdot \phi > 0$

In that case the good choice of gauge fixing corresponds to n^i =time-like. The condition (III.7) is expected to reduce the group $\text{SL}(2, \mathbb{R})$ to a two dimensional subgroup. A moment of reflection shows that this is given by the cartesian product of the $\text{U}(1) \subset \text{SL}(2, \mathbb{R})$ that leaves invariant n^i , and the little group (the boosts) leaving invariant the vector ϕ^i . All this will become transparent in the following.

The choice n^i =space-like leads to a degenerate situation for phase space points where $\phi^i \propto n^i$, as on these points the little groups associated to n^i and ϕ^i coincide. This leads to complications that we will not analyze in this work.

To simplify notation we can take $n^i = (1, 0, 0)$ which simply turns (III.7) into simply $g_3 = \phi^0 \approx 0$. With this choice, the matrix $G_{\alpha\beta} = \{g_\alpha, g_\beta\}$ becomes

$$G(x, y) = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\sigma\phi^2\delta(x-y) \\ 0 & 0 & 0 & \sigma\phi^1\delta(x-y) \\ 0 & \sigma\phi^2\delta(x-y) & -\sigma\phi^1\delta(x-y) & 0 \end{bmatrix} \quad (\text{III.8})$$

In order to isolate the second class part of the previous constraints we make the following redefinition

$$\begin{aligned}
C_0 &= g_0 = \partial\phi^0 + \sigma(\omega^1\phi^2 - \omega^2\phi^1) \approx 0, \\
C_1 &= \phi^1 g_1 + \phi^2 g_2 = \phi^1(\partial\phi^1 + \omega^2\phi^0 - \omega^0\phi^2) + \phi^2(\partial\phi^2 + \omega^0\phi^1 - \omega^1\phi^0) \approx 0, \\
C_2 &= \phi^1 g_1 - \phi^2 g_2 = \phi^1(\partial\phi^1 + \omega^2\phi^0 - \omega^0\phi^2) - \phi^2(\partial\phi^2 + \omega^0\phi^1 - \omega^1\phi^0) \approx 0, \\
C_3 &= g_3 = \phi^0 \approx 0.
\end{aligned} \tag{III.9}$$

With this definition the matrix $C = \{C_\alpha, C_\beta\}$ is block diagonal, namely, up to terms involving constraints:

$$C \approx \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2\sigma\phi^2\phi^1\delta(x-y) \\ 0 & 0 & 2\phi^2\sigma\phi^1\delta(x-y) & 0 \end{bmatrix} \tag{III.10}$$

This implies that the pair (C_2, C_3) is second class and can therefore explicitly solved. For instance they can be used to solve for ω^0 . Namely

$$\phi^0 = 0, \quad \omega^0 = \frac{1}{2} \left(\frac{\partial\phi^1}{\phi^2} - \frac{\partial\phi^2}{\phi^1} \right). \tag{III.11}$$

The first class ones become:

$$\begin{aligned}
C_0 &= \omega^1\phi^2 - \omega^2\phi^1 = \epsilon_{AB}\omega^A\phi^B \approx 0, \\
C_1 &= \frac{1}{2}\partial(\phi^A\phi^A) \approx 0,
\end{aligned} \tag{III.12}$$

where $A, B = 1, 2$. It is easy to see that the Dirac bracket among the remaining variables is

$$\{\omega^B(x), \phi^A(x)\}_D = \delta^{BA}\delta(x-y). \tag{III.13}$$

Direct computation shows that the algebra of the first class constraints is Abelian. More precisely, if we define

$$C_0(a) = \int_{S^1} dx a(x) C_0(x), \quad \text{and} \quad C_1(b) = \int_{S^1} dx b(x) C_1(x), \tag{III.14}$$

we have the following Dirac bracket algebra:

$$\{C_0(a), C_0(b)\}_D = 0 = \{C_1(a), C_1(b)\}_D \tag{III.15}$$

and

$$\{C_0(a), C_1(b)\}_D = -\frac{1}{2} \int_{S^1} dx \int_{S^1} dy a(x) \partial b(y) \epsilon_{AB} \phi^A(x) \{\omega^B(x), \phi^C \phi^C(y)\}_D = 0. \tag{III.16}$$

The gauge transformations generated by the first class constraints are

$$\begin{aligned}
\delta_{(0)}\omega^A &= \{\omega^A, C_0(a)\}_D = -\sigma a \epsilon^{BA} \omega^B, \\
\delta_{(0)}\phi^A &= \{\phi^A, C_0(a)\}_D = -\sigma a \epsilon^{BA} \phi^B,
\end{aligned} \tag{III.17}$$

which correspond to local U(1) internal rotations, and

$$\begin{aligned}
\delta_{(1)}\omega^A &= \{\omega^A, C_1(a)\}_D = -\phi^A \partial a, \\
\delta_{(1)}\phi^A &= \{\phi^A, C_1(a)\}_D = 0.
\end{aligned} \tag{III.18}$$

What is the geometric meaning of the transformation generated by C_1 ? Recall that $C_1 \equiv \phi^1 g_1 + \phi^2 g_2$ which is nothing else than the expression in the gauge (III.7) of $\phi^i g_i$. This is precisely the generator of the internal ‘‘boosts’’ leaving the ‘‘space-like’’ field ϕ^i invariant, as anticipated above.

The Riemannian theory is fully described by the equations of this section. The only change is that both C_0 and C_1 generate U(1) transformations in that case.

2. The “time-like” sector: $\phi \cdot \phi < 0$

The gauge fixing analogous to the previous case would now be defined with n^i spacelike and we would expect condition (III.7) to reduce the group $\text{SL}(2, \mathbb{R})$ to the cartesian product of the boosts leaving invariant n^i , and of the little group (the $\text{U}(1)$ rotations) leaving invariant the vector ϕ^i . It seems however difficult to control the positivity of $\phi \cdot \phi$, which is no more automatic. A more appropriate choice is to take $\phi^i(x) = \phi(x)u^i$ where u is some fixed time-like vector. With the choice $u = (1, 0, 0)$, this amounts to add to the constraints (III.2)—taken with $\sigma = -1$ —the new constraints

$$g_3 = \phi^1 \approx 0, \quad g_4 = \phi^2 \approx 0.$$

The 5×5 Poisson bracket matrix $G_{\alpha\beta} = \{g_\alpha, g_\beta\}$, $\alpha, \beta = 0, \dots, 4$ reads (up to constraints):

$$G(x, y) \approx \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \end{bmatrix} \phi(x) \delta(x - y). \quad (\text{III.19})$$

Only g_0 is first class, the remainder four constraints being of second class. Eliminating them through the Dirac procedure, we are left with the strong conditions

$$\phi^A = 0, \quad \omega^A = 0, \quad A = 1, 2,$$

and the Dirac bracket algebra (with $\omega := \omega^0$)

$$\{\omega(x), \phi(y)\}_D = -\delta(x - y). \quad (\text{III.20})$$

The first class constraint reads

$$g_0(x) = \partial\phi \quad \text{or, in integral form:} \quad g_0(a) = - \int_{S^1} dx \partial a \phi. \quad (\text{III.21})$$

and generates the $\text{U}(1)$ gauge rotations which leaves ϕ^i invariant:

$$\{g_0(a), \phi(y)\}_D = 0, \quad \{g_0(a), \omega(x)\}_D = \partial a.$$

ω plays the role of the $\text{U}(1)$ connection. We note that the connection Q (III.5) is equivalent to ω , since $Q = \omega/\phi$ and ϕ is constrained to be a constant.

3. The “null” sector: $\phi \cdot \phi = 0$

As in the previous case we take $\phi^i(x) = \phi(x)u^i$ where u is some fixed null vector. With the choice $u = (1, 1, 0)$, this amounts to add to the constraints (III.2) – taken with $\sigma = -1$ – the new constraints

$$g_3 = \phi^0 - \phi^1 \approx 0, \quad g_4 = \phi^2 \approx 0.$$

It turns out that the constraint system is reducible as one has that $g_0 - g_1 = \partial g_3 - \omega^2 g_3$. Following the standard procedure we drop the constraint g_0 . The 4×4 Poisson bracket matrix $G_{\alpha\beta} = \{g_\alpha, g_\beta\}$, $\alpha, \beta = 1, \dots, 4$ reads (up to constraints):

$$G(x, y) \approx \begin{bmatrix} 0 & 0 & 0 & \phi^0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\phi^0 & 0 & 0 & 0 \end{bmatrix} \delta(x - y). \quad (\text{III.22})$$

Therefore g_2 and g_3 are first class, while g_1 and g_4 are second class. Eliminating them through the Dirac procedure, we are left with the strong conditions

$$\phi^2 = 0, \quad \omega^2 = -\frac{\partial\phi^1}{\phi^0},$$

and the Dirac bracket algebra

$$\{\omega_A(x), \phi^A(y)\}_D = \delta_A^B \delta(x-y) \quad A = 0, 1. \quad (\text{III.23})$$

The first class constraints can be replaced by the equivalent pair

$$C_0 = \omega_0 + \omega_1 \quad C_1 = \phi^0 - \phi^1. \quad (\text{III.24})$$

The second one imposes the field to be of the form $\vec{\phi} = (\phi, \phi, 0)$, while first one generates boosts (translations of $\vec{\phi}$ along the null line $\phi^0 - \phi^1 = 0$). The system has no degree of freedom (not even global) and the null case is trivial.

4. Diffeomorphisms, Virasoro and Abelian generators

For simplicity the following analysis is performed in the $\phi \cdot \phi > 0$ sector. Let us define

$$\Gamma = \frac{\epsilon_{AB} \phi^B \partial \phi^A}{\phi^C \phi^C}, \quad (\text{III.25})$$

solution of the equation

$$\partial \phi^A - \epsilon^{AB} \Gamma \phi^B = 0., \quad (\text{III.26})$$

an analog to the torsion free connection of general relativity in the first order formulation. We may check that $\Gamma \approx \omega_0$ and that (III.26), with Γ replaced by ω_0 , is a constraint, a combination of the first class constraint C_1 and of the second class one C_2 .

One can introduce variables invariant under the gauge group generated by C_0 as follows:

$$\Pi \equiv \frac{1}{2} \phi^A \phi^A \quad \text{and} \quad Q \equiv \frac{\phi^A \omega^A}{\phi^C \phi^C}, \quad (\text{III.27})$$

obeying the following equation:

$$\int_{S^1} dx a(x) \{\Gamma(x), Q(y)\}_D = 0, \quad (\text{III.28})$$

for arbitrary $a(x) \in C^1(S^1)$. Π corresponds, in the $\phi^0 = 0$ gauge, to the invariant O_1 defined in (III.4), whereas Q is the Abelian connection (III.5). The meaning of these quantities will become clearer in the next section.

The constraints C_0 and C_1 are scalar densities of weight one. This is why they are naturally smeared with scalar functions a and b in order to produce coordinate independent quantities $C_0(a)$ and $C_1(b)$ respectively. We would like now to define an equivalent set of constraints that would be suitably smeared with vector fields of S^1 . In order to do this one needs first to define vector density constraints which can be achieved by multiplying the original ones by density one phase space functions. Without further motivation we redefine the constraints as:

$$\begin{aligned} V_1 &= -\Gamma C_0 = -\frac{\epsilon_{AB} \phi^B \partial \phi^A}{\phi^C \phi^C} \epsilon_{DE} \omega^D \phi^E \approx 0, \\ V_2 &= -Q C_1 = -\frac{1}{2} \frac{\phi^A \omega^A}{\phi^C \phi^C} \partial(\phi^B \phi^B) \approx 0. \end{aligned} \quad (\text{III.29})$$

These are vector densities of weight one (or scalar densities of weight two). As long as we are away from configurations for which $\Gamma = 0$ or $Q = 0$ the previous constraints define the same constraint surface. Assuming we have two vector fields α and β , we define the smeared versions of the previous constraints in the obvious manner. Then one has that

$$\{V_1(\alpha), V_2(\beta)\}_D = 0 \quad (\text{III.30})$$

and

$$\begin{aligned} \{V_1(\alpha), V_1(\beta)\}_D &= V_1([\alpha, \beta]), \\ \{V_2(\alpha), V_2(\beta)\}_D &= V_2([\alpha, \beta]). \end{aligned} \quad (\text{III.31})$$

where $[\alpha, \beta]$ is the vector field commutator. Therefore V_1 and V_2 commute with respect to each other, each of them satisfying a classical Virasoro algebra. They look like diffeomorphism generators, however none of the two generates diffeomorphisms of S^1 . The combination that does this is:

$$D = V_1 + V_2. \quad (\text{III.32})$$

The analog of the Hamiltonian constraint of gravity then is

$$H = V_1 - V_2. \quad (\text{III.33})$$

These satisfy the ‘gravity’ algebra

$$\begin{aligned} \{D(\alpha), D(\beta)\}_D &= D([\alpha, \beta]), \\ \{H(\alpha), H(\beta)\}_D &= D([\alpha, \beta]), \\ \{H(\alpha), D(\beta)\}_D &= H([\alpha, \beta]). \end{aligned} \quad (\text{III.34})$$

The constraint $D(\alpha)$ generate standard diffeomorphisms as can be checked by a direct calculation, namely

$$\{D(\alpha), \phi^A\}_D = \alpha \Gamma \epsilon^{AB} \phi^B \approx -\alpha \partial \phi^A = -\mathcal{L}_\alpha \phi^A, \quad (\text{III.35})$$

$$\{D(\alpha), \omega^A\}_D = \alpha \Gamma \epsilon^{AB} \omega_B - \partial(\alpha Q) \phi^A \approx -(\alpha \partial \omega^A + \partial \alpha \omega^A) = -\mathcal{L}_\alpha \omega^A, \quad (\text{III.36})$$

where the weak equalities in (III.35) and (III.36) mean the insertion of the constraint equations. Similarly, H generates time evolution up to space diffeomorphisms and field equations.

B. Full Reduction

Notice, from (III.27), that $(Q(x), \Pi)$ are a conjugate $U(1)$ -invariant fields, i.e., they strongly commute with C_0 , and

$$\{Q(x), \Pi(y)\}_D = \delta(x - y). \quad (\text{III.37})$$

Since Π also commutes with C_1 , it represents a strong Dirac observable of the model. So we introduce

$$\mathcal{P} \equiv \int_{S^1} \Pi. \quad (\text{III.38})$$

On the other hand Q does not commute with C_1 but it transforms as an Abelian connection—c.f. (III.6):

$$\delta_{(1)} Q = \{Q, C_1(a)\}_D = -\partial a. \quad (\text{III.39})$$

Therefore, it is quite easy to define a strong Dirac observable

$$\mathcal{X} \equiv \int_{S^1} Q \quad (\text{III.40})$$

so that

$$\{\mathcal{X}, \mathcal{P}\}_D = 1. \quad (\text{III.41})$$

The connection Q is an \mathbb{R} -connection (associated to the boost structure group) in the $\phi \cdot \phi > 0$ sector, while it becomes a $U(1)$ -connection in the $\phi \cdot \phi < 0$ sector.

C. Time-gauge reduction

In this section we describe the results obtained in [11] where the ‘temporal gauge’ was considered. This gauge fixing is based in the one taken in four dimensions, through which we can obtain a compact gauge group as the residual symmetry, as described in [8]. It consists in making the zweibein component $\chi := \omega_x^0$ vanish and it is implemented as an extra constraint $\chi \approx 0$. The action then reads

$$S = \int dt \int_{S^1} dx (\phi_i \dot{\omega}^i + \omega_t^i D\phi_i + B \chi), \quad (\text{III.42})$$

and again the Dirac method is used in order to eliminate the second class constraints. The remaining constraints, namely \mathcal{G}'_0 and \mathcal{G}'_1 are

$$\mathcal{G}'_0(x) = (\omega_x^1) \mathcal{G}_0(x) = \sigma \omega_x^1 \partial_x \left(\frac{\partial_x \phi^2}{\omega_x^1} \right) + k (\omega_x^1)^2 \phi^2 - \omega_x^1 \omega_x^2 \phi^1, \quad (\text{III.43})$$

$$\mathcal{G}'_1(x) = \omega_x^1 \mathcal{G}_1(x) = \omega_x^1 \partial_x \phi^1 + \omega_x^2 \partial_x \phi^2. \quad (\text{III.44})$$

The Dirac bracket algebra of these constraints is closed:

$$\begin{aligned} \{\mathcal{G}'_0(\epsilon), \mathcal{G}'_0(\eta)\}_D &= \sigma \mathcal{G}'_1([\epsilon, \eta]), \\ \{\mathcal{G}'_0(\epsilon), \mathcal{G}'_1(\eta)\}_D &= -\mathcal{G}'_0([\epsilon, \eta]), \\ \{\mathcal{G}'_1(\epsilon), \mathcal{G}'_1(\eta)\}_D &= -\mathcal{G}'_1([\epsilon, \eta]), \end{aligned} \quad (\text{III.45})$$

where $[\epsilon, \eta] = (\epsilon \partial_x \eta - \eta \partial_x \epsilon)$, which confirms that \mathcal{G}'_0 and \mathcal{G}'_1 are first class. In fact, time diffeomorphisms are generated by \mathcal{G}'_0 up to constraints, up to field equations (“on-shell realization”), and up to a compensating local Lorentz transformation which takes care of the time gauge condition. The second unbroken invariance is that of space diffeomorphisms, generated by \mathcal{G}'_1 .

Observe also that \mathcal{G}'_0 and \mathcal{G}'_1 are scalar densities of weight 1 and this ensures that they form a closed Lie algebra (III.45)—in contrast with gravity in higher dimensions where the algebra closes with field dependent structure “constants” [8, 22, 23]. Such a feature is characteristic of 2-dimensional theories with general covariance, such as the bosonic string in the approach of [24].

A new redefinition

$$\mathcal{C}_+ = \frac{\sqrt{-\sigma}}{2} \mathcal{G}'_0 - \frac{1}{2} \mathcal{G}'_1, \quad (\text{III.46})$$

$$\mathcal{C}_- = -\frac{\sqrt{-\sigma}}{2} \mathcal{G}'_0 - \frac{1}{2} \mathcal{G}'_1, \quad (\text{III.47})$$

leads to the algebra

$$\begin{aligned} \{\mathcal{C}_+(\epsilon), \mathcal{C}_+(\eta)\}_D &= \mathcal{C}_+([\epsilon, \eta]), \\ \{\mathcal{C}_-(\epsilon), \mathcal{C}_-(\eta)\}_D &= \mathcal{C}_-([\epsilon, \eta]), \\ \{\mathcal{C}_+(\epsilon), \mathcal{C}_-(\eta)\}_D &= 0. \end{aligned} \quad (\text{III.48})$$

which shows a factorization in two classical Virasoro algebras, as in the $\phi^0 = 0$ gauge of Subsection III A.

IV. QUANTIZATION

Quantization prescriptions take a classical theory as an input and are supposed to give us a quantum theory. As it is well known (and should be expected) this recipe is not complete and leads often to inequivalent quantum theories. Consequently, it is instructive to have explicit examples available to illustrate this point. Here we show that different natural choices lead to inequivalent quantum theories in our ultra simple gauge field theory.

A. Dirac quantization

The physical Hilbert space is given by gauge invariant square integrable functions of G . They are therefore class functions $f(g) = f(aga^{-1})$ for all $g, a \in G$. There are important differences between the

quantization of the Riemannian and Lorentzian cases. Therefore, we shall treat each case separately in this section.

1. *The Riemannian case, $G = \text{SU}(2)$*

This case is treated in great detail in [7, 25]. We briefly review the results here. As in the case of LQG one shifts emphasis from smooth connections to holonomies

$$g_\gamma[\omega] = \text{P exp}\left(-\int_\gamma \omega\right)$$

along oriented paths $\gamma \subset \Sigma$. In the context of the Dirac quantization program one first introduces an auxiliary Hilbert \mathcal{H}_{aux} space where the holonomy $g_\gamma[\omega]$ and the scalar fields ϕ^i are represented as operators. In the present case, once holonomies and ϕ^i have been chosen as fundamental variables, there is a unique choice of \mathcal{H}_{aux} where diffeomorphisms are unitarily implemented [26]. This Hilbert space is given by the Cauchy completion in a appropriate topology of the algebra of functionals of the connection that depend on the holonomy of ω along paths that are edges of arbitrary graphs in Σ (the details of this construction turn out not to be important for our simple model). In a second step one promotes the Gauss constraints (III.2) to self adjoint operators satisfying the appropriate quantum constraint algebra, and finally looks for the physical Hilbert space $\mathcal{H}_{phys} \subset \mathcal{H}_{aux}$ defined by the kernel of the quantum constraints.

Due to the simplicity of our model these steps can be shortcut and we can directly construct the physical Hilbert space \mathcal{H}_{phys} in one stroke. The logic is as follows: The Gauss constraints generate infinitesimal $\text{SU}(2)$ gauge transformations. Elements of \mathcal{H}_{phys} are gauge invariant functions of the Holonomy. As in LQG this restricts the set of possible graphs on which states are defined to closed ones. In our case there is a unique close graph corresponding to the entire initial value surface Σ . Therefore, the physical Hilbert space is given by functions of the holonomy $g[A] \in \text{SU}(2)$ around $\Sigma = S^1$ which are in the kernel of the Gauss constraints (III.2). Those are square integrable functions which are invariant under $\psi(g) = \psi(aga^{-1})$ for any $a \in \text{SO}(3)$ (this invariance is the residual gauge action on the based point from where the holonomy around the Universe is defined). The latter are the so-called class functions $\psi(g) \in \mathcal{L}^2(\text{SO}(3))/\mathcal{G} \subset \mathcal{L}^2(\text{SO}(3))$. Using the Peter-Weyl theorem they can be written as

$$\Psi(g) = \sum_j (2j+1) \psi_j \text{Tr}(D^j[g]), \quad (\text{IV.1})$$

where $D^j[g]$ are unitary irreducible representations of $\text{SU}(2)$, and $\psi_j = \int dg \overline{\text{Tr}(D^j[g])} \Psi[g]$ with dg the Haar measure. Consequently there is a natural construction of the physical Hilbert space based on the following choice of inner product

$$\langle \Psi, \Phi \rangle = \int dg \overline{\Psi(g)} \Phi(g).$$

Any gauge invariant function ($\psi(g) = \psi(aga^{-1}) \in \mathcal{H}_{phys}$) can be expanded in terms of the characters $\chi_j(g) = \text{Tr}[D^j(g)]$. This are the strict analog of the so-called spin-network states of LQG. A fact that will become more important in the Lorentzian case is that class functions can be thought of as functions of the (unique up to conjugation) Cartan subgroup $H \subset \text{SO}(3)$. More concretely, any $\text{SU}(2)$ rotation can be characterized by a $\text{U}(1)$ rotation by an angle $\nu \in [0, \pi]$ around an axis defined by a unit vector $\hat{n} \in R^3$. The latter is entirely defined by a point on the unit 2-sphere labelled by spherical coordinates $\theta \in [0, \pi]$ and $\phi \in [0, 2\pi]$. One can express the $\text{SU}(2)$ Haar measure in these coordinates as follows:

$$dg = \frac{1}{2\pi^2} \sin^2(\nu) (\sin(\theta) d\theta d\phi) d\nu.$$

Now any class function depends only on the coordinate ν , namely $\psi(g) = \psi(\nu)$. Therefore, the inner product above takes the form

$$\langle \psi, \phi \rangle = \frac{2}{\pi} \int_0^\pi d\nu \sin^2(\nu) \overline{\psi(\nu)} \phi(\nu), \quad (\text{IV.2})$$

where we have explicitly performed the θ and ϕ integration. In the Lorentzian case the analog of the last step (in a formal manipulation) leads to a trivial divergence due to the noncompactness of the gauge group. It is clear that one can regularize this divergence by simply dropping such integration. We will see this in detail in the following section.

Incidentally, notice that the characters can be expressed in terms of linear combinations of powers of $\chi_{1/2}(g)$ which shows that the most general gauge invariant functional of the generalized connection is the observable (III.4) $O_2 = \text{Tr}P \exp(-\int_{S^1} \omega)$. The Dirac observable O_1 is quantized by the self-adjoint operator $O_1 = -\hbar^2 \Delta$, where Δ denotes the Laplacian on $SU(2)$, the characters $\chi_j(g) = \text{Tr}[D^j(g)]$ are its eigenstates with eigenvalues $\hbar^2 j(j+1)$, namely

$$O_1|j\rangle = \hbar^2 j(j+1)|j\rangle, \quad (\text{IV.3})$$

where we have used Dirac bracket notation $\chi_j(g) \rightarrow |j\rangle$. The Dirac observable O_2 acts by multiplication, in the spin network basis its action is given by

$$O_2|j\rangle = c_j^{(+)}|j + \frac{1}{2}\rangle + c_j^{(-)}|j - \frac{1}{2}\rangle, \quad (\text{IV.4})$$

where $c_j^{(\pm)}$ are the corresponding Clebsh-Gordon coefficients. The gauge invariant combination O_3 of ϕ^i and $g_\Sigma[\omega]$ can be quantized using the commutator of O_1 and O_2 (Note that the direct quantization of O_3 would require dealing with nontrivial ordering issues).

2. The Lorentzian case, $G = \text{SL}(2, \mathbb{R})$

The Lorentzian case is more involved, and, to our knowledge, has not been described in the literature. The main technical complication is the noncompactness of the gauge group which implies that the standard nonperturbative techniques applicable to standard gauge theories with compact groups need to be revised due to the appearance of divergences in the naive treatment. The main technical complication is due to the fact that, unlike the compact case, class functions $\psi(g) = \psi(aga^{-1})$ are not square integrable functions in the Haar measure of $\text{SL}(2, \mathbb{R})$. The main difficulty with which one needs to deal is the definition of the physical inner product for functions of $\text{SL}(2, \mathbb{R})/\mathcal{G}$ so that appropriate reality conditions are satisfied by the Dirac observables. On the other hand the structure of the theory is richer.

As we saw in the Riemannian case, physical states are characterized by class functions which in turn can be thought off as functions of elements of the Cartan subgroups. The new feature in the Lorentzian sector is that there are two inequivalent (under conjugation) Cartan subgroups in $\text{SL}(2, \mathbb{R})$. On the one hand one has $H_1 = \text{U}(1) \in \text{SL}(2, \mathbb{R})$, given by rotations fixing an internal time axis, namely

$$H_1 = \left\{ g_\nu = \begin{bmatrix} \cos(\nu) & \sin(\nu) \\ -\sin(\nu) & \cos(\nu) \end{bmatrix} \right\}, \quad (\text{IV.5})$$

and $H_2 \in \text{SL}(2, \mathbb{R})$ given by the subgroup of boosts fixing some spacelike internal direction, explicitly

$$H_2 = \left\{ g_\eta = \begin{bmatrix} \exp(\eta) & 0 \\ 0 & \exp(-\eta) \end{bmatrix} \right\}. \quad (\text{IV.6})$$

This means that conjugation $g \rightarrow aga^{-1}$ for fixed $g \in \text{SL}(2, \mathbb{R})$ and arbitrary $a \in \text{SL}(2, \mathbb{R})$ generate orbits (gauge orbits) which are labelled by elements of H_1, H_2 , respectively¹.

Therefore, gauge invariant states $\Psi(g) = \Psi(aga^{-1})$ can be characterized by two functions

$$\Psi[g] = \begin{cases} \psi_1(\nu) & \text{for } [g] \in H_1 \\ \psi_2(\eta) & \text{for } [g] \in H_2 \end{cases}.$$

As mentioned above the noncompactness of the gauge group implies that gauge invariant states $\Psi(g) = \Psi(aga^{-1})$ are not square integrable with respect to the Haar measure. The reason is that physical states

¹ More precisely, this is true for the so called regular elements of $\text{SL}(2, \mathbb{R})$ which are an open subset of the full group with complement of measure zero with respect to the Haar measure [27].

are constant on noncompact adjoint orbits. However, an inner product can be introduced in such a way that these states are normalizable and the appropriate self-adjoint property of observables holds.

This is easily done by mimicking what we did in the previous section when finding an explicit parametrization of the Haar measure in terms of the Cartan subgroup. It is still true that one can write a regular element of $\text{SL}(2, \mathbb{R})$ as Abelian “rotations around an axis \hat{n} ”. The main difference is that the unit vector \hat{n} can be either time like or space like. In other words the (rotationally invariant) 2-sphere of directions is now replaced by the ($\text{SO}(2, 1)$ invariant) time like Hyperboloid h_0 , (given by the points in Minkowski internal spacetime $x^i x^j \eta_{ij} = 1$) together with the future and past space like Hyperboloid h_{\pm} (given by the points in Minkowski internal spacetime² $x^i x^j \eta_{ij} = -1$). The second difference is that “rotations around \hat{n} ” are now standard $\text{U}(1)$ rotations (elements of H_1) only when $\hat{n} \in h_{\pm}$, while they are replaced by boosts (elements of H_2) when $\hat{n} \in h_0$.

Elements of $g \in \text{SL}(2, \mathbb{R})$ can be modelled by points on three dimensional DeSitter spacetime (thought of as embedded in a four dimensional flat spacetime of signature $(++--)$) as follows

$$g = \begin{pmatrix} x^0 + x^3, x^1 + x^2 \\ x^1 - x^2, x^0 - x^3 \end{pmatrix} \quad (x^0)^2 - (x^1)^2 + (x^2)^2 - (x^3)^2 = 1.$$

In terms of these coordinates the invariant measure takes the simple form

$$dg = dx^0 dx^1 dx^2 dx^3 \delta((x^0)^2 - (x^1)^2 + (x^2)^2 - (x^3)^2 - 1).$$

Elements $g \in \text{SL}(2, \mathbb{R})$ equivalent by conjugation to elements in $H_1 \subset \text{SL}(2, \mathbb{R})$ are characterized by $|\text{Tr}[g]| \leq 2$. It is easy to see that these elements can be described in terms of hyperbolic coordinates as

$$\begin{aligned} x^0 &= \cos[\nu] \\ x^1 &= \sin[\nu] \sinh[\rho] \cos[\phi] \\ x^2 &= \sin[\nu] \cosh[\rho] \\ x^3 &= \sin[\nu] \sinh[\rho] \sin[\phi], \end{aligned}$$

where $\rho \in \mathbb{R}^+$ and $\phi \in [0, 2\pi]$ label points on h_{\pm} . For these regular elements the invariant measure becomes

$$dg = \sinh(\rho) \sin^2(\nu) d\nu d\rho d\phi, \quad \text{for } [g] \in H_1. \quad (\text{IV.7})$$

Elements $g \in \text{SL}(2, \mathbb{R})$ equivalent by conjugation to elements in $H_2 \subset \text{SL}(2, \mathbb{R})$ are characterized by $|\text{Tr}[g]| \geq 2$. In terms of hyperbolic coordinates they are characterized by

$$\begin{aligned} x^0 &= \cosh[\eta] \\ x^1 &= \sinh[\eta] \sin[\theta] \cosh[\rho] \\ x^2 &= \sinh[\eta] \sin[\theta] \sinh[\rho] \\ x^3 &= \sinh[\eta] \cos[\theta], \end{aligned}$$

where $\rho \in \mathbb{R}^+$ and $\theta \in [0, \pi]$ label points on h_0 . Explicitly we have $\text{Tr}[g] = 2 \cosh[\eta]$. The invariant measure becomes

$$dg = \sin(\theta) \sinh^2(\eta) d\eta d\rho d\phi, \quad \text{for } [g] \in H_2. \quad (\text{IV.8})$$

Indeed the physical Hilbert space $\mathcal{H}_{phys} = \mathcal{H}_1 \oplus \mathcal{H}_2$ is such that $\Psi(g) = (\psi_1(\nu), \psi_2(\eta)) \in \mathcal{H}_{phys}$ with $\psi_1(\nu) \in \mathcal{H}_1$ and $\psi_2(\eta) \in \mathcal{H}_2$. The inner product is respectively:

$$\langle \psi, \phi \rangle_1 = \int_0^{2\pi} d\nu \sin^2(\nu) \bar{\psi}(\nu) \phi(\nu), \quad \text{and} \quad \langle \psi, \phi \rangle_2 = \int_0^{\infty} d\eta \sinh^2(\eta) \bar{\psi}(\eta) \phi(\eta),$$

² Minkovski metric is $\eta_{ij} = \text{diag}(-1, 1, 1)$.

where the integration measure is defined by dropping the redundant integrations from the invariant measures (IV.7) and (IV.8) respectively. The two Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 are super-selection sectors. Harmonic analysis on $\text{SL}(2, \mathbb{R})$ implies that any function $f(g) \in \mathcal{L}^2(\text{SL}(2, \mathbb{R}))$ can be written as:

$$f(g) = \sum_{n \geq 1} (2n-1) \text{Tr}(f_n^+ D_n^+[g]) + \sum_{n \geq 1} (2n-1) \text{Tr}(f_n^- D_n^-[g]) + \int_0^\infty ds \mu(s) \text{Tr}(f_s D_s[g]). \quad (\text{IV.9})$$

As in the Euclidean case, physical states can be spanned in terms of characters. From the point of view of the space $\mathcal{L}^2(\text{SL}(2, \mathbb{R}))$ these states are distributional. The restriction of the characters to H_1 and H_2 give an orthonormal basis of \mathcal{H}_1 and \mathcal{H}_2 respectively of eigen-states of the Dirac observable $O_1 = \phi \cdot \phi = -\hbar^2 \Delta$. In \mathcal{H}_1 the Laplacian takes the explicit form $\Delta = \sin(\nu)^{-1} \partial_\nu^2 \sin(\nu) + 1/4$ and the characters are $\chi_n^\pm(\nu) = \pm \exp(\pm i(n-1)\nu)/(2i \sin(\nu))$ and the eigenvalues of O_1 are given by $\hbar^2(n - \frac{1}{2})(n - \frac{3}{2})$. Therefore, the spin network states $\chi_n^\pm(\nu) \rightarrow |n, \pm\rangle$, which satisfy

$$\langle n, +|m, +\rangle_1 = \langle n, -|m, -\rangle_1 = \delta_{nm} \quad \text{and} \quad \langle n, +|m, -\rangle_1 = 0,$$

are a basis for \mathcal{H}_1 diagonalizing O_1

$$O_1 |n\rangle = -\hbar^2(n - \frac{1}{2})(n - \frac{3}{2}) |n\rangle. \quad (\text{IV.10})$$

The Dirac observable O_2 acts by multiplication by $O_2 = 2 \cos(\nu)$ in \mathcal{H}_1 . In the spin network basis its action is given by

$$O_2 |n, \pm\rangle = |n \pm 1, \pm\rangle + |n \mp 1, \pm\rangle. \quad (\text{IV.11})$$

In \mathcal{H}_2 the Laplacian takes the explicit form $\Delta = -\sinh(\eta)^{-1} \partial_\eta^2 \sinh(\eta) + 1/4$ and the characters are $\chi_s(\eta) = \pm \cos(s\eta)/|\sinh(\eta)|$ and the eigenvalues of O_1 are given by $\hbar^2(s^2 + \frac{1}{4})$. Now spin network states are labelled by a real parameter $\chi_s(\nu) \rightarrow |s\rangle$. We have

$$O_1 |s\rangle = \hbar^2(s^2 + \frac{1}{4}) |s\rangle, \quad (\text{IV.12})$$

The Dirac observable $O_2 = \cosh(\eta)$ acts by multiplication. Its action on spin network states is not a spin network state.

B. Time-gauge quantization

The purpose of [11] and [12] is a quantization along the lines of loop quantization in 1+3 dimensions with a time-gauge fixing, in the presumably simpler case of the (1+1)-dimensional Jackiw-Teitelboim model. However, the procedure still remains somewhat more complicated than in the other cases studied in the present paper. The construction of the kinematical Hilbert space is based on the wave functionals defined on the configuration space spanned by the "holonomies" of the scalar fields ϕ^1 and ϕ^2 defined in Subsection III C, the "polymer-like" scalar product used there leading to nonseparability of the Hilbert space [28]. The conjugate fields ω^1 and ω^2 are represented as functional differential operators, which are diagonal in a spin-network-like orthonormal basis.

The construction of operators representing the classical constraints (III.44) goes through a cell regularization, and the hope is to check the algebra (III.45) at the quantum level, the final task being that of solving the quantum constraints.

C. Quantization in the $\phi \cdot n = 0$ gauge.

Recall that the $\phi \cdot n = 0$ gauge in the Lorentzian case, as described classically in Section (III A), lets us with two first class constraints C_0 and C_1 (III.12) in the $\phi \cdot \phi > 0$ case, C_0 generating U(1) internal rotations in the (ϕ^1, ϕ^2) plane, and C_1 a boost leaving the vector $\phi = (0, \phi^1, \phi^2)$ invariant. In the case $\phi \cdot \phi < 0$, we are left with one constraint g_0 (III.21) generating U(1) internal rotations leaving the vector $\phi = (\phi, 0, 0)$ invariant. In the former case, to the transformation generated by C_1 is associated the Abelian connection Q (III.5). In the latter case, to the U(1) invariance is associated the connection $\omega = \omega^0$. In the spirit of loop quantization, one must thus choose the holonomies H_I of Q or ω along intervals I of S^1

as the configuration space variables (from the connections we go to the generalized connections). These Abelian holonomies are given by

$$H_I = \exp\left(-\int_I dx \mathcal{T}(x) \tau_\star\right), \quad (\text{IV.13})$$

with

$$\begin{aligned} \mathcal{T} = \mathcal{Q}, \quad \tau_\star &= \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} && \text{for the sector } \phi \cdot \phi > 0, \\ \mathcal{T} = \omega, \quad \tau_\star &= \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} && \text{for the sector } \phi \cdot \phi < 0. \end{aligned}$$

Notice that these are the generators whose exponentiation leads to group elements of the form IV.6 (a boost) and IV.5 (a rotation), respectively. The holonomy $H = H_{S^1}$ is obviously a Dirac observable, element of the adjoint representation of the group generated by C_1 or g_0 .

Let us now discuss the two sectors separately.

1. Sector $\phi \cdot \phi > 0$

It will be convenient to use the invariant (under the action of C_0) variables (III.27), namely

$$\Pi \equiv \frac{1}{2} \phi^A \phi^A \quad \text{and} \quad Q \equiv \frac{\phi^A \omega^A}{\phi^C \phi^C}.$$

The advantage of these variables is that Π is nothing else but O_1 (III.4), one of the Dirac observables, and Q is the Abelian connection discussed above, transforming under the remaining Abelian gauge symmetry generated by the constraint C_1 (boosts) as:

$$\{Q, C_1(b)\}_D = -\partial b.$$

(See (III.12, III.14)). A finite transformation of H_I as given by (IV.13) reads:

$$H'_I = \exp(\tau_\star(b_t - b_s)) H_I,$$

where b_s and b_t are the values of b at the ends (source/target) of the interval I . Therefore, the holonomy around the space, $H := H_{S^1}$, being gauge invariant is a Dirac observable. It takes the form

$$H = \exp(-\tau_\star \eta) = \begin{bmatrix} \exp(\eta) & 0 \\ 0 & \exp(-\eta) \end{bmatrix}, \quad \text{with } \eta = \int_{S^1} dx Q(x) \text{ and } \tau_\star \equiv \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}. \quad (\text{IV.14})$$

From the classical Dirac bracket algebra (III.13) follows

$$\{\Pi(x), H\}_D = \tau_\star H, \quad (\text{IV.15})$$

and consequently

$$\{\Pi, \eta\}_D = -1. \quad (\text{IV.16})$$

This suggests to take Π and η as the classical phase space coordinates, with $\Pi \in \mathbb{R}_+$ (real positive) and $\eta \in \mathbb{R}$. (Owing to the constancy of Π following from the constraint C_1 , we can take for Π the value of $\Pi(x)$ at any point $x \in C_1$.)

We can quantize this sector in the conventional Schroedinger scheme. Here, elements of the physical Hilbert space are functions $\Phi(\eta)$ —with η defined in (IV.14)—belonging to $L_2(\mathbb{R}, d\eta)$, and Π is represented by the operator $\hat{\Pi} = -i\hbar d/d\eta$. Eigenfunctions of $\hat{\Pi}$ are unnormalizable “plane waves” $\exp(i\rho\eta/\hbar)$. Restricting to “wave packets” of positive frequency $\rho > 0$ we recover the set of positive real number as the (now continuous) spectrum of $\hat{\Pi}$. The spectrum of $\Pi = O_1$ is thus constituted by the positive real numbers in agreement with the results of the previous section, but it does not quite coincide with it for small values of s , as there is no discrete gap here (in contrast with (IV.12)). However there is agreement in the asymptotic regime.

Alternatively, as H is an element of the Abelian boost group, parametrized by the real number η one could explore the quantization based on the so-called polymer representations [17, 18]. Our task is to construct a physical Hilbert space \mathcal{H}_{phys} as a representation of the quantum algebra $[\hat{\Pi}, \hat{\eta}] = -i\hbar$ or, better:

$$[\hat{\Pi}, \hat{h}] = \hbar \hat{h}, \quad \text{where } h = \exp(i\eta), \quad (\text{IV.17})$$

corresponding to the classical algebra (IV.16). The elements of \mathcal{H}_{phys} will be taken as functions of the boost group, which we parametrize by real numbers ρ considered as elements of the Bohr compactification [29] \mathbb{R}_B of \mathbb{R} . Accordingly, the integration measure for the “almost periodic functions”

$$\Phi_s(\rho) = \exp\left(-\frac{i}{\hbar}s\rho\right), \quad s \in \mathbb{R},$$

is given by

$$\int_{\mathbb{R}_B} d\mu(\rho) \Phi_s(\rho) = \delta_{s0}. \quad (\text{IV.18})$$

The “cylindrical vector space” \mathcal{H}_{cyl} is defined as the set of all finite linear combinations of almost periodic functions, and a hermitean scalar product is defined with the help of the integration measure (IV.18). Hence, in particular:

$$\langle \Phi_s | \Phi_t \rangle = \delta_{st}. \quad (\text{IV.19})$$

The action of the operator $\hat{\eta}$ is defined through the action of its almost periodic counterparts $\Phi_t(\rho)$ for any $t \in \mathbb{R}$:

$$\hat{\Phi}_t \Phi_s(\rho) = \Phi_{t+s}(\rho),$$

whereas that of $\hat{\Pi}$ is defined by

$$\hat{\Pi} \Phi_s(\rho) = -i\hbar \frac{d}{d\rho} \Phi_s(\rho),$$

in agreement with the algebra (IV.17).

We observe that the Φ_s are eigenvectors of $\hat{\Pi}$ with eigenvalue s . Since this operator owes to be positive beyond of being self-adjoint, we define \mathcal{H}_{cyl+} as the space generated by the restricted basis $\{\Phi_s; s \geq 0\}$ —in analogy with the separation of the positive and negative frequency parts in the relativistic quantum theory of free fields. Finally, the physical Hilbert space \mathcal{H}_{phys} is defined as the Cauchy completion of \mathcal{H}_{cyl+} with respect to the norm induced by the scalar product (IV.19). Possessing an uncountable orthonormal basis, \mathcal{H}_{phys} is nonseparable. We note that the spectrum of $\hat{\Pi}$ coincide with the one found in the Schroedinger representation, yet it must be considered as “discrete” as the corresponding eigenvectors have finite norm.

2. Sector $\phi \cdot \phi < 0$

In this sector, as discussed in Subsection (III A 2), the classical phase space variables are $\omega = \omega^0$ and $\phi = \phi^0$, obeying the canonical Dirac bracket algebra (III.20). The constraint g_0 (III.21) lets ϕ to be a constant, and ω transforms under g_0 as the connection associated to the residual group $U(1)$ of gauge transformations preserving the gauge fixing conditions $\phi^1 = \phi^2 = 0$. Thus the classical Dirac observables are ϕ —taken at an arbitrary valor of the space coordinate x —and the holonomy along the space slice:

$$H = \exp(\tau, \xi) = \begin{bmatrix} \cos(\xi) & \sin(\xi) \\ -\sin(\xi) & \cos(\xi) \end{bmatrix}, \quad \xi = \int_{S^1} dx \omega(x), \quad \tau_* = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

It is convenient to perform the quantization in terms of the variables ϕ and ξ . From now on we switch emphasis from $H(\xi) \rightarrow h(\xi) := \exp(i\xi)$. From the classical Dirac bracket $\{\phi(x), h\}_D = ih$, we define the corresponding quantum commutator as

$$[\hat{\phi}, \hat{h}] = \hbar \hat{h}.$$

With this choice of variables, elements of the physical Hilbert space are continuous functions of $U(1)$, i.e., $\Phi(\theta)$, with $\theta \in [0, 2\pi]$, and $h(\theta)$ acts simply by multiplication. There is a spin network basis given by the unitary irreducible representations of $U(1)$, explicitly:

$$\Phi_n(\theta) = \frac{1}{\sqrt{2\pi}} \exp(in\theta), \quad \text{for all } n \in \mathbb{Z}.$$

The spectrum of $\hat{\phi}$ is discrete, given by its eigenvalues $n\hbar$. Thus the spectrum of $O_1 = \hat{\phi} \cdot \hat{\phi}$ is in agreement with the results of the previous section but has eigenvalues that differ from those corresponding to (IV.10) for small value of n . However, the eigenvalues of O_1 approach each other in the large (in Planck units) eigenvalue limit.

D. Quantization after totally reducing

The last equation tells us that the reduced phase space of the Jackiw-Teitelboim model corresponds to that of a system with a single degree of freedom. So this is just standard one dimensional quantum mechanics. The physical Hilbert space can be taken as $\mathcal{H}_{ph} \equiv \mathcal{L}^2(\mathbb{R})$. Wave functions are continuous functions $\Psi(\mathcal{X})$ upon which \mathcal{X} act by multiplication and $\mathcal{P} = -i\hbar\partial/\partial\mathcal{X}$. Notice that there is no discreteness to be found if we use the standard representation.

V. CONCLUSIONS

This paper is divided in two parts. In the first part we performed the canonical analysis of the JT model and studied the phase space structure of the theory. We showed that there are dynamically independent sectors corresponding to the cases $\phi \cdot \phi > 0$ (space-like), $\phi \cdot \phi = 0$ (null) and $\phi \cdot \phi < 0$ (time-like). The system has no local degrees of freedom. However, for $M = S^1 \times \mathbb{R}$ there is one global topological degree of freedom in the time-like and space-like sectors respectively. In the null sector no global degree of freedom is present no matter the spacetime topology.

A partial gauge fixing allows, in the space-like case, to reduce the DeSitter gauge symmetry of the JT model to the two dimensional diffeomorphism invariance of gravity. We explicitly showed in that case how, after partial gauge fixing, the remaining first class constraints relate to the generators of two Virasoro symmetries, and the familiar diffeomorphism and scalar constraints of gravitational theories.

In the second part we studied the quantization of the JT model using background independent techniques. We first performed the quantization of the model without the introduction of gauge fixing. Even though this was well known in the Riemannian case, the Lorentzian case presented some technical difficulties related to the noncompactness of the gauge group. The difficulties were overcome using group averaging techniques which naturally lead to two possible Hilbert space representation of the fundamental observables. These inequivalent quantum theories have both physical interpretation as they are in one-to-one correspondence with the superselection sectors $\phi \cdot \phi > 0$ (space-like), and $\phi \cdot \phi < 0$ (time-like).

Alternatively, one can partially (or totally) reduce the gauge freedom of the system at the classical level by introducing gauge conditions, and explore afterwards the quantization of the reduced system. We have explicitly shown that representations of the basic fields obtained following this alternative avenue are not equivalent to the previous ones. The main results of the second part are summarized in Table (I), which shows the various inequivalent sectors of the quantum Jackiw-Teitelboim theory. We have called

	Riemannian	Lorentzian	
	Spect. O_1	Spect. O_1 in Sector I	Spect. O_1 in Sector II
No gauge-fixing	$\hbar^2 n(n+1)$ for $n \in \mathbb{Z}/2$	$-\hbar^2(n - \frac{1}{2})(n - \frac{3}{2})$ for $n \in \mathbb{Z}$	$\hbar^2(s^2 + \frac{1}{4})$ for $s \in \mathbb{R}^+$
$n \cdot \phi = 0$ gauge	$\hbar^2 n^2$ for $n \in \mathbb{Z}$	$-\hbar^2 n^2$ for $n \in \mathbb{Z}$	$\hbar^2 s^2$ for $s \in \mathbb{R}^+$
Full reduction	\mathbb{R}^+ , continuous	\mathbb{R}^- , continuous	\mathbb{R}^+ , continuous
Time gauge	?	?	?

TABLE I: Spectrum of the observable O_1 defined in Eq. (III.4), according to various quantization schemes.

I and II the sectors (phases) of the Lorentzian theory which are described by the Hilbert spaces \mathcal{H}_I and

\mathcal{H}_2 in the no gauge fixing part (Section IV A), and by the $\phi \cdot \phi < 0$ and $\phi \cdot \phi > 0$ sectors in the $n \cdot \phi = 0$ gauge (Section IV C).

We observe that the spectrum of O_1 in the nongauge-fixed and $n \cdot \phi$ -gauge coincide in the large eigenvalue limit in both the Riemannian and the Lorentzian phase I. This is compatible with a common semi-classical limit. But, in the Lorentzian phase II the spectrum is that of all positive real numbers, “discrete” or “continuous” depending on the quantization being “polymeric” or of the “Schrodinger” type. Finally, the spectrum of O_1 is completely changed (continuous) in the fully reduced case, for the Riemannian and for both phases of the Lorentzian theory. The gauge group structure vanishing in this case implies that the kind of representations used in the former cases are not even available. As discussed in Section IV B, one could use a polymer like representation to recover a “discrete” spectrum but the microscopic Planckian structure is lost in the fully reduced setting.

-
- [1] R. Jackiw, in “Quantum Theory of Gravity”, edited by S. Christensen (Hilger, Bristol,1984); C. Teitelboim, Phys. Lett. B126198341; C. Teitelboim in “Quantum Theory of Gravity”, edited by S. Christensen (Hilger, Bristol,1984).
 - [2] T. Fukuyama and K. Kamimura, “Gauge Theory of Two-Dimensional Gravities”, Phys. Lett. 160B (1985) 259.
 - [3] Marc Henneaux, “Quantum Gravity in Two-Dimension: Exact Solution of Jackiw Model”, Phys. Rev. Lett. 54 (1985) 959.
 - [4] A.H. Chamseddine and D. Wyler, “Gauge theory of topological gravity in 1+1 dimensions”, Phys. Lett. B228198975.
 - [5] W. Kummer, H. Liebl, and D. V. Vassilevich, “Exact path integral quantization of generic 2-d dilaton gravity”, Nucl. Phys. B493 (1997) 491.
 - [6] W. Kummer, H. Liebl, and D. V. Vassilevich, “Integrating geometry in general 2d dilaton gravity with matter”, Nucl. Phys. B544 (1999) 403.
 - [7] E.R. Livine, Alejandro Perez, C. Rovelli, “2D manifold-independent spinfoam theory”, Class. Quantum Grav. 20 (2003) 4425, arXiv:gr-qc/0102051.
 - [8] A. Ashtekar and J. Lewandowski, “Background independent quantum gravity: A Status report”, Class. Quantum Grav. 21 (2004) R53, e-Print Archive: gr-qc/0404018; A. Perez, “Introduction to loop quantum gravity and spin foams,” Lectures at the 2nd International Conference on Fundamental Interactions, Brazil, arXiv:gr-qc/0409061.
 - [9] D. Grumiller, W. Kummer and D.V. Vassilevich, Phys. Rep. 369 (2002) 327
 - [10] L. Bergamin and R. Meyer, “Two-Dimensional Quantum Gravity with Boundary”, To appear in the proceedings of 4th Advanced Research Workshop: Gravity, Astrophysics, and Strings at the Black Sea, Kiten, Bourgas, Bulgaria, 10-16 Jun 2007, arXiv:0711.3595 [hep-th].
 - [11] C.P. Constantinidis, J.A. Lourenço, I. Morales, O. Piguet and Alex Rios, “Canonical Analysis of the Jackiw-Teitelboim Model in the Temporal Gauge. I. The Classical Theory”, Class. Quantum Grav. 25 (2008) 125003, arXiv:0802.0112 [gr-qc].
 - [12] C.P. Constantinidis, J.A. Lourenço and O. Piguet, work in progress.
 - [13] D. Birmingham, M. Blau, M. Rakowski and G.T. Thompson, “Topological Field Theory”, Phys.Rept. 209:129-340,1991. M. Blau and G. Thompson, “Topological Gauge Theories of Antisymmetric Tensor Field”, Ann. Phys. 205(1991) 130-172.
 - [14] L. Freidel and E.R. Livine, “Spin networks for noncompact groups”, J. Math. Phys. 44 (2003) 1322, arXiv:hep-th/0205268.
 - [15] A. Gomberoff and D. Marolf, “On group averaging for SO(n,1),” Int. J. Mod. Phys. D 8 (1999) 519 [arXiv:gr-qc/9902069].
 - [16] J.L. Willis, “On the Low-Energy Ramifications and a Mathematical Extension of Loop Quantum Gravity”, Ph. D. Thesis (2005), Penn State Univ. (available at <http://www.gravity.psu.edu/research/archives/thesis/index.html>).
 - [17] A. Ashtekar, S. Fairhurst, J.L. Willis, ”Quantum Gravity, Shadow States and Quantum Mechanics”, Class. Quantum Grav. 20 (2003) 1031, arXiv:gr-qc/0207106v3.
 - [18] A. Corichi, T. Vukasinac, J.A. Zapata, ”Hamiltonian Space in polymer quantum mechanics”, Class. Quantum Grav. 24 (2007) 1495, arXiv:gr-qc/0610072, A. Corichi, T. Vukasinac, J.A. Zapata, ”Polymer Quantum Mechanics and its Continuum Limit”, Phys. Rev. D76 (2207) 044016, arXiv:0704.0007 [gr-qc].
 - [19] P.A.M. Dirac, “Lectures on Quantum Mechanics”, Belfer Graduate School of Science, Yeshiva University, 1964.
 - [20] M. Henneaux and C. Teitelboim, “Quantization of Gauge Systems”, Princeton University Press, 1991.
 - [21] D.M. Gitman and I.V. Tyutin, “Quantization of Fields with Constraints”, Springer-Verlang Series in Nuclear

and Particle Physics, Berlin Heidelberg, 1990.

- [22] C. Rovelli, "Quantum Gravity", Cambridge Monography on Math. Physics (2004).
- [23] T. Thiemann, "Modern Canonical Quantum General Relativity", Cambridge Monographs on Mathematical Physics (2007).
- [24] T. Thiemann, "The LQG string: Loop quantum gravity quantization of string theory I: Flat target space", *Class. Quantum Grav.* 23 (2006) 1923, arXiv: hep-th/0401172.
- [25] K. Isler and C.A. Trugenberger, "Gauge Theory of Two-Dimensional Quantum Gravity", Phys. Rev. Lett. 631989834.
- [26] J. Lewandowski, A. Okołow, H. Sahlmann and T. Thiemann, "Uniqueness of diffeomorphism invariant states on holonomy flux algebras", *Commun. Math. Phys.* 267 (2006) 703, arXiv:gr-qc/0504147.
- [27] W. Rühl, "The Lorentz group and harmonic analysis", W.A. Benjamin Inc., New York, 1970.
- [28] A. Ashtekar, J. Lewandowski, H. Sahlmann, "Polymer and Fock representations for a scalar field", *Class. Quantum Grav.* 20 (2003) L11-1, e-Print: arXiv:gr-qc/0211012.
- [29] W. Rudin, "Functional Analysis", McGraw-Hill Book Company, 1973.