

The Uniqueness Problem of Sequence Product on Operator Effect Algebra $\varepsilon(H)$

Weihua Liu and Junde Wu

Department of Mathematics, Zhejiang University, Hangzhou 310027, P. R. China.

Let H be a complex Hilbert space. A quantum effect on H is a bounded linear self-adjoint operator A defined on H that satisfies $0 \leq A \leq I$. We denote the set of all quantum effects on H by $\varepsilon(H)$. The sequential product \circ defined on $\varepsilon(H)$ is an important concept because it can be used to describe the quantum measurements. A sequential product defined by $A \circ B = A^{\frac{1}{2}}BA^{\frac{1}{2}}$ for any two quantum effects A, B has recently been introduced and studied. In 2005, Professor Gudder presented an open problem as following: Is $A \circ B = A^{\frac{1}{2}}BA^{\frac{1}{2}}$ the only sequential product on $\varepsilon(H)$? In this paper, for $\dim H = 2$, we construct a new sequential product on $\varepsilon(H)$ which is different from $A \circ B = A^{\frac{1}{2}}BA^{\frac{1}{2}}$. Thus, we answer the Gudder's open problem negatively. In particular, our result shows the phase-changed phenomena of quantum measurement $A \circ B$, this is much more important in physics.

PACS numbers: 02.10-v, 02.30.Tb, 03.65.Ta