

Does a functional integral really need a Lagrangian?

Denis Kochan*

Department of Theoretical Physics,
FMFI, Comenius University,
842 48 Bratislava, Slovakia
(Dated: November 5, 2018)

Path integral formulation of quantum mechanics (and also other equivalent formulations) depends on a Lagrangian and/or Hamiltonian function that is chosen to describe the underlying classical system. The arbitrariness presented in this choice leads to a phenomenon called *Quantization ambiguity*. For example both $L_1 = \dot{q}^2$ and $L_2 = e^{\dot{q}}$ are suitable Lagrangians on a classical level ($\delta L_1 = \delta L_2$), but quantum mechanically they are diverse.

This paper presents a simple rearrangement of the path integral to a surface functional integral. It is shown that the surface functional integral formulation gives transition probability amplitude which is free of any Lagrangian/Hamiltonian and requires just the underlying classical equations of motion. A simple example examining the functionality of the proposed method is considered.

Dedicated to my friend and colleague Pavel Bóna.

PACS numbers: 03.65.Ca, 11.10.Ef, 31.15.xk

Keywords: quantization of (non-)Lagrangian systems, path vs. surface functional integral

I. A STANDARD PATH INTEGRAL LORE

According to Feynman [1], the probability amplitude of the transition of a system from the space-time configuration (q_0, t_0) to (q_1, t_1) is given as follows:

$$\mathbf{A}(q_1, t_1 | q_0, t_0) \propto \int_{\tilde{\gamma}} [\mathcal{D}\tilde{\gamma}] \exp \left\{ \frac{i}{\hbar} \int p_a dq^a - H dt \right\}. \quad (1)$$

Here the path-summation is taken over all trajectories $\tilde{\gamma}(t) = (\tilde{q}(t), \tilde{p}(t), t)$ in the extended phase space which are constrained as follows:

$$\begin{aligned} \tilde{\gamma}(t_0) &= (\tilde{q}(t_0) = q_0, \tilde{p}(t_0) - \text{arbitrary}, t_0), \\ \tilde{\gamma}(t_1) &= (\tilde{q}(t_1) = q_1, \tilde{p}(t_1) - \text{arbitrary}, t_1). \end{aligned}$$

To obtain a proper normalization of the Feynman propagator, one requires:

$$\begin{aligned} \delta(\tilde{q}_0 - q_0) &= \int_{\mathbf{R}^n[q_1]} dq_1 \mathbf{A}^*(q_1, t_1 | \tilde{q}_0, t_0) \mathbf{A}(q_1, t_1 | q_0, t_0), \\ \delta(q_1 - q_0) &= \lim_{t_1 \rightarrow t_0} \mathbf{A}(q_1, t_1 | q_0, t_0). \end{aligned}$$

The first equation asks for the conservation of the total probability and the second expresses the obvious fact that no evolution takes place whenever t_1 approaches t_0 .

It is a miraculous consequence (not a requirement!) of the propagator definition (1) that it satisfies an evolutionary chain rule (Chapman-Kolmogorov equation)

$$\mathbf{A}(q_1, t_1 | q_0, t_0) = \int_{\mathbf{R}^n[q]} dq \mathbf{A}(q_1, t_1 | q, t) \mathbf{A}(q, t | q_0, t_0),$$

whose infinitesimal version is the celebrated Schrödinger equation.

It is a curious fact that Formula (1) was not originally discovered by Feynman. In his pioneering paper [2] he arrives at a functional integral in the configuration space only

$$\mathbf{A}(q_1, t_1 | q_0, t_0) \propto \int [\mathcal{D}q] \exp \left\{ \frac{i}{\hbar} \int_{q(t)} L(q, \dot{q}, t) dt \right\}. \quad (2)$$

Later, however, it was shown that this formula represents a very special case of the most general prescription (1). Formula (1) is at the heart of our further discussion.

II. DEHAMILTONIANIZATION

A step beyond involves eliminating the Hamiltonian function H from Formula (1). The price to be paid for this will be to replace the path summation therein by a surface functional integration.

Our aim is the transition probability amplitude between (q_0, t_0) and (q_1, t_1) . Let us suppose that there exists a unique classical trajectory in the extended phase space $\gamma_{cl}(t) = (q_{cl}(t), p_{cl}(t), t)$ which connects these points (locally this assumption is always satisfied). Then for any curve $\tilde{\gamma}(t) = (\tilde{q}(t), \tilde{p}(t), t)$ which enters the path integration in (1), we can assign two auxiliary curves which we call $\lambda_0(s)$ and $\lambda_1(s)$. They are parameterized by $s \in [0, 1]$ and specified as follows:

$$\begin{aligned} \lambda_0(s) &= (q_0, \pi_0(s), t_0) \text{ where } \pi_0(0) = p_{cl}(t_0), \pi_0(1) = \tilde{p}(t_0), \\ \lambda_1(s) &= (q_1, \pi_1(s), t_1) \text{ where } \pi_1(0) = p_{cl}(t_1), \pi_1(1) = \tilde{p}(t_1). \end{aligned}$$

Let us emphasize that neither $\lambda_0(s)$ nor $\lambda_1(s)$ varies with respect to the q and t coordinates in the extended phase space. They are allowed to evolve only with respect to

*Electronic address: kochan@fmph.uniba.sk

the momentum variables. There are, of course, infinitely many of such curves, but as we will see nothing in the theory will be dependent on a particular choice of $\lambda_0(s)$ and $\lambda_1(s)$.

Using these curves one can write:

$$\int_{\tilde{\gamma}} p_a dq^a - H dt = \int_{\gamma_{cl}} p_a dq^a - H dt + \oint_{\partial\Sigma} p_a dq^a - H dt, \quad (3)$$

where $\partial\Sigma = \tilde{\gamma} - \lambda_1 - \gamma_{cl} + \lambda_0$ is a contour spanned by four curves $\tilde{\gamma}(t)$, $\gamma_{cl}(t)$, $\lambda_0(s)$, $\lambda_1(s)$ counting their orientations.

The first integral on the right is the classical action $\mathbf{S}_{cl}(q_1, t_1 | q_0, t_0)$. While the contour integral in (3) can be rearranged to represent a surface integral:

$$\begin{aligned} \oint_{\partial\Sigma} p_a dq^a - H dt = \\ \int_{\Sigma} dp_a \wedge \left(dq^a - \frac{\partial H}{\partial p_a} dt \right) - \frac{\partial H}{\partial q^a} dq^a \wedge dt =: \int_{\Sigma} \Omega. \end{aligned} \quad (4)$$

Surface Σ spanning the contour $\partial\Sigma$ is understood here as a map from the parametric space $(t, s) \in [t_0, t_1] \times [0, 1]$ to the extended phase space, i.e.

$$\Sigma : (t, s) \mapsto (q^a(t, s), p_a(t, s), t(t, s) = t).$$

Partial derivatives of the initial Hamiltonian function can be substituted using the velocity-momentum relations and classical equations of motion:

$$\frac{\partial H}{\partial p_a} = T^{ab} p_b (= \dot{q}^a) \quad \text{and} \quad \frac{\partial H}{\partial q^a} = -F_a (= -\dot{p}_a).$$

Here we consider the physically mostly relevant situation only. In this case the velocities and momenta become related linearly by the metric tensor $T_{ab}(q)$ (and its inverse) defined by the kinetic energy $T = \frac{1}{2} T_{ab} \dot{q}^a \dot{q}^b = \frac{1}{2} T^{ab} p_a p_b$ of the system, then

$$\Omega = dp_a \wedge dq^a - \left(T^{ab} p_a dp_b - F_a dq^a \right) \wedge dt. \quad (5)$$

This object represents a canonical two-form in the extended phase space. It is a straightforward generalization of the standard closed two-form $d\theta = dp \wedge dq - dH \wedge dt$ to the case when the forces are not potential-generated.

It is clear that for a given pair of trajectories $(\tilde{\gamma}, \gamma_{cl})$ there exists infinitely many Σ surfaces. They form a set which we call $\mathcal{U}_{\tilde{\gamma}}$. Since the surface integral $\int_{\Sigma} \Omega$ is only boundary dependent and Formulas (3) and (4) are satisfied, we can write:

$$\exp \left\{ \frac{i}{\hbar} \int_{\tilde{\gamma}} p_a dq^a - H dt \right\} = \frac{e^{\frac{i}{\hbar} \mathbf{S}_{cl}}}{\infty_{\tilde{\gamma}}} \int_{\mathcal{U}_{\tilde{\gamma}}} [\mathcal{D}\Sigma] \exp \left\{ \frac{i}{\hbar} \int_{\Sigma} \Omega \right\}.$$

Here $\infty_{\tilde{\gamma}}$ stands for the number of elements pertaining to the corresponding stringy set $\mathcal{U}_{\tilde{\gamma}}$. Assuming no topological obstructions from the side of the extended phase

space, $\infty_{\tilde{\gamma}}$ becomes an infinite constant independent of $\tilde{\gamma}$. Taking all of this into account we can rewrite (1) as follows:

$$\mathbf{A}(q_1, t_1 | q_0, t_0) \propto e^{\frac{i}{\hbar} \mathbf{S}_{cl}} \int_{\mathcal{U}} [\mathcal{D}\Sigma] \exp \left\{ \frac{i}{\hbar} \int_{\Sigma} \Omega \right\}. \quad (6)$$

In this formula the undetermined normalization constant ∞ was included into the integration measure $[\mathcal{D}\Sigma]$ and the path integral over $\tilde{\gamma}$'s was converted to the surface functional integral as was promised:

$$\int_{\mathcal{U}_{\tilde{\gamma}}} [\mathcal{D}\tilde{\gamma}] \int_{\mathcal{U}_{\tilde{\gamma}}} [\mathcal{D}\Sigma] \dots = \int_{\bigcup_{\tilde{\gamma}} \mathcal{U}_{\tilde{\gamma}}} [\mathcal{D}\Sigma] \dots =: \int_{\mathcal{U}} [\mathcal{D}\Sigma] \dots$$

The set $\mathcal{U} = \bigcup_{\tilde{\gamma}} \mathcal{U}_{\tilde{\gamma}}$ over which the functional integration is carried out contains all extended phase space strings which are anchored to the given classical trajectory γ_{cl} .

To eliminate Hamiltonian H completely we need to express $\mathbf{S}_{cl}(q_1, t_1 | q_0, t_0)$ in terms of the force field. Such a quantity may not exist in general, however we will see that in special cases one can recover an appropriate analog of $\mathbf{S}_{cl}(q_1, t_1 | q_0, t_0)$ requiring a certain behavior of

$$\mathbf{A}(q_1, t_1 | q_0, t_0) \propto e^{\frac{i}{\hbar} \mathbf{S}_{cl}} \int_{\mathcal{U}} [\mathcal{D}\Sigma] \exp \left\{ \frac{i}{\hbar} \int_{\Sigma} \Omega \right\}.$$

in the limit $\hbar \rightarrow 0$.

III. FUNCTIONALITY

A major advantage of the surface functional integral formulation rests in its explicit independence on Hamiltonian H . From the point of view of classical physics, dynamical equations and the force fields entering them seem to be more fundamental than the Hamiltonian and/or Lagrangian function, which provide these equations in a relatively compact but ambiguous way, see [3]. Therefore from the conceptual point of view, Formula (6) gives us transition probability amplitude from a different and hopefully new perspective. It is clear that for the potential generated forces the surface functional integral formula (6) gives nothing new compared to (1), since in this case Ω is closed and can be represented as $\Omega = d(p_a dq^a - H dt)$. There are, of course, some hidden subtleties which we pass over either quickly or in silence, however, all of them are discussed in [4].

To show functionality we need to analyze either a strongly non-Lagrangian system [5] or a weakly non-Lagrangian one. For the sake of simplicity let us focus on the second case. To this end, let us consider a system consisting of a free particle with unit mass affected by friction:

$$\ddot{q} = -\kappa \dot{q} \quad \Leftrightarrow \quad \Omega = dp \wedge (dq - p dt) - \kappa p dq \wedge dt.$$

In the considered example, the surface functional integral can be carried out explicitly (for details see [4]). At the

end one arrives at the path integral in the configuration space with a surprisingly trivial result:

$$\int_{\mathcal{U}} [\mathcal{D}\Sigma] \exp \left\{ \frac{i}{\hbar} \int_{\Sigma} \Omega \right\} \propto \exp \left\{ -\frac{i}{\hbar} \int_{t_0}^{t_1} \left(\frac{1}{2} \dot{q}_{cl}^2 - \kappa p_{cl} q_{cl} \right) dt \right\} \times \int [\mathcal{D}q] \exp \left\{ \frac{i}{\hbar} \int_{t_0}^{t_1} \left(\frac{1}{2} \dot{q}^2 - \kappa p_{cl} q \right) dt \right\}.$$

If we define \mathbf{S}_{cl} to be

$$\mathbf{S}_{cl}(q_1, t_1 | q_0, t_0) = \int_{t_0}^{t_1} \left(\frac{1}{2} \dot{q}_{cl}^2 - \kappa p_{cl} q_{cl} \right) dt, \quad (7)$$

then

$$\mathbf{A}(q_1, t_1 | q_0, t_0) \propto \int [\mathcal{D}q] \exp \left\{ \frac{i}{\hbar} \int_{t_0}^{t_1} \left(\frac{1}{2} \dot{q}^2 - \kappa p_{cl} q \right) dt \right\}$$

and in the classical limit $\hbar \rightarrow 0$ we arrive at the saddle point equation which is specified by the functional term in the exponent above:

$$\ddot{q} = -\kappa \dot{q}_{cl}.$$

This differential equation is different from the equation $\ddot{q} = -\kappa \dot{q}$ that we started with initially, but both of them coincide when a solution $q(t)$ satisfying $q(t_0) = q_0$ and $q(t_1) = q_1$ is looking for. In the present situation we gain:

$$\mathbf{S}_{cl} = \frac{\kappa}{4} (q_1 - q_0) \frac{(q_0 + 3q_1)e^{-\kappa t_1} - (q_1 + 3q_0)e^{-\kappa t_0}}{e^{-\kappa t_0} - e^{-\kappa t_1}}$$

and

$$\mathbf{A}(q_1, t_1 | q_0, t_0) = \sqrt{\frac{\kappa}{4\pi i \hbar \tanh \frac{\kappa}{2}(t_1 - t_0)}} e^{\frac{i}{\hbar} \mathbf{S}_{cl}}. \quad (8)$$

Here we have already employed the normalization conditions specified in the first paragraph. One can immediately verify that in the frictionless limit ($\kappa \rightarrow 0$) the

transition probability amplitude $\mathbf{A}(q_1, t_1 | q_0, t_0)$ matches the Schrödinger propagator for a single free particle.

IV. CONCLUSION

Quantization of dissipative systems has been very attractive problem from the early days of quantum mechanics. It has been revived again and again across the decades. Many phenomenological techniques and effective methods have been suggested. References [6] and [7] provide a very basic list of papers dealing with this point.

We have developed here a new quantization method that generalizes the conventional path integral approach. We have focused only on the nonrelativistic quantum mechanics of spinless systems. However, the generalization to the field theory is straightforward.

Let us stress that the proposed method represents an alternative approach to [7] and possesses several qualitative advantages. For example, propagator (8) is invariant with respect to time translations, the same symmetry property which is possessed by the underlying equation of motion. Moreover, it is reasonable to expect that the “dissipative quantum evolution” will not remain Markovian. This fact is again confirmed, since the probability amplitude under consideration does not satisfy the memoryless Chapman-Kolmogorov equation mentioned in the first paragraph.

Finally, let us believe that the simple geometrical idea behind the surface functional integral quantization will fit within the Ludwig Faddeev dictum *quantization is not a science, quantization is an art!*

Acknowledgments

This research was supported in part by MŠMT Grant LC06002, VEGA Grant 1/1008/09 and by the MŠ SR program for CERN and international mobility.

\mathcal{A} . \mathcal{M} . \mathcal{D} . \mathcal{S} .

-
- [1] R. P. Feynman, A. R. Hibbs: *Quantum Mechanics and Path Integrals*, McGraw-Hill Inc., New York, 1965.
[2] R. P. Feynman: *Space-Time Approach to Non-Relativistic Quantum Mechanics*, Rev. of Mod. Phys. **20** (1948), 367-387.
[3] E. P. Wigner: *Do the Equations of Motion Determine the Quantum Mechanical Commutation Relations?*, Phys. Rev. **77** (1950), 711-712.
S. Okubo: *Does the Equation of Motion determine Commutation Relations?*, Phys. Rev. D **22** (1980), 919-923.
M. Henneaux: *Equations of motion, Commutation Relations*

- and Ambiguities in the Lagrangian Formalism*, Annals Phys. **140** (1982), 45-64.
[4] D. Kochan: *Direct quantization of equations of motion*, Acta Polytechnica **47** No. 2-3 (2007), 60-67, arXiv:hep-th/0703073.
D. Kochan: *How to Quantize Forces(?): An Academic Essay on How the Strings Could Enter Classical Mechanics*, J. Geom. Phys. **60** (2010), 219-229, arXiv:hep-th/0612115.
D. Kochan: *Quantization of Non-Lagrangian systems: some irresponsible speculations* AIP Conf. Proc. **956** (2007), 3-8.

- D. Kochan: *Quantization of Non-Lagrangian Systems*, Int. J. Mod. Phys. A **24** Nos. 28 & 29 (2009), 5319-5340.
- D. Kochan: *Functional integral for non-Lagrangian systems*, Phys. Rev. A **81**, 022112 (2010), arXiv:1001.1863.
- [5] J. Douglas: *Solution of the inverse problem in the calculus of variations*, Trans. Am. Math. Soc. **50** (1941), 71-128.
- [6] H. Beteman: *On Dissipative Systems and Related Variational Principles*, Phys. Rev. **38** (1938), 815-819.
- R. P. Feynman, F. L. Vernon: *The Theory of a General Quantum System Interacting with a Linear Dissipative System*, Annals of Physics **24** (1963), 118-173.
- M. D. Kostin: *On the Schrödinger-Langevin Equation*, J. Chem. Phys. **57** (1972), 3589-3591.
- K. K. Kan, J. J. Griffin: *Quantized Friction and the Correspondence Principle: Single Particle with Friction*, Phys. Lett. **50B** (1974), 241-243.
- R. W. Hasse: *On the Quantum Mechanical Treatment of Dissipative Systems*, J. Math. Phys. **16** (1975), 2005-2011.
- H. Dekker: *Classical and quantum mechanics of the damped harmonic oscillator*, Phys. Rep. **80** (1981), 1-112.
- R. Alicki: *Path integrals and stationary phase approximation for quantum dynamical semigroups. Quadratic systems*, J. Math. Phys. **23** (1982), 1370-1375.
- A. O. Caldeira, A. J. Leggett: *Path integral approach to quantum Brownian motion*, Physica **121A** (1983), 587-616.
- J. Geicke: *Semi-classical quantisation of dissipative equations*, J. Phys. A: Math. Gen. **22** (1989) 1017-1025.
- U. Weiss: *Quantum Dissipative Systems* (2nd ed.), Series in Modern Condensed Matter Physics-Vol.10, World Scientific, Singapore, 1999.
- V. E. Tarasov: *Quantization of non-Hamiltonian and Dissipative Systems*, Phys. Lett. A **288** (2001), 173-182, quant-ph/0311159.
- H. P. Breuer, F. Petruccione: *The Theory of Open Quantum Systems*, Oxford University Press, Oxford, 2002.
- M. Razavy: *Classical and Quantum Dissipative Systems*, Imperial Colleges Press, London, 2005.
- D. Chruściński, J. Jurkowski: *Quantum damped oscillator I: dissipation and resonances*, Ann. Phys. **321** (2006), 854-874, quant-ph/0506007.
- D. M. Gitman, V. G. Kupriyanov: *Canonical quantization of so-called non-Lagrangian systems*, Eur. Phys. J. C **50** (2007), 691-700, arXiv: hep-th/0605025.
- V. E. Tarasov: *Quantum Mechanics of Non-Hamiltonian and Dissipative Systems*, Elsevier Science, 2008.
- [7] P. Caldirola: *Forze non conservative nella meccanica quantistica*, Nuovo Cim. **18** (1941), 393-400.
- E. Kanai: *On the Quantization of the Dissipative Systems*, Prog. Theor. Phys. **3** (1948), 440-442.
- I. C. Moreira: *Propagators for the Caldirola-Kanai-Schrödinger Equation*, Lett. Nuovo Cim. **23** (1978), 294-298.
- A. D. Jannussis, G. N. Brodimas, A. Streclas: *Propagator with friction in Quantum Mechanics*, Phys. Lett. **74A** (1979), 6-10.
- U. Das, S. Ghosh, P. Sarkar, B. Talukdar: *Quantization of Dissipative Systems with Friction Linear in Velocity*, Physica Scripta. **71** (2005), 235-237.