

HOLOMORPHIC EXTENSION OF EIGENFUNCTIONS

BERNHARD KRÖTZ AND HENRIK SCHLICHTKRULL

ABSTRACT. Let $X = G/K$ be a Riemannian symmetric space of non-compact type. We prove a theorem of holomorphic extension for eigenfunctions of the Laplace-Beltrami operator on X , by techniques from the theory of partial differential equations.

1. Introduction

Let X be a Riemannian symmetric space of non-compact type. Then $X = G/K$, where G is a connected semisimple Lie group and K a maximal compact subgroup. We choose the group G such that it is contained in a complexification $G_{\mathbb{C}}$, and we denote by $K_{\mathbb{C}} \subset G_{\mathbb{C}}$ the complexification of K . The symmetric space $X_{\mathbb{C}} = G_{\mathbb{C}}/K_{\mathbb{C}}$ carries a natural complex structure, and it contains X as a totally real submanifold.

We are interested in eigenfunctions of the Laplace-Beltrami operator Δ on X . Since this operator is elliptic and G -invariant, every eigenfunction admits a holomorphic extension to some open G -invariant neighborhood of X in $X_{\mathbb{C}}$. The G -orbits in $X_{\mathbb{C}}$ are generally difficult to parametrize, but let us recall that a particular G -invariant open neighborhood Ξ of X , for which the orbit structure is compellingly simple, has been proposed in [1]. It is commonly called the *complex crown* of X , and it has been thoroughly investigated in recent years. See for example [?, ?, 3, 4, 10, 11, 12]. In the present paper we show that every eigenfunction for Δ extends holomorphically to Ξ .

Our result generalizes a result from [11] that every joint eigenfunction for the full set of invariant differential operators on X extends holomorphically to Ξ . The proof given in [11] invokes the Helgason conjecture, affirmed in [9] by micro-local analysis. Our proof is considerably simpler. The crucial step is an application of a theorem from the theory of analytic partial differential equations. This theorem asserts the existence of a holomorphic extension to solutions which are holomorphic on one side of a non-characteristic surface.

At the end of the paper a further generalization is given to functions on G , which are eigenfunctions for the Casimir operator and right- K -finite.

2. Notation

We denote by $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ the Lie algebra of G and its Cartan decomposition. We choose a maximal abelian subspace \mathfrak{a} in \mathfrak{p} and denote by $\Sigma \subset \mathfrak{a}^*$ the corresponding system of restricted roots. The root spaces in \mathfrak{g} are denoted by \mathfrak{g}^{α} , where $\alpha \in \Sigma$, and by $\Sigma^+ \subset \Sigma$ we denote a positive system. The centralizer of \mathfrak{a} in K is $M = Z_K(\mathfrak{a})$, and the Weyl group is $W = N_K(\mathfrak{a})/M$, where $N_K(\mathfrak{a})$ is the normalizer.

Recall the definition of the complex crown Ξ of X . We set

$$\Omega := \{Y \in \mathfrak{a} \mid |\alpha(Y)| < \frac{\pi}{2}, \forall \alpha \in \Sigma\}.$$

Then

$$\Xi := G \exp(i\Omega) K_{\mathbb{C}} = \{g \exp(iY) \cdot x_0 \mid g \in G, Y \in \Omega\} \subset X_{\mathbb{C}}.$$

Here x_0 denotes the standard base point $eK_{\mathbb{C}}$ in $X_{\mathbb{C}}$.

3. Results

Lemma 3.1. *The G -invariant crown Ξ is an open subset of $X_{\mathbb{C}}$. The surjective map*

$$\Phi : G \times \Omega \rightarrow \Xi, \quad (g, Y) \mapsto g \exp(iY) \cdot x_0$$

is real analytic, and the topology of Ξ is identical to the quotient topology with respect to this map.

Let Ω^+ be the intersection of Ω with the positive open chamber in \mathfrak{a} , and let $\Xi' = \Phi(G \times \Omega^+)$. Then Ξ' is open and dense in Ξ , and

$$\Phi' : G/M \times \Omega^+ \rightarrow \Xi', \quad (gM, Y) \mapsto \Phi(g, Y)$$

is a diffeomorphism.

Proof. Apart from the statement about the topology of Ξ , this can be found in [11], §4. For the topological statement we need to prove that a subset of Ξ is open if its preimage is open. It suffices to prove the following. Let $z_n \rightarrow z \in \Xi$ be a converging sequence. Then there exists a subsequence of the form $z_j = \Phi(g_j, Y_j)$ with converging sequences $g_j \rightarrow g \in G$ and $Y_j \rightarrow Y \in \Omega$. It follows from [1], Propositions 1 and 7, that there exist sequences g_n in G , k_n in K and Y_n in Ω such that $z_n = \Phi(g_n k_n, Y_n)$, and such that g_n and $k_n \exp(iY_n) \cdot x_0$ both converge. By passing to a subsequence, we may assume that k_j converges, and since $Y \mapsto \exp(iY) \cdot x_0$ is a diffeomorphism of Ω onto its image, it then follows that Y_j converges in Ω . \square

Theorem 3.2. *Let $f \in C^\infty(X)$ be an eigenfunction for Δ . Then f extends to a holomorphic function on Ξ .*

Proof. As Δ is elliptic, the regularity theorem for elliptic differential operators (see [6], Theorem 7.5.1) implies that f is an analytic function. As such it has an extension to a holomorphic function on some open neighborhood U_0 of x_0 in Ξ . It follows from the proof in [6], that U_0 can be chosen independently of f , that is, *every* eigenfunction can be holomorphically extended to U_0 (the radius of convergence obtained in the proof depends only on the corresponding radii for the coefficients of the differential operator). In particular, it follows from the fact that Δ is G -invariant, that $L_g f$ extends to U_0 for all $g \in G$. The union U of the

G -translated sets $L_{g^{-1}}(U_0)$ is then a G -invariant open neighborhood of X in Ξ , to which f extends.

We now consider the open dense subset $\Xi' \subset \Xi$ from Lemma 3.1. The intersection $U \cap \Xi'$ is non-empty, open and G -invariant. Let $Y_0 \in \Omega^+$, and for $r > 0$ let B_r denote the open ball in \mathfrak{a} of radius r , centered at Y_0 . If $B_r \subset \Omega^+$, then we define an open set

$$T_r = G \exp(iB_r) \cdot x_0 \subset \Xi',$$

which we regard as a G -invariant ‘circular tube’ in Ξ' . We claim that if f extends holomorphically to a set containing some circular tube $T_r \subset \Xi'$ centered at Y_0 , then it extends to all circular tubes in Ξ' centered at Y_0 . Since Y_0 was arbitrary, and since Ω^+ is simply connected, it follows from this claim that f extends holomorphically from $U \cap \Xi'$ to Ξ' .

In order to establish the claim we use Theorem 9.4.7 of [8], due to Zerner [13]. We write $\Delta_{\mathbb{C}}$ for the extension of Δ to a $G_{\mathbb{C}}$ -invariant holomorphic differential operator on $X_{\mathbb{C}}$. Obviously, the holomorphic extension that we seek will be an eigenfunction for $\Delta_{\mathbb{C}}$ on Ξ . It follows from Lemma 3.1 that each circular tube T_r , for which the closure is contained in Ξ' , has real-analytic boundary ∂T_r . In order to apply Zerner’s theorem it suffices to establish that ∂T_r is non-characteristic for $\Delta_{\mathbb{C}}$, for all such tubes. By G -invariance, it suffices to consider boundary points $x \in \partial T_r$ with $x \in \exp(i\Omega) \cdot x_0$. Recall from [11], p. 207, that when $x \in \exp(i\Omega) \cdot x_0$ we have a complex-linear isomorphism

$$(3.1) \quad \mathfrak{p}_{\mathbb{C}} \ni Z \mapsto \tilde{Z}_x \in T_x \Xi,$$

where \tilde{Z} is the holomorphic vector field on $X_{\mathbb{C}}$ given by

$$\tilde{Z}_x \varphi = L_Z \varphi(x) = \frac{d}{dz} \varphi(\exp(-zZ)x)|_{z=0}.$$

In this isomorphism the tangent space at x of the boundary ∂T_r will then be a real hyperplane given by an equation $\operatorname{Re} \zeta(Z) = 0$ for some cotangent vector $\zeta \in \mathfrak{p}_{\mathbb{C}}^*$. Since the tube T_r is G -invariant, it follows that $\operatorname{Re} \zeta$ annihilates Z for all $Z \in \mathfrak{p}$, so ζ is purely imaginary on \mathfrak{p} .

Let $(X_j^\alpha)_{\alpha \in \Sigma^+, 1 \leq j \leq m_\alpha}$ together with $Y_1, \dots, Y_r \in \mathfrak{a}$ be an orthonormal basis for \mathfrak{p} such that $X_j^\alpha \in \mathfrak{p}_\alpha := [\mathfrak{g}^\alpha + \mathfrak{g}^{-\alpha}] \cap \mathfrak{p}$. Here $m_\alpha = \dim \mathfrak{p}_\alpha$ as usual. In the universal enveloping algebra we have

$$\Delta = \sum_{\alpha \in \Sigma^+} \sum_{j=1}^{m_\alpha} (X_j^\alpha)^2 + \sum_{i=1}^r Y_i^2$$

with respect to the right action, where functions on G/K are regarded as a right K -invariant function on G . Observe that modulo \mathfrak{k} ,

$$\operatorname{Ad}(a)^{-1}(X_j^\alpha) = \cosh(\alpha(\log a))X_j^\alpha$$

for $a \in A$ (see [11] p. 207), and hence

$$R_{X_j^\alpha} f(a) = -[\cosh \alpha(\log a)]^{-1} L_{X_j^\alpha} f(a).$$

It follows that

$$(3.2) \quad \Delta = \sum_{\alpha \in \Sigma^+} \sum_{j=1}^{m_\alpha} [\cosh \alpha(\log a)]^{-2} (X_j^\alpha)^2 + \sum_{i=1}^r Y_i^2$$

at $z = a \cdot x_0 \in X$, with respect to the left action.

By analytic continuation, the same equation holds as well for $\Delta_{\mathbb{C}}$ and $a \in A_{\mathbb{C}}$. In particular, at $x = \exp(iY) \cdot x_0$ we obtain

$$\Delta_{\mathbb{C}} = \sum_{\alpha \in \Sigma^+} \sum_{j=1}^{m_\alpha} [\cos \alpha(Y)]^{-2} [(\tilde{X}_j^\alpha)_x]^2 + \sum_{i=1}^r [(\tilde{Y}_i)_x]^2.$$

Note that the condition that x belongs to the crown precisely ensures that $\cos \alpha(Y) \neq 0$, so that the expression makes sense. As ζ is purely imaginary on \mathfrak{p} , it follows that all terms in the above sum are ≤ 0 when applied to ζ . Thus the principal symbol of $\Delta_{\mathbb{C}}$ is non-zero at ζ , and the boundary of T_r is non-characteristic. It follows that Zerner's theorem can be applied, so that f extends holomorphically to Ξ' .

For the extension to the full set Ξ we shall apply Bochner's theorem (see [7], Theorem 2.5.10). From what we have seen so far, for all $g \in G$ the function

$$f_g : \mathfrak{a} \rightarrow \mathbb{C}, \quad Y \mapsto f(g \exp(Y) \cdot x_0)$$

extends to a holomorphic function on a tubular neighborhood $\mathfrak{a} + i\omega$ of \mathfrak{a} in $\mathfrak{a}_{\mathbb{C}}$, and also to $\mathfrak{a} + i\Omega^+$. For elements $w \in N_K(\mathfrak{a})$ we have

$$f_g(\text{Ad}(w)Y) = f_{gw}(Y).$$

It follows that f_g extends to a holomorphic function on each Weyl conjugate of $\mathfrak{a} + i\Omega^+$, hence to $\mathfrak{a} + i\Omega'$, where $\Omega' = \cup \text{Ad}(w)(\Omega^+)$ is the set of regular elements in Ω . Now Bochner's theorem implies that f_g extends to a holomorphic function on the tube over the convex hull of $\omega \cup \Omega'$, that is, to $\mathfrak{a} + i\Omega$. Furthermore, $g \mapsto f_g$ is continuous into $H(\mathfrak{a} + i\Omega)$ (with standard topology), since it is continuous into the space $H(\mathfrak{a} + i(\omega \cup \Omega'))$ which by Bochner's theorem is topologically isomorphic.

Recall that for all $g, g' \in G$ and $Y, Y' \in \Omega$ we have

$$g \exp(iY) \cdot x_0 = g' \exp(iY') \cdot x_0$$

if and only if there exists $w \in N_K(\mathfrak{a})$ and $k \in Z_K(Y)$ with $g' = gkw$ and $Y' = \text{Ad}(w^{-1})Y$. It follows easily that by

$$g \exp(iY) \cdot x_0 \mapsto f_g(Y)$$

we obtain a well-defined extension of f on Ξ . The topological statement in the first part of Lemma 3.1 implies that this extension is continuous. Since the extension is holomorphic on Ξ' , it must be holomorphic everywhere. \square

We list some easy consequences of the preceding theorem and its proof. From the Iwasawa decomposition $G = NAK$ associated to the positive system Σ^+ , we obtain the familiar horospherical projection $x \mapsto H(x) \in \mathfrak{a}$, defined by $x \in N \exp H(x) \cdot x_0$ for $x \in X$. For each $\lambda \in \alpha_{\mathbb{C}}^*$ the function

$$x \mapsto e^{\lambda(H(x))}$$

on X is an eigenfunction for Δ , hence extends to a holomorphic function on Ξ . We obtain:

Corollary 3.3. *The projection $H : X \rightarrow \mathfrak{a}$ extends to a holomorphic map $\Xi \rightarrow \mathfrak{a}_{\mathbb{C}}$. Moreover, $\Xi \subset N_{\mathbb{C}} A_{\mathbb{C}} K_{\mathbb{C}}$.*

Proof. Let $h_{\lambda}(z)$ denote the analytic continuation of $e^{\lambda(H(x))}$. Since $h_{-\lambda}(z) = h_{\lambda}(z)^{-1}$ we conclude that $h_{\lambda}(z) \neq 0$. As Ξ is simply connected, the analytic continuation of $\lambda(H(x))$ is obtained by taking logarithms, and the first statement follows. For the last statement, we note that once the Iwasawa A -component allows an analytic continuation, then so does the N -component. Indeed, knowing the A -component, we can determine the N -component of $x \in X$ from $\theta(x)x^{-1}$, where θ denotes the Cartan involution. \square

The preceding corollary was obtained for classical groups in [10]. The general case follows from results established in [12], [2] and [4] with [5].

Let $\omega \subset \Omega$ be open, convex and W -invariant, and let $T_{\omega} \subset \Xi$ denote the open set

$$T_{\omega} = G \exp(i\omega) \cdot x_0.$$

Corollary 3.4. *Let $f \in C^{\infty}(X)$ and let P be a non-trivial polynomial of one variable. If $P(\Delta)f$ extends to a holomorphic function on T_{ω} , then so does f .*

Proof. By treating the factors of P successively we may assume that $P(\Delta) = \Delta - \lambda$. The proof of Theorem 3.2 can then be repeated. \square

The following generalization is more far-reaching. We denote by $C \in \mathcal{Z}(\mathfrak{g})$ the Casimir element of \mathfrak{g} .

Theorem 3.5. *Let $f \in C^{\infty}(G)$ be a right K -finite eigenfunction of C . Then f extends to a holomorphic function on*

$$\tilde{\Xi} := G \exp(i\hat{\Omega}) K_{\mathbb{C}} \subset G_{\mathbb{C}}.$$

Proof. Recall that f being K -finite means that the translates $R_k f$ by $k \in K$ span a finite dimensional space, which is then a representation space for K . We may assume that it is irreducible, and then f is an eigenfunction for the Casimir element $C_{\mathfrak{k}}$ of \mathfrak{k} , acting from the right. The operator $C + 2C_{\mathfrak{k}}$ is elliptic, so it follows that f is real analytic. The proof of Theorem 3.2 can now be repeated, with the following changes.

In Lemma 3.1 we replace the map Φ by

$$\tilde{\Phi} : G \times \Omega \times K_{\mathbb{C}} \rightarrow \tilde{\Xi}, \quad (g, Y, k) \mapsto g \exp(iY)k,$$

and we define $\tilde{\Phi}'$ as before, but now on $(G \times \Omega^+ \times K_{\mathbb{C}})/M$, where M acts on the first and last factor, from the right and left, respectively.

The G -invariant tubes $T_r \subset \Xi$ are replaced by their $G \times K_{\mathbb{C}}$ -invariant preimages $\tilde{T}_r = T_r K_{\mathbb{C}} \subset \tilde{\Xi}$. The map $Z \mapsto L_Z$ from $\mathfrak{p}_{\mathbb{C}}$ onto $T_x \Xi$ in (3.1) is replaced by $Z \oplus U \mapsto L_Z + R_U$ from $\mathfrak{g}_{\mathbb{C}} = \mathfrak{p}_{\mathbb{C}} \oplus \mathfrak{k}_{\mathbb{C}}$ onto $T_x \tilde{\Xi}$. Here $x \in \exp(i\Omega)$. The cotangent vector ζ , normal to \tilde{T}_r at x is zero on $\mathfrak{k}_{\mathbb{C}}$ and purely imaginary on \mathfrak{p} , by the same argument as before.

Since f is a $C_{\mathfrak{k}}$ -eigenfunction, the action of C on it differs only by a constant from that of the operator Δ described in (3.2). Hence $\partial \tilde{T}_r$ is non-characteristic for C , and the application of Zerner's theorem goes through. The rest of the argument is essentially unchanged. \square

References

- [1] Akhiezer, D. and Gindikin, S., *On Stein extensions of real symmetric spaces*, Math. Ann. **286** (1990), 1–12.
- [2] Burns, D., Halverscheid, S. and R. Hind, *The geometry of Grauert tubes and complexification of symmetric spaces*, Duke Math. J. **118** (2003), no. **3**, 465–491.
- [3] Gindikin, S. and Krötz, B. *Invariant Stein domains in Stein symmetric spaces and a nonlinear complex convexity theorem*, Int. Math. Res. Not. **2002**, 959–971.
- [4] Huckleberry, A. *On certain domains in cycle spaces of flag manifolds*, Math. Ann. **323** (2002), 797–810.
- [5] ———, Erratum: *On certain domains in cycle spaces of flag manifolds*, to be submitted
- [6] Hörmander, L., *Linear Partial Differential Operators*, Springer Grundlehren 116, 1976.
- [7] ———, *An introduction to complex analysis in several variables*, Van Nostrand, 1966.
- [8] ———, *The Analysis of Linear Partial Differential Operators I*, Springer Grundlehren 256, 1983.
- [9] Kashiwara, M. et al, *Eigenfunctions of invariant differential operators on a symmetric space*, Ann. of Math. **107** (1978), 1–39.

- [10] Krötz, B. and R. Stanton, *Holomorphic extensions of representations. I. Automorphic functions*, Ann. of Math. **(2) 159** (2004), no. **2**, 641–724.
- [11] ———, *Holomorphic extensions of representations. II. Geometry and harmonic analysis*, Geom. Funct. Anal. **15** (2005), no. **1**, 190–245.
- [12] Matsuki, T. *Stein extensions of Riemann symmetric spaces and some generalization*, J. Lie Theory **13** (2003), 565–572.
- [13] Zerner, M., *Domaine d'holomorphie des fonctions vérifiant une équation aux dérivées partielles*, C. R. Acad. Sci. Paris **272** (1971), 1646.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, D-53111 BONN,
EMAIL: KROETZ@MPIM-BONN.MPG.DE

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN 5, DK-2100 KØPENHAVN, EMAIL: SCHLICHT@MATH.KU.DK