

Some applications of the Poisson summation formula

M. L. Glasser

Department of Physics, Clarkson University
Potsdam, NY 13699-5820 (USA)
laryg@clarkson.edu

Nikos Bagis

Department of Informatics, Aristotle University
Thessaloniki Greece
bagkis@hotmail.com

Abstract

In this note we derive two special forms of the Poisson's summation formula, for even and odd functions, which we apply to obtain representations for some Euler-type numbers Q_n , as well as to sum various series related to elliptic functions and elementary transcendental constants.

1. Introduction

One of the fundamental results of Fourier analysis is the Poisson Summation Formula, which can be written (see [Ch] pg. 43-47):

$$a \sum_{k=-\infty}^{\infty} f(ak) = \sum_{k=-\infty}^{\infty} f^{\wedge}(bk) \quad (1)$$

where $a > 0$, $a = 2\pi / a$ and

$$f^{\wedge}(x) = \int_{-\infty}^{\infty} f(t)e^{-ixt} dt$$

is the Fourier Transform of f . Eq.(1) is valid under the relatively weak condition

$$f(x) = O\left[\left(1+|x|^c\right)^{-1}\right], \text{ for } x \in \mathbb{R} \text{ and for some } c > 0.$$

It is also known that if $f_c^{\wedge}(\gamma) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(t) \cos(t\gamma) dt$ is the Cosine Transform of a

function f then someone can get the Poisson's Cosine Summation Formula (See [T] pg. 60-68):

$$\sqrt{a} \left(\frac{f(0)}{2} + \sum_{n=1}^{\infty} f(ka) \right) = \sqrt{b} \left(\frac{f_c^{\wedge}(0)}{2} + \sum_{k=1}^{\infty} f_c^{\wedge}(kb) \right) \quad (2)$$

where $a > 0$, $b > 0$, $ab = 2\pi$.

2. Lemmas

Lemma 1.

Let f is analytic in the upper half plane $\text{Im}(z) > 0$ and continuous in $\text{Im}(z) \geq 0$ and there exists $C, N > 0$ and $0 \leq b < \pi$ such that

$$|f(z)| \leq C(1+|z|)^N e^{b|\text{Re}(z)|}, \text{ for every } z \text{ in } \text{Im}(z) \geq 0 \quad (3)$$

then the series $\sum_{k=0}^{\infty} (-1)^k f(i(k+1/2))$ is Abel summable and holds the following relation:

$$\int_{-\infty}^{\infty} \frac{f(t)}{\cosh(\pi t)} dt = 2 \lim_{r \rightarrow 1^-} \sum_{k=0}^{\infty} (-1)^k f(i(k+1/2)) r^k \quad (4)$$

If also $|f(ik)| \leq \frac{C'}{k+1}$, $k = 1, 2, 3, \dots$ then the series converge and its sum is equal with the integral in Eq.(4).

Lemma 2.

Let f is as in Lemma 1 then the series $\sum_{k=0}^{\infty} (-1)^k k f(ik)$ is Abel summable and holds the following relation:

$$\int_{-\infty}^{\infty} \frac{f(t)t}{\sinh(\pi t)} dt = 2 \lim_{r \rightarrow 1^-} \sum_{k=1}^{\infty} (-1)^{k-1} k f(ik) r^k \quad (5)$$

If also $|f(ik)| \leq \frac{C''}{k^2}$, $k = 1, 2, 3, \dots$ then the series converge and its sum is equal with the integral in Eq.(5). (For a more general result of Lemma's 1, 2 see also [Bag]).

3. Theorems

Theorem 1.

Let $a > 0$, f is even and as in Lemma 1, then

$$\sqrt{a} \left(\frac{f(0)}{2} + \sum_{k=1}^{\infty} \frac{f(ka)}{\cosh(\pi ka)} \right) = \sqrt{\frac{2b}{\pi}} \left(\frac{ce}{2} + \sum_{k=0}^{\infty} \frac{(-1)^k f(i(k+1/2))}{e^{b(k+1/2)} - 1} \right) \quad (6)$$

where $ab = 2\pi$ and $ce = \lim_{x \rightarrow 0^+} \left(\sum_{k=0}^{\infty} (-1)^k f(i(k+1/2)) e^{-(k+1/2)x} \right)$.

Theorem 2.

Let $a > 0$, f is odd and as in Lemma 2, then

$$\sqrt{a} \left(\frac{f'(0)}{2\pi} + \sum_{k=1}^{\infty} \frac{f(ka)}{\sinh(\pi ka)} \right) = \sqrt{\frac{b2}{\pi}} \left(c_o/2 + i \sum_{k=1}^{\infty} \frac{(-1)^k f(ki)}{e^{bk} - 1} \right) \quad (7)$$

when $ab = 2\pi$ and $c_o = \lim_{x \rightarrow 0^+} \left(\sum_{k=1}^{\infty} (-1)^k i f(ki) e^{-kx} \right)$.

4. Proof of Lemmas 1,2

Proof of Lemma 1.

Let $a > 0$. We define the function

$$g(z) = e^{iza} f(z) \frac{1}{\cosh(\pi z)}$$

g is meromorphic in $\text{Im}(z) > 0$ and has simple poles at the points $z = i(k + 1/2)$ with

$$\text{Res}(g; i(k + 1/2)) = e^{-a(k+1/2)} f(i(k + 1/2)) \frac{(-1)^k}{\pi i}. \text{ Thus if } m > 1 \text{ natural, } R = m \text{ and } \gamma_R$$

is the upper half circle with diameter $[-R, R]$ we get:

$$\frac{1}{2\pi i} \int_{\gamma_{R_m}} g(z) dz = \sum_{k=0}^{m-1} \frac{(-1)^k}{\pi i} e^{-a(k+1/2)} f(i(k + 1/2)) \quad (8)$$

If m natural number we have $|\cosh(\pi \text{Re}^{i\theta})| \geq ce^{\pi R|\cos(\theta)|}$, where $c > 0$ is a positive constant. From (3) we get for $0 < a \leq \pi - b$:

$$\left| \int_0^{\pi} f(R \cdot e^{i\theta}) \frac{e^{ia \text{Re}^{i\theta}}}{\cosh(\pi(\text{Re}^{i\theta}))} i \text{Re}^{i\theta} d\theta \right| \leq \left| \int_0^{\pi} (1+R)^N \text{Re}^{(b-\pi)R|\cos(\theta)|} e^{-aR|\cos(\theta)|} d\theta \right| \leq$$

$$\leq \int_0^{\pi} (1+R)^{N+1} e^{-aR} d\theta \rightarrow 0 \text{ as } R \rightarrow \infty \text{ (} R = m \text{ natural number).}$$

From (3) we have $\int_{-\infty}^{\infty} \left| f(t) \frac{1}{\cosh(\pi t)} dt \right| < \infty$ and $\sum_{k=0}^{\infty} e^{-ak} |f(i(k + 1/2))| < \infty$

Thus we get $R = m \rightarrow \infty$ in Eq.(8) and we have:

$$\int_{-\infty}^{\infty} \frac{f(t) e^{ita}}{\cosh(\pi t)} dt = 2 \sum_{k=0}^{\infty} (-1)^k f(i(k + 1/2)) e^{-a(k+1/2)}, \text{ for every } a > 0 \quad (9)$$

The main result follows easily setting $a \rightarrow 0^+$ in Eq.(9).

Proof of Lemma 2.

As in Lemma 1.

5. Proof of Theorems

Proof of Theorem 1.

For f even and as in Lemma 1 we have

$$\int_0^{\infty} \frac{f(t) \cos(t\gamma)}{\cosh(\pi t)} dt = \sum_{k=0}^{\infty} (-1)^k f(i(k+1/2)) e^{-(k+1/2)\gamma} \quad (10)$$

From Eq.(10) and the Poisson Cosine Formula Eq.(2):

$$\sqrt{a} \left(\frac{f(0)}{2} + \sum_{k=1}^{\infty} f(ka) \right) = \sqrt{b} \left(\frac{f_c \wedge (0)}{2} + \sum_{k=1}^{\infty} f_c \wedge (kb) \right)$$

or

$$\sqrt{a} \left(\frac{f(0)}{2} + \sum_{k=1}^{\infty} \frac{f(ka)}{\cosh(\pi ka)} \right) = \sqrt{b} \left(\frac{ce}{2} + \sum_{k=0}^{\infty} \frac{(-1)^k f(i(k+1/2))}{e^{b(k+1/2)} - 1} \right)$$

Where we have used that $\sum_{l=0}^{\infty} e^{-(k+1/2)lb} = \frac{1}{e^{b(k+1/2)} - 1}$.

Proof of Theorem 2.

From Lemma 2 we have $\int_{-\infty}^{\infty} \frac{f(t) t e^{ita}}{\sinh(\pi t)} dt = 2 \sum_{k=1}^{\infty} (-1)^{k-1} k f(ik) e^{-ka}$, $a > 0$

If g is even then

$$\int_0^{\infty} \frac{g(t) t}{\sinh(\pi t)} \cos(ta) dt = \sum_{k=1}^{\infty} (-1)^{k-1} k g(ik) e^{-ka} \quad (11)$$

From (1) and the Poisson cosine formula we have

$$\sqrt{a} \left(\frac{1}{2} \lim_{h \rightarrow 0} \left(\frac{g(h)h}{\sinh(\pi h)} \right) + \sum_{n=1}^{\infty} \frac{g(ka)ka}{\sinh(ka)} \right) = \sqrt{b} \left(\sqrt{\frac{2}{\pi}} c_e / 2 - \sum_{m=1}^{\infty} 2 \sqrt{\frac{2}{\pi}} \sum_{k=1}^{\infty} (-1)^{k-1} k g(ki) e^{-kmb} \right)$$

or

$$\sqrt{a} \left(\frac{1}{2} \lim_{h \rightarrow 0} \left(\frac{g(h)h}{\sinh(\pi h)} \right) + \sum_{n=1}^{\infty} \frac{g(ka)ka}{\sinh(ka)} \right) = \sqrt{\frac{b2}{\pi}} \left(c_e / 2 - 2 \sum_{k=1}^{\infty} \frac{(-1)^{k-1} k g(ki)}{e^{bk} - 1} \right)$$

or

$$\sqrt{a} \left(\frac{f(0)}{2\pi} + \sum_{n=1}^{\infty} \frac{g(ka)ka}{\sinh(ka)} \right) = \sqrt{\frac{b2}{\pi}} \left(c_e / 2 - 2 \sum_{k=1}^{\infty} \frac{(-1)^{k-1} k g(ki)}{e^{bk} - 1} \right)$$

The result follows replacing $f(x) = x g(x)$

6. Applications of Theorems 1,2

1) As a first application we take $f(t) = t^{2m+1}$ in Theorem 2. Since

$$\begin{aligned}
c_o &= (-1)^m \lim_{x \rightarrow 0^+} \sum_{k=1}^{\infty} (-1)^{k+1} k^{2m+1} e^{-kx} = \\
&(-1)^{m+1} \lim_{x \rightarrow 0^+} \frac{d^{2m+1}}{dx^{2m+1}} \sum_{k=1}^{\infty} (-1)^{k+1} e^{-kx} = (-1)^{m+1} \lim_{x \rightarrow 0} \frac{d^{2m+1}}{dx^{2m+1}} \left(\frac{1}{e^x + 1} \right) = \\
&(-1)^{m+1} \lim_{x \rightarrow 0} \frac{d^{2m+1}}{dx^{2m+1}} \left(\sum_{k=0}^{\infty} \frac{Q_k}{k!} x^k \right) = \\
&\frac{(-1)^{m+1}}{2} Q_{2m+1}
\end{aligned} \tag{12}$$

Theorem 2 therefore yields

$$\begin{aligned}
\frac{a}{2} \left[\frac{\delta_{m,0}}{2\pi} + a^{2m+1} \sum_{k=1}^{\infty} \frac{k^{2m+1}}{\sinh(\pi a k)} \right] = \\
\frac{(-1)^{m+1}}{4} Q_{2m+1} + (-1)^m \sum_{k=1}^{\infty} \frac{(-1)^{k+1} k^{2m+1}}{e^{2\pi k/a} - 1}.
\end{aligned} \tag{13}$$

In particular, for $m = 0$ the hyperbolic series on the right hand side of (13) can be expressed in terms of elliptic functions [Z] which yields

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1} k}{e^{2\pi k/a} - 1} = \frac{1}{4} \left(\frac{1}{2} - \frac{a}{\pi} \right) + \frac{K}{\pi^2} [E - K] \tag{14}$$

Where K and E are the complete elliptic integrals of the first and second kind of modulus k given by $K'/K = a$. When a is the square root of a rational number, k is one of the so-called singular moduli and the elliptic integrals can be expressed in terms of the Gamma function.

2) Next, we select $f(t) = t^{4n}$, $\mathbb{N} \ni n > 1$, for which, as above, we find $c_e - 1/2Q_{4n+1}$.

Then, taking $a = 1$, Theorem 2 gives

$$\sum_{k=1}^{\infty} \frac{k^{4n+1} (e^{k\pi} + (-1)^k)}{(e^{k\pi} - 1)(e^{k\pi} + 1)} = -1/4Q_{4n+1} \tag{15}$$

By using partial fractions and the identity ([Be2] pg. 262):

$$\sum_{k=1}^{\infty} \frac{k^{4n+1}}{e^{2\pi k} - 1} = \frac{B_{4n+2}}{8n+4}, \tag{16}$$

where B_n , $n = 0, 1, 2, 3, \dots$ denotes the Bernoulli numbers, we find

$$\sum_{k=1}^{\infty} \frac{(2k-1)^{4n+1}}{e^{\pi(2k-1)} + 1} = -1/4Q_{4n+1} - \frac{2^{4n-1}}{2n+1} B_{4n+2}. \tag{17}$$

3) Other identities are:

a) Setting $f(t) = \frac{\sinh\left(\frac{t\pi}{6}\right)^6}{t^6}$ in Theorem 2, for $a = \pi$, $b = 2$ we get the result:

$$\frac{41\zeta(5)}{6912} = \frac{\pi^6}{93312} + 2 \sum_{k=1}^{\infty} \frac{(-1)^k \sin^6\left(\frac{k\pi}{6}\right)}{k^5(e^{2k} - 1)} + \frac{1}{\pi^4} \sum_{k=1}^{\infty} \frac{\sinh^6\left(\frac{k\pi}{6}\right)}{k^5 \sinh(\pi^2 k)}$$

Which is a formula for $\zeta(5)$.

$$\text{b) } \frac{7\zeta(3)}{128} = \frac{\pi^3}{512} + \sum_{k=1}^{\infty} \frac{(-1)^k \sin^4\left(\frac{k\pi}{4}\right)}{k^3(e^{\pi k} - 1)} + \frac{1}{8} \sum_{k=1}^{\infty} \frac{\sinh^4\left(\frac{k\pi}{2}\right)}{k^3 \sinh(2\pi k)}$$

c) For $f(t) = \left(\frac{\cos(tc) - 1}{t}\right)^3$, $a = 1$, $b = 2\pi$ we get:

$$\begin{aligned} 8 \sum_{k=1}^{\infty} \frac{(\cos(ck) - 1)^3}{k^3 \sinh(k\pi)} + 16 \sum_{k=1}^{\infty} \frac{(-1)^k (\cosh(ck) - 1)^3}{k^3 (e^{2k\pi} - 1)} = \\ = -c^3 + c\pi^2 - 30Li_3(-e^{-c}) + 12Li_3(-e^{-2c}) - 2Li_3(-e^{-3c}) - 15\zeta(3) \quad : (18) \end{aligned}$$

where $Li_v(x) = \sum_{k=1}^{\infty} \frac{x^k}{k^v}$, see [Be1] pg. 246).

Set now

$$f(c) := -30Li_3(-e^{-c}) + 12Li_3(-e^{-2c}) - 2Li_3(-e^{-3c}), \quad c \geq 0$$

Then we have the analytic continuation of f :

$$f(-c) = -2c^3 + 2c\pi^2 + f(c)$$

$$\text{d) } \frac{1}{8} \sec^2\left(\frac{1}{2}\right) \tan\left(\frac{1}{2}\right) = 4 \sum_{k=1}^{\infty} \frac{k^2 \sinh(2k)}{\sinh(2k\pi)} - \sum_{k=1}^{\infty} \frac{(-1)^k k^2 \sin(k)}{e^{\pi k} - 1}$$

$$\text{e) } \frac{1}{16} (2 - \cos(1)) \sec^4\left(\frac{1}{2}\right) = 8 \sum_{k=1}^{\infty} \frac{k^3 \sinh(2k)}{\sinh(2k\pi)} - \sum_{k=1}^{\infty} \frac{(-1)^k k^3 \cos(k)}{e^{\pi k} - 1}$$

$$\text{f) } \tan\left(\frac{1}{2}\right) = \frac{2}{\pi} + 4 \sum_{k=1}^{\infty} \frac{\sinh(2k)}{\sinh(2k\pi)} + 4 \sum_{k=1}^{\infty} \frac{(-1)^k \sin(k)}{e^{\pi k} - 1}$$

4) If we set $f(t) = t^{4\nu+1} \cos(\pi t)$, $\nu = 1, 2, 3, \dots$ and $a = 1$, $b = 2\pi$ Theorem 2 then

$$\sum_{k=1}^{\infty} \frac{(-1)^k k^{4\nu+1} \cosh(k\pi)}{e^{2k\pi} - 1} = -\frac{c_o}{2}$$

i) For $\nu = 1$ we get

$$\sum_{k=1}^{\infty} \frac{(-1)^k k^5 \cosh(k\pi)}{e^{2k\pi} - 1} = \frac{1}{64} (33 - 26 \cosh(\pi) + \cosh(2\pi)) \operatorname{sech}^6\left(\frac{\pi}{2}\right)$$

ii) For $\nu = 2$ we get

$$\sum_{k=1}^{\infty} \frac{(-1)^k k^9 \cosh(k\pi)}{e^{2k\pi} - 1} = \frac{-e^\pi (1 - 502e^\pi + 14608e^{2\pi} - 88234e^{3\pi} + 156190e^{4\pi} - 88234e^{5\pi} + 14608e^{6\pi} - 502e^{7\pi} + e^{8\pi})}{(e^\pi + 1)^{10}}$$

iii) For $\nu = 3 \dots$ etc. (All these sums are rational functions of e^π with integer coefficients)

5) If we set $f(t) = \frac{\sin(\pi t \nu)}{t}$, $a = 1/\nu$, $b = 2\pi\nu$ in Theorem 1 then with $\nu > 0$, ν -real, we get:

$$\operatorname{arccot}\left(e^{\nu\pi/2}\right) = 2 \sum_{k=0}^{\infty} \frac{(-1)^k \sinh(\nu\pi(k+1/2))}{(e^{(2k+1)\nu\pi} - 1)(2k+1)}$$

6) If we set $f(t) = \left(\frac{\sin(\pi t/\nu)}{t}\right)^2$, $a = 1/\nu$, $b = 2\pi\nu$ in Theorem 1 then we get:

$$\frac{1}{4} e^{\pi/\nu} \varphi(-e^{2\pi/\nu}, 2, \frac{1}{2}) + \frac{1}{4} e^{-\pi/\nu} \varphi(-e^{-2\pi/\nu}, 2, 1/2) = 2G + \frac{\pi^2}{2\nu^3} - 8 \sum_{k=1}^{\infty} \frac{(-1)^k \sinh^2\left(\pi\left(\frac{k}{\nu} + \frac{1}{2\nu}\right)\right)}{(e^{(2k+1)\pi\nu} - 1)(2k+1)^2} + \nu \sum_{k=1}^{\infty} \frac{\sin^2\left(\frac{k\pi}{\nu}\right)}{\cosh(k\pi/\nu) k^2}$$

Where G is Catalan's constant: $G = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2}$ and $\varphi(z, s, a) = \sum_{k=0}^{\infty} \frac{z^k}{(a+k)^s}$ is LerchPhi function (see [Be1]).

6. References

[A,S]: Milton Abramowitz, Irene A. Stegun .Handbook of Mathematical Functions. Dover.

[Ba,Gl]: N. Bagis, M. L. Glasser. "Integrals and Series Resulting from Two Sampling Theorems". Journal of Sampling Theory in Signal and Image Processing. Vol. 5, No. 1, Jan. 2006, pp. 89-97. ISSN: 1530-6429.

[Bag]: N. Bagis. "On Series Integrals and Continued Fractions" Hardy Ramanujan Journal, Vol 26, pg 23-29. 2003.

[Be1]: B. C. Berndt. "Ramanujan`s Notebooks Part I". Springer-Verlag, New York Inc.(1985)

[Be2]: B.C. Berndt. "Ramanujan`s Notebooks Part II". Springer-Verlag, New York Inc.(1989) p. 262

[Ch]: C.K.Chui., "An Introduction to Wavelets", Academic Press, New York, 1992.

[KK]: Konrad Knopp. "Theory and Application of Infinite Series". Dover Publications, Inc. New York. 1990.

[T]: E.C.Titchmarsh, "Introduction to the theory of Fourier integrals". Oxford University Press, Amen House, London, 1948.

[Z]: I.J. Zucker, SIAM J. Math. Anal.**10**.192 (1979)