

q -ANALOG OF TABLEAU CONTAINMENT

JANG SOO KIM

ABSTRACT. We prove a q -analog of the following result due to McKay, Morse and Wilf: the probability that a random standard Young tableau of size n contains a fixed standard Young tableau of shape $\lambda \vdash k$ tends to $f^\lambda/k!$ in the large n limit, where f^λ is the number of standard Young tableaux of shape λ . We also consider the probability that a random pair (P, Q) of standard Young tableaux of the same shape contains a fixed pair (A, B) of standard Young tableaux.

1. INTRODUCTION


In 2002, McKay, Morse and Wilf [7] computed the large n limit of the probability that a random standard Young tableau of size n contains a fixed standard Young tableau. In this paper we obtain a q -analog of this result. In order to state the McKay–Morse–Wilf theorem and our generalization, we need to introduce several notations on permutations and standard Young tableaux.

We denote by \mathfrak{S}_n the set of permutations of $[n] = \{1, 2, \dots, n\}$. For a permutation $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$, let $\pi_{\leq k}$ (resp. $\pi_{>k}$) denote the permutation in \mathfrak{S}_k which is order-isomorphic to the subword of π consisting of the integers $i \leq k$ (resp. $i > k$). For example, if $\pi = 513697428$, then $\pi_{\leq 4} = 1342$ and $\pi_{>4} = 12534$. Similarly, let $\pi^{\leq k}$ (resp. $\pi^{>k}$) denote the permutation in \mathfrak{S}_k which is order-isomorphic to $\pi_1\pi_2 \cdots \pi_k$ (resp. $\pi_{k+1}\pi_{k+2} \cdots \pi_n$). For example, if $\pi = 513697428$, then $\pi^{\leq 4} = 3124$ and $\pi^{>4} = 53214$. If $\sigma = \pi_{\leq k}$ for some k , we say that π *contains* σ . Note that our permutation containment is different from the usual pattern containment, see [1].

We denote by \mathfrak{I}_n the set of involutions in \mathfrak{S}_n . For a permutation $\sigma \in \mathfrak{S}_k$, let $\mathfrak{I}_n(\sigma)$ denote the set of involutions in \mathfrak{I}_n containing σ . In other words,

$$\mathfrak{I}_n(\sigma) = \{\pi \in \mathfrak{I}_n : \pi_{\leq k} = \sigma\}.$$


A *partition* $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ is a weakly decreasing sequence of positive integers called *parts*. The sum of the parts of a partition λ is denoted by $|\lambda|$. If $|\lambda| = n$, we say that λ is a partition of n , also written as $\lambda \vdash n$. We denote by \emptyset the unique partition of 0. The *Ferrers diagram* of a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ is the left justified array of squares such that the i th row has λ_i squares. For example, the Ferrers diagram of $(4, 3, 1)$ is the following:



Key words and phrases. q -analog, tableau containment, permutation containment.

The author was supported by the SRC program of Korea Science and Engineering Foundation (KOSEF) grant funded by the Korea government (MEST) (No. R11-2007-035-01002-0).

We will identify a partition with its Ferrers diagram. Thus $\mu \subset \lambda$ means that the Ferrers diagram of μ is contained in the Ferrers diagram of λ . For two partitions λ and μ with $\mu \subset \lambda$, the *skew shape* λ/μ is the set theoretic difference $\lambda - \mu$. If $|\lambda| - |\mu| = n$, we write $\lambda/\mu \vdash n$. For example, the skew shape $\lambda/\mu \vdash 4$ for $\lambda = (4, 3, 2)$ and $\mu = (3, 2)$ is the following:



Occasionally we view a partition λ as the skew shape λ/\emptyset .

A *skew standard Young tableau* T , or a *skew SYT* T for short, is a filling of a skew shape $\lambda/\mu \vdash n$ with the integers in $[n]$ such that the integers are increasing along rows and columns. In this case we say that the *shape* of T is λ/μ , denoted by $\text{sh}(T) = \lambda/\mu$, and the *size* of T is n . If the shape T is a partition, T is called a *standard Young tableau* or *SYT* for short. We denote by \mathcal{T}_n the set of all SYTs of size n .

For a SYT T , let $T_{\leq k}$ denote the SYT obtained by removing all squares containing integers greater than k . Similarly let $T_{> k}$ denote the skew SYT obtained by removing all squares containing integers less than or equal to k and by decreasing all remaining integers by k . For example, if

$$(1) \quad T = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 4 & 7 \\ \hline 3 & 5 & 6 & \\ \hline 8 & 9 & & \\ \hline \end{array}$$

then $T_{\leq 5}$ and $T_{> 5}$ are the following:

$$T_{\leq 5} = \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \quad T_{> 5} = \begin{array}{|c|c|c|} \hline & & 2 \\ \hline & 1 & \\ \hline 3 & 4 & \\ \hline \end{array}$$

For two SYTs U and T , if $U = T_{\leq k}$ for some k , we say that T *contains* U . For a SYT A , we denote by $\mathcal{T}_n(A)$ the set of all SYTs of size n containing A .

We assume the reader is familiar with the Robinson–Schensted correspondence, see [8, 11]. If π corresponds to the pair of SYTs (P, Q) in the Robinson–Schensted correspondence, we write $\pi \xrightarrow{\text{RS}} (P, Q)$, and also $P(\pi) = P$ and $Q(\pi) = Q$. If π is an involution, we write $\pi \xleftrightarrow{\text{RS}} P$.

For given $A \in \mathcal{T}_m$, $\pi \in \mathcal{I}_n$ and $T \in \mathcal{T}_n$ with $\pi \xleftrightarrow{\text{RS}} T$, we can easily see that T contains A if and only if π contains σ for some $\sigma \in \mathfrak{S}_m$ with $P(\sigma) = A$. Thus the Robinson–Schensted correspondence induces the following bijection:

$$(2) \quad \text{RS} : \mathcal{T}_n(A) \rightarrow \bigcup_{\sigma: P(\sigma)=A} \mathcal{I}_n(\sigma).$$

McKay, Morse and Wilf [7] proved that for $\sigma \in \mathfrak{S}_m$,

$$(3) \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{I}_n(\sigma)|}{|\mathcal{I}_n|} = \frac{1}{m!}.$$

As a corollary, they obtained that the probability that a random SYT of size n contains A of shape $\alpha \vdash m$ tends to $f^\alpha/m!$ in the large n limit, where f^α denotes

the number of SYTs of shape α . In other words,

$$(4) \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A)|}{|\mathcal{T}_n|} = \frac{f^\alpha}{m!}.$$

Jaggard [4] defined the j -set (see Section 2) $J(\pi)$ of a permutation π and found the following exact formula for $|\mathcal{J}_{n+m}(\sigma)|$ for $\sigma \in \mathfrak{S}_m$:

$$(5) \quad |\mathcal{J}_{n+m}(\sigma)| = \sum_{\substack{j \in J(\sigma) \\ k = n - m + j}} \binom{n}{k} t_k,$$

where $t_k = |\mathcal{J}_k|$, the number of involutions in \mathfrak{S}_k . Using (5), Jaggard [4] found another proof of (3).

In this paper we find q -analogs of (3), (4) and (5). To state these, we need the following definitions.

For a permutation $\pi = \pi_1 \pi_2 \cdots \pi_n$, a *descent* of π is an integer $i \in [n - 1]$ such that $\pi_i > \pi_{i+1}$. For a skew SYT T of size n , a *descent* of T is an integer $i \in [n - 1]$ such that $i + 1$ is in a row lower than the row containing i . Let $D(\pi)$ (resp. $D(T)$) denote the set of all descents of π (resp. T). Let $\text{maj}(\pi)$ (resp. $\text{maj}(T)$) denote the sum of all descents of π (resp. T). For example, if $\pi = 513697428$, we have $D(\pi) = \{1, 5, 6, 7\}$ and $\text{maj}(\pi) = 19$. For the SYT T in (1), we have $D(T) = \{2, 4, 7\}$ and $\text{maj}(T) = 13$.

We define

$$\text{imaj}(\pi) = \text{maj}(\pi^{-1}), \quad A_n(p, q) = \sum_{\pi \in \mathfrak{S}_n} p^{\text{imaj}(\pi)} q^{\text{maj}(\pi)},$$

$$t_n(q) = \sum_{\pi \in \mathcal{J}_n} q^{\text{maj}(\pi)}, \quad f^{\lambda/\mu}(q) = \sum_{\text{sh}(T) = \lambda/\mu} q^{\text{maj}(T)},$$

$$[n]_q! = (1 + q)(1 + q + q^2) \cdots (1 + q + \cdots + q^{n-1}), \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n - k]_q!}.$$

Finally we can state our main results.

Theorem 1.1. *For $\sigma \in \mathfrak{S}_m$, we have*

$$\sum_{\pi \in \mathcal{J}_{n+m}(\sigma)} q^{\text{maj}(\pi^{>m})} = \sum_{\substack{j \in J(\sigma) \\ k = n - m + j}} q^{\text{maj}(\sigma^{>j})} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q).$$

For a real number $r > 0$, we define $\bar{r} = \min(r, r^{-1})$.

Theorem 1.2. *For $\sigma \in \mathfrak{S}_m$ and a real number $q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\sum_{\pi \in \mathcal{J}_n(\sigma)} q^{\text{maj}(\pi^{>m})}}{\sum_{\pi \in \mathcal{J}_n} q^{\text{maj}(\pi^{>m})}} = \frac{q^{\text{maj}(\sigma)} + (1 - \bar{q})C}{[m]_q! + (1 - \bar{q})D},$$

where C and D are polynomials in q and \bar{q} . (We refer the reader to Theorem 2.5 for their exact form.)

Theorem 1.3. *For a SYT A of shape $\alpha \vdash m$ and a real number $q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\sum_{T \in \mathcal{T}_n(A)} q^{\text{maj}(T_{>m})}}{\sum_{T \in \mathcal{T}_n} q^{\text{maj}(T_{>m})}} = \frac{f^\alpha(q) + (1 - \bar{q})E}{[m]_q! + (1 - \bar{q})D},$$

where E and D are polynomials in q and \bar{q} . (We refer the reader to Theorem 3.5 for their exact form.)

We can obtain similar results by considering pairs of SYTs. For pairs (P, Q) and (A, B) of SYTs, we say that (P, Q) *contains* (A, B) if P and Q contain A and B respectively. We denote by $\mathcal{T}_n(A, B)$ the set of pairs (P, Q) of SYTs of the *same shape* of size n containing (A, B) .

Given $A \in \mathcal{T}_a$, $B \in \mathcal{T}_b$ and $\pi \xrightarrow{\text{RS}} (P, Q)$, it is easy to see that (P, Q) contains (A, B) if and only if $\pi_{\leq a} = \sigma$ and $\pi^{\leq b} = \tau$ for some $\sigma \in \mathfrak{S}_a$ and $\tau \in \mathfrak{S}_b$ with $P(\sigma) = A$ and $Q(\tau) = B$. We denote by $\mathfrak{S}_n(\sigma, \tau)$ the set of $\pi \in \mathfrak{S}_n$ such that π contains σ and π^{-1} contains τ^{-1} . In other words, for $\sigma \in \mathfrak{S}_a$ and $\tau \in \mathfrak{S}_b$,

$$\mathfrak{S}_n(\sigma, \tau) = \{\pi \in \mathfrak{S}_n : \pi_{\leq a} = \sigma, \pi^{\leq b} = \tau\}.$$

Thus the Robinson–Schensted correspondence induces the following bijection:

$$(6) \quad RS : \mathcal{T}_n(A, B) \rightarrow \bigcup_{\substack{\sigma: P(\sigma)=A \\ \tau: Q(\tau)=B}} \mathfrak{S}_n(\sigma, \tau).$$

In Section 2 we define the j_2 -set $J(\sigma, \tau)$ of a pair (σ, τ) of permutations, and prove the following (p, q) -analogs of Theorems 1.1, 1.2 and 1.3.

Theorem 1.4. *Let a, b, n, m and ℓ be integers with $a + m = b + n = \ell$. Then, for $\sigma \in \mathfrak{S}_a$ and $\tau \in \mathfrak{S}_b$, we have*

$$\sum_{\pi \in \mathfrak{S}_\ell(\sigma, \tau)} p^{\text{imaj}(\pi_{>a})} q^{\text{maj}(\pi^{>b})} = \sum_{\substack{j \in J(\sigma, \tau) \\ k = n - a + j}} p^{\text{imaj}(\tau_{>j})} q^{\text{maj}(\sigma^{>j})} \begin{bmatrix} m \\ k \end{bmatrix}_p \begin{bmatrix} n \\ k \end{bmatrix}_q A_k(p, q).$$

In particular, when $p = q = 1$ we have

$$|\mathfrak{S}_\ell(\sigma, \tau)| = \sum_{\substack{j \in J(\sigma, \tau) \\ k = n - a + j}} \binom{m}{k} \binom{n}{k} k!.$$

Theorem 1.5. *For $\sigma \in \mathfrak{S}_a$, $\tau \in \mathfrak{S}_b$ and real numbers $p, q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\sum_{\pi \in \mathfrak{S}_n(\sigma, \tau)} p^{\text{imaj}(\pi_{>a})} q^{\text{maj}(\pi^{>b})}}{\sum_{\pi \in \mathfrak{S}_n} p^{\text{imaj}(\pi_{>a})} q^{\text{maj}(\pi^{>b})}} = \frac{p^{\text{imaj}(\tau)} q^{\text{maj}(\sigma)} + (1 - \bar{p})(1 - \bar{q})C'}{[b]_p! [a]_q! + (1 - \bar{p})(1 - \bar{q})D'},$$

where C' and D' are polynomials in p, \bar{p}, q and \bar{q} . (We refer the reader to Theorem 2.6 for their exact form.) In particular, when $p = 1$ or $p = q = 1$ we have

$$\lim_{n \rightarrow \infty} \frac{\sum_{\pi \in \mathfrak{S}_n(\sigma, \tau)} q^{\text{maj}(\pi^{>b})}}{\sum_{\pi \in \mathfrak{S}_n} q^{\text{maj}(\pi^{>b})}} = \frac{q^{\text{maj}(\sigma)}}{b! [a]_q!},$$

$$\lim_{n \rightarrow \infty} \frac{|\mathfrak{S}_n(\sigma, \tau)|}{|\mathfrak{S}_n|} = \frac{1}{a!b!}.$$

Theorem 1.6. *Let A, B be SYTs of shape $\alpha \vdash a, \beta \vdash b$ respectively. Then, for real numbers $p, q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\sum_{(P,Q) \in \mathcal{T}_n(A,B)} p^{\text{maj}(P_{>a})} q^{\text{maj}(Q_{>b})}}{\sum_{(P,Q) \in \mathcal{T}_n(\emptyset, \emptyset)} p^{\text{maj}(P_{>a})} q^{\text{maj}(Q_{>b})}} = \frac{f^\beta(p) f^\alpha(q) + (1 - \bar{p})(1 - \bar{q})E'}{[b]_p! [a]_q! + (1 - \bar{p})(1 - \bar{q})D'},$$

where E' and D' are polynomials in p, \bar{p}, q and \bar{q} . (We refer the reader to Theorem 3.6 for their exact form.) In particular, when $p = 1$ or $q = 1$ we have

$$\lim_{n \rightarrow \infty} \frac{\sum_{(P,Q) \in \mathcal{T}_n(A,B)} q^{\text{maj}(Q_{>b})}}{\sum_{(P,Q) \in \mathcal{T}_n(\emptyset, \emptyset)} q^{\text{maj}(Q_{>b})}} = \frac{f^\beta f^\alpha(q)}{b! [a]_q!},$$

$$\lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A, B)|}{|\mathcal{T}_n(\emptyset, \emptyset)|} = \frac{f^\alpha f^\beta}{a!b!}.$$

The rest of this paper is organized as follows. In Section 2, we define j_2 -sets and prove Theorems 1.1, 1.2, 1.4 and 1.5. In Section 3, we prove Theorems 1.3 and 1.6. In Section 4, we find a simple method to determine whether a given set is a j_2 -set. In Section 5, we prove a lemma which plays an important role in proving the limit theorems. In Section 6, we propose some open problems.

2. PERMUTATION CONTAINMENT

Recall the definitions of $\pi^{\leq k}, \pi^{>k}, \pi_{\leq k}$ and $\pi_{>k}$. These are easy to remember using the following argument. We can consider a permutation $\pi = \pi_1 \pi_2 \cdots \pi_n$ as a collection of *bi-letters* $\begin{smallmatrix} i \\ j \end{smallmatrix}$ as follows:

$$\pi = \left\{ \begin{array}{cccc} 1 & 2 & \cdots & n \\ \pi_1 & \pi_2 & & \pi_n \end{array} \right\}.$$

Then $\pi_{\leq k}$ (resp. $\pi_{>k}, \pi^{\leq k}$ and $\pi^{>k}$) is the permutation obtained from π by taking the bi-letters $\begin{smallmatrix} i \\ j \end{smallmatrix}$ with $j \leq k$ (resp. $j > k, i \leq k$ and $i > k$) and by relabeling them if necessary. It is easy to see that $(\pi^{\leq k})^{-1} = (\pi^{-1})_{\leq k}$.

In this section we prove Theorems 1.1, 1.2, 1.4 and 1.5 which are related to permutation containment. To do this, we start with a decomposition of permutation matrices.

For $\pi \in \mathfrak{S}_n$, the *permutation matrix* $M(\pi)$ is the $n \times n$ matrix whose (i, j) -entry is 1 if $\pi_i = j$, and 0 otherwise. For example,

$$M(4132) = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

Consider a 0-1 matrix M such that each row and column contains at most one 1. There exists a unique permutation π whose permutation matrix is obtained from

M by removing the rows and columns consisting of zeroes. In this case, we write $\pi \sim M$. If $\pi \sim M$ and $\pi \sim N$, we also write $M \sim N$. For example,

$$(7) \quad \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix} \sim 213.$$

For an $n \times m$ matrix M , we denote by $\text{row}(M)$ (resp. $\text{col}(M)$) the word $\mathbf{r} = r_1 r_2 \cdots r_n$ (resp. $\mathbf{c} = c_1 c_2 \cdots c_m$) of integers such that r_i (resp. c_i) is the sum of elements in the i th row (resp. column) of M . For example, if M is the second matrix in (7), then $\text{row}(M) = 1011$ and $\text{col}(M) = 11010$.

Let a, b, m, n , and ℓ be fixed integers with $a + m = b + n = \ell$. Consider a permutation $\pi \in \mathfrak{S}_\ell$. We divide the permutation matrix of π as follows:

$$M(\pi) = \begin{matrix} & a & m \\ b & \begin{pmatrix} M_{(1,1)} & M_{(1,2)} \\ M_{(2,1)} & M_{(2,2)} \end{pmatrix} \\ n & \end{matrix}$$

where the numbers outside the matrix indicate the sizes of the block matrices. Note that

$$\begin{aligned} \pi_{\leq a} &\sim \begin{pmatrix} M_{(1,1)} \\ M_{(2,1)} \end{pmatrix}, & \pi_{>a} &\sim \begin{pmatrix} M_{(1,2)} \\ M_{(2,2)} \end{pmatrix}, \\ \pi^{\leq b} &\sim \begin{pmatrix} M_{(1,1)} & M_{(1,2)} \end{pmatrix}, & \pi^{>b} &\sim \begin{pmatrix} M_{(2,1)} & M_{(2,2)} \end{pmatrix}. \end{aligned}$$

Assume that $M_{(1,1)}$ contains j 1's. Then $M_{(1,2)}$, $M_{(2,1)}$ and $M_{(2,2)}$ contain $b - j$, $a - j$ and $n - a + j$ 1's respectively. Then we define

$$\phi_{a,b}(\pi) = (\pi_{(1,1)}, \pi_{(1,2)}, \pi_{(2,1)}, \pi_{(2,2)}, \mathbf{c}_1, \mathbf{r}_1, \mathbf{c}_2, \mathbf{r}_2),$$

where $\pi_{(r,s)}$ is the permutation satisfying $\pi_{(r,s)} \sim M_{(r,s)}$ for $r = 1, 2$ and $s = 1, 2$, and $\mathbf{c}_1 = \text{col}(M_{(2,1)})$, $\mathbf{c}_2 = \text{col}(M_{(2,2)})$, $\mathbf{r}_1 = \text{col}(M_{(1,2)})$ and $\mathbf{r}_2 = \text{col}(M_{(2,2)})$. For example, if $\pi = 7152436$ with

$$M(\pi) = \left(\begin{array}{cc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right),$$

then

$$\phi_{2,3}(\pi) = (1, 21, 1, 213, 01, 101, 11010, 0111).$$

It is easy to see that the following is a bijection:

$$\phi_{a,b} : \mathfrak{S}_\ell \rightarrow \bigcup_{\substack{0 \leq j \leq a \\ k = n - a + j}} \mathfrak{S}_j \times \mathfrak{S}_{b-j} \times \mathfrak{S}_{a-j} \times \mathfrak{S}_k \times \binom{[a]}{a-j} \times \binom{[b]}{b-j} \times \binom{[m]}{k} \times \binom{[n]}{k},$$

where $\binom{[n]}{k}$ denotes the set of words consisting of k 1's and $n - k$ 0's.

For $\sigma \in \mathfrak{S}_a$, $\tau \in \mathfrak{S}_b$ and $\mathbf{r} \in \binom{[a+b]}{b}$, the *shuffle* $\text{sf}(\sigma, \tau; \mathbf{r})$ is the permutation in \mathfrak{S}_{a+b} obtained from \mathbf{r} by replacing the i th 0 to σ_i and the j th 1 to $a + \tau_j$ for $1 \leq i \leq a$ and $1 \leq j \leq b$. For example, $\text{sf}(3142, 231; 0010110) = 3164752$. The following lemma is due to Garsia and Gessel [3].

Lemma 2.1. For $\sigma \in \mathfrak{S}_a$, $\tau \in \mathfrak{S}_b$, we have

$$\sum_{\mathbf{r} \in \binom{[a+b]}{b}} q^{\text{maj}(\text{sf}(\sigma, \tau; \mathbf{r}))} = q^{\text{maj}(\sigma) + \text{maj}(\tau)} \begin{bmatrix} a+b \\ b \end{bmatrix}_q.$$

Jaggard [4] defined the j -set as follows. For a permutation π , the j -set $J(\pi)$ of π is defined to be the set of integers $j \geq 0$ such that $\pi^{\leq j}$ is an involution. In other words,

$$J(\pi) = \{j : \pi^{\leq j} = (\pi^{-1})_{\leq j}\}.$$

Note that we always have $0 \in J(\pi)$ because $\pi^{\leq 0} = (\pi^{-1})_{\leq 0}$ is the empty permutation.

As we have seen in (5), the j -set is useful to express the number of elements in $\mathfrak{J}_n(\sigma)$. We define the j_2 -set which is useful to express the number of elements in $\mathfrak{S}_n(\sigma, \tau)$.

Definition 2.1. The j_2 -set $J(\sigma, \tau)$ of a pair (σ, τ) of permutations is defined to be

$$J(\sigma, \tau) = \{j : \sigma^{\leq j} = \tau_{\leq j}\}.$$

Now we are ready to prove the following two theorems from which Theorems 1.1 and 1.4 follow.

Theorem 2.2. For $\sigma \in \mathfrak{S}_a$, we have

$$\begin{aligned} \sum_{\pi \in \mathfrak{J}_{n+a}} q^{\text{maj}(\pi^{>a})} &= \sum_{\substack{0 \leq j \leq a \\ k = n-a+j}} t_j \binom{a}{j} [a-j]_q! \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q), \\ \sum_{\pi \in \mathfrak{J}_{n+a}(\sigma)} q^{\text{maj}(\pi^{>a})} &= \sum_{\substack{j \in J(\sigma) \\ k = n-a+j}} q^{\text{maj}(\sigma^{>j})} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q). \end{aligned}$$

Proof. We omit the proof since it is similar to the proof of the following theorem. \square

Theorem 2.3. Let a, b, n, m and ℓ be integers with $a + m = b + n = \ell$. Then, for $\sigma \in \mathfrak{S}_a$ and $\tau \in \mathfrak{S}_b$, we have

$$\begin{aligned} \sum_{\pi \in \mathfrak{S}_\ell} p^{\text{imaj}(\pi^{>a})} q^{\text{maj}(\pi^{>b})} &= \sum_{\substack{0 \leq j \leq a \\ k = n-a+j}} j! \binom{a}{j} \binom{b}{j} [b-j]_p! [a-j]_q! \begin{bmatrix} m \\ k \end{bmatrix}_p \begin{bmatrix} n \\ k \end{bmatrix}_q A_k(p, q), \\ \sum_{\pi \in \mathfrak{S}_\ell(\sigma, \tau)} p^{\text{imaj}(\pi^{>a})} q^{\text{maj}(\pi^{>b})} &= \sum_{\substack{j \in J(\sigma, \tau) \\ k = n-a+j}} p^{\text{imaj}(\tau^{>j})} q^{\text{maj}(\sigma^{>j})} \begin{bmatrix} m \\ k \end{bmatrix}_p \begin{bmatrix} n \\ k \end{bmatrix}_q A_k(p, q). \end{aligned}$$

Proof. Fix an integer j with $0 \leq j \leq a$ and consider a permutation $\pi \in \mathfrak{S}_\ell$ such that $\pi_{(1,1)} \in \mathfrak{S}_j$, where

$$\phi_{a,b}(\pi) = (\pi_{(1,1)}, \pi_{(1,2)}, \pi_{(2,1)}, \pi_{(2,2)}, \mathbf{c}_1, \mathbf{r}_1, \mathbf{c}_2, \mathbf{r}_2).$$

Then $(\pi^{>a})^{-1} = \text{sf}(\pi_{(1,2)}^{-1}, \pi_{(2,2)}^{-1}; \mathbf{c}_2)$ and $\pi^{>b} = \text{sf}(\pi_{(2,1)}, \pi_{(2,2)}; \mathbf{r}_2)$. Thus

$$(8) \quad p^{\text{imaj}(\pi^{>a})} q^{\text{maj}(\pi^{>b})} = p^{\text{maj}(\text{sf}(\pi_{(1,2)}^{-1}, \pi_{(2,2)}^{-1}; \mathbf{c}_2))} q^{\text{maj}(\text{sf}(\pi_{(2,1)}, \pi_{(2,2)}; \mathbf{r}_2))}.$$

Let $k = n - a + j$. By Lemma 2.1, the sum of (8) over all $\pi_{(2,2)} \in \mathfrak{S}_k$, $\mathbf{c}_2 \in \binom{[m]}{k}$ and $\mathbf{r}_2 \in \binom{[n]}{k}$ equals

$$(9) \quad p^{\text{maj}(\pi_{(1,2)})} q^{\text{maj}(\pi_{(2,1)})} \begin{bmatrix} m \\ k \end{bmatrix}_p \begin{bmatrix} n \\ k \end{bmatrix}_q A_k(p, q).$$

Summing (9) over all j , $\pi_{(1,1)} \in \mathfrak{S}_j$, $\pi_{(1,2)} \in \mathfrak{S}_{b-j}$, $\pi_{(2,1)} \in \mathfrak{S}_{a-j}$, $\mathbf{c}_1 \in \binom{[a]}{a-j}$ and $\mathbf{r}_1 \in \binom{[b]}{b-j}$, and using the well known result $\sum_{\pi \in \mathfrak{S}_n} q^{\text{maj}(\pi)} = [n]_q!$, see [1, 10], we get the first identity.

If $\pi \in \mathfrak{S}_\ell(\sigma, \tau)$, then $\text{sf}(\pi_{(1,1)}, \pi_{(1,2)}; \mathbf{r}_1) = \tau$ and $\text{sf}(\pi_{(1,1)}^{-1}, \pi_{(2,1)}^{-1}; \mathbf{c}_1) = \sigma^{-1}$, which implies $\pi_{(1,1)} = \sigma^{\leq j} = \tau_{\leq j}$, $\pi_{(1,2)} = \tau_{> j}$ and $\pi_{(2,1)} = \sigma^{> j}$. Thus we have $j \in J(\sigma, \tau)$, and j determines $\pi_{(1,1)}$, $\pi_{(1,2)}$, $\pi_{(2,1)}$, \mathbf{c}_1 and \mathbf{r}_1 . Then we get the second identity by summing (9) over all $j \in J(\sigma, \tau)$. \square

To prove Theorems 1.2 and 1.5 we need the following lemma whose proof is given in Section 5. Recall that for a real number $r > 0$, we denote $\bar{r} = \min(r, r^{-1})$.

Lemma 2.4. *For real numbers $p, q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\frac{t_{n+1}(q)}{[n+1]_q!}}{\frac{t_n(q)}{[n]_q!}} = 1 - \bar{q}, \quad \lim_{n \rightarrow \infty} \frac{\frac{A_{n+1}(p, q)}{[n+1]_p! [n+1]_q!}}{\frac{A_n(p, q)}{[n]_p! [n]_q!}} = (1 - \bar{p})(1 - \bar{q}).$$

Now we prove the following theorem which implies Theorem 1.2.

Theorem 2.5. *For $\sigma \in \mathfrak{S}_m$ and a real number $q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\sum_{\pi \in \mathcal{J}_n(\sigma)} q^{\text{maj}(\pi^{> m})}}{\sum_{\pi \in \mathcal{J}_n} q^{\text{maj}(\pi^{> m})}} = \frac{\sum_{j \in J(\sigma)} q^{\text{maj}(\sigma^{> j})} \begin{bmatrix} m \\ j \end{bmatrix}_q [j]_q! (1 - \bar{q})^j}{\sum_{j=0}^m [m]_q! t_j \binom{m}{j} (1 - \bar{q})^j}.$$

Proof. By Theorem 2.2, the left hand side is equal to

$$(10) \quad \lim_{n \rightarrow \infty} \frac{\sum_{\pi \in \mathcal{J}_{n+m}(\sigma)} \frac{q^{\text{maj}(\pi^{> m})}}{[n]_q!}}{\sum_{\pi \in \mathcal{J}_{n+m}} \frac{q^{\text{maj}(\pi^{> m})}}{[n]_q!}} = \lim_{n \rightarrow \infty} \frac{\sum_{j \in J(\sigma)} \frac{q^{\text{maj}(\sigma^{> j})}}{[m-j]_q!} \frac{t_{n-m+j}(q)}{[n-m+j]_q!}}{\sum_{j=0}^m t_j \binom{m}{j} \frac{t_{n-m+j}(q)}{[n-m+j]_q!}}.$$

By Lemma 2.4, we have

$$\lim_{n \rightarrow \infty} \frac{\frac{t_{n-m+j}(q)}{[n-m+j]_q!}}{\frac{t_{n-m}(q)}{[n-m]_q!}} = (1 - \bar{q})^j.$$

Then, we get the theorem by dividing the numerator and the denominator of the right hand side of (10) by $t_{n-m}(q)/[n-m]_q!$, and by multiplying them by $[m]_q!$. \square

Similarly, we can prove the following theorem, which implies Theorem 1.5.

Theorem 2.6. For $\sigma \in \mathfrak{S}_a$, $\tau \in \mathfrak{S}_b$ and real numbers $p, q > 0$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\sum_{\pi \in \mathfrak{S}_n(\sigma, \tau)} p^{\text{maj}(\pi > a)} q^{\text{maj}(\pi > b)}}{\sum_{\pi \in \mathfrak{S}_n} p^{\text{maj}(\pi > a)} q^{\text{maj}(\pi > b)}} \\ = \frac{\sum_{j \in J(\sigma, \tau)} p^{\text{maj}(\tau > j)} q^{\text{maj}(\sigma > j)} \begin{bmatrix} b \\ j \end{bmatrix}_p \begin{bmatrix} a \\ j \end{bmatrix}_q [j]_p! [j]_q! (1 - \bar{p})^j (1 - \bar{q})^j}{\sum_{j=0}^a [b]_p! [a]_q! j! \binom{a}{j} \binom{b}{j} (1 - \bar{p})^j (1 - \bar{q})^j}. \end{aligned}$$

Note that when $p = q = 1$ in Theorem 2.6 we have

$$(11) \quad \lim_{\ell \rightarrow \infty} \frac{|\mathfrak{S}_n(\sigma, \tau)|}{|\mathfrak{S}_\ell|} = \frac{1}{a!b!},$$

which means that the probability that a random permutation $\pi \in \mathfrak{S}_\ell$ contains σ and π^{-1} contains τ^{-1} tends to $1/a!b!$ as ℓ approaches infinity. This statement fits well with the following intuition. If we divide $M(\pi)$ as

$$M(\pi) = \begin{matrix} & a & m \\ b & \begin{pmatrix} M_{(1,1)} & M_{(1,2)} \\ M_{(2,1)} & M_{(2,2)} \end{pmatrix} \end{matrix},$$

where $m = \ell - a$ and $n = \ell - b$, then the probability that $M_{(1,1)}$ contains a 1 will be very low when ℓ is very large. Thus the probability that $M_{(2,1)} \sim \sigma$ and $M_{(1,2)} \sim \tau$ will be very close to $1/a!$ and $1/b!$ respectively, which is consistent with (11).

3. TABLEAU CONTAINMENT

Jaggard [4] proved that for a SYT A of shape α and an integer j ,

$$(12) \quad \#\{\sigma : P(\sigma) = A, j \in J(\sigma)\} = \sum_{\mu \vdash j} f^{\alpha/\mu}.$$

We have the following analog of (12): given SYTs A and B of shape α and β respectively and an integer j , we have

$$(13) \quad \#\{(\sigma, \tau) : P(\sigma) = A, Q(\tau) = B, j \in J(\sigma, \tau)\} = \sum_{\mu \vdash j} f^{\beta/\mu} f^{\alpha/\mu}.$$

Using (12), Jaggard [4] gave another proof of the following formula due to Sagan and Stanley [9]:

$$(14) \quad \sum_{\lambda/\alpha \vdash n} f^{\lambda/\alpha} = \sum_{k \geq 0} \binom{n}{k} t_k \sum_{\alpha/\mu \vdash n-k} f^{\alpha/\mu}.$$

Similarly using (13) we can prove the following formula also due to Sagan and Stanley [9]:

$$(15) \quad \sum_{\substack{\lambda/\alpha \vdash m \\ \lambda/\beta \vdash n}} f^{\lambda/\alpha} f^{\lambda/\beta} = \sum_{k \geq 0} \binom{m}{k} \binom{n}{k} k! \sum_{\substack{\beta/\mu \vdash m-k \\ \alpha/\mu \vdash n-k}} f^{\beta/\mu} f^{\alpha/\mu}.$$

In this section we generalize these results and prove Theorems 1.3 and 1.6.

We will use the following well known fact: for given $\pi \xleftrightarrow{\text{RS}} (P, Q)$ we have $D(\pi^{-1}) = D(P)$ and $D(\pi) = D(Q)$, see [11]. Thus $q^{\text{maj}((\pi^{-1})^{>a})} = q^{\text{maj}(P_{>a})}$ and $q^{\text{maj}(\pi^{>b})} = q^{\text{maj}(Q_{>b})}$ for all nonnegative integers a and b .

We start with the following two lemmas which are respectively a q -analog of (12) and a (p, q) -analog of (13).

Lemma 3.1. *For a SYT A of shape α and an integer j , we have*

$$\sum_{\sigma: \begin{cases} P(\sigma)=A \\ j \in J(\sigma) \end{cases}} q^{\text{maj}(\sigma^{>j})} = \sum_{\mu \vdash j} f^{\alpha/\mu}(q).$$

Proof. Since it is similar to the proof of the following lemma, we omit it. \square

Lemma 3.2. *For SYTs A and B of shape α and β respectively and an integer j , we have*

$$\sum_{(\sigma, \tau): \begin{cases} P(\sigma)=A \\ Q(\tau)=B \\ j \in J(\sigma, \tau) \end{cases}} p^{\text{imaj}(\tau_{>j})} q^{\text{maj}(\sigma^{>j})} = \sum_{\mu \vdash j} f^{\beta/\mu}(p) f^{\alpha/\mu}(q).$$

Proof. Let $X = \{(\sigma, \tau) : P(\sigma) = A, Q(\tau) = B, j \in J(\sigma, \tau)\}$ and $Y = \{(U, V) : \mu \vdash j, \text{sh}(U) = \beta/\mu, \text{sh}(V) = \alpha/\mu\}$. It is sufficient to find a bijection $\xi : X \rightarrow Y$ such that if $\xi(\sigma, \tau) = (U, V)$, then $\text{imaj}(\tau_{>j}) = \text{maj}(U)$ and $\text{maj}(\sigma^{>j}) = \text{maj}(V)$.

We define $\xi(\sigma, \tau) = (U, V)$ by $U = P(\tau)_{>j}$ and $V = Q(\sigma)_{>j}$. Then we have $\text{imaj}(\tau_{>j}) = \text{maj}(U)$ and $\text{maj}(\sigma^{>j}) = \text{maj}(V)$.

To prove ξ is a bijection, it is sufficient to show that for $(U, V) \in Y$ there exists a unique pair $(\sigma, \tau) \in X$ satisfying $\xi(\sigma, \tau) = (U, V)$. Let $\alpha \vdash a$. Since $P(\sigma) = A, Q(\sigma)_{>j} = V$, by reversing the insertion algorithm $a-j$ times, we can find $\sigma_a, \sigma_{a-1}, \dots, \sigma_{j+1}$ and $P(\sigma^{\leq j})$. Since $P(\tau)_{\leq j} = P(\tau_{\leq j}) = P(\sigma^{\leq j})$ and $P(\tau)_{>j} = U$, we can determine $P(\tau)$. Thus we get τ , from which we can determine $\sigma^{\leq j} = \tau_{\leq j}$. Thus we get σ , and there is a unique pair $(\sigma, \tau) \in X$ with $\xi(\sigma, \tau) = (U, V)$. \square

Now we have the following q -analog of (14) and (p, q) -analog of (15).

Theorem 3.3. *For a partition α and an integer n , we have*

$$\sum_{\lambda/\alpha \vdash n} f^{\lambda/\alpha}(q) = \sum_{k \geq 0} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q) \sum_{\alpha/\mu \vdash n-k} f^{\alpha/\mu}(q).$$

Proof. Let $\alpha \vdash a$ and let A be a SYT of shape α . Then the left hand side is equal to

$$\begin{aligned}
 \sum_{T \in \mathcal{T}_{n+a}(A)} q^{\text{maj}(T_{>a})} &= \sum_{\sigma: P(\sigma)=A} \sum_{\pi \in \mathcal{J}_{n+a}(\sigma)} q^{\text{maj}(\pi^{>a})} \\
 &= \sum_{\sigma: P(\sigma)=A} \sum_{\substack{j \in J(\sigma) \\ k=n-a+j}} q^{\text{maj}(\sigma^{>j})} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q) \quad (\text{by Theorem 2.2}) \\
 &= \sum_{\substack{0 \leq j \leq a \\ k=n-a+j}} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q) \sum_{\sigma: \begin{cases} P(\sigma)=A \\ j \in J(\sigma) \end{cases}} q^{\text{maj}(\sigma^{>j})} \\
 &= \sum_{\substack{0 \leq j \leq a \\ k=n-a+j}} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q) \sum_{\mu \vdash j} f^{\alpha/\mu}(q) \quad (\text{by Lemma 3.1}) \\
 &= \sum_{k \geq 0} \begin{bmatrix} n \\ k \end{bmatrix}_q t_k(q) \sum_{\alpha/\mu \vdash n-k} f^{\alpha/\mu}(q).
 \end{aligned}$$

□

Theorem 3.4. *For partitions α, β and integers m, n , we have*

$$\sum_{\substack{\lambda/\alpha \vdash m \\ \lambda/\beta \vdash n}} f^{\lambda/\alpha}(p) f^{\lambda/\beta}(q) = \sum_{k \geq 0} \begin{bmatrix} m \\ k \end{bmatrix}_p \begin{bmatrix} n \\ k \end{bmatrix}_q A_k(p, q) \sum_{\substack{\beta/\mu \vdash m-k \\ \alpha/\mu \vdash n-k}} f^{\beta/\mu}(p) f^{\alpha/\mu}(q).$$

Proof. This can be done similarly as in the proof of Theorem 3.3. □

We note that Theorems 3.3 and 3.4 can also be proved using the following identities of skew Schur functions:

$$\begin{aligned}
 \sum_{\lambda} s_{\lambda/\alpha}(\mathbf{x}) &= \sum_{\mu} s_{\alpha/\mu}(\mathbf{x}) \prod_i (1 - x_i)^{-1} \prod_{i < j} (1 - x_i x_j)^{-1}, \\
 \sum_{\lambda} s_{\lambda/\alpha}(\mathbf{x}) s_{\lambda/\beta}(\mathbf{y}) &= \prod_{i,j} (1 - x_i y_j)^{-1} \sum_{\mu} s_{\beta/\mu}(\mathbf{x}) s_{\alpha/\mu}(\mathbf{y}).
 \end{aligned}$$

Now we can prove the following theorem, which implies Theorem 1.3.

Theorem 3.5. *For a SYT A of shape $\alpha \vdash m$ and a real number $q > 0$, we have*

$$\lim_{n \rightarrow \infty} \frac{\sum_{T \in \mathcal{T}_n(A)} q^{\text{maj}(T_{>m})}}{\sum_{T \in \mathcal{T}_n} q^{\text{maj}(T_{>m})}} = \frac{\sum_{j=0}^m \begin{bmatrix} m \\ j \end{bmatrix}_q [j]_q! (1 - \bar{q})^j \sum_{\mu \vdash j} f^{\alpha/\mu}(q)}{\sum_{j=0}^m [m]_q! t_j \binom{m}{j} (1 - \bar{q})^j}.$$

Proof. By the Robinson–Schensted correspondence (2), we have

$$\lim_{n \rightarrow \infty} \frac{\sum_{T \in \mathcal{T}_n(A)} q^{\text{maj}(T_{>m})}}{\sum_{T \in \mathcal{T}_n} q^{\text{maj}(T_{>m})}} = \lim_{n \rightarrow \infty} \frac{\sum_{\sigma: P(\sigma)=A} \sum_{\pi \in \mathcal{J}_n(\sigma)} q^{\text{maj}(\pi^{>m})}}{\sum_{\pi \in \mathcal{J}_n} q^{\text{maj}(\pi^{>m})}}.$$

By Theorem 2.5, the above limit is equal to

$$\frac{\sum_{\sigma: P(\sigma)=A} \sum_{j \in J(\sigma)} q^{\text{maj}(\sigma^{>j})} \begin{bmatrix} m \\ j \end{bmatrix}_q [j]_q! (1-\bar{q})^j}{\sum_{j=0}^m [m]_q! t_j \binom{m}{j} (1-\bar{q})^j} = \frac{\sum_{j=0}^m \begin{bmatrix} m \\ j \end{bmatrix}_q [j]_q! (1-\bar{q})^j \sum_{\sigma: \begin{cases} P(\sigma)=A \\ j \in J(\sigma) \end{cases}} q^{\text{maj}(\sigma^{>j})}}{\sum_{j=0}^m [m]_q! t_j \binom{m}{j} (1-\bar{q})^j}.$$

By Lemma 3.1, we are done. \square

Similarly, we get the following, which implies Theorem 1.6.

Theorem 3.6. *Let A and B be SYTs of shape $\alpha \vdash a$ and $\beta \vdash b$ respectively. Then for real number $p, q > 0$, we have*

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\sum_{(P,Q) \in \mathcal{T}_n(A,B)} p^{\text{maj}(P_{>a})} q^{\text{maj}(Q_{>b})}}{\sum_{(P,Q) \in \mathcal{T}_n(\emptyset, \emptyset)} p^{\text{maj}(P_{>a})} q^{\text{maj}(Q_{>b})}} \\ = \frac{\sum_{j=0}^a \begin{bmatrix} b \\ j \end{bmatrix}_p \begin{bmatrix} a \\ j \end{bmatrix}_q [j]_p! [j]_q! (1-\bar{p})^j (1-\bar{q})^j \sum_{\mu \vdash j} f^{\beta/\mu}(p) f^{\alpha/\mu}(q)}{\sum_{j=0}^a [b]_p! [a]_q! j! \binom{a}{j} \binom{b}{j} (1-\bar{p})^j (1-\bar{q})^j}. \end{aligned}$$

Let us consider the special cases $q = 1$ and $p = q = 1$ of Theorems 3.5 and 3.6, which are

$$(16) \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A)|}{|\mathcal{T}_n|} = \frac{f^\alpha}{a!}.$$

$$(17) \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A, B)|}{|\mathcal{T}_n(\emptyset, \emptyset)|} = \frac{f^\alpha f^\beta}{a! b!}.$$

As mentioned in the introduction, (16) is first proved by McKay et al. [7]. To our knowledge (17) is new. For the rest of this section we give another proofs of (16) and (17) using (14) and (15).

Using the well known asymptotic behavior of $t_n \sim \frac{1}{\sqrt{2}} n^{n/2} \exp(-\frac{n}{2} + \sqrt{n} - \frac{1}{4})$, we can easily see that

$$\lim_{n \rightarrow \infty} \frac{t_n}{t_{n+1}} = 0.$$

Thus

$$\lim_{n \rightarrow \infty} \frac{nt_{n-1}}{t_{n+1}} = \lim_{n \rightarrow \infty} \frac{t_{n+1} - t_n}{t_{n+1}} = 1.$$

Using induction one can easily prove that

$$(18) \quad \lim_{n \rightarrow \infty} \frac{n^a t_{n-a}}{t_{n+a}} = \lim_{n \rightarrow \infty} \frac{n^a t_n}{t_{n+2a}} = 1.$$

By (14), we have

$$\lim_{n \rightarrow \infty} \frac{|\mathcal{J}_n(\sigma)|}{|\mathcal{J}_n|} = \lim_{n \rightarrow \infty} \frac{\sum f^{\lambda/\alpha}}{t_{n+a}} = \lim_{n \rightarrow \infty} \frac{\sum_{k \geq 0} \binom{n}{k} t_k \sum_{\alpha/\mu \vdash n-k} f^{\alpha/\mu}}{t_{n+a}}.$$

Since $\alpha/\mu \vdash n - k$, we only need to consider k with $n - a \leq k \leq n$. Using (18), it is not difficult to see that for $k > n - a$, we have

$$\lim_{n \rightarrow \infty} \frac{\binom{n}{k} t_k}{t_{n+a}} = 0,$$

and

$$\lim_{n \rightarrow \infty} \frac{\binom{n}{n-a} t_{n-a}}{t_{n+a}} = \frac{1}{a!}.$$

This gives another proof of (16).

The same argument can be applied to (17). More precisely, by (15),

$$\lim_{\ell \rightarrow \infty} \frac{|\mathcal{T}_\ell(A, B)|}{|\mathcal{T}_\ell(\emptyset, \emptyset)|} = \lim_{\ell \rightarrow \infty} \frac{\sum_{\lambda/\alpha \vdash m} f^{\lambda/\alpha} f^{\lambda/\beta}}{\ell!} = \lim_{\ell \rightarrow \infty} \frac{\sum_{k \geq 0} \binom{m}{k} \binom{n}{k} k! \sum_{\substack{\beta/\mu \vdash m-k \\ \alpha/\mu \vdash n-k}} f^{\beta/\mu} f^{\alpha/\mu}}{\ell!},$$

where $\ell = a + m = b + n$, $\text{sh}(A) = \alpha \vdash a$ and $\text{sh}(B) = \beta \vdash b$. By the condition $\beta/\mu \vdash m - k$, we only need to consider the integers k with $k \geq m - b = n - a = \ell - a - b$. It is easy to see that for $k > m - b$, we have

$$\lim_{\ell \rightarrow \infty} \frac{\binom{m}{k} \binom{n}{k} k!}{\ell!} = 0,$$

and

$$\lim_{\ell \rightarrow \infty} \frac{\binom{m}{m-b} \binom{n}{n-a} (\ell - a - b)!}{\ell!} = \frac{1}{a!b!}.$$

This gives another proof of (17).

We note that in fact the above two proofs are special cases of the proofs of Theorems 3.5 and 3.6.

4. CRITERION FOR j_2 -SETS

In this section we find a criterion for a set to be a j_2 -set. First, we review some previous results on j -sets. Throughout this section we assume that n , m and k are positive integers.

Kim and Kim [5] proved the following theorem.

Theorem 4.1. *Let J be a j -set with largest element $m \geq 2$. Then, for $n > m$, $J \cup \{n\}$ is a j -set if and only if $n = m + 1$ or $n - m \geq m - \max(J \cap [m - 2])$.*

Using Theorem 4.1 we can easily determine whether a given set is a j -set. To do this we need the following definition introduced by Corteel and Lovejoy [2]. An *overpartition* is a weakly decreasing sequence of positive integers in which the last occurrence of an integer may be overlined.

For a set $S = \{s_0, s_1, s_2, \dots, s_n\}$ of nonnegative integers with $s_0 < s_1 < \dots < s_n$, we define $\Delta(S)$ to be the sequence (a_1, a_2, \dots, a_n) where $a_i = s_{n-i+1} - s_{n-i}$ for $i \in [n]$. For example, if

$$(19) \quad S = \{0, 1, 2, 3, 5, 6, 9, 13, 17, 18, 19, 20, 22\},$$

then

$$(20) \quad \Delta(S) = (2, 1, 1, 1, 4, 4, 3, 1, 2, 1, 1, 1).$$

And then we define $\overline{\Delta}(S)$ to be the sequence obtained from $\Delta(S) = (a_1, a_2, \dots, a_n)$ by applying the following algorithm.

First we set $k = 1$. Find the smallest index i such that $a_i = 1$ and $k \leq i < n$. If we have no such i , then we finish the algorithm. Otherwise we replace a_i and a_{i+1} with $\overline{a_{i+1} + 1}$. Set $k = i + 2$ and repeat this process.

For example, for $\Delta(S)$ in (20), we have

$$(21) \quad \overline{\Delta}(S) = (2, \overline{2}, \overline{5}, 4, 3, \overline{3}, \overline{2}, 1).$$

Assume that we have $\overline{\Delta}(S) = (b_1, \dots, b_m)$. Let $\mathbf{i} = \{i_1, i_2, \dots, i_k\}$ be the set of integers with $1 \leq i_1 < i_2 < \dots < i_k \leq m$ such that $b_j = \overline{2}$ if and only if $j \in \mathbf{i}$. We define $\psi(S)$ to be the sequence $(\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_{k+1})$, where $\mathbf{s}_j = (b_{i_{j-1}+1}, b_{i_{j-1}+2}, \dots, b_{i_j})$ for $j \in [k+1]$, and $i_0 = 0$ and $i_{k+1} = n$.

For example, for S in (19), we have

$$(22) \quad \psi(S) = ((2, \overline{2}), (\overline{5}, 4, 3, \overline{3}, \overline{2}), (1)).$$

With the above observation Kim and Kim [5] proved the following.

Corollary 4.2. *Let S be a set of nonnegative integers with $\psi(S) = (\mathbf{s}_1, \dots, \mathbf{s}_{k+1})$. Then S is a j -set if and only if $0 \in S$, \mathbf{s}_i is an overpartition ending with $\overline{2}$ for all $i \in [k]$ and $\mathbf{s}_{k+1} = (1)$ or $\mathbf{s}_{k+1} = \emptyset$.*

Hence, the set S in (19) is a j -set because $\psi(S)$ in (22) satisfies the condition in Corollary 4.2.

Note that since $J(\pi) = J(\pi, \pi^{-1})$, every j -set is also a j_2 -set. However, the converse is not true. For example, the j_2 -set $J(312, 312) = \{0, 1, 3\}$ is not a j -set.

We have the following theorem analogous to Theorem 4.1.

Theorem 4.3. *Let J be a j_2 -set such that the two largest elements of J are $n - k$ and n . Then $J \cup \{n + m\}$ is a j_2 -set if and only if $m = 1$ or $m \geq k$.*

Before proving Theorem 4.3 we will see how to determine whether a given set is a j_2 -set.

Consider a set S of integers with $\Delta(S) = (a_1, a_2, \dots, a_n)$. Let $\mathbf{i} = \{i_1, i_2, \dots, i_k\}$ be the set of integers with $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $a_j = 1$ if and only if $j \in \mathbf{i}$. We define $\psi_2(S)$ to be the sequence $(\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_k)$, where $\mathbf{s}_j = (a_{i_{j-1}+1}, a_{i_{j-1}+2}, \dots, a_{i_j})$ for $j \in [k]$ and $i_0 = 0$. For example, if

$$(23) \quad S = \{0, 1, 3, 6, 7, 8, 12, 13, 14, 15, 17\},$$

then

$$\begin{aligned} \Delta(S) &= (2, 1, 1, 1, 4, 1, 1, 3, 2, 1), \\ \psi_2(S) &= ((2, 1), (1), (1), (4, 1), (1), (3, 2, 1)). \end{aligned}$$

It is easy to see that Theorem 4.3 implies the following.

Corollary 4.4. *Let S be a set of nonnegative integers with $\psi_2(S) = (\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_k)$. Then S is a j_2 -set if and only if $0 \in S$ and \mathbf{s}_i is a partition with exactly one part equal to 1 for all $i \in [k]$.*

By Corollary 4.4, the set S in (23) is a j_2 -set. Using Corollary 4.4, we can easily get the following generating function for the number $j_2(n)$ of j_2 -sets with largest element n .

Corollary 4.5. *We have*

$$\begin{aligned} \sum_{n \geq 0} j_2(n)x^n &= \frac{1}{1-x \prod_{i \geq 2} \frac{1}{1-x^i}} \\ &= 1 + x + x^2 + 2x^3 + 4x^4 + 8x^5 + 15x^6 + 29x^7 + 55x^8 + 105x^9 \\ &\quad + 200x^{10} + 381x^{11} + 725x^{12} + 1381x^{13} + 2629x^{14} + 5005x^{15} + \dots \end{aligned}$$

For the rest of this section we prove Theorem 4.3. Our proof is similar to, but simpler than, the proof of Theorem 4.1 in [5].

Note that if J is a j_2 -set, then $J \cap [k]$ is also a j_2 -set for all integers k .

Proposition 4.6. *Let J be a j_2 -set with largest element n . Then there is a permutation in \mathfrak{S}_n such that $J(\pi, \pi) = J$.*

Proof. Let (σ, τ) be a pair with $J = J(\sigma, \tau)$. Since $\sigma^{\leq n} = \tau_{\leq n}$, if we set $\pi = \sigma^{\leq n}$, we have $J(\pi, \pi) = J$. \square

Recall that, for 0-1 matrices M and N , we write $M \sim N$ if the matrices obtained from M and N by removing rows and columns consisting of zeroes are the same.

To prove Theorem 4.3 we need the following four lemmas.

Lemma 4.7. *Let J be a j_2 -set such that the three largest elements of J are $n - k$, n and $n + m$ with $m \geq 2$. Then $m \geq k$.*

Proof. Consider $\pi \in \mathfrak{S}_{n+m}$ with $J(\pi, \pi) = J$. We divide $M(\pi)$ as follows:

$$M(\pi) = \begin{matrix} & n & m \\ n & \begin{pmatrix} A & B \\ C & D \end{pmatrix} \\ m & \end{matrix}$$

Since $n \in J(\pi, \pi)$, we have $\begin{pmatrix} A & B \\ C \end{pmatrix} \sim \begin{pmatrix} A \\ C \end{pmatrix}$. Let $\sigma = \pi^{\leq n} = \pi_{\leq n}$. Assume that B has s nonzero entries. Then C also has s nonzero entries. If $s = 0$, then we get $n + 1 \in J(\pi, \pi)$ because $\pi^{\leq n+1} \sim \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} \sim \pi_{\leq n+1}$. But this is a contradiction to $m \geq 2$. Thus $s \geq 1$. Since $\sigma^{\leq n-s} \sim A \sim \sigma_{\leq n-s}$, we get $\sigma^{\leq n-s} = \sigma_{\leq n-s}$. Thus

$$\pi^{\leq n-s} = (\pi^{\leq n})^{\leq n-s} = \sigma^{\leq n-s} = \sigma_{\leq n-s} = (\pi_{\leq n})_{\leq n-s} = \pi_{\leq n-s}$$

and we get $n - s \in J(\pi, \pi)$. Since $n - k$ is the largest element in $J \cap [n - 1]$, we get $n - s \leq n - k$. Since B has at most m nonzero entries, we get $k \leq s \leq m$. \square

Lemma 4.8. *Let J be a j_2 -set such that the two largest elements of J are $n - k$ and n . Then $J \cup \{n + k\}$ is a j_2 -set.*

Proof. If $k = 1$, then it is clear. Assume $k \geq 2$. Consider $\sigma \in \mathfrak{S}_n$ with $J(\sigma, \sigma) = J$. Let π be the unique permutation in \mathfrak{S}_{n+k} satisfying

$$(24) \quad M(\pi) = \begin{matrix} & n & k \\ n & \begin{pmatrix} A & C \\ B & \mathbf{0} \end{pmatrix}, \\ k & \end{matrix}$$

where A, B and C are the matrices of size $n \times n$, $k \times n$ and $n \times k$ respectively such that $M(\sigma) \sim \begin{pmatrix} A \\ B \end{pmatrix} \sim \begin{pmatrix} A & C \end{pmatrix}$. The condition $n - k \in J(\sigma, \sigma)$ guarantees that there is a unique π . Since $J(\pi, \pi) \cap [n] = J$ and $n + k \in J(\pi, \pi)$, it is sufficient to show that $n + s \notin J(\pi, \pi)$ for all $1 \leq s < k$. Suppose $n + s \in J(\pi, \pi)$ for some $1 \leq s < k$. Then we have

$$(25) \quad \begin{pmatrix} A & C \\ B' & \mathbf{0} \end{pmatrix} \sim \begin{pmatrix} A & C' \\ B & \mathbf{0} \end{pmatrix},$$

where B' (resp. C') is the matrix consisting of the first s rows of B (resp. columns of C). Removing the last $k - s$ nonzero rows and columns of the matrices in both sides of (25), we get $\sigma^{\leq n-k+s} = \sigma_{\leq n-k+s}$, i.e., $n - k + s \in J(\sigma, \sigma) = J$, which is a contradiction to the assumption that $n - k$ and n are the two largest elements of J . \square

Lemma 4.9. *Let J be a j_2 -set such that the two largest elements of J are $n - k$ and n . If $k \geq 2$, then $(J \setminus \{n\}) \cup \{n + 1\}$ is a j_2 -set.*

Proof. Consider $\sigma = \sigma_1 \sigma_2 \cdots \sigma_n \in \mathfrak{S}_n$ with $J(\sigma, \sigma) = J$. Since $n - 1 \notin J(\sigma, \sigma)$, we have $\sigma_n \neq n$. Let $\pi \in \mathfrak{S}_{n+1}$ be the permutation such that

$$\pi_i = \begin{cases} \sigma_i & \text{if } i < n \text{ and } \sigma_i < n, \\ n + 1 & \text{if } i < n \text{ and } \sigma_i = n, \\ n & \text{if } i = n, \\ \sigma_n & \text{if } i = n + 1. \end{cases}$$

Then $M(\sigma)$ and $M(\pi)$ are decomposed as follows:

$$M(\sigma) = \begin{array}{|c|c|} \hline A & C \\ \hline B & \mathbf{0} \\ \hline \end{array}, \quad M(\pi) = \begin{array}{|c|c|c|} \hline & 0 & \\ \hline A & \vdots & C \\ \hline & 0 & \\ \hline 0 & \cdots & 0 & 1 & 0 \\ \hline B & & 0 & 0 & \\ \hline \end{array},$$

where A, B and C are $(n - 1) \times (n - 1)$, $1 \times (n - 1)$ and $(n - 1) \times 1$ matrices respectively. It is not difficult to see that $J(\pi, \pi) = (J \setminus \{n\}) \cup \{n + 1\}$. \square

Lemma 4.10. *Let J be a j_2 -set such that the two largest elements of J are $n - 1$ and n . Then $J \cup \{n + k\}$ is a j_2 -set for all positive integers k .*

Proof. It is clear if $k = 1$. Assume $k \geq 2$. Consider $\sigma \in \mathfrak{S}_{n-1}$ with $J(\sigma, \sigma) = J \setminus \{n\}$. Let π be the permutation in \mathfrak{S}_{n+k} such that $M(\pi) = \begin{pmatrix} M(\sigma) & \mathbf{0} \\ \mathbf{0} & A \end{pmatrix}$,

where $A = \begin{pmatrix} \mathbf{0} & I_{k-1} \\ 1 & \mathbf{0} \end{pmatrix}$ and I_{k-1} is the $(k - 1) \times (k - 1)$ identity matrix. Then $J(\pi, \pi) = J \cup \{n + k\}$. \square

Now we can prove Theorem 4.3.

Proof of Theorem 4.3. Assume that $J' = J \cup \{n + m\}$ is a j_2 -set. Then the three largest integers of J' are $n - k, n$ and $n + m$. If $m = 1$, we are done. If $m \geq 2$, then by Lemma 4.7 we have $m \geq k$.

Now assume that $m = 1$ or $m \geq k$. If $m = 1$, then clearly $J \cup \{n + m\}$ is a j_2 -set. Assume that $m \geq k$. If $k = 1$, then by Lemma 4.10 $J \cup \{n + m\}$ is a j_2 -set. If $k \geq 2$, then by Lemmas 4.8 and 4.9 we have that $J \cup \{n + i\}$ is a j_2 -set for all integers $i \geq k$. In particular $J \cup \{n + m\}$ is a j_2 -set. \square

5. PROOF OF LEMMA 2.4

In this section we prove Lemma 2.4. First we prove the following three lemmas.

Lemma 5.1. *We have*

$$\frac{t_n(q^{-1})}{[n]_{q^{-1}}!} = \frac{t_n(q)}{[n]_q!},$$

$$\frac{A_n(p^{-1}, q)}{[n]_{p^{-1}}! [n]_q!} = \frac{A_n(p, q^{-1})}{[n]_p! [n]_{q^{-1}}!} = \frac{A_n(p^{-1}, q^{-1})}{[n]_{p^{-1}}! [n]_{q^{-1}}!} = \frac{A_n(p, q)}{[n]_p! [n]_q!}.$$

Proof. For $T \in \mathcal{T}_n$ we have $\text{maj}(T) + \text{maj}(T') = \binom{n}{2}$, where T' is the transpose of T . Thus

$$t_n(q^{-1}) = \sum_{T \in \mathcal{T}_n} q^{-\text{maj}(T)} = q^{-\binom{n}{2}} \sum_{T \in \mathcal{T}_n} q^{\text{maj}(T')} = q^{-\binom{n}{2}} t_n(q).$$

Since $q^{-\binom{n}{2}} / [n]_{q^{-1}}! = 1 / [n]_q!$, we get the first identity. The rest identities can be proved similarly. \square

Lemma 5.2. *For a real number $0 < q < 1$, we have*

$$\log \left(\prod_{i \geq 1} (1 - q^i)^{-i} \right) < \left(1 + \frac{q}{(1 - q)^2} \right) \left(1 + \log \frac{1}{1 - q} \right).$$

Proof. The left hand side is equal to

$$\begin{aligned} \sum_{i \geq 1} i \log \left(\frac{1}{1 - q^i} \right) &= \sum_{i \geq 1} i \sum_{j \geq 1} \frac{q^{ij}}{j} \\ &= \sum_{i, j \geq 2} \frac{iq^{ij}}{j} + \sum_{i \geq 1} iq^i + \sum_{j \geq 1} \frac{q^j}{j} - 1 \\ &< \sum_{i, j \geq 2} \frac{iq^{i+j}}{j} + \sum_{i \geq 1} iq^i + \sum_{j \geq 1} \frac{q^j}{j} \\ &< \left(1 + \sum_{i \geq 1} iq^i \right) \left(1 + \sum_{j \geq 1} \frac{q^j}{j} \right) = \left(1 + \frac{q}{(1 - q)^2} \right) \left(1 + \log \frac{1}{1 - q} \right). \end{aligned}$$

\square

Let $\mathbf{x} = (x_1, x_2, \dots)$ and $\mathbf{y} = (y_1, y_2, \dots)$ be two infinite sequences of independent variables. Let $s_\lambda(\mathbf{x})$ denote the Schur function in the variables \mathbf{x} . The following formulas are well known, see [6, 11]:

$$(26) \quad \sum_{n \geq 0} \sum_{\lambda \vdash n} s_\lambda(\mathbf{x}) z^n = \prod_{i \geq 1} (1 - x_i z)^{-1} \prod_{1 \leq i < j} (1 - x_i x_j z^2)^{-1},$$

$$(27) \quad \sum_{n \geq 0} \sum_{\lambda \vdash n} s_\lambda(\mathbf{x}) s_\lambda(\mathbf{y}) z^n = \prod_{i, j \geq 1} (1 - x_i y_j z)^{-1}.$$

It is also known, see [11, Proposition 7.19.11], that

$$s_\lambda(1, q, q^2, \dots) = \frac{f^\lambda(q)}{(1-q)^n [n]_q!}.$$

Thus we get the following:

$$(28) \quad \sum_{\lambda \vdash n} s_\lambda(1, q, q^2, \dots) = \frac{t_n(q)}{(1-q)^n [n]_q!},$$

$$(29) \quad \sum_{\lambda \vdash n} s_\lambda(1, p, p^2, \dots) s_\lambda(1, q, q^2, \dots) = \frac{A_n(p, q)}{(1-p)^n (1-q)^n [n]_p! [n]_q!}.$$

Lemma 5.3. *For real numbers $0 < p, q < 1$, we have*

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{\lambda \vdash n} s_\lambda(1, q, q^2, \dots) &= \prod_{i \geq 1} (1 - q^i)^{-1} \prod_{0 \leq i < j} (1 - q^{i+j})^{-1}, \\ \lim_{n \rightarrow \infty} \sum_{\lambda \vdash n} s_\lambda(1, p, p^2, \dots) s_\lambda(1, q, q^2, \dots) &= \prod_{\substack{i, j \geq 0 \\ i+j > 0}} (1 - p^i q^j)^{-1}. \end{aligned}$$

Proof. Let $\xi_n(q) = \sum_{\lambda \vdash n} s_\lambda(1, q, q^2, \dots)$. By (26), we have

$$\sum_{n \geq 0} \xi_n(q) z^n = \prod_{i \geq 0} (1 - q^i z)^{-1} \prod_{0 \leq i < j} (1 - q^{i+j} z^2)^{-1},$$

equivalently,

$$\sum_{n \geq 0} (\xi_n(q) - \xi_{n-1}(q)) z^n = \prod_{i \geq 1} (1 - q^i z)^{-1} \prod_{0 \leq i < j} (1 - q^{i+j} z^2)^{-1},$$

where $\xi_{-1}(q) = 0$.

Then

$$\lim_{N \rightarrow \infty} \xi_N(q) = \lim_{N \rightarrow \infty} \sum_{n=0}^N (\xi_n(q) - \xi_{n-1}(q)) = \prod_{i \geq 1} (1 - q^i)^{-1} \prod_{0 \leq i < j} (1 - q^{i+j})^{-1}$$

converges, because

$$\prod_{i \geq 1} (1 - q^i)^{-1} < \prod_{0 \leq i < j} (1 - q^{i+j})^{-1} = \prod_{i \geq 1} (1 - q^i)^{-\lceil \frac{i}{2} \rceil} < \prod_{i \geq 1} (1 - q^i)^{-i},$$

where $\prod_{i \geq 1} (1 - q^i)^{-i}$ converges by Lemma 5.2. Thus we get the first limit. Similarly, we can prove the second limit. \square

Proof of Lemma 2.4. We will only prove the first limit. The second can be proved similarly. Using the well known asymptotic behavior of $t_n \sim \frac{1}{\sqrt{2}}n^{n/2} \exp(-\frac{n}{2} + \sqrt{n} - \frac{1}{4})$, we can easily see that it holds for $q = 1$.

Assume $q \neq 1$. By Lemma 5.1, it is sufficient to show that for $0 < q < 1$,

$$\lim_{n \rightarrow \infty} \frac{\frac{t_{n+1}(q)}{[n+1]_q!}}{\frac{t_n(q)}{[n]_q!}} = 1 - q.$$

Since

$$\frac{t_n(q)}{[n]_q!} = (1 - q)^n \sum_{\lambda \vdash n} s_\lambda(1, q, q^2, \dots),$$

we are done by Lemma 5.3. □

6. FURTHER STUDY

In this paper we have found a q -analog and a (p, q) -analog of the following:

$$(30) \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A)|}{|\mathcal{T}_n|} = \frac{f^\alpha}{a!},$$

$$(31) \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A, B)|}{|\mathcal{T}_n(\emptyset, \emptyset)|} = \frac{f^\alpha f^\beta}{a!b!}.$$

In a probabilistic point of view (30) means that the probability that a random SYT T of size n contains a given SYT A of shape $\alpha \vdash a$ tends to $\frac{f^\alpha}{a!}$ as n approaches infinity. This clearly implies that the probability that a random pair (P, Q) of SYTs of size n contains a given pair (A, B) of SYT of shape $\alpha \vdash a$ and $\beta \vdash b$ tends to $f^\alpha f^\beta / a!b!$ as n approaches infinity. However, (31) implies that adding the additional condition that we must have $\text{sh}(P) = \text{sh}(Q)$ does not change the probability, which is nontrivial. With this observation we conjecture the following.

Conjecture 6.1. For given SYTs A_1, \dots, A_k with $\text{sh}(A_i) = \alpha_i \vdash a_i$ for $i \in [k]$, we denote by $\mathcal{T}_n(A_1, \dots, A_k)$ the set of k -tuples (T_1, \dots, T_k) such that $\text{sh}(T_1) = \dots = \text{sh}(T_k)$ and $T_i \in \mathcal{T}_n(A_i)$ for all $i \in [k]$. Then

$$\lim_{n \rightarrow \infty} \frac{|\mathcal{T}_n(A_1, \dots, A_k)|}{|\mathcal{T}_n(E_1, \dots, E_k)|} = \frac{f^{\alpha_1} \dots f^{\alpha_k}}{a_1! \dots a_k!},$$

where $E_1 = \dots = E_k = \emptyset$.

Problem 6.1. Find a q -analog of Conjecture 6.1 if it is true.

In our q -analog and (p, q) -analog of (30) and (31) we have the assumption $p, q > 0$. We can extend the ranges of p and q by proving Lemma 2.4 for more general p and q . Thus we propose the following problem.

Problem 6.2. Find negative real numbers or complex numbers p and q satisfying the following:

$$\lim_{n \rightarrow \infty} \frac{\frac{t_{n+1}(q)}{[n+1]_q!}}{\frac{t_n(q)}{[n]_q!}} = 1 - \bar{q}, \quad \lim_{n \rightarrow \infty} \frac{\frac{A_{n+1}(p, q)}{[n+1]_p! [n+1]_q!}}{\frac{A_n(p, q)}{[n]_p! [n]_q!}} = (1 - \bar{p})(1 - \bar{q}).$$

ACKNOWLEDGEMENTS

I would like to thank Professor Richard Stanley for helpful discussion. I would also like to thank Professor Ron King for pointing out an error in Lemma 5.3 in an earlier version of this paper. I am grateful to the anonymous referees for their very careful reading and helpful comments which have improved the readability of this paper significantly.

REFERENCES

- [1] Miklós Bóna. *Combinatorics of permutations*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [2] Sylvie Corteel and Jeremy Lovejoy. Overpartitions. *Trans. Amer. Math. Soc.*, 356(4):1623–1635 (electronic), 2004.
- [3] A. M. Garsia and I. Gessel. Permutation statistics and partitions. *Adv. in Math.*, 31(3):288–305, 1979.
- [4] Aaron D. Jaggard. Subsequence containment by involutions. *Electron. J. Combin.*, 12:Research Paper 14, 15 pp. (electronic), 2005.
- [5] Dongsu Kim and Jang Soo Kim. The initial involution patterns of permutations. *Electron. J. Combin.*, 14(1):Research Paper 2, 15 pp. (electronic), 2007.
- [6] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [7] Brendan D. McKay, Jennifer Morse, and Herbert S. Wilf. The distributions of the entries of Young tableaux. *J. Combin. Theory Ser. A*, 97(1):117–128, 2002.
- [8] Bruce E. Sagan. *The symmetric group*, volume 203 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001.
- [9] Bruce E. Sagan and Richard P. Stanley. Robinson-Schensted algorithms for skew tableaux. *J. Combin. Theory Ser. A*, 55(2):161–193, 1990.
- [10] Richard P. Stanley. *Enumerative Combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997.
- [11] Richard P. Stanley. *Enumerative Combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999.

E-mail address: kimjs@math.umn.edu