

Problems on Minkowski sums of convex lattice polytopes

Tadao Oda

odatadao@math.tohoku.ac.jp

Abstract submitted at the Oberwolfach Conference

“Combinatorial Convexity and Algebraic Geometry” 26.10–01.11, 1997

Throughout, we fix the notation $M := \mathbf{Z}^r$ and $M_{\mathbf{R}} := \mathbf{R}^r$.

Given convex lattice polytopes $P, P' \subset M_{\mathbf{R}}$, we have


$$(M \cap P) + (M \cap P') \subset M \cap (P + P'),$$

where $P + P'$ is the Minkowski sum of P and P' , while the left hand side means $\{m + m' \mid m \in M \cap P, m' \in M \cap P'\}$.

Problem 1 For convex lattice polytopes $P, P' \subset M_{\mathbf{R}}$ when do we have the equality

$$(M \cap P) + (M \cap P') = M \cap (P + P')?$$

We always have the equality if $r = 1$. This need not be the case, however, if $r \geq 2$ as the following example shows:



In this example, each of P and P' is nice (known as basic or unimodular), but their relative position is not.

We may regard the case $P' = \nu P$ for a positive integer $\nu > 0$ as a special case of *nice relative position*. We have $P + \nu P = (\nu + 1)P$, and

$$(M \cap P) + (M \cap \nu P) \subset M \cap (\nu + 1)P.$$

Problem 2 When do we have the equality

$$(M \cap P) + (M \cap \nu P) = M \cap (\nu + 1)P$$

for all $\nu \in \mathbf{Z}_{>0}$?

This problem is related to the projective normality of projective toric varieties.

We obviously have the equality if $r = 1$. Koelman [9] showed that the equality always holds if $r = 2$.

More generally, Sturmfels [10] and others showed that the equality holds if P has a basic (also known as unimodular) triangulation.

In view of toric geometry, the following could be a reasonable formulation of the problem as to when P and P' are in nice relative position: We fix an r -dimensional convex lattice polytope P , and let P' to be obtained from P by *independent parallel translation* of facets (codimension one faces) of P . The combinatorial face structure of P' might differ from that of P .

By toric geometry, the convex lattice polytope P corresponds to an r -dimensional projective toric variety X over the complex number field \mathbf{C} together with an ample divisor D on X , while P' gives rise to an effective divisor D' on X . D' is ample if the combinatorial face structure of P' coincides with that of P . When D' is merely nef, the combinatorial face structure of P' could be slightly degenerate.

Problem 3 If D' is nef, do we have the surjectivity of the canonical multiplication map

$$H^0(X, \mathcal{O}_X(D)) \otimes_{\mathbf{C}} H^0(X, \mathcal{O}_X(D')) \longrightarrow H^0(X, \mathcal{O}_X(D + D'))?$$

We know that $H^1(X, \mathcal{O}_X(D')) = 0$ when D' is nef, hence

$$H^1(X \times X, \mathcal{O}_{X \times X}(p_1^{-1}D + p_2^{-1}D')) = 0$$

by Künneth formula. Consequently, Problem 3 is equivalent to the following:

Problem 4 Let \mathcal{I} be the $\mathcal{O}_{X \times X}$ -ideal corresponding to the diagonal subvariety $\Delta(X)$ of $X \times X$. If D' is nef, do we have

$$H^1(X \times X, \mathcal{I} \otimes_{\mathcal{O}_{X \times X}} \mathcal{O}_{X \times X}(p_1^{-1}D + p_2^{-1}D')) = 0?$$

Hopefully, we might have an affirmative answer at least when X is *smooth* and D' is *ample*. There have been unsuccessful attempts in this direction by means of Frobenius splittings in characteristic $p > 0$.

Without assuming X to be smooth nor D' to be ample, let us give another formulation for the problem.

Let $N := \text{Hom}_{\mathbf{Z}}(M, \mathbf{Z})$ with the canonical bilinear pairing $\langle \cdot, \cdot \rangle : M \times N \rightarrow \mathbf{Z}$. Consider the finite complete fan Σ for N corresponding to X . As usual, denote by

$$\Sigma(1) = \{\rho_1, \rho_2, \dots, \rho_l\}$$

the set of one-dimensional cones in Σ , and let $n_j \in N$ be the primitive generator for $\rho_j \in \Sigma(1)$. Let us introduce a free \mathbf{Z} -module $\tilde{N} := \bigoplus_{j=1}^l \mathbf{Z}\tilde{n}_j$ with the basis consisting of the symbols $\{\tilde{n}_1, \dots, \tilde{n}_l\}$ corresponding to $\Sigma(1)$, and the \mathbf{Z} -linear map

$$\pi : \tilde{N} \rightarrow N \quad \text{with } \pi(\tilde{n}_j) := n_j \text{ for } j = 1, \dots, l.$$

Let $\tilde{M} := \text{Hom}_{\mathbf{Z}}(\tilde{N}, \mathbf{Z})$ with the dual basis $\{\tilde{m}_1, \dots, \tilde{m}_l\}$. Since π has finite cokernel, the dual \mathbf{Z} -linear map

$$\pi^* : M \rightarrow \tilde{M} \quad \text{with } \pi^*(m) := \sum_{j=1}^l \langle m, n_j \rangle \tilde{m}_j \text{ for any } m \in M$$

is injective. Let $\mathcal{M} := \text{coker}(\pi^*)$ and denote by $\mu_j \in \mathcal{M}$ the image of $\tilde{m}_j \in \tilde{M}$. We call $(\mathcal{M}, \{\mu_1, \dots, \mu_l\})$ the linear Gale transform of $(N, \{n_1, \dots, n_l\})$.

\mathcal{M} is canonically isomorphic to the Weil divisor class group of X (modulo linear equivalence). Let

$$\tilde{M} \supset \tilde{M}_{\geq 0} := \sum_{j=1}^l \mathbf{Z}_{\geq 0} \tilde{m}_j \quad \text{and} \quad \mathcal{M} \supset \mathcal{M}_{\geq 0} := \sum_{j=1}^l \mathbf{Z}_{\geq 0} \mu_j.$$

$\tilde{M}_{\geq 0}$ is canonically isomorphic to the semigroup of torus-invariant effective Weil divisors on X . For $j = 1, 2, \dots, l$ we will use D_j and \tilde{m}_j interchangeably to denote the torus-invariant irreducible Weil divisor corresponding to the one-dimensional cone ρ_j .

The homogeneous coordinate ring introduced by Cox, Audin, Delzant, et al. (cf. [4]) is the semigroup algebra

$$S := \mathbf{C}[\tilde{M}_{\geq 0}] = \mathbf{C}[x_1, x_2, \dots, x_l] \quad \text{with } x_j := \mathbf{e}(\tilde{m}_j) \in S \text{ for } j = 1, \dots, l.$$

We endow the polynomial ring S with the $(\mathcal{M}_{\geq 0})$ -grading defined by

$$\deg x_j := \mu_j \quad \text{for } j = 1, \dots, l.$$

For $\alpha \in \mathcal{M}$, we denote by S_α the homogeneous part of degree α .

Note that the $(\mathcal{M}_{\geq 0})$ -graded ring S depends only on the 1-skeleton $\Sigma(1)$ of Σ . Problem 1 is more or less equivalent to the following:

Problem 5 Given $\alpha, \beta \in \mathcal{M}_{\geq 0}$, when is the multiplication map

$$S_\alpha \otimes_{\mathbf{C}} S_\beta \longrightarrow S_{\alpha+\beta}$$

surjective?

The fan Σ determines the polyhedral cone

$$C \subset \mathcal{M}_{\mathbf{R}} := \mathcal{M} \otimes_{\mathbf{Z}} \mathbf{R}$$

spanned by nef divisor classes. The intersection $\mathcal{M} \cap C^\circ$ of \mathcal{M} with the interior C° of C is the semigroup of ample divisor classes on X . Then Problems 3 and 4 are almost equivalent to the following:

Problem 6 Is the multiplication map $S_\alpha \otimes_{\mathbf{C}} S_\beta \rightarrow S_{\alpha+\beta}$ surjective if $\alpha \in \mathcal{M} \cap C^\circ$ and $\beta \in \mathcal{M} \cap C$? What if $\alpha, \beta \in \mathcal{M} \cap C^\circ$?

The study of the diagonal ideal sheaf $\mathcal{I} \subset \mathcal{O}_{X \times X}$ is important not only in connection with Problem 4 but in its own right. In explaining a possible approach to the study, let us follow the notation of Cox [4].

We denote

$$x^D := \prod_{j=1}^l x_j^{a_j} \in S \quad \text{for } D = \sum_{j=1}^l a_j \tilde{m}_j \in \tilde{M}_{\geq 0}$$

and

$$\deg x^D := \sum_{j=1}^l a_j \mu_j =: [D].$$

By our convention $D_j = \tilde{m}_j$, we have

$$x^{D_j} = x_j \quad \text{and} \quad [D_j] = \mu_j \quad \text{for } j = 1, 2, \dots, l.$$

For each $\alpha \in \mathcal{M}$ we denote by $\mathcal{O}_X(\alpha)$ the \mathcal{O}_X -module corresponding to the degree-shifted graded S -module $S(\alpha)$. We also need the following notation later: For each $m \in M$ we denote the zero and polar divisors of the character $\mathbf{e}(m)$ of the torus regarded as a rational function on X by

$$D^+(m) := \sum_{\substack{1 \leq j \leq l \\ \langle m, n_j \rangle > 0}} \langle m, n_j \rangle D_j \quad \text{and} \quad D^-(m) := \sum_{\substack{1 \leq j \leq l \\ \langle m, n_j \rangle < 0}} (-\langle m, n_j \rangle) D_j,$$

hence $\pi^*(m) = D^+(m) - D^-(m)$.

We have a canonical homomorphism of $(\mathcal{M}_{\geq 0} \times \mathcal{M}_{\geq 0})$ -graded \mathbf{C} -algebras

$$S \otimes_{\mathbf{C}} S \longrightarrow \mathbf{C}[\mathcal{M}_{\geq 0}] \otimes_{\mathbf{C}} S$$

defined by

$$x^D \otimes x^E \mapsto \mathbf{e}([D]) \otimes x^{D+E} \quad \text{for } D, E \in \tilde{M}_{\geq 0}.$$

The ideal $\mathcal{I} \subset \mathcal{O}_{X \times X}$ for the diagonal subvariety $\Delta(X) \subset X \times X$ obviously corresponds to the $(\mathcal{M}_{\geq 0} \times \mathcal{M}_{\geq 0})$ -homogeneous ideal

$$I := \ker(S \otimes_{\mathbf{C}} S \rightarrow \mathbf{C}[\mathcal{M}_{\geq 0}] \otimes_{\mathbf{C}} S).$$

Problems 5 and 6 ask the surjectivity of the (α, β) -component $S_{\alpha} \otimes_{\mathbf{C}} S_{\beta} \rightarrow \mathbf{e}(\alpha) \otimes_{\mathbf{C}} S_{\alpha+\beta}$ of this homomorphism under various conditions on $\alpha, \beta \in \mathcal{M}_{\geq 0}$.

I is a homogeneous binomial ideal in the $(\mathcal{M}_{\geq 0} \times \mathcal{M}_{\geq 0})$ -graded \mathbf{C} -algebra $S \otimes_{\mathbf{C}} S$ depending only on the 1-skeleton $\Sigma(1)$ of the fan Σ . We may try to find nice $(\mathcal{M}_{\geq 0} \times \mathcal{M}_{\geq 0})$ -graded $S \otimes_{\mathbf{C}} S$ -free resolutions of I to consider Problem 4.

Identifying the logarithmic derivatives dx_j/x_j with \tilde{m}_j for $j = 1, 2, \dots, l$ as usual, we get a canonical injective homomorphism of graded S -modules

$$\Omega_S^1 \longrightarrow S \otimes_{\mathbf{Z}} \tilde{M}, \quad dx_j \mapsto x_j \otimes \tilde{m}_j \quad \text{for } j = 1, 2, \dots, l.$$

Denote by

$$\Omega := \ker(\Omega_S^1 \rightarrow S \otimes_{\mathbf{Z}} \tilde{M} \rightarrow S \otimes_{\mathbf{Z}} \mathcal{M})$$

the kernel of the composite of this homomorphism with the canonical projection $S \otimes_{\mathbf{Z}} \tilde{M} \rightarrow S \otimes_{\mathbf{Z}} \mathcal{M}$. It is not hard to show that the sheaf Ω_X^1 of Zariski differential 1-forms (resp. $\bigoplus_{j=1}^l \mathcal{O}_X(-D_j)$) is the \mathcal{O}_X -module associated to the graded S -module Ω (resp. Ω_S^1). In this way, we get the following well-known result:

Proposition 7 (Generalized Euler exact sequence. cf. Batyrev-Cox [1])

We have an exact sequence of \mathcal{O}_X -modules

$$0 \rightarrow \Omega_X^1 \rightarrow \bigoplus_{j=1}^l \mathcal{O}_X(-D_j) \rightarrow \mathcal{O}_X \otimes_{\mathbf{Z}} \mathcal{M} \rightarrow 0.$$

Remark 8 The graded S -module Ω is generated over S by

$$\left\{ x^D dx^E - x^E dx^D \mid D, E \in \tilde{M}_{\geq 0}, D \sim E \right\},$$

hence by

$$\left\{ x^{D^+(m)} dx^{D^-(m)} - x^{D^-(m)} dx^{D^+(m)} \mid m \in M \right\}.$$

The vectors $n_1, n_2, \dots, n_l \in N$ give rise to an arrangement \mathcal{A} of hyperplanes $\{n_j\}^{\perp} \subset M_{\mathbf{R}}$. A chamber Γ for \mathcal{A} is one of the top-dimensional polyhedral cones appearing in the partition of $M_{\mathbf{R}}$ induced by the arrangement \mathcal{A} . If we choose for each chamber Γ a set Ξ_{Γ} of generators of the semigroup $M \cap \Gamma$, then Ω is generated over S by

$$\left\{ x^{D^+(m)} dx^{D^-(m)} - x^{D^-(m)} dx^{D^+(m)} \mid m \in \bigcup_{\text{chambers } \Gamma} \Xi_{\Gamma} \right\}.$$

Remark 9 As in Remark 8, we see that the “diagonal” ideal $I \subset S \otimes_{\mathbf{C}} S$ is generated over $S \otimes_{\mathbf{C}} S$ by

$$\left\{ x^D \otimes x^E - x^E \otimes x^D \mid D, E \in \tilde{M}_{\geq 0}, D \sim E \right\},$$

hence by

$$\left\{ x^{D^+(m)} \otimes x^{D^-(m)} - x^{D^-(m)} \otimes x^{D^+(m)} \mid m \in \bigcup_{\text{chambers } \Gamma} \Xi_{\Gamma} \right\}.$$

Example 10 When $X = \mathbf{P}^r$ is the projective space, the corresponding polytope P is a unimodular simplex, and $l = r + 1$. We have $\mathcal{M} = \mathbf{Z}$, and $S_{\alpha} \otimes S_{\beta} \rightarrow S_{\alpha+\beta}$ in Problem 5 is obviously surjective for all $\alpha, \beta \in \mathcal{M}_{\geq 0}$. Nevertheless, the description of the diagonal ideal $\mathcal{I} \subset \mathcal{O}_{X \times X}$ is nontrivial. By Beilinson [2] we have an exact sequence

$$0 \rightarrow \mathcal{O}_X(-r) \boxtimes \Omega_X^r(r) \rightarrow \cdots \rightarrow \mathcal{O}_X(-j) \boxtimes \Omega_X^j(j) \rightarrow \cdots \rightarrow \mathcal{O}_X(-1) \boxtimes \Omega_X^1(1) \rightarrow \mathcal{I} \rightarrow 0,$$

where $\mathcal{F} \boxtimes \mathcal{F}' := (p_1^* \mathcal{F}) \otimes_{\mathcal{O}_{X \times X}} (p_2^* \mathcal{F}')$ is the external tensor product on $X \times X$ of \mathcal{O}_X -modules \mathcal{F} and \mathcal{F}' . (Thanks are due to Miles Reid for pointing out this result to the author.) One way of proving this is to note that the $S \otimes_{\mathbf{C}} S$ -module homomorphism

$$S \otimes_{\mathbf{C}} \Omega_S^1 \longrightarrow S \otimes_{\mathbf{C}} S, \quad 1 \otimes dx_j \mapsto x_j \otimes 1 \quad \text{for } j = 1, 2, \dots, r + 1$$

induces a surjection $S \otimes_{\mathbf{C}} \Omega \rightarrow I$. Thus the Koszul complex arising out of the corresponding $\mathcal{O}_{X \times X}$ -homomorphism (cf., e.g., Eisenbud [6])

$$\mathcal{O}_X(-1) \boxtimes \Omega_X^1(1) \longrightarrow \mathcal{O}_{X \otimes X},$$

whose cokernel is $\mathcal{O}_{\Delta(X)}$, gives the exact sequence above.

Similarly, the Koszul complex arising out of the S -homomorphism $\Omega_S^1 \rightarrow S$ which sends dx_j to x_j for $j = 1, 2, \dots, r + 1$ is nothing but the complex (Ω_S^1, d) . By Remark 8, we see that Ω is the image of $d : \Omega_S^2 \rightarrow \Omega_S^1$. Consequently, we get an exact sequence of \mathcal{O}_X -modules

$$0 \rightarrow \mathcal{O}_X(-r - 1)^{\oplus_{r+1} C_{r+1}} \rightarrow \cdots \rightarrow \mathcal{O}_X(-j)^{\oplus_{r+1} C_j} \rightarrow \cdots \rightarrow \mathcal{O}_X(-2)^{\oplus_{r+1} C_2} \rightarrow \Omega_X^1 \rightarrow 0,$$

where

$${}_{r+1}C_j := \binom{r+1}{j}$$

are the binomial coefficients.

Alternatively, we could use the Eagon-Northcott complex (cf. Eagon-Northcott [5], Kirby [8] and Eisenbud [6]) for the 2×2 -minors of

$$\begin{pmatrix} x_1 \otimes 1 & x_2 \otimes 1 & \cdots & x_{r+1} \otimes 1 \\ 1 \otimes x_1 & 1 \otimes x_2 & \cdots & 1 \otimes x_{r+1} \end{pmatrix}$$

to get an exact sequence

$$0 \rightarrow \mathcal{E}_r \rightarrow \cdots \rightarrow \mathcal{E}_p \rightarrow \cdots \rightarrow \mathcal{E}_1 \rightarrow \mathcal{I} \rightarrow 0,$$

where

$$\mathcal{E}_p := \bigoplus_{\substack{j+k=p+1 \\ j \geq 1, k \geq 1}} \bigwedge^{p+1} \tilde{M} \otimes_{\mathbf{Z}} \mathcal{O}_{X \times X}(-j, -k)$$

with

$$\mathcal{O}_{X \times X}(\alpha, \beta) := \mathcal{O}_X(\alpha) \boxtimes \mathcal{O}_X(\beta) \text{ for } \alpha, \beta \in \mathcal{M} = \mathbf{Z}.$$

References

- [1] V. Batyrev and D. Cox, On the Hodge structure of projective hypersurfaces in toric varieties, *Duke Math. J.* 75 (1994), 293–338.
- [2] A. A. Beilinson, Coherent sheaves on \mathbf{P}^n and problems of linear algebra, *Func. Anal. App.* 12, No. 3 (1978), 214–216.
- [3] I. N. Bernshtein, I. M. Gelfand and S. I. Gelfand, Algebraic bundles over \mathbf{P}^n and problems of linear algebra, *Func. Anal. App.* 12, No. 3 (1978), 212–214.
- [4] D. Cox, The homogeneous coordinate ring of a toric variety, *J. Algebraic Geometry* 4 (1995), 17–50.
- [5] J. A. Eagon and D. G. Northcott, Ideals defined by matrices and a certain complex associated with them, *Proc. Royal Soc. London* 269 (1962), 188–204.
- [6] D. Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Graduate Texts in Math. 150, Springer-Verlag, New York, Berlin, Heidelberg, London, Paris, Tokyo, Hong Kong, Barcelona, Budapest, 1995
- [7] A. G. Khovanskii, Sums of finite sets, orbits of commutative semigroups and Hilbert functions, *Func. Anal. App.* 29, No. 2 (1995), 102–112.
- [8] D. Kirby, A sequence of complexes associated with a matrix, *J. London Math. Soc.* (2) 7 (1973), 523–530.
- [9] R. Koelman, Generators for the ideal of a projectively embedded toric surface, *Tohoku Math. J.* 45 (1993), 385–392.
- [10] B. Sturmfels, Equations defining toric varieties, alg-geom/9610018, AMS Summer Inst. in Algebraic Geometry, Santa Cruz, 1995.