

BRIEF COMMUNICATION. AN INDEFINITE STURM THEORY.

ALESSANDRO PORTALURI

INTRODUCTION

Sturm theory for second order differential equations was generalized to systems and higher order equations with positive leading coefficient by several authors. (See for instance [1], [2] and [6] among the others). Here we propose a Sturm oscillation theorem for indefinite systems with Dirichlet boundary conditions of the form

$$(0.1) \quad l(x, D)v := p_{2m} \frac{d^{2m}v}{dx^{2m}} + p_{2m-2}(x) \frac{d^{2m-2}v}{dx^{2m-2}} + \dots + p_1(x) \frac{dv}{dx} + p_0(x)v = 0,$$

where p_i is a smooth path of matrices on the complex n -dimensional vector space \mathbb{C}^n , p_{2m} is the symmetry of the form $\text{diag}(I_{n-\nu}, -I_\nu)$ for some $\nu \geq 0$. This generalization was obtained along the lines of [4] and [5] in which the second order equations were considered. Full proofs and the relation with Maslov index will appear elsewhere.

1. VARIATIONAL SET UP AND MAIN RESULTS

We will use the variational approach to (0.1) as described in [2] and we will stick to the notations of that paper. Given the complex n -dimensional Hermitian space $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$, for any $m \in \mathbb{N}$ let $\mathcal{H}^m := H^m(J, \mathbb{C}^n)$ be the Sobolev space of all H^m -maps from $J := [0, 1]$ into \mathbb{C}^n .

A *derivative dependent Hermitian form* is the form $\Omega(x)[u] = \sum_{i,j=0}^m \langle D^i u(x), \omega_{i,j}(x) D^j u(x) \rangle$, where, each $\omega_{i,j}$ is a smooth path of x -dependent Hermitian matrices with constant leading coefficient $\omega_{m,m} := p_{2m}$ and such that $\omega_{i,2m-1-i} = 0$ for each $i = 0, \dots, m$. Each derivative dependent Hermitian form, defines a Hermitian form $q: \mathcal{H}^m \rightarrow \mathbb{R}$ by setting $q(u) := \int_J \Omega(x)[u] dx$. If $v \in \mathcal{H}^m$ and $u \in \mathcal{H}^{2m}$ then, using integration by parts, the corresponding sesquilinear form $q(u, v)$ can be written as

$$(1.1) \quad q(u, v) = \int_J \langle v(x), l(x, D)u(x) \rangle dx + \phi(u, v),$$

where $l(x, D)$ is a differential operator of the form of (0.1) and $\phi(u, v)$ is a bilinear form depending only on the $(m-1)$ -jet, $j^{m-1}v(x) := (v(x), \dots, v^{(m-1)}(x))$ and on the $(2m-1)$ -jet $j^{2m}u(x)$ at the boundary $x = 0, 1$.

Let $\mathcal{H}_0^m := \mathcal{H}_0^m(J) := \{u \in \mathcal{H}^m : j^{m-1}u(0) = 0 = j^{m-1}u(1)\}$ and let q_Ω be the restriction of the Hermitian form q to \mathcal{H}_0^m . For each $\lambda \in J$, let us consider the space $\mathcal{H}_0^m([0, \lambda])$ with the form $\int_{[0, \lambda]} \Omega(x) dx$. Via the substitution $x \mapsto \lambda x$, we transfer this form to $\mathcal{H}_0^m(J)$, so, we come to the forms Ω_λ and q_λ defined respectively by

$$(1.2) \quad \Omega_\lambda(x)[u] := \sum_{i,j=0}^m \langle D^i u(x), \lambda^{2m-(i+j)} \omega_{i,j}(\lambda x) D^j u(x) \rangle \quad \text{and we let} \quad q_\lambda(u) := \int_J \Omega_\lambda(x)[u] dx.$$

Then $\lambda \mapsto q_\lambda$ is a smooth path of Hermitian forms acting on \mathcal{H}_0^m with $q_1 = q_\Omega$ and with $q_0(u) = \int_J \langle p_{2m} D^m u, D^m u \rangle dx$. Now we introduce the following definition.

DEFINITION 1.1. A conjugate instant for q_Ω is any point $\lambda \in (0, 1]$ such that $\ker q_\lambda \neq \{0\}$.

Let C_λ be the path of bounded selfadjoint Fredholm operators associated to q_λ via the Riesz representation theorem.

LEMMA 1.2. The Hermitian form q_0 is non-degenerate. Moreover each q_λ is a Fredholm Hermitian form. (i.e. C_λ is a Fredholm operator). In particular $\dim \ker q_\lambda < +\infty$ and q_λ is non degenerate if and only if $\ker q_\lambda = \{0\}$.

Date: December 10, 2008. Revised version.

2000 Mathematics Subject Classification. 34B24.

The author was partially supported by MIUR project *Variational Methods and Nonlinear Differential Equations*.

Proof. That the operator C_0 is an isomorphism can be proven exactly as in [4, Proposition 3.1]. On the other hand each q_λ is a weakly continuous perturbation of q_0 since it differs from q_0 only by derivatives of u of order less than m . Therefore $C_\lambda - C_0$ is compact for all $\lambda \in J$ and hence C_λ is Fredholm of index 0. The last assertion follows from this. \square

When the form q_Ω is non degenerate, i.e. when 1 is not a conjugate instant, we introduce the following definition.

DEFINITION 1.3. *The (regularized) Morse index of q_Ω is defined by*

$$(1.3) \quad \mu_{\text{Mor}}(q_\Omega) := -\text{sf}(q_\lambda, J),$$

where sf denotes the spectral flow of the path q_λ i.e., the number of positive eigenvalues of C_λ at $\lambda = 0$ which become negative at $\lambda = 1$ minus the number of negative eigenvalues of C_λ which become positive. (See, for instance [4], for a more detailed exposition).

Now we define the conjugate index. By (1.1) $q(u, v) = \int_J \langle v(x), l(x, D)u(x) \rangle dx$ for $u \in \mathcal{H}_0^{2m}$ and $v \in \mathcal{H}_0^m$ and therefore elements of the kernel of q_λ are weak solutions of the Dirichlet boundary value problem

$$\begin{cases} l(x, D)u = 0 \\ j^{m-1}u(0) = 0 = j^{m-1}u(1). \end{cases}$$

Let $\mathcal{O} = \{z := \lambda + is \in \mathbb{C} : 0 < \lambda < 1, -1 < s < 1\}$. For each $z \in \overline{\mathcal{O}}$ let us consider the closed unbounded Fredholm operator A_z on $L^2(J, \mathbb{C}^{mn})$ with domain $\mathcal{D}(A_z) = \{u \in \mathcal{H}^{2m} : j^{m-1}u(0) = 0 = j^{m-1}u(1)\}$ defined by $A_z u := l_\lambda(x, D)u + isu$. Since $A_{\bar{z}} = \bar{A}_z$, it follows that A_z has a continuous inverse for $s = \Im(z) \neq 0$. Let us consider the splitting $\mathbb{C}^{2mn} := (\mathbb{C}^n)^m \times (\mathbb{C}^n)^m$. For each $w = (w_1, \dots, w_m) \in (\mathbb{C}^n)^m$, let $u_z(w, x)$ be the unique solution of the Cauchy problem

$$\begin{cases} l(x, D)u + isu = 0 \\ u(0) = u'(0) = \dots = u^{m-1}(0) = 0, \quad u^m(0) = w_1, \dots, u^{2m-1}(0) = w_m, \end{cases}$$

and let $\mathcal{R}_z : \mathbb{C}^{nm} \rightarrow \mathbb{C}^{nm}$ be the endomorphism defined by $\mathcal{R}_z(w) := j^{m-1}(u_z(w, 1))$. Clearly $u \in \ker A_z$ if and only if $(u^m(0), \dots, u^{2m-1}(0)) \in \ker \mathcal{R}_z$. From this and by using regularity of the weak solutions, we get that the following three statements are equivalent:

(i) $\ker A_z \neq \{0\}$; (ii) $\Im(z) = 0$ and $\lambda = \Re(z)$ is a conjugate instant; (iii) $\det \mathcal{R}_z = 0$. Because of these three equivalent statements, the function $\rho(z) := \det \mathcal{R}_z$ does not vanishes on the boundary $\partial \mathcal{O}$.

DEFINITION 1.4. *The conjugate index $\mu_{\text{con}}(q_\Omega)$, of q_Ω is minus the winding number of $\rho|_{\partial \Omega} \rightarrow \mathbb{C} - \{0\}$ or equivalently minus the Brouwer degree, i.e. $-\deg(\rho, \mathcal{O}, 0)$.*

With this said our main results are

THEOREM 1. (Generalized Sturm Oscillation theorem). *With the notation above, we have*

$$\mu_{\text{con}}(q_\Omega) = \mu_{\text{Mor}}(q_\Omega).$$

THEOREM 2. (Generalized Sturm comparison theorem). *If Ω_0, Ω_1 are two derivative dependent Hermitian forms with $\Omega_0[u](x) \leq \Omega_1[u](x)$ for all $x \in J$, then we have*

$$\mu_{\text{Mor}}(q_{\Omega_0}) \leq \mu_{\text{Mor}}(q_{\Omega_1}).$$

2. PROOFS

Proof. (of Theorem 1). We split the proof of this result into some steps.

First step. Notice that the family A_z has the form $A + B_z$ where A is a fixed unbounded closed operator and B_z is an A -bounded perturbation of A depending on z and that the operator valued one-form $dA_z A_z^{-1}$ is well-defined on $\partial \mathcal{O}$. Exactly as in Proposition 4.4 in [4] one proves the following Lemma.

LEMMA 2.1. *The form $dA_z A_z^{-1}$ takes values in operators of the trace class and*

$$(2.1) \quad \mu_{\text{con}}(q_\Omega) = -\frac{1}{2\pi i} \int_{\partial \mathcal{O}} \text{Tr} dA_z A_z^{-1}.$$

Let us assume now that the path $\{A_\lambda\}_{\lambda \in J}$ has only regular crossing points with the variety of singular operators. This means that the quadratic form $\Gamma(A, \lambda)$ given by the restriction to the kernel of A_λ of $\langle A'_\lambda, \cdot \rangle$ is nondegenerate. Since regular crossing points are isolated, there are only finite number of them. In what follows we will show that if \mathcal{D}_j is a small enough neighborhood of a crossing point $\lambda_j \in \mathcal{O}$, then

$$(2.2) \quad \frac{1}{2\pi i} \int_{\partial \mathcal{D}_j} \text{Tr} dA_z A_z^{-1} = \text{sign} \Gamma(A, \lambda_j).$$

For a fixed j , choose a positive number $\mu > 0$ such that the only point in the spectrum of A_{λ_j} in the interval $[-\mu, \mu]$ is 0 and then choose η small enough such that neither μ nor $-\mu$ lies in the spectrum of A_λ for $|\lambda - \lambda_j| < \eta$. For such a λ , let P_λ be the orthogonal projection in H onto the spectral subspace associated

to the part of the spectrum of A_λ lying in the interval $[-\mu, \mu]$. Then $A_\lambda P_\lambda = P_\lambda A_\lambda$ on the domain of A_λ . By [3, Chapter II, Section 6] there exist a smooth path U of unitary operators of H defined in $[\lambda_j - \eta, \lambda_j + \eta]$, such that $U_{\lambda_j} = \text{Id}_H$ and such that $P_\lambda U_\lambda = U_\lambda P_{\lambda_j}$. Let us consider the smooth operator valued function $N_z = U_\lambda^{-1} A_z U_\lambda$ defined on some open neighborhood of $\mathcal{D}_j = [\lambda_j - \eta, \lambda_j + \eta] \times [-1, 1]$ together with the differential one-form $\theta = dN_z N_z^{-1}$. Clearly θ is in the trace class and $\text{Tr } dA_z A_z^{-1} = \text{Tr } dN_z N_z^{-1}$.

Second step. If $H_j = \text{Im } P_{\lambda_j} = \ker A_{\lambda_j}$, under the splitting $H = H_j \oplus H_j^\perp$, the one-form θ splits into $\theta_0 = dN_z N_z^{-1}|_{H_j}$ and $\theta_1 = dN_z N_z^{-1}|_{H_j^\perp}$. Thus by taking traces we have

$$(2.3) \quad \frac{1}{2\pi i} \int_{\partial \mathcal{D}_j} \text{Tr } \theta = \frac{1}{2\pi i} \int_{\partial \mathcal{D}_j} \text{Tr } \theta_0 + \frac{1}{2\pi i} \int_{\partial \mathcal{D}_j} \text{Tr } \theta_1$$

where the last term in (2.3) vanishes.

In fact, $N_z|_{H_j^\perp}$ is invertible on $\overline{\mathcal{O}}$ and hence the one form θ_1 is exact. Now, let us consider the path of symmetric endomorphisms $M: [\lambda_j - \eta, \lambda_j + \eta] \rightarrow \mathcal{L}(H_j)$ given by $M_\lambda = N_\lambda|_{H_j}$. By elementary calculations it follows that $\text{sign } \Gamma(M_\lambda, \lambda_j) = \text{sign } \Gamma(A, \lambda_j)$.

Third step. If M is the path defined above and if $M_z = M_\lambda + is\text{Id}$, for $z \in \mathcal{D}_j$, then

$$\text{sign } \Gamma(M, \lambda_j) = \frac{1}{2\pi i} \int_{\partial \mathcal{D}_j} \text{Tr } dM_z M_z^{-1}.$$

Using [3, Chapter II, Theorem 6.8], one reduces M_λ to a diagonal form and the result follows by direct integration. This together with (2.3) proves (2.2).

Fourth step. Using regularity it is easy to see that any regular crossing point for the path A is also a regular crossing point for the path q i.e., the crossing form defined as the restriction $\Gamma(q, \lambda)$ of \dot{q}_λ to $\ker q_\lambda$ is non-degenerate. Moreover the crossing forms $\Gamma(A, \lambda)$ and $\Gamma(q, \lambda)$ coincide. Since for path with only regular crossings the spectral flow is given by the sum of the signatures of the crossing forms it follows from Lemma 2.1 and formula (2.2) that theorem 1 holds for paths with only regular crossings. In order to conclude remains to show that it is possible to extend the above calculation to general paths having only regular crossings. To do so we will apply a perturbation argument of Robbin and Salamon to the path of operators obtained by the restriction of the family A_z to the real line. By Theorem 4.22 in [7], we can find a $\delta > 0$ such that $A_\lambda^\delta := A_\lambda + \delta \text{Id}$ is a path of self-adjoint Fredholm operators with only regular crossing points. Let \mathcal{R}_z^δ be the matrix associated to the perturbed family A_z^δ and let $\rho^\delta(z) := \det \mathcal{R}_z^\delta$. Taking closed disjoint neighborhoods \mathcal{D}_j of $(\lambda_j, 0)$ in \mathcal{O} and summing over all crossing points, we obtain

$$\mu_{\text{con}}(q_\Omega) = - \sum_{j=1}^k \text{deg}(\rho^\delta, \mathcal{D}_j, 0) = \sum_{j=1}^k - \frac{1}{2\pi i} \int_{\partial \mathcal{D}_j} \text{Tr } dA_z^\delta (A_z^\delta)^{-1} = - \sum_{j=1}^k \text{sign } \Gamma(A, \lambda_j) = -\text{sf}(q_\lambda, J)$$

by the homotopy invariance of the spectral flow. Now Theorem 1 is proved. \square

Proof. of Theorem 2. Since the top order terms coincide, the difference between the two Fredholm Hermitian forms q^1 and q^2 is weakly semi-continuous. Thus in order to conclude it is enough to show that if q^1 and q^2 are two admissible families of Fredholm Hermitian forms on \mathcal{H}_0^m whose difference is weakly semi-continuous and such that $q_0^1 = q_0^2$, $q_1^2 \leq q_1^1$, then $\text{sf}(q^2) \leq \text{sf}(q^1)$. The rest of the proof is devoted to show this.

It is easy to see that there exists a small $\eta > 0$ such that $\tilde{q}_\lambda^1 = q_\lambda^1 + \lambda\eta \|\cdot\|^2$ is a family of Fredholm Hermitian forms that is homotopic to q^1 by $(t, \lambda) \rightarrow q_\lambda^1 + t\lambda\eta \|\cdot\|^2$ and therefore $\text{sf}(\tilde{q}^1) = \text{sf}(q^1)$. But now $\tilde{q}_0^1 = q_0^2$ and $\tilde{q}_1^1 < \tilde{q}_1^2$. Given the family of Fredholm Hermitian forms defined on T by $\phi(s, \lambda) = s\tilde{q}_\lambda^1 + (1-s)q_\lambda^2$ and by using the free homotopy property of the spectral flow for closed paths, it follows that the restriction of ϕ to ∂T has spectral flow zero. Furthermore, $\text{sf}(\tilde{q}^1) - \text{sf}(q^2) = \text{sf}(\rho)$, where $\rho(s) = (1-s)\tilde{q}_1^2 + s\tilde{q}_1^1$. Since $\dot{\rho} = \tilde{q}_1^1 - \tilde{q}_1^2$ is positive definite at each crossing point, all crossing points are regular, and each gives a positive contribute $\dim \ker \rho(s)$ to the spectral flow of ρ . Thus $\text{sf}(\rho) \geq 0$ and hence $\text{sf}(q^2) \leq \text{sf}(q^1)$. \square

Acknowledgements. I would like to thank Jacobo Pejsachowicz for many useful conversations. I am also very grateful to the anonymous referee for his numerous suggestions, criticisms and for having pointed out some crucial mistakes in the original version.

REFERENCES

- [1] V.I. Arnol'd, *Sturm theorems in Symplectic geometry*, Funk. Anal. i Prilozhen. **19**, n.4, 1–10 (1985).
- [2] H. Edwards, *A generalized Sturm Theorem*, Ann. of Math. **80** (1964), 2–57.
- [3] T. Kato, *Perturbation Theory for Linear Operators*, Grundlehren der Mathematischen Wissenschaften, vol. 132, Springer-Verlag, New York/Berlin, 1980.
- [4] M. Musso, J. Pejsachowicz, A. Portaluri, *A Morse Index Theorem for Perturbed Geodesics on semi-Riemannian Manifolds*, Topological Methods in Nonlinear Analysis, **25** (2005), no. 1, 69–99.
- [5] M. Musso, J. Pejsachowicz, A. Portaluri, *Bifurcation for Perturbed Geodesics on semi-Riemannian Manifolds*. Esaim Control Optim. Calc. Var. **13** (2007), no. 3, 598–621.
- [6] V. Y. Ovsienko, *Selfadjoint differential operators and curves on a Lagrangian Grassmannian that are subordinate to a loop*. Math. Zametki **47** (1990), no. 3, 65–73; translation in Math. Notes **47** (1990), no. 3–4, 270–275.
- [7] J. Robbin, D. Salamon, *The spectral flow and the Maslov index*, Bull. London Math. Soc. **27** (1995), 1–33.

DIPARTIMENTO DI MATEMATICA "ENNIO DE GIORGI",
COLLEGIO FIORINI, UNIVERSITÀ DEL SALENTO,
VIA PER ARNESANO, LECCE, LE
ITALY
E-mail address: `alessandro.portaluri@unile.it`