#### AVERAGE-CASE PERTURBATIONS AND SMOOTH CONDITION NUMBERS

DIEGO ARMENTANO

Centro de Matemática Facultad de Ciencias Universidad de la República Calle Igua 4225 11400 Montevideo, Uruguay diego@cmat.edu.uy

> ABSTRACT. In this paper we define a new condition number adapted to directionally uniform perturbations. The definitions and theorems can be applied to a large class of problems. We show the relation with the classical condition number, and study some interesting examples.

*Keywords:* Condition Numbers, Random Matrices, Systems of Linear Equations, Finding Kernels, Eigenvalue and Eigenvector Problems, Solving Polynomial Systems of Equations.

AMS subject classifications: 65F35; 65Y05

## 1. INTRODUCTION

Let X and Y be two real (or complex) Riemannian manifolds of real dimensions m and n $(m \ge n)$  respectively associated to some computational problem, where X is the space of *inputs* and Y is the space of *outputs*. Let  $V \subset X \times Y$  be the *solution variety*, i.e. the subset of pairs (x, y) such that y is an output corresponding to the input x. Let  $\pi_1 : V \to X$  and  $\pi_2 : V \to Y$  be the canonical projections, and  $\Sigma \subset V$  be the set of critical points of the projection  $\pi_1$ .

In case dim  $V = \dim X$ , for each  $(x, y) \in V \setminus \Sigma$  there is a differentiable function locally defined from some neighborhoods  $U_x$  and  $U_y$  of  $x \in X$  and  $y \in Y$  respectively, namely

$$G := \pi_2 \circ \pi_1^{-1}|_{U_x} : U_x \to U_y$$

Let us denote by  $\langle \cdot, \cdot \rangle_x$  and  $\langle \cdot, \cdot \rangle_y$  the Riemannian (or Hermitian) inner product in the tangent spaces  $T_x X$  and  $T_y Y$  at x and y respectively. The derivative  $DG(x) : T_x X \to T_y Y$  is called the *condition matrix* at (x, y). The *classical condition number* at  $(x, y) \in V \setminus \Sigma$  is defined as

(1) 
$$\kappa(x,y) := \max_{\substack{\dot{x} \in T_x X \\ \|\dot{x}\|^2_x = 1}} \|DG(x)\dot{x}\|_y$$

This number is an upper-bound -to first-order approximation- of the *worst-case* sensitivity of the output error with respect to small perturbations of the input. It plays an important role to understand the behavior of algorithms and, as a consequence, appears in the usual bounds of the running time of execution. There exist an extensive literature about the role of the condition

#### DIEGO ARMENTANO

number in numerical analysis and complexity of algorithms, see for example [13] and references therein.

In many practical situations, there exist a discrepancy between theoretical analysis and observed performance of an algorithm. There exist several approaches that attempt to rectify this discrepancy. Among them we find *average-case analysis* (see [7, 10]) and *smooth analysis* (see [11, 4, 15]). For a comprehensive review on this subject with historical notes see [5].

In this paper, averaging is performed in a different form. In many problems, the space of inputs has a much larger dimension than the one of the space of outputs  $(m \gg n)$ . Then, it is natural to assume that infinitesimal perturbations of the input will produce drastic changes in the output, only when they are performed in a few directions. Then, a possibly different approach to analyze complexity of algorithms is to replace "worst direction" by a certain mean over all possible directions. This alternative was already suggested and studied in [14] in the case of linear system solving Ax = b, and more generally, in [12] in the case of matrix perturbation theory where the first-order perturbation expansion is assumed to be random.

In this paper we extend this approach to a large class of computational problems, restricting ourselves to the case of directionally uniform perturbations.

Generalizing the concept introduced in [14] and [12], we define the *pth-average condition number* at (x, y) as

(2) 
$$\kappa_{av}^{[p]}(x,y) := \left[\frac{1}{\operatorname{vol}(S_x^{m-1})} \int_{\dot{x} \in S_x^{m-1}} \|DG(x)\dot{x}\|_y^p \, dS_x^{m-1}(\dot{x})\right]^{1/p} \quad (p = 1, 2, \ldots)$$

where  $\operatorname{vol}(S_x^{m-1}) = \frac{2\pi^{m/2}}{\Gamma(m/2)}$  is the measure of the unit sphere  $S_x^{m-1}$  in  $T_xX$ , and  $dS_x^{m-1}$  is the induced volume element. We will be mostly interested in the case p = 2, which we simply write  $\kappa_{av}$  and call it average condition number.

Before the statement of the main theorem, we define the Frobenius condition number as

$$\kappa_F(x,y) := \|DG(x)\|_F = \sqrt{\sigma_1^2 + \dots + \sigma_n^2}$$

where  $\|\cdot\|_F$  is the Frobenius norm and  $\sigma_1, \ldots, \sigma_n$  are the singular values of the condition matrix.

# Theorem 1.

$$\kappa_{av}^{[p]}(x,y) = \frac{1}{\sqrt{2}} \left[ \frac{\Gamma\left(\frac{m}{2}\right)}{\Gamma\left(\frac{m+p}{2}\right)} \right]^{1/p} \mathbb{E}(\|\eta_{\sigma_1,\dots,\sigma_n}\|^p)^{1/p}$$

where  $\|\cdot\|$  is the Euclidean norm in  $\mathbb{R}^n$  and  $\eta_{\sigma_1,\ldots,\sigma_n}$  is a centered Gaussian vector in  $\mathbb{R}^n$  with diagonal covariance matrix  $Diag(\sigma_1^2,\ldots,\sigma_n^2)$ . In particular,

(3) 
$$\kappa_{av}(x,y) = \frac{\kappa_F(x,y)}{\sqrt{m}}.$$

This result is most interesting when  $n \ll m$ , for in that case

$$\kappa_{av}(x,y) \le \sqrt{\frac{n}{m}} \cdot \kappa(x,y) \ll \kappa(x,y),$$

as  $\kappa_F(x, y) \leq \sqrt{n} \cdot \kappa(x, y)$ . Thus, in these cases one may expect much better stability properties than those predicted by classical condition numbers.

In numerical analysis, many authors are interested in relative errors. Thus, in the case  $(X, \|\cdot\|_X)$ and  $(Y, \|\cdot\|_Y)$  are normed vector spaces, instead of consider the (absolute) condition number (1), one could take the *relative condition number* defined as

$$\kappa_{rel}(x) := \frac{\|x\|_X}{\|y\|_Y} \kappa(x, y),$$

and the relative Frobenius condition number as

$$\kappa_{relF}(x) := \frac{\|x\|_X}{\|y\|_Y} \kappa_F(x, y),$$

(for simplicity we drop the y in the argument).

In the same way, we define the relative pth-average condition number as

(4) 
$$\kappa_{rel}{}^{[p]}_{av}(x) := \frac{\|x\|_X}{\|y\|_Y} \kappa_{av}{}^{[p]}(x,y), \quad (p = 1, 2, \ldots)$$

For the case p = 2 we simply write  $\kappa_{relav}$  and call it relative average condition number.

Theorem 1 remains true if one change the (absolute) condition number by the relative condition number. In particular,

$$\kappa_{relav}(x) := \frac{\kappa_{relF}(x,y)}{\sqrt{m}}.$$

# 2. Componentwise Analysis

In the case  $Y = \mathbb{R}^n$  we define the *kth-componentwise condition number* at  $(x, y) \in V$  as:

(5) 
$$\kappa(x, y, k) := \max_{\substack{\dot{x} \in T_x X \\ \|\dot{x}\|^2_x = 1}} |(DG(x)\dot{x})_k| \qquad (k = 1, \dots, n),$$

where  $|\cdot|$  is the absolute value and  $w_k$  indicates the kth-component of the vector  $w \in \mathbb{R}^n$ .

Following [14] for the linear case, we define the average componentwise condition number as

(6) 
$$\kappa_{av}^{[p]}(x,y,k) := \left[\frac{1}{\operatorname{vol}(S_x^{m-1})} \int_{\dot{x}\in S_x^{m-1}} |(DG(x)\dot{x})_k|^p \, dS_x^{m-1}(\dot{x})\right]^{1/p} \quad (p=1,2,\ldots).$$

Then we have:

## Proposition 1.

$$\kappa_{av}^{[p]}(x,y,k) = \left[\frac{1}{\sqrt{\pi}} \cdot \frac{\Gamma\left(\frac{m}{2}\right)}{\Gamma\left(\frac{m+p}{2}\right)} \cdot \Gamma\left(\frac{p+1}{2}\right)\right]^{1/p} \cdot \kappa_F(x,y,k).$$

In particular,

$$\kappa_{av}(x, y, k) = \frac{\kappa_F(x, y, k)}{\sqrt{m}}.$$

*Proof.* Observe that  $\kappa_{av}^{[p]}(x, y, k)$  is the *p*th-average condition number for the problem of finding the *k*th-component of  $G = (G_1, \ldots, G_n)$ . Theorem 1 applied to  $G_k$  yields

$$\kappa_{av}^{[p]}(x,y,k) = \frac{1}{\sqrt{2}} \left[ \frac{\Gamma\left(\frac{m}{2}\right)}{\Gamma\left(\frac{m+p}{2}\right)} \right]^{\frac{1}{p}} \mathbb{E}(|\eta_{\sigma_1}|^p)^{1/p}$$

where  $\sigma_1 = \|DG_k(x)\| = \kappa(x, y, k)$ . Then,

$$\mathbb{E}(|\eta_{\sigma_1}|^p)^{1/p} = \kappa(x, y, k) \cdot \mathbb{E}(|\eta_1|^p)^{1/p}$$

where  $\eta_1$  is a standard normal in  $\mathbb{R}$ . Finally,

$$\mathbb{E}(|\eta_1|^p) = \frac{2}{\sqrt{2\pi}} \int_0^\infty \rho^p e^{-\rho^2/2} \, d\rho = \frac{2}{\sqrt{2\pi}} 2^{\frac{p-1}{2}} \Gamma(\frac{p+1}{2})$$

and the proposition follows.

#### 3. Examples

In this section we will compute the average condition number for different problems: systems of linear equations, eigenvalue and eigenvector problems, finding kernels of linear transformations and solving polynomial systems of equations. The first two have been computed in [12] and are an easy consequence of *Theorem 1* and the usual condition number.

3.1. Systems of Linear Equations. We consider the problem of solving the system of linear equations Ay = b, where  $A \in \mathcal{M}_n(\mathbb{R})$  the space of  $n \times n$  matrices with the Frobenius inner product, i.e.  $\langle A, B \rangle_F = \text{trace}(B^t A)$  ( $B^t$  is the transpose of B), and  $b \in \mathbb{R}^n$ .

If we assume that b is fixed, then the input space  $X = \mathscr{M}_n(\mathbb{R})$  with the Frobenius inner product,  $Y = \mathbb{R}^n$  with the Euclidean inner product, and  $\Sigma$  equals the subset of non-invertible matrices. Then the map  $G : \mathscr{M}_n(\mathbb{R}) \setminus \Sigma \to \mathbb{R}^n$  is globally defined and differentiable, namely

$$G(A) = A^{-1}b \ (=y).$$

By implicit differentiation,

(7) 
$$DG(A)(\dot{A}) = -A^{-1}\dot{A}y.$$

Is easy to see from (7) that  $\kappa(A, y) = ||A^{-1}|| \cdot ||y||$  and  $\kappa_F(A, y) = ||A^{-1}||_F \cdot ||y||$ . Then from *Theorem 1* we get

(8) 
$$\kappa_{av}(A,y) = \frac{\|A^{-1}\|_F \cdot \|y\|}{n} \le \frac{\kappa(A,y)}{\sqrt{n}}.$$

A similar result was proved in [12].

For the general case, we have  $X = \mathscr{M}_n(\mathbb{R}) \times \mathbb{R}^n$  with the Frobenius inner product in  $\mathscr{M}_n(\mathbb{R})$ and the Euclidean inner product in  $\mathbb{R}^n$ . Then,  $G : \mathscr{M}_n(\mathbb{R}) \setminus \Sigma \times \mathbb{R}^n \to \mathbb{R}^n$  satisfies  $G(A, b) = A^{-1}b$ . Is easy to see that  $\kappa((A, b), y) = ||A^{-1}|| \cdot \sqrt{1 + ||y||^2}$  and  $\kappa_F((A, b), y) = ||A^{-1}||_F \cdot \sqrt{1 + ||y||^2}$ . Again from *Theorem 1* we get

$$\kappa_{av}((A,b),y) = \frac{\|A^{-1}\|_F \cdot \sqrt{1+\|y\|^2}}{\sqrt{n^2+n}} \le \frac{\kappa((A,b),y)}{\sqrt{n+1}}.$$

For the kth-componentwise condition number, we have that

$$\kappa_{av}((A,b),y,k) = \left[\frac{1}{\sqrt{\pi}} \cdot \frac{\Gamma\left(\frac{n^2}{2}\right)}{\Gamma\left(\frac{n^2+p}{2}\right)} \cdot \Gamma\left(\frac{p+1}{2}\right)\right]^{1/p} \cdot \kappa((A,b),y).$$

A similar result was proved in [14].

In [7], it is proved that the expected value of the relative condition number  $\kappa_{rel}(A) = ||A|| \cdot ||A^{-1}||$ of a random matrix A whose elements are i.i.d standard normal, satisfies:

$$\mathbb{E}(\log \kappa_{rel}(A)) = \log n + c + o(1),$$

as  $m \to \infty$ , where  $c \approx 1.537$ . If we consider the relative average condition number defined in (4), we get from (8)

$$\mathbb{E}(\log \kappa_{relav}(A)) = \frac{1}{2}\log n + c + o(1),$$

as  $m \to \infty$ .

3.2. Eigenvalue and Eigenvector Problem. Let  $X = \mathscr{M}_n(\mathbb{C})$  be the space of  $n \times n$  complex matrices with the Frobenius Hermitian inner product,  $Y = \mathbb{P}(\mathbb{C}^{n+1}) \times \mathbb{C}$  and  $V = \{(A, v, \lambda) : Av = \lambda v\}$ . Then for  $(A, v, \lambda) \in V \setminus \Sigma$  the condition matrices  $DG_1$  and  $DG_2$  associated with the eigenvector and eigenvalue problem are

$$DG_1(A)\dot{A} = (\pi_{v^{\perp}}(\lambda I - A)|_{v^{\perp}})^{-1} \left(\pi_{v^{\perp}}\dot{A}v\right) \quad \text{and} \quad DG_2(A)\dot{A} = \frac{\langle u, Av \rangle}{\langle u, v \rangle}$$

where u is some left eigenvector associated with  $\lambda$ , i.e.  $u^*A = \overline{\lambda}u^*$  (see [Bez IV]). The associated condition numbers are:

(9) 
$$\kappa_1(A, v) = \left\| (\pi_{v^{\perp}}(\lambda I - A)|_{v^{\perp}})^{-1} \right\| \text{ and } \kappa_2(A, \lambda) = \frac{\|u\| \cdot \|v\|}{|\langle u, v \rangle|}.$$

From our *Theorem 1*, we get the respective average condition numbers:

$$\kappa_{av1}(A,v) = \frac{1}{n} \left\| (\pi_{v\perp}(\lambda I - A)|_{v\perp})^{-1} \right\|_F \le \frac{1}{\sqrt{n}} \kappa_1(A,v),$$
$$\kappa_{av2}(A,\lambda) = \frac{1}{n} \kappa_2(A,\lambda).$$

A similar result for  $\kappa_{av2}(A, \lambda)$  was proved in [12].

3.3. Finding Kernels of Linear Transformations. Let  $\mathscr{M}_{k,p}(\mathbb{C})$  be the linear space of  $k \times p$  complex matrices with the Frobenius Hermitian inner product, i.e.  $\langle A, B \rangle_F = \operatorname{trace}(B^*A)$  ( $B^*$  is the adjoint of B), and  $\mathcal{R}_r \subset \mathscr{M}_{k,p}(\mathbb{C})$  the subset of matrices of rank r. We consider the problem of solving the system of linear equations Ax = 0. For this purpose, we introduce the *Grassmannian*  $\mathbb{G}_{p,\ell}$  of complex subspaces of dimension  $\ell$  in  $\mathbb{C}^p$ , where  $\ell = \dim(\ker A) = p - r$ .

The input space  $X = \mathcal{R}_r$  is a smooth submanifold of  $\mathscr{M}_{k,p}(\mathbb{C})$  of complex dimension  $(k+p)r-r^2$ (see [6]). Thus, it has a natural Hermitian structure induced by the Frobenius metric on  $\mathscr{M}_{k,p}(\mathbb{C})$ .

In what follows, we identify  $\mathbb{G}_{p,\ell}$  with the quotient  $\mathbb{S}_{p,\ell}/\mathcal{U}_{\ell}$  of the *Stiefel* manifold

$$\mathbb{S}_{p,\ell} := \{ M \in \mathscr{M}_{p,\ell}(\mathbb{C}) : M^*M = I \}$$

#### DIEGO ARMENTANO

by the unitary group  $\mathcal{U}_{\ell} \subset \mathscr{M}_{\ell}(\mathbb{C})$ , which acts on the right of  $\mathbb{S}_{p,\ell}$  in the natural way (see [6]). Then, the complex dimension of the output space  $Y = \mathbb{G}_{p,\ell}$  is (p-r)r.

We will use the same symbol to represent an element of  $\mathbb{S}_{p,\ell}$  and its class in  $\mathbb{G}_{p,\ell}$ . The manifold  $\mathbb{S}_{p,\ell}$  has a canonical Hermitian structure induced by the Frobenius norm in  $\mathscr{M}_{\ell}(\mathbb{C})$ . On the other hand,  $\mathcal{U}_{\ell}$  is a Lie group of isometries acting on  $\mathbb{S}_{p,\ell}$ . Therefore,  $\mathbb{G}_{p,\ell}$  is a Homogeneous space (see [8]), with a natural Riemannian structure that makes the projection  $\mathbb{S}_{p,\ell} \to \mathbb{G}_{p,\ell}$  a Riemannian submersion. The orbit of  $M \in \mathbb{S}_{p,\ell}$  under the action of the unitary group  $\mathcal{U}_{\ell}$ , namely,  $o_{\ell}(M) = \{MU : U \in \mathcal{U}_{\ell}\}$ , defines a smooth submanifold of  $\mathbb{S}_{p,\ell}$ . In this form we can define a local chart of a small neighborhood of  $M \in \mathbb{G}_{p,\ell}$  from the affine spaces

$$M + T_M o_\ell(M)^{\perp}$$

where  $T_M o_\ell(M)^{\perp}$  is the orthogonal complement of  $T_M o_\ell(M)$  in  $T_M \mathbb{S}_{p,\ell}$ . Implicit differentiation in local coordinates of the input-output map G at the point  $G(A) = M \in \mathbb{S}_{p,\ell}$  yields

(10) 
$$DG(A)(\dot{A}) = -A^{\dagger}\dot{A}M,$$

where  $\dot{A} \in T_A \mathcal{R}_r$ ,  $A^{\dagger}$  is the Moore-Penrose inverse of A and  $A^{\dagger} \dot{A} M \in T_M o_{\ell}(M)^{\perp}$ .

One way to compute the singular values of the condition matrix described in (10), is to take an orthonormal basis in  $\mathscr{M}_{k,p}(\mathbb{C})$  which diagonalizes A. From the singular value decomposition, there exists orthonormal basis  $\{u_1, \ldots, u_k\}$  of  $\mathbb{C}^k$ , and  $\{v_1, \ldots, v_p\}$  of  $\mathbb{C}^p$ , such that  $Av_i = \sigma_i u_i$ for  $i = 1, \ldots, r$ , and  $Av_i = 0$  for  $i = r + 1, \ldots, p$ . Thus,  $\{u_i v_j^* : i = 1, \ldots, k; j = 1, \ldots, p\}$  is an orthonormal basis of  $\mathscr{M}_{k,p}(\mathbb{C})$  which diagonalizes A. In this basis the tangent space  $T_A \mathcal{R}_r$  is the orthogonal complement of the subspace generated by  $\{u_i v_j^* : i = r + 1, \ldots, k; j = r + 1, \ldots, p\}$ . From where we conclude that

$$\kappa(A, M) = \|DG(A)\| = \|A^{\dagger}\|, \quad \kappa_F(A, M) = \sqrt{p - r} \cdot \|A^{\dagger}\|_F.$$

From our Theorem 1,

$$\kappa_{av}(A,M) = \frac{\sqrt{p-r}}{\sqrt{(k+p-r)r}} \cdot \|A^{\dagger}\|_F \le \sqrt{\frac{p(p-r)}{(k+p-r)r}} \cdot \kappa(A,M).$$

In [1], it is proved that

$$\mathbb{E}(\log \kappa_{rel}(A): A \in \mathcal{R}_r) \le \log \left[\frac{k+p-r}{k+p-2r+1}\right] + 2.6,$$

where the expected value is computed with respect to the normalized naturally induced measure in  $\mathcal{R}_r$ . Our *Theorem 1* immediately yields a bound for the average relative condition number, namely,

$$\mathbb{E}(\log \kappa_{relav}(A): A \in \mathcal{R}_r) \le \frac{1}{2} \log \left[\frac{(k+p-r)r}{(k+p-2r+1)^2 p(p-r)}\right] + 2.6$$

3.4. Finding Roots Problem I: Univariate Polynomials. We start with the case of one polynomial in one complex variable. Let  $X = \mathcal{P}_d = \{f : f(z) = \sum_{i=0}^d f_i z^i, f_i \in \mathbb{C}\}$ . Identifying  $\mathcal{P}_d$  with  $\mathbb{C}^{d+1}$  we can define two standard inner products in the space  $\mathcal{P}_d$ : - Weyl inner product:

(11) 
$$\langle f,g\rangle_W := \sum_{i=0}^d f_i \overline{g_i} \binom{d}{i}^{-1};$$

- Canonical Hermitian inner product:

(12) 
$$\langle f, g \rangle_{\mathbb{C}^{d+1}} := \sum_{i=0}^{d} f_i \overline{g_i}$$

The solution variety is given by  $V = \{(f, z) : f(z) = 0\}$ . Thus, by implicit differentiation

$$DG(f)(\dot{f}) = -(f'(\zeta))^{-1}\dot{f}(\zeta).$$

We denote by  $\kappa_W$  and  $\kappa_{\mathbb{C}^{d+1}}$  the condition numbers with respect to the Weyl and Euclidean inner product. The reader may check that

$$\kappa_W(f,\zeta) = \frac{(1+|\zeta|^2)^{d/2}}{|f'(\zeta)|} \quad \text{and} \quad \kappa_{\mathbb{C}^{d+1}}(f,\zeta) = \frac{\sqrt{\sum_{i=0}^d |\zeta|^{2i}}}{|f'(\zeta)|},$$

(for a proof see [3], p. 228). From *Theorem 1*, we get:

$$\kappa_{avW}^{[2]}(f,\zeta) = \frac{1}{\sqrt{2(d+1)}} \kappa_W(f,\zeta), \qquad \kappa_{av\mathbb{C}^{d+1}}^{[2]}(f,\zeta) = \frac{1}{\sqrt{2(d+1)}} \kappa_{\mathbb{C}^{d+1}}(f,\zeta).$$

3.5. Finding Roots Problem II: Systems of Polynomial Equations. We now study the case of complex homogeneous polynomial systems. Let  $\mathcal{H}_{(d)}$  the space of systems  $f : \mathbb{C}^{n+1} \to \mathbb{C}^n$ ,  $f = (f_1, \ldots, f_n)$  where each  $f_i$  is a homogeneous polynomial of degree  $d_i$ . We consider  $\mathcal{H}_{(d)}$  with the homogeneous analogous of the Weyl structure defined above (see Chapter 12 of [3] for details).

Let  $X = \mathbb{P}(\mathcal{H}_{(d)})$  and  $Y = \mathbb{P}(\mathbb{C}^{n+1})$  and  $V = \{(f, \zeta) : f(\zeta) = 0\}$ . We denote by  $N = \sum_{i=1}^{n} {d_i+n \choose n} - 1$  the complex dimension of X. We may think of 2N as the size of the input. Then,

$$DG(f)(\dot{f}) = -\left(Df(\zeta)_{\zeta^{\perp}}\right)^{-1}\dot{f}(\zeta),$$

and the condition number is

$$\kappa_W(f,\zeta) = \left\| \left( Df(\zeta)_{\zeta^\perp} \right)^{-1} \right\|,\,$$

where some norm 1 affine representatives of f and  $\zeta$  have been chosen (cf.[3]). Associated with this quantity, we consider

(13) 
$$\kappa_W(f) := \sqrt{\frac{1}{\mathcal{D}} \sum_{\{\zeta : f(\zeta) = 0\}} \kappa_W(f, \zeta)^2}$$

where  $\mathcal{D} = d_1 \cdots d_n$  is the number of projective solutions of a generic system.

The expected value of  $\kappa_W^2(f)$  is an essential ingredient in the complexity analysis of pathfollowing methods (cf. [9], [2]). In [2] the authors proved that

(14) 
$$\mathbb{E}_f\left[\kappa_W(f)\right)^2\right] \le 8nN,$$

where f is chosen at random with the Weyl distribution.

The relation between complexity theory and  $\kappa_{av}$  is not clear yet. However, it is interesting to study the expected value of the  $\kappa_{av}$ -analogous of equation (14), namely

$$\kappa_{avW}(f) := \sqrt{\frac{1}{\mathcal{D}}} \sum_{\{\zeta: f(\zeta)=0\}} \kappa_{avW}(f,\zeta)^2.$$

From our *Theorem 1* we get,

$$\kappa_{avW}(f,\zeta) \leq \frac{\kappa_W(f,\zeta)}{\sqrt{N/n}}, \qquad \mathbb{E}_f\left[\kappa_{avW}(f)\right)^2\right] \leq 8n^2.$$

Note that the last bound depends on the number of unknowns n, and not on the size of the input  $n \ll N$ .

## 4. PROOF OF THE MAIN THEOREM

In the case of complex manifolds, the condition matrix turns to be an  $n \times n$  complex matrix. In what follows, we identify it with the associated  $2n \times 2n$  real matrix. We center our attention in the real case.

The main theorem follows immediately from Lemma 1 and Proposition 2 below.

**Lemma 1.** Let  $\eta$  be a Gaussian standard random vector in  $\mathbb{R}^m$ . Then

$$\kappa_{av}^{[p]}(x,y) = \frac{1}{\sqrt{2}} \left[ \frac{\Gamma\left(\frac{m}{2}\right)}{\Gamma\left(\frac{m+p}{2}\right)} \right]^{\frac{1}{p}} \cdot \left[ \mathbb{E}(\|DG(x)\eta\|^p) \right]^{1/p},$$

where  $\mathbb{E}$  is the expectation operator and  $\|\cdot\|$  is the Euclidean norm in  $\mathbb{R}^n$ .

*Proof.* Let  $f : \mathbb{R}^m \to \mathbb{R}$  be the continuous function given by

$$f(v) = \|DG(x)v\|.$$

Then

$$\left[\mathbb{E}(\|DG(x)\eta\|^p)\right]^{1/p} = \left[\frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m} f(v)^p \cdot e^{-\|v\|^2/2} \, dv\right]^{1/p}.$$

Integrating in polar coordinates we get that

(15) 
$$\mathbb{E}(\|DG(x)\eta\|^p) = \frac{I_{m+p-1}}{(2\pi)^{m/2}} \cdot \int_{S^{m-1}} f^p \, dS^{m-1}$$

where

$$I_j = \int_0^{+\infty} \rho^j \, e^{-\rho^2/2} \, d\rho, \quad j \in \mathbb{N}.$$

Making the change of variable  $u = \rho^2/2$  we obtain

$$I_j = 2^{\frac{j-1}{2}} \Gamma(\frac{j+1}{2}),$$

therefore

(16) 
$$I_{m+p-1} = 2^{\frac{m+p-2}{2}} \cdot \Gamma\left(\frac{m+p}{2}\right).$$

Then joining together (15) and (16) we obtain the result.

## **Proposition 2.**

$$\mathbb{E}(\|DG(x)\eta\|^p) = \mathbb{E}(\|\eta_{\sigma_1,\dots,\sigma_n}\|^p)$$

where  $\eta_{\sigma_1,\ldots,\sigma_n}$  is a centered Gaussian vector in  $\mathbb{R}^n$  with diagonal covariance matrix  $Diag(\sigma_1^2,\ldots,\sigma_n^2)$ .

*Proof.* Let DG(x) = UDV be a singular value decomposition of DG(x). By the invariance of the Gaussian distribution under the action of the orthogonal group in  $X, V\eta$  is again a standard Gaussian random vector. Then,

$$\mathbb{E}(\|DG(x)\eta\|^p) = \mathbb{E}(\|UD\eta\|^p),$$

and by the invariance under the action of the orthogonal group of the Euclidean norm, we get

$$\mathbb{E}(\|DG(x)\eta\|^p) = \mathbb{E}(\|D\eta\|^p).$$

Finally  $D\eta$  is a centered Gaussian vector in  $\mathbb{R}^n$  with covariance matrix  $Diag(\sigma_1^2, \ldots, \sigma_n^2)$ , and the proposition follows. For the case p = 2

$$\kappa_{av}(x,y) = \left[\mathbb{E}\left(\sigma_1^2\eta_1^2 + \ldots + \sigma_n^2\eta_n^2\right)\right]^{1/2}$$

where  $\eta_1, \ldots, \eta_n$  are i.i.d. standard normal in  $\mathbb{R}$ . Then

$$\kappa_{av}(x,y) = \left(\sum_{i=1}^{n} \sigma_i^2\right)^{1/2} = \kappa_F(x,y).$$

Acknowledgments

The author thanks Professors Carlos Beltrán, Michael Shub and Mario Wschebor for useful discussions.

#### References

- [1] C. Beltrán: Estimates On The Condition Number Of Random, Rank-Deficient Matrices. (to appear)
- [2] C. Beltrán and L.M. Pardo: Fast linear homotopy to find approximate zeros of polynomial systems. (to appear)
- [3] L. Blum, F. Cucker, M. Shub and S. Smale: Complexity and real computation. Springer-Verlag, New York, 1998.
- [4] P. Bürgisser, F. Cucker and M. Lotz: Smoothed analysis of complex conic condition numbers, J. Math. Pures et Appl. 86, 293-309, 2006.
- [5] P. Bürgisser: Smoothed Analysis of Condition Numbers, (to appear).
- [6] J.P. Dedieu: Points Fixes, Zéros et la Méthode de Newton. Mathématiques & Applications SMAI-Springer Verlag, 54, 2006.
- [7] A. Edelman: Eigenvalues and condition numbers of random matrices, SIAM J. of Matrix Anal. and Applic., 9, 543-560, 1988.
- [8] S. Gallot, D. Hulin and J. Lafontaine: Riemannian geometry. Third edition. Universitext. Springer-Verlag, Berlin, 2004.
- M. Shub and S. Smale: Complexity of Bezout's theorem. IV. Probability of success; extensions. SIAM J. Numer. Anal. 33, no. 1, 128-148, 1996.
- [10] S. Smale: On the efficiency of Algorithms of analysis, Bull. Amer. Math. Soc., 13, 87-121, 1985.
- [11] D.A. Spielman and S.H. Teng: Smoothed analysis of algorithms, ICM 2002, Beijing, Vol. I, pp. 597-606, 2002.
- [12] G. W. Stewart: Stochastic perturbation theory, SIAM Rev. 32, no. 4, 579-610, 1990.
- [13] L.N. Trefethen and D. Bau, III: Numerical linear algebra, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1997.

- [14] N. Weiss, G.W. Wasilkowski, H. Wozniakowski, and M. Shub: Average condition number for solving linear equations. *Linear Algebra Appl.* 83, 79-102, 1986.
- [15] M. Wschebor: Smoothed Analysis of  $\kappa(a)$ , J. of Complexity, 20, 97-107, 2004.