

1 Efficacy of Put and Spd sprayed on leaves from *Brassica juncea* plants against Cd<sup>2+</sup>-induced  
2 oxidative stress.

3

4

5 Michel AOUN<sup>a</sup>, Annick HOURMANT<sup>a</sup>, Jean-Yves CABON<sup>b</sup>

6

7 Laboratoire de Toxicologie alimentaire et cellulaire, Université de Bretagne Occidentale  
8 (Brest-France)<sup>a</sup>.

9 Laboratoires de Chimie, électrochimie moléculaire et chimie analytique-UMR 6521,  
10 Université de Bretagne Occidentale (Brest-France)<sup>b</sup>.

11 \*Dr. Michel AOUN, UBO

12 6, Avenue Le Gorgeu

13 29200 BREST

14 michel.aoun@univ-brest.fr\*

15

## 16 **Abstract**

17

18 The protective effect exerted by polyamines (Put and Spd) against cadmium (Cd)  
19 stress was investigated in *Brassica juncea* plants. Treatment with CdCl<sub>2</sub> (75 μM)  
20 resulted in a rise of Cd accumulation, a decrease of fresh and dry weights in every  
21 plant organ, an increase of free polyamine content at limb and stem levels as well as a  
22 decrease at root level. On the other hand, the total conjugated polyamine levels in the stem  
23 tissues were unaffected by Cd. In the leaf tissues, this metal caused a reduction of chlorophyll  
24 a content, a rise of guaiacol peroxidase (GPOX) activity and an increase of malondialdehyde  
25 (MDA), soluble glucide, proline and amino acid contents.

26 Exogenous application, by spraying, of putrescine (Put) and spermidine (Spd) to leaf tissues  
27 reduced CdCl<sub>2</sub>-induced stress. These polyamines proved to exert a partial,  
28 though significant, protection of the foliar fresh weight and to alleviate the oxidative stress  
29 generated by Cd through reductions of MDA amounts and GPOX (E.C.1.11.1.7) activity.  
30 The enhancement of chlorophyll a content in plants by Put and those of Chl a and Chl b by  
31 Spd both constitute evidences of their efficacy against the Cd<sup>2+</sup>-induced loss of pigments.  
32 Conversely to Put, Spd caused a decrease of Cd content in leave tissues and a rise in the stems  
33 and roots; these findings are in favour of a stimulation of Cd uptake by Spd. The proline  
34 stimulation observed with Cd was reduced further to the spraying of Put onto tissues, but the  
35 decrease induced by Spd was more limited. In the plants treated with Cd, the amino acid  
36 contents in the leaves were unaffected by Put and Spd spraying; on the other hand, Cd<sup>2+</sup>  
37 disturbe polyamine levels (free and acido-soluble conjugated-forms); we notice the rise of  
38 total free PAs and the decrease of their conjugated-ones.  
39 The results of the present study suggest that Put and Spd can efficiently protect *Brassica*  
40 *juncea* plants against Cd-induced oxidative damage likely through a strong reduction of H<sub>2</sub>O<sub>2</sub>  
41 generation and pigment protection. Moreover, the use of Spd in phytoremediation processes is  
42 briefly discussed.

43 **Key words:** *Brassica juncea*, CdCl<sub>2</sub>, oxidative stress, amino acids, PAs, proline.

44

#### 45 **Abbreviations**

46

47	GPOX	Guaiacol peroxidase
48	MDA	Malondialdehyde
49	MES	2-(N-morpholino ethane sulfonic) acid
50	PAs	Polyamines

51	PCA	Perchloric acid
52	Put	Putrescine
53	Spd	Spermidine
54	Spm	Spermine
55	SSAT	Spermidine / Spermine N1-Acetyltransferase

56

## 57 **Introduction**

58

59 The contamination of soils, water and the atmosphere further to the dispersal of industrial and  
60 urban wastes generated by human activities is, nowadays, a major environmental concern. It  
61 is caused by a wide range of organic contaminants (combustible and putrescible substances,  
62 hazardous wastes, explosives and petroleum products) and inorganic pollutants (heavy  
63 metals) (Adriano, 1986; Alloway, 1990).

64 Among these metals, Cd as well as Zn, Cu, Pb, .... are widespread contaminants  
65 known to be toxic to humans, animals and plants. Furthermore, Cd<sup>2+</sup> is easily taken up by  
66 plants, where its toxicity becomes manifest through growth reduction (Chen and Kao, 1995),  
67 decrease in chlorophyll contents (Stobart et al. 1985; Larsson et al. 1998) and decline in  
68 transpiration rate (Haag-Kerwer et al. 1999). Moreover, lipid peroxidation (Ferrat *et al.*, 2003)  
69 and guaiacol peroxidase activity (Shah *et al.*, 2001; Srivastava *et al.*, 2004; Smeets *et al.*,  
70 2005; Mishra *et al.*, 2006; Hsu et Kao, 2007; Semane *et al.*, 2007) are enhanced by this metal  
71 and others, whereas carbohydrate metabolism (Moya *et al.*, 1993; Costa and Spitz, 1997),  
72 amino acid content (Sharma and Dietz, 2006), proline (Lesko and Simon-Sarkadi, 2002;  
73 Singh and Tewari, 2003; Sharma and Dietz, 2006) and polyamine one (Lesko and Simon-  
74 Sarkadi, 2002; Groppa et al. 2003; Kuthanova et al. 2004; Taulavuori *et al.*, 2005 ; Sharma  
75 and Dietz, 2006; Groppa *et al.*, 2007a, b; Groppa and Benavides, 2008) are altered.

76 Oxygen is essential for aerobic organisms, but the reactive oxygen species (ROS), including  
77 the superoxide anion ( $O_2^{\cdot-}$ ), hydroxyl radicals ( $OH^{\cdot}$ ) and hydrogen peroxide ( $H_2O_2$ ) (Foyer et  
78 al. 1994, 1997; Asada, 1999), which are generated in the course of metabolic reactions within  
79 the cells, are known to be toxic. Since long, ROS have been considered as damaging to cell  
80 membranes (Apel and Hirt, 2004) and implicated in ageing and wounding (Thompson et al.  
81 1987). It is only recently that they have emerged as ubiquitous signalling molecules involved  
82 in the recognition of stress factors as well as in the response to them (Foyer and Noctor,  
83 2005). Plants cope with the oxidative stress by using antioxidant enzymes such as superoxide  
84 dismutase (SOD, E.C. 1.15.1.1), ascorbate peroxidase (APX, E.C. 1.11.11.1), catalase (CAT,  
85 EC 1.11.1.6), glutathione reductase (GR, E.C. 1.6.4.2), peroxidase (POX, E.C.1.11.1.6),  
86 guaiacol peroxidase (GPOX, E.C. 1.11.1.7), glutathione peroxidase (GSH-PX, E.C. 1.11.1.9)  
87 and many NADP-dehydrogenases as well as antioxidant compounds of low molecular weight,  
88 such as ascorbate and glutathione (Noctor and Foyer, 1998; Asada, 1999; Del Rio et al. 1998,  
89 2002).

90 The toxicity of Cd to plant cells is related to the oxidative stress caused through the  
91 generation of ROS (Piqueras et al. 1999; Sandalio et al. 2001; Schützendübel and Polle, 2002;  
92 Olmos et al. 2003; Kuo and Kao, 2004; Romero- Puertas et al. 2004), the stimulation, or  
93 inhibition, of the activities by antioxidant enzymes (Shaw, 1995; Gallego et al. 1996; Chaoui  
94 et al. 1997; Dixit et al. 2001; Shah et al. 2001; Innelli et al. 2002; Kuo and Kao, 2004), the  
95 production of oxidative damage, the induction of lipid peroxidation (Shaw, 1995; Gallego et  
96 al. 1996; Chaoui et al. 1997; Lozano-Rodriguez et al. 1997; Dixit et al. 2001; Shah et al.  
97 2001, Chien et al. 2002; Kuo and Kao, 2004) and protein oxidation (Pena et al. 2006).

98

99 Aliphatic polyamines are synthesised in both Prokaryotes and Eukaryotes. Among them,  
100 putrescine (Put), spermidine (Spd) and spermine (Spm) are the most common ones. Moreover,

101 Put and Spd are usually more abundant than Spm, which is often present at trace amounts. In  
102 plants, these polyamines are not only involved in numerous cellular and molecular processes  
103 (Slocum et al. 1984; Bouchereau et al. 1999; Wallace et al. 2003), but also in their responses  
104 to environmental stresses by salt (Sun et al. 2002; Zhao et al. 2003), water (Xing et al. 2000),  
105 chilling (Lee, 1997; Zheng et al. 2000), acid rain (Velikova et al. 2000), ozone (Ormord and  
106 Beckerson, 1986; Navakoudis et al. 2003), paraquat (Ye et al. 1997) and heavy metal(s)  
107 (Groppa et al. 2001; Wang et al., 2004; Taulavuori et al. 2005; Sharma and Dietz, 2006;  
108 Groppa et al. 2007a, b). Moreover, in engineered plants, tolerance to multiple environmental  
109 stresses is enhanced by an overproduction of polyamines (Roy and Wu, 2001; Capell et al.  
110 2004; Kasukabe et al. 2004). Less is known about the biosynthetic activities that lead to their  
111 accumulation in response to heavy metals (Weinstein et al. 1986; Wettlaufer et al. 1991;  
112 Groppa et al. 2003; Balestrasse et al. 2005).

113 Evidences of the protection exerted by exogenous polyamines against oxidative stress of  
114 various origins are available in the literature: they are about paraquat, a viologen often used as  
115 herbicide (Bors et al. 1989; Minton et al. 1990; Chang and Kao, 1997; Kurepa et al. 1998;  
116 Benavides et al. 2000), acid rain (Velikova et al. 2000), salt (Tang 123 and Newton, 2005)  
117 and heavy metals, e.g. Cd and Cu, (Groppa et al. 2001; Urano et al. 2003; Tang et al. 2005;  
118 Franchin et al. 2007; Groppa et al. 2007a, b; Hsu and Kao, 2007). Drolet and co-workers  
119 (1986) showed that polyamines are effective radical scavengers in a number of chemical- and  
120 *in vitro* enzyme-systems. Under osmotic stress, ROS production is enhanced by a reduction of  
121 polyamine content in the leaves of *Glycyrrhiza inflata* (Li and Wang, 2004).

122 These considerations led us to carry out experiments aimed at determining whether the  
123 toxicity by Cd was affected by spraying Put and Spd onto the leaves of *Brassica juncea* plants  
124 issued from seedlings grown hydroponically. They showed that polyamine metabolism was

125 disturbed by Cd and that, further to an exogenous application of Put and Spd by spraying at  
126 the leaf tissues, the oxidative damage caused by CdCl<sub>2</sub> was reduced.

127

## 128 **Material and methods**

129

### 130 Plant material

131

132 The cultivar, *Brassica juncea* I39/1, is a pure spring line, genetically fixed and  
133 obtained by autofertilisation.

134

### 135 Culture conditions

136

137 Indian mustard (*Brassica juncea*) seeds were grown on humidified vermiculite at 22°C for  
138 3 days under dark conditions. At day 3, uniformly germinated seeds were selected and  
139 cultivated again on vermiculite for 15 days. Then, at day 18, plantlets were extracted from  
140 vermiculite and their roots were carefully washed under water flow. The extracted plants were  
141 grown, for 84 hours, in a hydroponic system composed of a half-strength Hoagland solution  
142 (Hoagland and Arnon, 1950) in MES buffer (1 mM) and enriched with CdCl<sub>2</sub> (75 µM). the  
143 final pH was 6.60. Solutions of Put (1 mM) and Spd (1 mM) were prepared in Teepol (0.01  
144 %, v/v) and further sprayed at the surface of leaves. The control plants were sprayed with only  
145 Teepol. All experiments were carried out in a growth chamber maintained at 22°C  
146 (light)/20°C (dark) under a photoperiod (16 h) with 60 µmol photon.m<sup>-2</sup>.s<sup>-1</sup>) provided by cool  
147 white fluorescent lamps.

148

149 Determination of pigment contents

150

151 The contents in chlorophylls a, b and carotenoids were determined by using the procedure  
152 developed by Lichtenthaler (Lichtenthaler, 1987). Briefly, about 1 g of limb fresh weight was  
153 extracted at 4°C with acetone 100% (25 ml), then the mixture was centrifuged 10 min at  
154 5 000 g and 4°C prior to the recording of absorbance at 470, 662 and 645 nm, respectively,  
155 with a UV-1605 spectrophotometer from Shimadzu.

156

157 Measurement of Cd in the different organs

158

159 Cadmium concentrations in the different solutions were measured by electrothermal atomic  
160 absorption spectrometry with a Perkin-Elmer, SIMAA 6100, spectrometer operated in the  
161 single-element mode. At harvest, the leaves, stems and roots were washed twice with distilled  
162 water for 5 min, weighed and then oven-dried for 4 days at 80°C prior to grounding. All  
163 solutions were prepared with high-purity water from a MilliQ-system (Millipore, Milford,  
164 MA, USA). Aliquots of about 200 mg of grounded matter were transferred to Teflon vessels  
165 prior to the addition of nitric acid, hydrogen peroxide and hydrofluoric acid in mixture (4/3/1,  
166 v/v/v). Then, the vessels were closed and exposed to microwave digestion as detailed  
167 elsewhere (Weiss et al. 1999).

168

169 Assessment of lipid peroxidation

170

171 Lipid peroxidation was assessed from the amount of produced malondialdehyde (MDA) able  
172 to react with thiobarbituric acid and was measured according to the method developed by  
173 Minotti and Aust (1987) and improved by Iturbe-Ormaetxe and co-workers (1998).

174 Approximately 1 g of leaves (fresh weight) was extracted by grinding with 5 mL of a mixture  
175 of meta-phosphoric acid (5%, w/v) and butylhydroxy-toluene (50/1, v/v). After centrifugation  
176 at 5 000 g for 30 min, 4 ml of the supernatant were homogenized with 200  $\mu$ L of  
177 butylhydroxytoluene (2%, w/v), 1 mL of HCl (25%, v/v) and 1 mL of 2-thiobarbituric acid  
178 (1%, w/v) prepared in NaOH (50 mM). The resulting mixture was incubated at 95°C for 30  
179 min, then quickly cooled in an ice bath to stop the reaction. Finally, the chromogen was  
180 extracted by addition of 3 ml of 1-butanol to the mixture. Then, after centrifugation at 500 g  
181 for 5 min, the absorbance was read at 532 nm. The extinction coefficient of MDA ( $\epsilon = 155$   
182  $\text{mmol L}^{-1} \text{cm}^{-1}$ ) was used to calculate the sample content in MDA.

183

184 Determination of guaiacol peroxidase activity (GPOX)

185

186 GPOX activity in leaves was measured by the procedure developed by MacAdam and co-  
187 workers (1992) after slight changes: briefly the use of guaiacol (4  $\mu$ mol) as substrate by  
188 peroxidases in the presence of  $\text{H}_2\text{O}_2$  leads to the formation of tetraguaiacol (1  $\mu$ mol), whose  
189 absorption, read at 470 nm, gives the amount of enzyme. The extinction coefficient of  
190 tetraguaiacol used to calculate GPOX activity was  $26.2 \text{ mM}^{-1} \text{ cm}^{-1}$ . One unit of GPOX was  
191 defined as the amount of enzyme that caused the formation of 1  $\mu$ mol tetraguaiacol per min.  
192 GPOX activity was expressed as the amount of oxidized guaiacol.

193

194 Amino acid determination

195

196 This determination is based on the fact that every amino acid reacts with ninhydrin at 100°C  
197 to form the Ruheman purple (Yemm and Cooking, 1955). Approximately, 1 g of leaves (fresh  
198 weight) was mixed in a glass tube with distilled water (7 mL) and heated at 100°C for 1 hour.

199 After cooling on ice, the mixture was ground and centrifuged at 5000 g for 10 min at 4°C. The  
200 pellet was submitted to a second extraction and centrifugation under the same conditions.  
201 Then, after pooling, 200 µl of supernatant were mixed with 0.5 ml of citrate buffer (160 mM,  
202 pH 4.6), before addition of 1 mL of ninhydrin (54 mM, w/v) dissolved in ethanol 70 % and  
203 2 mL of ascorbic acid (1%, w/v). After strong agitation (vortex), the tubes were placed in a  
204 water bath at 100°C for 20 min, then cooled on ice prior to the addition of 3 mL of ethanol  
205 70%. Then absorbance was read at 570 nm. The standard curve was constructed from a  
206 dilution series of a stock solution of L-leucine (5 mM). The amounts of amino acids were  
207 expressed as µmol equivalent leucine per g (fresh weight).

208

209

210 Measurement of free proline

211

212 The sample content in free proline was determined by spectrophotometry at 520 nm according  
213 to the procedure by Bates and co-workers. (1973): 1 g of leaves (fresh weight) was extracted  
214 with 10 mL of sulfosalicylic acid (3%, w/v). After filtration, 0.5 mL of the filtrate were  
215 incubated at 100°C with 1 mL ninhydrin (1%, w/v). The reaction was let to develop for 1  
216 hour, and then stopped by cooling in an ice bath prior to the addition of 3 mL of toluene.  
217 After vortexing of the final mixture for 2 min, it was incubated for 3 h in the dark and at room  
218 temperature before reading of the absorbance at 520 nm. The standard curve was constructed  
219 from a dilution series of a stock solution of L-proline (5 mM). The amounts of proline were  
220 expressed in µmol.g<sup>-1</sup> (fresh weight).

221

222 Sugar determination

223

224 Soluble sugars were determined according to the method by Cerning-Beroard (1975): under  
225 certain conditions of pH and heat, soluble sugars are degraded into furfural, which further  
226 reacts with anthrone to form a blue compound whose absorbance is read at 625 nm. About 1 g  
227 (F.W) of leaf was mixed in a tube with 7 ml distilled water before heating at 100°C for one  
228 hour. After cooling on ice, the mixture was ground, then centrifuged for 10 min at 5 000 g and  
229 4°C. The pellet was subjected to a second extraction and centrifugation under the same  
230 conditions. Then, the two supernatants were pooled, mixed and filtered on a 45- $\mu$ m filter  
231 paper.

232 An aliquot (500  $\mu$ l) was vortexed in glass tubes with 5 ml of a solution of anthrone (92 mg  
233 anthrone and 92 mg de thiourea dissolved in 100 ml sulfuric acid (70 %, v/v), then the  
234 mixture was heated at 100°C for 10 min. After cooling in an ice bath, absorbance was read at  
235 625 nm. The results are expressed as  $\mu$ moles equivalent glucose.g<sup>-1</sup> F.W.

236

237 Polyamine determination

238

239 Free PAs were extracted from leaf, stem and root tissues by the procedure described by Flores  
240 and Glaston (1982). To determine the amount of conjugated PAs in the acid soluble fraction,  
241 200  $\mu$ l of the supernatant were mixed with an equal volume of 12 N HCl and hydrolysed for a  
242 night at 110°C in flame-sealed ampoules. The hydrolysates were dried under a flow of air and  
243 resuspended in 200  $\mu$ l PCA (5 %, v/v). About PAs, they were derivatised according to the  
244 method by Flores and Glaston (1982), slightly modified by Féray and co-workers (1992). The  
245 dansyl derivatives of putrescine (Put), spermidine (Spd) and spermine (Spm) were separated  
246 by HPLC and quantified by fluorescence spectrophotometry. The HPLC apparatus consisted  
247 of two solvent pumps (Kontron 422) coupled to a high performance mixer, a 7125 Rheodyne  
248 injection valve fitted with 20  $\mu$ l loop and a stainless precolumn spheri ODS 5 mm

249 (Brownlee (30 X 4.6 mm) in conjunction with a stainless steel analytical column  
250 (Ultrasphere). The mobile phase consisted of 0.01 M NaH<sub>2</sub>PO<sub>4</sub>, pH 4.4 as solvent (A) and of  
251 methanol/acetonitrile in mixture (50/50, v/v) as solvent B. Both solvents were mixed at  
252 ambient temperature to produce a gradient elution at a flow-rate of 1 mL.min<sup>-1</sup> starting with B  
253 80% for 5 min, followed with B 80-89% for 2 min, then B 89-100% for 5 min, B 100-80% for  
254 3 min and finally B 80% for 10 min. The gradient and data analysis were controlled by a  
255 microcomputer data system (Kontron Inst.). Detection by spectrofluorimetry was achieved  
256 with a SFM 25 apparatus (Kontron) set at 360 and 510 nm wavelengths for excitation and  
257 emission, respectively.

258 The retention times (min) for Put, Spd and Spm were 7.6, 13.5 and 15.4, respectively.

259

260 Statistical analysis

261

262 The different results are expressed as the mean ± SD (standard deviation). Three independent  
263 experiments were performed and each measurement was made in triplicate. The statistical  
264 differences between measurements were assessed by one-way ANOVA and LSD tests.

265

## 266 **Results**

267

268 By comparison with the controls, Figure 1 shows clearly the significant reductions in the fresh  
269 weight of organs from 18-day-old *Brassica juncea* plants (leaves: 34%, roots: 30% and stems:  
270 20%) induced by the presence of CdCl<sub>2</sub> (75 µM) in the culture medium. One should note that  
271 the spraying of leaf tissues with Put or Spd in plants subjected to CdCl<sub>2</sub> had a positive effect  
272 by reducing the loss of fresh weight for the leaves (22-23% against 34%). On the other hand,  
273 this beneficial action was not observed for roots and stems (Fig. 1).

274 Table 1 highlight that 84 h-exposure of plants to CdCl<sub>2</sub> (75 μM) caused significant reductions  
275 in the dry weight (17% whatever the organ) and total water content of leaves (35%), roots  
276 (32%) and stems (24%). The dry weights of leaves and stems proved to be unaffected by  
277 spraying of Put or Spd upon the leaf tissues of the Cd-exposed plants conversely to that of the  
278 roots where the loss was only 9% against 17%.

279 Further to their exposure to CdCl<sub>2</sub>, the plants accumulated substantial amounts in their organs.  
280 As expected, this accumulation was the greatest in the roots (4506 μg.g<sup>-1</sup> D.W.) and far much  
281 lower in the stems and leaves (933 and 355 μg.g<sup>-1</sup> D.W, respectively) (Table 2). Spraying of  
282 Put on leaf tissues caused no significant difference 303 in the Cd contents of organs,  
283 conversely to that of Spd: indeed, Cd contents was significantly decreased in the leaves (20%)  
284 and significantly increased in the stems 305 and roots (22 and 86 %, respectively).

285

286 Figure 2 shows that, by comparison to controls, exposure of plants to CdCl<sub>2</sub> (75 μM) caused a  
287 significant reduction of the leave contents in chlorophyll a (15%), but not in chlorophyll b and  
288 carotenoids (P < 0.05). Then the Chl a/b ratio in Cd-exposed plants was 2.84 against 3.04.  
289 Spraying of Put or Spd upon control plants had no significant effect upon the pigment  
290 contents (Fig. 2). It is worth noting that spraying of Put upon Cd-exposed plants resulted in an  
291 elevation of Chl a content. That of spd caused significant rises of Chl a and Chl b contents (p  
292 < 0.05), and, thus, the Chl a/b ratio was significantly lowered (2.34 against 2.84, control  
293 value: 3.04). However, the carotenoid contents was unaffected by spraying of Put or Spd.

294

295 Figure 3 evidences the enhancement of lipid peroxidation (+ 157%) by exposure to Cd. On  
296 the other hand, MDA contents were unaffected by spraying of Put or Spd onto control plants  
297 (Fig. 3). The MDA content of Cd-exposed plants was lowered further to the spraying with Put

298 to control value but Spd proved to be less efficient (reduction of MDA content by 33% ,  
299 though still 71% higher than in controls.

300 Exposure to CdCl<sub>2</sub> induced a significant enhancement of GPOX activity (2.4-fold with  
301 respect to controls,  $p < 0.05$ ) (Fig. 4). On the other hand, it was unaffected by spraying of Put  
302 or Spd upon controls conversely to Cd-exposed plants where Put and Spd alleviated partially,  
303 but significantly the oxidative stress generated by Cd (1.9-fold reduction of GPOX activity).

304

305 Concerning the contents in free amino-acids, it was significantly increased (65%) by exposure  
306 to CdCl<sub>2</sub> (Fig. 5), but spraying of controls and Cd-exposed plants with Put or Spd had no  
307 significant effect upon the content in free amino-acids.

308

309 Table 3 evidences the rise in the proline content of leaf tissues (228%) induced by  
310 exposure to CdCl<sub>2</sub>. This content was unaffected by spraying of control plants with Put or Spd,  
311 conversely to the case of Cd-exposed plants, where further to Put spraying the proline content  
312 was markedly lowered (113% against 228%). However, Spd proved to be less efficient (1.86  
313 fold less than in Cd-exposed plants, but 80% higher than in control plants.

314

315 Figure 6 evidences the marked rise (117%) of soluble sugar levels observed further to the  
316 exposure of *Brassica juncea* seedlings to CdCl<sub>2</sub>. Spraying of Put or Spd upon the leaves of  
317 control plants had no effect on their soluble sugar content. The high sugar levels observed in  
318 Cd-exposed plants were greatly lowered by Spd, but not by Put: indeed, they were 56%  
319 higher than in controls.

320

321 In comparison to control plants, exposure to CdCl<sub>2</sub> (75  $\mu$ M) induced a significant  
322 accumulation of Put, Spd and Spm in the leaves (62, 33 and 31%) and the stems (13, 18 and

323 45%) (Fig. 7 A, B). On the other hand, in the roots, free polyamine levels were decreased by  
324 22, 58 and 71% for Put, Spd and Spm, respectively (Fig. 7C). Put amounts ( $P < 0.05$ ) were  
325 significantly increased at leaf and stem levels (Fig 7 A, B) and decreased in roots (Fig. 7 C)  
326 further to the spraying of Put (1 mM) on Cd-treated plants; moreover, Put (1 mM) decreased  
327 significantly Spd and Spm contents in leaf and root tissues (Fig. 7 A, C), but Spd and Spm  
328 stem levels were similar to those reported in Cd-treated plants (Fig. 7 B).

329 Spd (1 mM) applied exogenously increased significantly free Put contents at foliar, stem and  
330 root levels in comparison to Cd treated plants (Fig. 7 A, B, C). Moreover, Spd contents were  
331 increased significantly at foliar and root levels. However, Spd did not modified Spd stem  
332 levels and Spm contents at different organs in comparison to Cd treated plants (Fig. 7).

333

334  $\text{CdCl}_2$  (75  $\mu\text{M}$ ) treatment enhance significantly endogenous conjugated Put levels of 101 and  
335 10 % at foliar and stem levels respectively (Fig. 8 A, B), but not Put root levels in comparison  
336 to control plants (Fig. 8 C). Cd did not affected conjugated Spd levels but increase conjugated  
337 Spm levels of 36 % at foliar levels (Fig. 8 A). However, Cd decrease Spd and Spm levels of  
338 25 % at stem level, and 35 and 30 % at root levels in comparison to control values (Fig. 8 B,  
339 C). Put (1 mM) applied exogenously did not showed any significant effect on Put, Spd and  
340 Spm contents at foliar level ( $P < 0.05$ ). However, Put and Spd contents were increased but not  
341 Spm at stem level in comparison to Cd treated plants. Moreover, Put contents were unchanged  
342 at roots level, but Spd and Spm contents were decreased in comparison to Cd treated plants.

343 Spd (1 mM) applied exogenously did not show any significant effect on Spm contents but it  
344 was responsible for Put decrease and Spd increase at foliar level. At stem level, Spd  
345 application did not modify the contents in endogenous Spd and Spm, but cause a significant  
346 decrease of Put level ( $P < 0.05$ ). At root level, Spd application didn't have any effect on Spm

347 contents, but decrease significantly Put and Spd contents at room levels in comparison to Cd-  
348 exposed plants.

349

### 350 **Discussion and conclusions**

351

352 The results showed that the application of CdCl<sub>2</sub> (75 µM) in hydroponic conditions, causing a  
353 decline in overall growth of plants due to reduced masses of fresh and dry materials of the  
354 various organs. The percentages of water are also down. These results agree with previous  
355 work on *Brassicacea* (Larsson et al. 1998; Haag-Kerwer et al. 1999; Singh and Tewari, 2003).  
356 The foliar application of the Put or the Spd does not affect the biomass of various organs of *B.*  
357 *juncea* but produced a slight increase in weight of dry matter accompanied by a slight  
358 decrease in the percentage of water from aerial parts.

359 When the foliar spraying with one or other of PAs was associated with exposure to  
360 CdCl<sub>2</sub> (75 µM), it reduces the reduction of the mass of fresh leaf produced by the  
361 metal and cancel its effect on the mass of dry matter without changing the reduction in the  
362 percentage of water content. No changes by PAs inhibitory effects of fresh and dry masses  
363 due to cadmium on the stems and roots were observed. Only the water content of these  
364 organs, reduced by cadmium, was restored to the value of the control by the PAs.

365 The extent of pigment used to evaluate the toxicity of metals. The Cd exposure for 84 h led to  
366 a significant decrease of chlorophyll a. This effect is consistent with earlier work (Padmaja et  
367 al. 1990; Larsson et al. 1998; Groppa et al. 2007a, b). and follows, according Mysliwa-  
368 Kurdziel and Strzalka (2002), an inhibition by Cd, the biosynthesis of chlorophyll or  
369 activation of chlorophyllase (Abdel-Basset et al. 1995). Cadmium does not alter the content of  
370 chlorophyll b and carotenoids and, again, chlorophyll b appears to be less sensitive than  
371 chlorophyll a to Cd stress. At the toxicity of Cd, a slight lowering of the ratio of chl a / b,

372 agreement with the work of Baszynski et al. (1980) and Larsson et al. (1998). The application  
373 of exogenous Put or Spd does not alter the levels of pigments. However, when the foliar spray  
374 of one or other of PAs was associated with the treatment by cadmium, it helps to keep  
375 chlorophyll a and, in the case of Spd, to increase the content of chlorophylls a and b. The PAs  
376 are present in the membranes of thylacoides, the PSII and light-harvesting complex II  
377 (Kotzabatsis et al. 1993; Legocka and Zajchert, 1999) and known for their antisenescent  
378 effect (Galston and Kaur-Sawhney, 1987; Pandey et al. 2000). The fact that the only  
379 exogenous application of Put or Spd didn't produce any effect on pigments suggests that the  
380 Put and Spd foremost protect the pigments by a trapping of ROS produced under the effect of  
381 Cd, ROS that lead to the destruction pigments.

382 Indeed, many works related a protective effect of exogenous PAs against oxidative stress  
383 generated by another environmental factors such ozone (Bors et al. 1989; Navakoudis et al.  
384 2003), UV radiation (Sfichi et al. 2004), paraquat (Chang and Kao, 1997; Benavides et al.  
385 2000), acid rain (Velikova et al. 2000), high salinity (Tang and Newton, 2005), water deficit  
386 and chilling (Nayyar and Chander, 2004), and metals (Groppa et al., 2001, 2007 a, b).  
387 Furthermore, the protection of the photosynthetic apparatus by exogenous Put against ozone  
388 and UV radiation has been reported by Navakoudis et al. (2003) and Sfichi et al. (2004).

389

390 The results showed a Cd absorption by plant roots exposed for 84 h to CdCl<sub>2</sub> (75 μM)  
391 followed by transportation to parts, indicating transport by xylem as shown by salt et al.  
392 (1995) and Gong et al. (2003). The application of the Put did not alter the Cd content of  
393 various organs but a slight increase in the amount of Cd was observed in the leaves, because  
394 an increase by the diamine of the dry mass of these organs.

395 The spraying Spd changes the distribution pattern of Cd between different parts of the plant.

396 Thus, Spd induced a very strong stimulation of root uptake followed by an

397 increase of transport in the stems and this is true that the results are expressed in  $\mu\text{g Cd.g}^{-1}$   
398 D.W. or in  $\mu\text{g Cd}$  per organ. A slight decline in Cd was produced by the Spd in the leaves  
399 could explain the higher content of chlorophyll (less production of ROS and trapping by Spd).  
400 However, given the increase by the Spd of the leaf D.W., the amount of Cd accumulated was  
401 ultimately identical to that found in untreated plants to the Spd.

402 In an attempt to explain the effect of the root-level, Spd can assume a transport by the phloem  
403 of Spd to these bodies, transport demonstrated by earlier work (Béraud et al. 1992; Antognoni  
404 et al. 1998). A change in membrane permeability could arise and it is known that polyamine  
405 can influence the root absorption of a cation such a potassium (De Agazio et al. 1988). These  
406 results differ from those obtained by Hsu and Kao (2007) who report a reduction in the  
407 absorption of Cd by detached leaves of rice, after pretreatment with the Spd. This simplified  
408 system can probably explain this difference in effect observed.

409 It is now well known that abiotic stress including heavy metals cause molecular damage  
410 caused by overproduction of reactive species of oxygen (ROS). The overproduction of ROS  
411 cause lipid peroxidation which leads to the formation of degradation products such as alkanes  
412 and aldehydes (malondialdehyde) (Ferrat et al. 2003).

413 The malondialdehyde (MDA) which is a good marker for the lipoperoxidation, is significantly  
414 increased in the leaves of plants treated with cadmium. The foliar spary with either PAs can  
415 cancel (Put) or Sharply reduce (Spd) the production of MDA caused by the metaland the  
416 implementation of Put or Spd alone did not have any impact on the formation of this  
417 aldehyde. The results therefore reflect oxidative stress generated by cadmium and show a  
418 protective effect of PAs against metal stress. These data are consistent with the work of  
419 Groppa et al. (2001), Wang et al. (2007) and Zhao and Yang (2008) who report a reduction of  
420 lipid peroxidation induced by Cd and Cu by the exogenous application of Spd and Spm,

421 respectively in leaf discs of *Helianthus annuus* and in the leaves of *Nymphoides peltatum* and  
422 *Malus hupehensis*.

423 To fight against oxidative damage, plants set up systems for defense and non-enzymatic  
424 antioxidant enzyme that plays a role in regulating the levels of ROS. Among the enzymes for  
425 the destruction of radical species include peroxidase.

426 A sharp increase in the activity of the guaiacol peroxidase (GPOX) is induced in leaves by  
427 treatment with Cd and this suggests a role for this enzyme in the elimination of H<sub>2</sub>O<sub>2</sub>  
428 produced in excess. Several studies have suggested the involvement of guaiacol peroxidase in  
429 oxidative stress imposed by metals (Shah et al. 2001; Schützenbüdel and Polle, 2002;  
430 Sirvastava et al. 2004; Smeets et al., 2005; Mishra et al. 2006; Hsu and Kao, 2007; Semane et  
431 al. 2007). The foliar application of Spd or Put combined with Cd treatment can mitigate the  
432 increased activity of guaiacol peroxidase induced by the metal. Contrary to what is observed  
433 by Tang et al. (2005), the only application of exogenous Put and Spd produces no change in  
434 the peroxidase activity, we can not conclude to an induction by Spd of *de novo* synthesis of  
435 this enzyme. It is more likely that exogenous Pas (Spd, Spm) have antioxidant function in  
436 protecting the tissues of oxidative damage induced by copper and cadmium as reported by  
437 Groppa et al. (2007 a, b) and Wang et al. (2007).

438 Exposure to heavy metals is often accomplished by a synthesis of various metabolites that  
439 accumulate in concentrations of the order of millimoles, which include some amino acids  
440 (Costa and Spitz, 1997; Sharma and Dietz, 2006; Lei et al. 2007), especially proline,  
441 carbohydrates (Samarakoon and Rauser, 1979; Jha and Dubey, 2004), and polyamines  
442 (Taulavuori et al. 2005; Sharma and Dietz, 2006; Groppa et al. 2007 a, b; Groppa and  
443 Benavides, 2008). The exposure of plants to cadmium produced a doubling of the content of  
444 total free amino acids in leaves. The foliar application of Put or Spd, whether conducted with  
445 or without treatment with Cd, did not affect the amino acid content of leaves. The

446 accumulation of amino acids in response to metal stress can be explained by the degradation  
447 of certain proteins or by the *de novo* synthesis of amino acids (Berlet and Stadtman, 1997;  
448 Siedlecka and Krupa, 2002; Hsu and Kao, 2003; Pena et al. 2006; Sharma and Dietz, 2006).  
449 Among the amino acids, proline, while constituting less than 5 % of total free amino acids  
450 (Matysik et al. 2002; Lei et al. 2007), is probably one of the most common metabolites that is  
451 synthesized in response to stress. Its content, which constituted 4.40 % of total amino acids,  
452 was significantly increased in leaves after treatment with cadmium but was unaffected by the  
453 only foliar application of Put or Spd. However, When the foliar application of one or other of  
454 PAs was associated with Cd treatment, it suppressed (with Put), or reduced (with Spd) the  
455 proline accumulation induced by Cd.

456 An increase in proline under the effect of cadmium has been reported for various plants such  
457 as lupine (Costa and Spitz, 1997), rice (Shah and Dubey, 1998), radish (Chen et al. 2003) and  
458 soybean (Balestasse et al. 2005). Such an increase in proline was also observed under the  
459 effect of other metals such as Mn (Lei et al. 2007), Zn, Pb, Co, Cu (Alia et al. 1995; Schat et  
460 al. 1997; Sharmila and Pardha Saradi, 2002; Sharma and Dietz, 2006), but also with other  
461 types of stress: saline (Demiral and Turkana, 2005), water (Taulavuori et al. 2005), UV  
462 radiation (Pardha Saradi et al. 1995), thermal (Taulavuori et al. 2005). Proline is considered  
463 part of a general adaptation syndrome to unfavorable environmental conditions. Different  
464 functions are attributed to the accumulation of fluid compatible osmoregulation, chelation and  
465 detoxification of metals, protection of enzymes, regulating cytosolic acidity, stabilizing the  
466 machinery of protein synthesis and trapping of reactive oxygen species (hydroxyl radicals,  
467 singlet oxygen) (Sharmila and Pardha Saradi, 2002).

468 The results of treatment with Cd combined with foliar application of Spd suggest that the  
469 proline could be as osmoticum or trapping of ROS. Indeed, a slight decrease in the percentage  
470 of water / D.W. accompanied by an increase in proline is observed, suggesting a role of

471 osmoregulation of proline. However, the combination of two treatments (Cd and Put) did not  
472 affect the percentage of water / leaf D.W. or their content of proline (identical to that of  
473 control). In addition, the foliar spray of Spd partially reduced the production of MDA and  
474 proline induced by exposure to cadmium. This effect is consistent with a protective membrane  
475 against the attack of free radicals, agreeing with the results of Smirnoff and Cumbe (1989)  
476 which suggest that proline reacts with hydroxyl radicals to generate non-toxic hydroxyproline  
477 and those of Alia et al. (2001) and Matysik et al. (2002) who showed that proline was able to  
478 scavenge singlet oxygen. In addition, the increased content of chlorophylls could result from  
479 protection by proline of thylacoidal membranes against the attack of ROS as reported by Kavi  
480 Kishor et al. (2005).

481 In addition to changes in amino acid contents, accumulation of carbohydrates has been  
482 reported in response to different environmental stresses especially metals (Samarakoon and  
483 Rauser, 1979; Moya et al. 1993; Costa and Spitz, 1997; Dubey and Singh, 1999).

484 The results showed that the treatment with Cd produced a doubling of the soluble  
485 carbohydrate contents of leaves but the only foliar spray with Put or Spd did not change in  
486 these metabolites. However, the combination of two treatments: Cd-PA, permits with the Spd,  
487 a total abolition of the accumulation caused by Cd and only a mitigation with Put. This  
488 accumulation of carbohydrates as a result of Cd could, like what has been reported for arsenic  
489 by Jha and Dubey (2004), reflect a deterioration in enzyme activities including invertase acid,  
490 the sucrose synthase and starch phosphorylase. Costa and Spitz (1997) also observed a  
491 disruption of soluble carbohydrate under the effect of Cd resulting from the action of free  
492 radicals produced. An increase in photosynthesis appears to be unlikely to reflect the increase  
493 in soluble sugars observed, Cd producing a reduction of pigment. The foliar application of Put  
494 and Spd to plants exposed to Cd, resulting in the maintenance of pigments and therefore likely  
495 to photosynthesis. It can therefore be assumed by a railroad polyamine, free radicals produced

496 under the effect of Cd, preventing (With Spd) or mitigating (With Put) and the disruption of  
497 carbohydrate metabolism.

498 Many studies showed the involvement of polyamines in defence mechanisms during the biotic  
499 and abiotic stresses (Flores, 1991; Galston et al. 1997; Bouchereau et al. 1999; Taulavuori et  
500 al. 2005; Groppa and Benavides, 2008), and can operate according to Rhee et al. (2007) as  
501 stress molecules.

502 The exposure of plants to Cd product, in leaves, an increase of PA contents, free and  
503 conjugated forms together, the effect is especially marked for Put. These results agreed with  
504 previous results obtained under the effect of Cd and Cu for leaf tissues of wheat, oats, and  
505 bean that show an increase in free Put due to the stimulation of ADC (Weinstein et al. 1986;  
506 Groppa et al. 2003, 2007 a, b) and also that of ODC (Groppa et al. 2003, 2007 a, b).  
507 Moreover, a reduction of diamine oxidase activity by Cd and Cu, as reported by Groppa et al.  
508 (2003) and Balestrasse et al. (2005), could help to increase the content of free Put. An  
509 elevated content of conjugated Put was also reported by Weinstein et al. (1986) suggesting  
510 that conjugated PAs can be used like forms of storage and can be exchanged with the free  
511 forms. The treatment of plants by Cd product, however, reduced levels of free and conjugated  
512 PAs in the roots. The lowering of free PA forms was linked to the reduction of Put and more  
513 to the reduction of Spd and Spm contents. The sharp rise compared Put / Spd that results from  
514 1.08 to 2.00, evidenced by an inhibition of Spd synthase by Cd and therefore the content of  
515 Spd, causing therefore a reduction of the Spd synthesis. However, the lowering of root PAs by  
516 Cd, could result, not only of reducing its synthesis but also an inhibition of the transport by  
517 the phloem. Indeed, the accumulation of Pas observed in the leaves could be partly due to the  
518 inhibition of transport.

519 The decrease in root PAs combined results of a reduction of conjugated forms of the Spd and  
520 the Spm and agrees with the notion that conjugated PAs can constitute forms of storage. It can

521 not reflect this reduction of conjugated PAs by Cd inhibition of their transport by phloem,  
522 contrary to free PAs, the conjugated PAs are transported either by the phloem or xylem  
523 (Antognoni et al. 1998).

524 The foliar spray of Put, surprisingly, change a little the content of free Put of the different  
525 organs, with the exception of a slight increase in the leaves and roots. However, a slight  
526 increase of the content of conjugated Put was observed in the different parts of plant. These  
527 results suggested that conjugated Put could help to keep constant the contents of free Put as  
528 shown by Kuthanova et al. (2004). Treatment with the Put did not alter or a little the contents  
529 of Spd and Spm of different organs but reduced sharply the content of their conjugated forms  
530 in the leaves and roots. These data showed that exogenous Put did not alter the content of  
531 endogenous free Put, Spd and Spm, that may imply a lack of penetration of diamine due to its  
532 cationic properties which leads to bind wall anionic sites (Evans and Malmberg, 1989;  
533 Tiburcio et al. 1990). However, the foliar spray of Put causing in hand, an increase of  
534 conjugated Put, and in the other hand, a decrease of conjugated forms of Spd and Spm in the  
535 leaves and roots, indicated that, Put had not only been absorbed at foliar level but also  
536 transported through the phloem to the roots. Indeed, the absorption of PAs had been identified  
537 in various parts of plants (Theiss et al. 2004) without being able to conclude the existence of a  
538 high affinity general transport system as has been demonstrated for animal cells (Hubert et al.  
539 1996; Seiler et al. 1996; Sakata et al. 2000). Furthermore, we know that PAs are transported  
540 by the xylem (Friedman et al. 1985; 1986; Antognoni et al. 1998). We can suppose that free  
541 PAs homeostasis can be maintained and associated with the disruption of conjugated forms.  
542 The combination of Put and Cd led to keep the contents of total PAs at leaf levels  
543 similar to those of control leaves with a change in the distribution pattern of PAs: increase of  
544 Put and decrease of Spd and Spm. It therefore appeared that treatment with Put did not alter  
545 the effect of Cd on the foliar Put content but slightly lowered those of Spd and Spm than those

546 of control. Moreover, the combination of Put and Cd resulted in a reduction (conjugated Put)  
547 or deletion (conjugated Spd and Spm) effect of Cd on the contents of foliar conjugated forms.  
548 The stem contents of total free and conjugated PAs were respectively increased by 28% and  
549 15 % by combination treatment of Put and Cd, mainly because of an accumulation of free and  
550 conjugated Put.

551 Therefore, it appeared that the application of Put permitted some protection to leaves, reducing  
552 the stimulatory effects of Cd on their levels of free and conjugated Spd and Spm, but did not  
553 change the disruption caused by the metal alone on the levels of free and conjugated Put of  
554 leaves and those of free Put of roots.

555 The foliar spray of Spd produced an increase in total free PAs in the various organs, resulting  
556 from an increase in Spd and especially Put. This effect was particularly marked in leaf,  
557 because there was an increase of Spd, after a local spraying of this triamine but a doubling of  
558 the content of Put. Such a rise in Put has been reported by De Agazio et al. (1995) and  
559 Duhazé et al. (2002) after contribution of exogenous Spd whose degradation by polyamine  
560 oxidase generated Put; this allowed to adjust the excess of Spd. We can not exclude also the  
561 participation of an interconversion pathway of Spd in Put, catalyzed by spermidine / spermine  
562 N1-Acetyltransferase (SSAT), highlighted among animal cells (Pegg, 1986; Rhee et al. 2007).  
563 Indeed, the contribution of exogenous acetylspermidine to germination of *Arabidopsis*  
564 *thaliana*, resulting in a sharp increase in free Put (Tassoni et al. 2000) and moreover, the  
565 presence of Spd and Spm was acetylated reported by Torrigiani et al. (1993) and Del Duca et  
566 al. (1995).

567 Treatment with Spd induced a decrease in total conjugated PAs in different parts of the  
568 plant, due to lower conjugated Put in aerial parts, and for stems and roots, to decrease of  
569 conjugated forms of Spd and Spm which can only be explained by a reduction of the transferase  
570 activities to the combination with hydroxycinnamic acids (Sun et al. 1991).

571 When Spd was combined with Cd, a sharp increase of free PAs was observed in aerial parts of  
572 plants, resulting mainly from a very significant increase of free Put, significantly higher than  
573 those observed with the single application of Spd. This accumulation of Put could result both  
574 of stimulating the synthesis induced by the metal but also to the interconversion of Spd in Put.  
575 At root levels, the combination treatment Spd-Cd greatly increased the content of Put  
576 opposing the inhibitory effect produced by Cd alone and amplified the action of treatment  
577 with exogenous Spd.

578 The combination treatment of Spd with Cd, will maintain conjugated PAs sheets at values  
579 similar to those found in control plants, negating the effect of Cd.

580 These results suggested that Spd might play a protective role with respect to the toxicity of  
581 metal. It was likely that the accumulation of Put observed in different parts of the plant  
582 resulted from an interconversion of Spd to Put, which can confer a certain tolerance to the  
583 stress induced by Cd.

584

#### 585 **Acknowledgement**

586

587 We thank Dr Thierry Guinet from 'Ecole Nationale d'Enseignement Supérieure Agronomique  
588 (ENESA) (Dijon-France) for providing us the seeds of the spring line *Brassica juncea*  
589 AB79/1.

590

#### 591 **References**

592

593 Abdel-Basset, R., Issa, AA., Adam, M.S., 1995. Chlorophyllase activity: Effects of heavy  
594 metals and calcium. *Photosynth.* 31, 421-425.

595 Adriano, D.C., 1986. Trace elements in the terrestrial environment Springer-Verlag, New  
596 York, pp, 533.

597 Alia, Prasad, K.V.S.K., Pardha Saradi, P., 1995. Effect of Zinc on free-radicals and proline in  
598 *Brassica juncea* and *Cajanus cajan*. Phytochem. 39, 45-47.

599 Alia, Mohanty, P., Matysik, J., 2001. Effect of proline on the production of singlet oxygen.  
600 Amino Acids 21, 195-200.

601 Alloway, B.J., 1990. In Heavy Metals in Soils (ed Alloway, B.J.), Blackie, Glasgow

602 Antognoni, F., Fornalè, S., Grimmer, C., Komor, E., Bagni, N., 1998. Long-distance  
603 translocation of polyamines in phloem and xylem of *Ricinus communis* L. plants. Planta 204,  
604 520-527.

605 Apel, K., Hirt, H., 2004. Reactive oxygen species metabolism, oxidative stress, and signal  
606 transduction. Annu. Rev. Plant Biol. 85, 235-241.

607 Antognoni, F., Fornalè, S., Grimmer, C., Komor, E., Bagni, N., 1998. Long-distance  
608 translocation of polyamines in phloem and xylem of *Ricinus communis* L. plants. Planta 204,  
609 520-527.

610 Asada, K., 1999. The water-water cycle in chloroplasts: Scavenging of reactive oxygen and  
611 dissipation of excess photons. Annu. Rev. Plant Physiol. Plant Mol. Biol. 50, 501-639.

612 Balestrasse, K.B., Gallego S.M., Benavides M.P., Tomaro M.L., 2005. Polyamines and  
613 proline are affected by cadmium stress in nodules and roots of soybean plants. Plant Soil 270,  
614 343-353.

615 Baszynski, T., Wajda, L., Krol, M., Wolinska, D., Krupa, Z., Tukendorf, A., 1980.  
616 Photosynthetic activities of cadmium-treated tomato plants. Physiol. Plant. 48, 365-370.

617 Bates, L.S., Waldren, R.P., Teare, I.D., 1973. Rapid determination of free proline for water-  
618 stress studies. Plant Soil 39, 205-207.

619 Benavides, M.P., Gallego, S.M., Comba, M.Z., Tomaro, M.L., 2000. Relationship between  
620 polyamines and paraquat toxicity in sunflower leaf discs. *Plant Growth Regul.* 31, 215-224.

621 Béraud, J., Brun, A., Féray, A., Hourmant, A., Penot, M., 1992. Long-distance transport of  
622 <sup>14</sup>C-putrescine in potato plantlets (*Solanum tuberosum* cv. Bintje). *Biochem. Physiol.*  
623 *Pflanzen.* 188, 169-176.

624 Berlett, B.S., Stadtman, E.R., 1997. Protein oxidation in aging, disease and oxidative stress.  
625 *Am. Soc. Biochem. Biol. Mol.* 272, 20313-20316.

626 Bors, W., Langebartels, C., Michel, C., Sandermann, H., 1989. Polyamines as radical  
627 scavengers and protectants against ozone damage *Phytochem.* 28, 1589-1595.

628 Bouchereau, A., Aziz, A., Martin-Tanguy, J., 1999. Polyamines and environmental  
629 challenges: recent development *Plant Sci.* 140, 103-125.

630 Capell, T., Bassie, L., Christou, P., 2004. Modulation of polyamine biosynthetic pathway in  
631 transgenic rice confers tolerance to drought stress. *Proc. Natl. Acad. Sci. USA* 101, 9909-  
632 9914.

633 Cerning-Beroard, J., 1975. A note on sugar determination by anthrone method. *Cereal Chem.*  
634 52, 857-860.

635 Chang, C.J., Kao, C.H., 1997 Paraquat toxicity is reduced by polyamines in rice leaves. *Plant*  
636 *Growth Regul.* 22, 163-168.

637 Chaoui, A., Mazhoudi, S., Ghorbal, M.H., Ferjani, E.E., 1997. Cadmium and zinc induction  
638 of lipid peroxidation and effects on antioxidant enzyme activities in bean (*Phaseolus vulgaris*  
639 L) *Plant Sci.* 127, 139-147.

640 Chen S.L., Kao, C.H., 1995. Cd induced changes in proline level and peroxidase activity in  
641 roots of rice seedlings. *Plant Growth Regul.* 17, 67-71.

642 Chen, Y.X., He, Y.F., Luo, Y.M., Yu, Y.L., Lin, Q., Wong, M.H., 2003. Physiological  
643 mechanism of plant roots exposed to cadmium. *Chem.* 50, 789-793.

644 Chien, H.F., Lin, C.C., Wang, J.W., Chen, C.T., Kao, C.H., 2002. Changes in ammonium ion  
645 content and glutamine synthetase activity in rice leaves caused by excess cadmium are a  
646 consequence of oxidative damage. *Plant Growth Regul.* 36, 41-47.

647 Costa, G., Spitz, E., 1997. Influence of cadmium on soluble carbohydrate, free amino acids,  
648 protein content of *in vitro* cultured *Lupinus albus*. *Plant Sci.* 128, 131-140.

649 De Agazio, M., Giardina, M.C., Grego, S., 1988. Effect of exogenous putrescine, spermidine  
650 and spermine on K<sup>+</sup> uptake and H<sup>+</sup> extrusion through plasma membrane in maize root  
651 segments. *Plant Physiol.* 87, 176-178.

652 De Agazio, M., Zacchini, M., Federico, R., Grego, S., 1995. Putrescine accumulation in maize  
653 roots treated with spermidine : Evidence for spermidine to putrescine conversion. *Plant Sci.*  
654 111, 181-185.

655 Delauney, A.J., Verma, D.P.S., 1993. Proline biosynthesis and osmoregulation in plants. *Plant*  
656 *J* 4, 215-223.

657 Das, P., Samantaray, S., Rout, G.R., 1997. Studies on cadmium toxicity: a review. *Environ*  
658 *Pollut.* 98, 29-36.

659 Del Duca, S., Beninati, S., Serafini-Fracassini, D., 1995. Polyamines in chloroplasts :  
660 identification of their glutamyl and acetyl derivatives. *Biochem. J.* 305, 233-237.

661 Del Rio, L.A., Corpas, F.J., Sandalio, L.M., Palma, J.M., Gomez, M., Borroso, B., 2002.  
662 Reactive oxygen species, antioxidant systems and nitric oxide in peroxisomes. *J. Exp. Bot.*  
663 53, 1255-1272.

664 Demiral, T., Türkan, I., 2005. Comparative lipid peroxidation, antioxidant defense systems  
665 and proline content in roots of two rice cultivars differing in salt tolerance. *Environ. Exp. Bot.*  
666 53, 247-257.

667 Dixit, V., Pandey, V., Shyam, R., 2001. Differential antioxidative responses to cadmium in  
668 roots and leaves of pea (*Pisum sativum* L. cv. Azad). *J. Exp. Bot.* 52, 1101-1109.

669 Drolet, G., Dumbroff, E.B., Legge, R.L., Thompson, J.E., 1986. Radical scavenging  
670 properties of polyamines Phytochem. 25, 367-371.

671 Dubey, R.S., Singh, A.K., 1999. Salinity induces accumulation of soluble sugars and alters  
672 the activity of sugar metabolizing enzymes in rice plants. Biol. Plant. 42, 233-239.

673 Duhazé, C., Gouzerh, G., Gagneul, D., Larher, F., Bouchereau, A., 2002. The conversion of  
674 spermidine to putrescine and 1,3-diaminopropane in the roots of *Limonium tataricum*. Plant  
675 Sci. 163, 639-646.

676 Evans, P.T., Malmberg, R.L., 1989. Do polyamines have roles in plant development? Annu.  
677 Rev. Plant Physiol. Plant Mol. Biol. 40, 235-269.

678 Féray, A., Hourmant, A., Penot, M., Cann-Moisan, C., Caroff, J., 1992. Effects of interaction  
679 between polyamines and benzyladénine on betacyanine synthesis in *Amaranthus* seedlings. J.  
680 Plant Physiol. 139, 680-684.

681 Ferrat, L., Pergent-Martini, C., Roméo, M., 2003. Assessment of the use of biomarkers in  
682 aquatic plants for the evaluation of environmental quality : Application to seagrasses. Aquatic  
683 Toxicol. 65, 187-204.

684 Flores, H.E., 1991. Changes in polyamine metabolism in response to abiotic stress. *In* :  
685 Slocum, R.D., Flores, H.E. (Eds.), Biochemistry and Physiology in plants. CRC Press, pp.  
686 213-225.

687 Flores, H.E., Galston; A.W., 1982. Analysis of polyamines in higher plants by high  
688 performance liquid chromatography. Plant Physiol. 69, 701-706.

689 Foyer, C.H., Descourvies, P., Kunert, K.J., 1994. Protection against oxygen radicals: an  
690 important defence mechanism studies in transgenic plants. Plant Cell. Environ. 17, 507-523.

691 Foyer, C.H., Noctor, G., 2005. Oxidant and antioxidant signalling in plants: re-evaluation of  
692 the concept of oxidative stress in a physiological context. Plant Cell. Environ. 28, 1056-1071.

693 Franchin, C., Fossati, T., Pasquini, E., Lingua, G., Castiglione, S., Torrigiani, P., Biondi, S.,  
694 2007. High concentrations of zinc and copper induce differential polyamine responses in  
695 micropropagated white poplar (*Populus alba*). *Physiol. Plant.* 130, 77-90.

696 Friedman, R., Levin, N., Altman, A., 1986. Presence and identification of polyamines in  
697 xylem and phloem exudates of plants. *Plant Physiol.* 82, 1154-1157.

698 Gallego, S.M., Benavides, M.P., Tomaro, M.L., 1996. Effect of heavy metal ion excess on  
699 sunflower leaves: evidence for involvement of oxidative stress. *Plant Sci.* 121, 151-159.

700 Galston, A.W., Kaur-Sawhney, R., 1987. Polyamines as endogenous growth regulators. *In* :  
701 *Plant hormones and their role in plant growth and development.* Ed. Davies, P.J., Martinus  
702 Nijhoff publisher. pp. 280-295.

703 Galston, A.W., Kaur-Sawhney, R., Altabella, T., Tiburcio, A.F., 1997. Plant polyamines in  
704 reproductive activity and response to abiotic stress. *Bot. Acta* 110, 197-207.

705 Gong, J.-M., Lee, D.A., Schroeder, J.I., 2003. Long-distance root-to-shoot transport of  
706 phytochelatins and cadmium in *Arabidopsis*. *Proc. Natl. Acad. Sci.* 100, 10118-10123.

707 Groppa, M.D., Beanvides, M.P., Tomaro, M.L., 2003. Polyamine metabolism in sunflower  
708 and wheat leaf discs under cadmium and copper stress. *Plant Sci.* 164: 293-299.

709 Groppa, M.D., Tomaro, M.L., Benavides, M.P. 2001. Polyamines as protectors against  
710 cadmium or copper-induced oxidative damage in sunflower leaf discs. *Plant Sci.* 161, 481-  
711 488.

712 Groppa, M.D., Ianuzzo, M.P., Tomaro, M.L., Benavides, M.P. 2007a. Polyamine metabolism  
713 in sunflower plants under long-terme cadmium or copper stress. *Amino Acids* 32, 265-275.

714 Groppa, M.D., Tomaro, M.L. Benavides, M.P., 2007b. Polyamines and heavy metal stress :  
715 the antioxidant behavior of spermine in cadmium- and copper-treated wheat leaves.  
716 *BioMetals* 20, 185-195.

717 Groppa, M.D., Benavides, M.P., 2008. Polyamines and abiotic stress : recent advances.  
718 Amino Acids 34, 35-45.

719 Haag-Kerwer, A., Schäfer, H.J., Heiss, S., Walter, C., Rausch, T., 1999. Cadmium exposure  
720 in *Brassica juncea* causes decline in transpiration rate and leaf expansion without effect on  
721 photosynthesis. J. Exp. Bot; 50, 1827-1835.

722 Hoagland, D.R., Arnon, D.I., 1950. The water-culture for growing plants without soil. Cal.  
723 Agric. Exp. Sta Cir. 347, 1-32.

724 Hsu, Y.T., Kao, C.H., 2003. Changes in protein and amino acid contents in two cultivars of  
725 rice seedlings with different apparent tolerance to cadmium. Plant Growth Regul. 40, 147-  
726 155.

727 Hsu, Y.T., Kao, C.H., 2007. Cadmium-induced oxidative damage in rice leaves is reduced by  
728 polyamines. Plant Soil 291, 27-37.

729 Hubert, M., Pelletier, J.G., Torossian, K., Dionne, P., Gamache, I., Charest-Gaudreault, R.,  
730 Audette, M., Poulin, R., 1996. 2,2'-Dithiobis(N-ethyl-spermine-5,5-carboamide) is a high  
731 affinity, membrane-impermeant antagonist of the mammalian polyamine transport system. J.  
732 Biol. Chem. 271, 27556-27563.

733 Innelli, M.A., Pietrni, R., Fiore, L., Petrilli, L., Massacci, A., 2002. Antioxidant response to  
734 cadmium in *Phragmites australis* plants. Plant Physiol. Biochem. 40, 977-982.

735 Iturbe-Ormaetxe, I., Escuredo, P.R., Arrese-Igor, C., Becana, M., 1998. Oxidative damage in  
736 pea plants exposed to water deficit or paraquat. Plant Physiol. 116, 173-181.

737 Jha, A.B., Dubey, R.S., 2004. Carbohydrate metabolism in growing rice seedlings under  
738 arsenic toxicity. Plant Physiol. 123, 1029-1036.

739 Kasukabe, Y., He, I., Nada, K., Misawa, S., Ihara, I., Kashibana, S., 2004. Overexpression of  
740 spermidine synthase enhances tolerance to multiple environmental stresses and up-regulates

741 the expression of various stress-regulated genes in transgenic *Arabidopsis thaliana*. Plant Cell  
742 Physiol. 45, 712-722.

743 Kavi Kishor, P.B., Sangam, S., Amruth, R.N., Sri Laxmi, P., Naidu, K.R., Rao, K.R.S.S.,  
744 Sreenath Rao, Reddy, K.J., Theriappan, P., Sreenivasulu, N., 2005. Regulation of proline  
745 biosynthesis, degradation, uptake and transport in higher plants : its implications in plant  
746 growth and abiotic stress tolerance. Curr. Sci. 88, 424-438.

747 Kotzabasis, K., Fotinou, C., Roubelakis-Angelakis, K.A., Ghanotakis, D., 1993. Polyamines  
748 in the photosynthetic apparatus. Photosystem II highly resolved subcomplexes are enriched in  
749 spermine. *Photosynth. Res.* 38, 83-88.

750 Kuo, M.C., Kao, C.H. 2004. Antioxidant enzyme activities are upregulated in response to  
751 cadmium in sensitive, but not in tolerant rice (*Oryza sativa* L) seedlings. Bot. Bull. Acad. Sin.  
752 45, 291-299.

753 Kurepa, J., Smalle, J, Van Montagu, M., Inzé, D., 1998. Polyamines and paraquat toxicity in  
754 *Arabidopsis thaliana*. Plant Cell Physiol. 39, 987-992.

755 Kusano, T., Berberich, T., Tateda, C., Takahashi, Y., 2008. Polyamines : essential factors for  
756 growth and survival. Planta 228, 367-381.

757 Kuthanova, A., Gemperlova, L., Zelenkova, S., Eder, J., Machackova, I., Opatrny, Z.,  
758 Cvikrova, M., 2004. Cytological changes and alterations in polyamine contents induced by  
759 cadmium in tobacco BY-2 cells. Plant Physiol. Biochem. 42, 149-156.

760 Larsson, E.H., Bordman, J.F., Asp, H., 1998. Influence of UV-B radiation and Cd<sup>2+</sup> on  
761 chlorophyll fluorescence, growth and nutrient content in *Brassica napus*. J. Exp. Bot. 49,  
762 1031-1039.

763 Lee, T.M., 1997. Polyamine regulation of growth and chilling tolerance of rice (*Oryza sativa*  
764 L.) roots cultured *in vitro*. Plant Sci. 122, 111-117.

765 Legocka, J., Zajchert, I., 1999. Role of spermidine in the stabilization of the apoprotein of the  
766 light-harvesting chlorophyll a/b-protein complex of photosystem II during leaf senescence  
767 process. *Acta Physiol. Plant.* 21, 127-132.

768 Lei, Y.B., Korpelainen, H., Li, C.Y., 2007. Physiological and biochemical responses to high  
769 Mn concentrations in two contrasting *Populus cathayana* populations. *Chem.* 68, 686-694.

770 Lesko, K., Simon-Sarkadi, L., 2002. Effect of cadmium stress on amino acid and polyamine  
771 content of wheat seedlings. *Period. Polytech. Ser. Chem; Eng;* 46, 65-71.

772 Li, C-Z, Wang, G-X, 2004. Interaction between reactive oxygen species, ethylene and  
773 polyamines in leaves of *Glycyrrhiza inflata* seedlings under root osmotic stress. *Plant Growth*  
774 *Regul.* 42, 55-60.

775 Lichtenthaler, H.K., 1987. *Methods in enzymology*, Vol 148 Chlorophylls and carotenoids:  
776 Pigments of photosynthetic biomembranes Academic Press, Inc.

777 MacAdam, J.W., Nelson, C.J., Sharpe, R.E., 1992. Peroxidase activity in the leaf elongation  
778 zone of tall fescue. *Plant Physiol.* 99, 872-878.

779 Matysik, J., Alia, Bhalu, B., Mohanty, P., 2002. Molecular mechanisms of quenching of  
780 reactive oxygen species by proline under stress in plants. *Curr. Sci.* 82, 525-532.

781 Minotti, G., Aust, S.D., 1987. The requirement for iron (III) in the initiation of lipid  
782 peroxidation by iron (II) and hydrogen peroxide. *J. Biol. Chem.* 262, 1098-1104.

783 Minton, K.W., Tabor, H., Tabor, C.W., 1990. Paraquat toxicity is increased in *Escherichia*  
784 *coli* defective in the synthesis of polyamines *Proc. Natl. Acad. Sci. USA* 87, 2851-2855

785 Mishra, S., Srivastava, S., Tripathi, R.D., Govindarajan, R., Kuriakose, S.V., Prasad, M.N.V.,  
786 2006. Phytochelatin synthesis and response of antioxidants during cadmium stress in *Bacopa*  
787 *monnieri* L. *Plant Physiol. Biochem.* 44, 25-37.

788 Moya, J.L., Ros, R., Picazo, I., 1993. Influence of cadmium and nickel on growth, net  
789 photosynthesis and carbohydrate distribution in rice plants. *Photosynth. Res.* 36, 75-80.

790 Mysliwa-Kurdziel, B., Strzalka, K., 2002. Influence of metals on biosynthesis of  
791 photosynthetic pigments. *In* : Physiology and Biochemistry of Metal Toxicity and Tolerance  
792 in Plants, eds Prasad, M.N.V., and Strzalka, K., Kluwer Academic Publishers, Netherlands,  
793 pp. 201-227.

794 Navakoudis, E., Lutz, C., Langebartels, C., Lutz-Meindl, U., Kotzabasis, K., 2003. Ozone  
795 impact on the photosynthetic apparatus and the protective role of polyamines. *Biochim.*  
796 *Biophys. Acta* 1621, 160-169.

797 Nayyar, H., Chander, S., 2004. Protective effects of polyamines against oxidative stress  
798 induced by water and cold stress in chickpea. *J. Agron. Crop. Sci.* 190, 355-365.

799 Noctor, G., Foyer, C.H., 1998. Ascorbate and glutathione: keeping active oxygen under  
800 control. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 49, 249-279.

801 Olmos, E.O., Martinez-Solano, J.R., Piqueras, A., Hellin, E., 2003. Early steps in the  
802 oxidative burs induced by Cd in cultured tobacco cells (BY-2 line). *J. Exp. Bot.* 54, 291-301.

803 Ormrod, D.P., Beckerson, D.W., 1986. Polyamines as antiozonants for tomato *Hortsci.* 21,  
804 1070-1071.

805 Padmaja, K., Prasad, D.D.K., Prasad, A.R.K., 1990. Inhibition of chlorophyll synthesis in  
806 *Phaseolus vulgaris* L. seedlings by cadmium acetate. *Phytosynt.* 24, 399-405.

807 Pandey, S., Ranade, S.A., Nagar, P.K., Kumar, N., 2000. Role of polyamines and ethylene as  
808 modulators of plant senescence. *J. Biosci.* 25, 291-299.

809 Pardha Saradhi, P., Alia, Arora, S., Prasad, K.V.S.K., 1995. Proline accumulate in plants  
810 exposed to UV radiation and protects them against UV induced peroxidation. *Biochem.*  
811 *Biophys. Res. Commun.* 209, 1-5.

812 Pegg, A.E., 1986. Recent advances in the biochemistry of polyamines in eukaryotes.  
813 *Biochem. J.* 234, 249-262.

814

815 Pena, L.B., Tomaro, M.L., Gallego, S.M., 2006. Effect of different metals on protease activity  
816 in sunflower cotyledons. *Electronic. J. Biotechnol.* ISSN: 0717- 3458 DOI: 10.2225/Vol. 9:  
817 issue 3-18.

818 Piqueras, A., Olmos, E., Martinez-Solano, J.R.; Hellin, E., 1999. Cd induced oxidative burst  
819 in tobacco BY-2 cell: time-course, subcellulaire location and antioxidant response. *Free Rad.*  
820 *Res.* 31: S25-S31.

821 Rhee, H.J., Kim, E.-J., Lee, J.K., 2007. Physiological polyamines : Simple primordial stress  
822 molecules. *J. Cell. Mol. Med.* 11, 685-703.

823 Romero-Puertas, M.C., Rodriguez-Serrano, M., Corpas, F.J., Gomez, M., Del Rio, L.A.,  
824 Sandalio, L.M., 2004. Cadmium-induced subcellular accumulation of  $O_2^-$  And  $H_2O_2$  in pea  
825 leaves *Plant Cell Environ.* 27, 1122-1134.

826 Roy, M., Wu, R., 2001. Arginine decarboxylase transgene expression and analysis of  
827 environmental stress tolerance in transgenic rice. *Plant Sci.* 160, 869-875.

828 Sakata, K., Kashiwagi, K., Igarashi, K., 2000. Properties of a polyamine transporter regulated  
829 by antizyme. *Biochem. J.* 347, 297-303.

830 Salt, D.E., Prince, R.C., Pickering, I.J., Raskin, I., 1995. Mechanisms of cadmium mobility  
831 and accumulation in Indian mustard. *Plant Physiol.* 109, 1427-1433.

832 Samarakoon, A.B., Rauser, W.E., 1979. Carbohydrate levels and phosphoassimilate export  
833 from leaves of *Phaseolus vulgaris* exposed to excess cobalt, nickel and zinc. *Plant Physiol.*  
834 63, 1165-1169.

835 Sandalio, L.M., Dalurzo, H.C., Gomez, M., Romero-Puertas, M.C., Del Rio, L.A., 2001.  
836 Cadmium-induced changes in the growth and oxidative metabolism of pea plant *J. Exp. Bot.*  
837 52, 2115-212.

838 Schat, H., Sharma, S.S., Vooijs, R., 1997. Heavy metal induced accumulation of free proline  
839 in a metal tolerant and a non tolerant ecotype of *Silene vulgaris*. *Physiol. Plant.* 101, 477-482.

840 Schützendübel, A., Polle, A., 2002. Plant responses to biotic stresses: heavy metal-induced  
841 oxidative stress and protection by mycorrhization J. Exp. Bot. 53, 1351-1366.

842 Seiler, N., Delcros, J.G., Moulinoux, J.P., 1996. Polyamine transport in mammalian cells. An  
843 update. Int. J. Biochem. Cell. Biol. 28, 843-861.

844 Semane, B., Cuypers, A., Smeets, K., Van Belleghem, F., Horemans, N., Schat, H.,  
845 Vangronsveld, J., 2007. Cadmium responses in *Arabidopsis thaliana*: glutathione metabolism  
846 and antioxidative defence system. Physiol. Plant. 129, 519-528.

847 Sfichi, L., Ioannidis, N., Kotzabasis, K., 2004. Thylakoid-associated polyamines adjust the  
848 UV-B sensitivity of the photosynthetic apparatus by means of light-harvesting complex II  
849 changes. Photochem. Photobiol. 80, 499-506.

850 Shah, K., Dubey, R.S., 1998. Effect of cadmium on proline accumulation and ribonuclease  
851 activity in rice seedlings: role of proline as a possible enzyme protectant. Biol. Plant. 40, 121-  
852 130.

853 Shah, K., Kumar, R.G., Verma, S., Dubey, R.S., 2001. Effect of cadmium on lipid  
854 peroxidation, superoxide anion generation and activities of antioxidant enzymes in growing  
855 rice seedlings. Plant Sci. 161: 1135-1144.

856 Sharma, S.S., Dietz, K-J., 2006. The significance of amino acids and amino acid-derived  
857 molecules in plant response and adaptation to heavy metal stress. J. Exp. Bot. 57: 711-726.

858 Sharmila, P., Pardha Saradhi, P., 2002. Proline accumulation in heavy metal stressed plants:  
859 An adaptative strategy. In : Prasad, M.N.V., and Strzalka, K., (eds.), Physiology and  
860 Biochemistry of Metal Toxicity and Tolerance in plants, pp. 179-199.

861 Shaw, B.P., 1995. Effect of mercury and cadmium on the activities of antioxidative enzymes  
862 in the seedlings of *Phaseolus aureus*. Biol. Plant. 37, 587-596.

863 Siedlecka, A., Krupa, Z., 2002. Functions of enzymes in heavy metal treated plants. *In* :  
864 Physiology and Biochemistry of Metal Toxicity and Tolerance in Plants eds Prasad, M.N.V.,  
865 and Strzalka, K., Kluwer Academic Publishers, Netherlands, pp. 303-324.

866 Singh, P.K., Tewari, K.K., 2003. Cadmium toxicity induced changes in plant water relations  
867 and oxidative metabolism of *Brassica juncea* L. plants. *J. Environ. Biol.* 24, 107-112.

868 Slocum, R.D., Kaur-Sawhney, R., Galston, A.W., 1984. The physiology and biochemistry of  
869 polyamines in plants. *Arch. Biochem. Biophys.* 235, 283-303.

870 Smeets, K., Cuypers, A., Lambrechts, A., Semane, B., Hoet, P., Van Laere, A., Vangronsveld,  
871 J., 2005. Induction of oxidative stress and antioxidative mechanisms in *Phaseolus vulgaris*  
872 after Cd application. *Plant Physiol. Biochem.* 43, 437-444.

873 Smirnoff, N., Cumbes, O.J., 1989. Hydroxyl radical scavenging activity of compatible  
874 solutes. *Phytochem.* 28, 1057-1060.

875 Srivastava, S., Tripathi, R.D., Dwivedi, U.N., 2004. Synthesis of phytochelatins and  
876 modulation of antioxidants in response to cadmium stress in *Cuscuta reflexa* - an  
877 angiospermic parasite. *J. Plant Physiol.* 161, 665-675.

878 Stobart, A.K., Griffiths, W.T., Ameen-Bukhari, I., Sherwood, R.P., 1985. The effect of Cd<sup>2+</sup>  
879 on the biosynthesis of chlorophyll of barley. *Physiol. Plant.* 63, 293-298.

880 Sun, C., Liu, Y.L., Zhang, W.H., 2002. Mechanism of the effect of polyamines on the activity  
881 of tonoplast of barley roots under salt stress. *Acta Bot. Sin.* 44, 1167-1172.

882 Sun, L.Y., Monneuse, M.O., Martin-Tanguy, J., Tepfer, D., 1991. Changes in flowering and  
883 the accumulation of polyamines and hydroxycinnamic acid-polyamine conjugates in tobacco  
884 plants transformed by the *rol-A* locus from the Ri TL-DNA of *Agrobacterium rhizogenes*.  
885 *Plant Sci.* 80, 145-156.

886 Tang, C-F., Liu, Y-G., Zeng, G-M., Li, X., Xu, W-H., Li, C-F., Yuan, X-Z., 2005. Effects of  
887 exogenous spermidine on antioxidant system responses of *Typha latifolia* L. under Cd<sup>2+</sup> stress  
888 J. Integr. Plant Biol. 47, 428-434.

889 Tang, W., Newton, R.J., 2005. Polyamines reduce salt-induced oxidative damage by increasing  
890 the activities of antioxidant enzymes and decreasing lipid peroxidation in Virginia Pine. Plant  
891 Growth Regul. 46: 31-43.

892 Tassoni, A., Van Buuren, M., Franceschetti, M., Fornalè, S., Bagni, N., 2000. Polyamine  
893 content and metabolism in *Arabidopsis thaliana* and effect of spermidine on plant  
894 development. Plant Physiol. Biochem. 38, 383-393.

895 Taulavuori, K., Prasad, M.N.V., Taulavuori, E., Laine, K., 2005. Metal stress consequences  
896 on frost hardiness of plants at northern high latitudes: a review and hypothesis. Environ.  
897 Pollut. 135, 209-220.

898 Theiss, C., Bohley, P., Bisswanger, H., Voigt, J., 2004. Uptake of polyamines by the  
899 unicellular green alga *Chlamydomonas reinhardtii* and their effect on ornithine decarboxylase  
900 activity. J. Plant Physiol. 161, 3-14.

901 Thompson, J.E., Legge, R.L., Barber, R.F., 1987. The role of free radicals in senescence and  
902 wounding. New Phytol. 105, 317-344.

903 Tiburcio, A.F., Kaur-Sawhney, R., Galston A.W., 1990. Polyamine metabolism. *In* : The  
904 Biochemistry of Plants. Mifflin, B.J., and Lea, P.J., (Eds) Vol. 16, pp. 283-325. San diego,  
905 Academic Press.

906 Torrigiani, P., Altamura, M.M., Scaramagli, S., Capitani, F., Falasca, G., Bagni, N., 1993.  
907 Regulation of rhizogenesis by polyamine in tobacco thin layers. J. Plant Physiol. 142, 81-87.

908 Urano, K., Yoshiba, Y., Nanjo, T., Igarashi, Y., Seki, M., Sekiguchi, F., Yamaguchi-  
909 Shinozaki, K., Shinozaki, K., 2003. Characterization of Arabidopsis genes involved in

910 biosynthesis of polyamines in abiotic stress responses and developmental stages. *Plant Cell*  
911 *Environ.* 26, 1917-1926.

912 Velikova, V., Yordanov, I., Edreva, A., 2000. Oxidative stress and some antioxidant systems  
913 in acid rain-treated bean plants Protective role of exogenous polyamines. *Plant Sci.* 151, 59-  
914 66.

915 Wallace, H.M., Fraser, A.V., Hughes, A., 2003. A perspective of polyamine metabolism  
916 *Biochem. J.* 376, 1-14.

917 Wang, X., Shi, G.X., Ma, G.Y., Xu, Q.S., Rafeek, K., Hu, J.Z., 2004. Effects of exogenous  
918 spermidine on resistance of *Nymphoides peltatum* to Hg<sup>2+</sup> stress. *J. Plant Physiol. Mol. Biol.*  
919 30: 69-74 (in Chinese with an English abstract).

920 Wang, X., Shi, G., Xu, Q., Hu jinzhao, H., 2007. Exogenous polyamines enhance copper  
921 tolerance of *Nymphoides pelatum*. *J. Plant Physiol.* 164, 1062-1070.

922 Weinstein, L.H., Kaur-Sawhney, R., Rajam, M.V., Wettlaufer, S.H., Galston, A.W., 1986.  
923 Cadmium-induced accumulation of putrescine in oat and bean leaves. *Plant Physiol.* 82, 641-  
924 645.

925 Weiss, D.J., Shotyk, W., Schafer, J., Loyall, U., Grollmund, E., Gloor, M., 1999. Microwave  
926 digestion of ancient peat and lead determination by voltammetry. *Fresenius J. Anal. Chem.*  
927 363, 300-305.

928 Wettlaufer, S.H., Osmeloski, J., Weinstein, L.H., 1991. Response of polyamines to heavy  
929 metal stress in oat seedlings. *Environ. Toxicol. Chem.* 10, 1083-1088.

930 Xing, G.S., Zhou, G.K., Li, Z.X., Cui, K.R., 2000. Studies of polyamine metabolism and  $\beta$ -N-  
931 oxalyl-L- $\alpha$ ,  $\beta$ - diamino-propionic acid accumulation in grass pea (*Lathyrus sativus*) under  
932 water stress. *Acta Bot. Sin.* 42, 1039-1044 (in Chinese with an English abstract).

933 Ye, B., Muller, H.H., Zhang, J., Gressel, J., 1997. Constitutively elevated levels of putrescine  
934 and putrescine generating enzymes correlated with oxidant stress resistance in *Conyza*  
935 *Canadensis* and wheat. *Plant Physiol.* 115, 1443-1451.

936 Yemm, E.W., Cocking, E.C., 1955. The determination of amino acids with ninhydrin.  
937 *Analyst.* 80, 209-213.

938 Zhao, F.G., Liu, Y.L., Zhang, W.H., 2003. Relationship between polyamine metabolism in  
939 roots and salt tolerance of barley seedlings. *Acta Bot. Sin.* 45, 295-300.

940 Zhao, H., Yang, H., 2008. Exogenous polyamines alleviate the lipid peroxidation induced by  
941 cadmium chloride stress in *Malus hupehensis*. *Rehd. Sci. Hort.* 116, 442-447.

942 Zheng, Y.H., Li, S.Y., Xi, Y.F., Su, X.G., Yi, Y.B., 2000. Polyamine changes and chilling  
943 injury in cold-stored loquat fruits. *Acta Bot. Sin.* 42, 824-827 (in Chinese with English  
944 abstract).

Figure




Figure 1: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the fresh weight of organs from *Brassica juncea* plants grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 μM) for 84 h. The results, expressed as means ± S.D., were calculated from 3 independent experiments made in triplicate and conducted on, at least, 30 plants per experiment. For each organ, the different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).




Figure 2: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the Chl a, Chl b and carotenoid contents of leaves from *Brassica juncea* exposed to CdCl<sub>2</sub> (75 µM) for 84 h. The results, expressed as means ± S.D., were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).




Figure 3: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the MDA contents of leaves from *Brassica juncea* plants grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 μM) for 84 h. The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).




Figure 4: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the guaiacol activity in the leaves of *Brassica juncea* plants exposed to CdCl<sub>2</sub> (75 µM) for 84 h. The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).




Figure 5: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the amino acid contents of leaves from *Brassica juncea* plants grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 μM) for 84 h. The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).






Figure 6: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the soluble sugar content of leaves from *Brassica juncea* plants grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 µM) for 84 h. The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).








Figure 7: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the free polyamine content of leaves: (A), stems: (B) and roots: (C) from *Brassica juncea* plants exposed to CdCl<sub>2</sub> (75 μM) for 84 h. The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).






Figure 8: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the conjugated polyamine content of leaves: (A), stems: (B) and roots: (C) from *Brassica juncea* plants exposed to CdCl<sub>2</sub> (75 μM) for 84 hours. The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test)

**Table**[Click here to download Table: Tables.doc](#)

Table 1: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the dry weight of organs of *Brassica juncea* seedlings grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 μM) for 84 h.

Treatments	Organs		
	Leaves	Stems	Roots
C	96.13 ± 3.03 <sup>b</sup>	12.73 ± 0.40 <sup>e</sup>	11.69 ± 0.62 <sup>ef</sup>
C + Put	110.94 ± 2.26 <sup>a</sup>	14.56 ± 0.40 <sup>d</sup>	11.74 ± 0.50 <sup>ef</sup>
C+ Spd	118.92 ± 2.58 <sup>a</sup>	14.73 ± 0.51 <sup>d</sup>	14.12 ± 0.51 <sup>d</sup>
Cd	79.47 ± 1.99 <sup>c</sup>	11.66 ± 0.60 <sup>ef</sup>	9.69 ± 0.44 <sup>g</sup>
Cd + Put	95.95 ± 2.29 <sup>b</sup>	10.93 ± 0.34 <sup>f</sup>	8.60 ± 0.37 <sup>g</sup>
Cd + Spd	97.67 ± 2.54 <sup>b</sup>	11.05 ± 0.26 <sup>f</sup>	8.59 ± 0.18 <sup>g</sup>

The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate significant differences at  $p = 0.05$  (One-way ANOVA and LSD test).

Table 2: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on Cd accumulation by organs of *Brassica juncea* plants grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 μM) for 84 h.

Treatments	Organs		
	Leaves	Stems	Roots
C	24.19 ± 1.26 <sup>g</sup>	22.11 ± 2.07 <sup>g</sup>	21.67 ± 2.47 <sup>g</sup>
C + Put	23.24 ± 1.23 <sup>g</sup>	22.78 ± 2.30 <sup>g</sup>	22.50 ± 2.18 <sup>g</sup>
C + Spd	23.33 ± 1.18 <sup>g</sup>	22.89 ± 2.18 <sup>g</sup>	22.83 ± 3.47 <sup>g</sup>
Cd	355.24 ± 22.31 <sup>e</sup>	932.67 ± 63.57 <sup>d</sup>	4506.00 ± 307.98 <sup>b</sup>
Cd + Put	336.67 ± 26.67 <sup>e</sup>	943.33 ± 64.67 <sup>d</sup>	4771.83 ± 336.48 <sup>b</sup>
Cd + Spd	285.09 ± 15.13 <sup>f</sup>	1139.67 ± 57.94 <sup>c</sup>	8378.33 ± 459.39 <sup>a</sup>

The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).

Table 3: Effect of an exogeneous application, by spraying, of either Put (1 mM), or Spd (1 mM), on the proline content of leaves from *Brassica juncea* plants grown under hydroponic conditions and exposed to CdCl<sub>2</sub> (75 μM) for 84 h.

Treatments	Proline Contents (μmol.g <sup>-1</sup> M.F.)	%
C	0.55 ± 0.02 <sup>a</sup>	100
C + Put	0.592 ± 0.04 <sup>a</sup>	107
C + Spd	0.56 ± 0.04 <sup>a</sup>	101
Cd	1.26 ± 0.06 <sup>c</sup>	228
Cd + Put	0.63 ± 0.03 <sup>a</sup>	113
Cd + Spd	0.91 ± 0.05 <sup>b</sup>	165

The results, expressed as means ± S.D, were calculated from 3 independent experiments made in triplicate. The different letters indicate differences significant at  $p = 0.05$  (One-way ANOVA and LSD test).