

Interplay between magnetism and superconductivity in Fe-pnictides

A. B. Vorontsov, M. G. Vavilov, and A. V. Chubukov

Department of Physics, University of Wisconsin, Madison, Wisconsin 53706, USA

(Dated: December 12, 2008)

We consider phase transitions and potential co-existence of spin-density-wave (SDW) magnetic order and extended s -wave (s^+) superconducting order within a two-band itinerant model of iron pnictides, in which SDW magnetism and s^+ superconductivity are competing orders. We show that depending on parameters, the transition between these two states is either first order, or involves an intermediate phase in which the two orders co-exist. We demonstrate that such co-existence is possible when SDW order is incommensurate.

PACS numbers: 74.25.Dw, 74.25.Ha

Introduction. Iron-based pnictide superconductors – oxygen containing 1111 materials RFeAsO (R = La, Nd, Sm) and oxygen free 122 materials AFe₂As₂ (A = Ba, Sr, Ca) are at the center of experimental and theoretical activities at the moment because of high potential for applications and for the discovery of new mechanisms of superconductivity. Most of parent compounds of Fe-pnictides are magnetically ordered. Upon doping, magnetism disappears and superconductivity emerges, but the nature of this transition remains unclear. Some experiments on fluoride-doped 1111 materials indicate that the transition is first-order,¹ some show behavior more consistent with a quantum-critical point separating magnetic and superconducting (SC) states², and some show a co-existence of magnetism and SC in both classes of materials.^{3,4,5}

It is by now rather firmly established that Fe-pnictides are metals in a paramagnetic phase for all dopings, with two sets of (almost) doubly degenerate Fermi surface (FS) pockets – a hole pocket centered at (0,0) and an electron pocket centered at $\boldsymbol{\pi} = (\pi, \pi)$ in the folded Brillouin zone. To a good approximation, hole and electron FS are circular and at zero doping have nearly identical sizes.^{6,7,8,9,10,11} Like in chromium (Ref. 12), this nesting geometry is favorable to a spin-density-wave (SDW) ordering at $\boldsymbol{\pi}$ as the corresponding susceptibility logarithmically diverges at $T = 0$,¹² and a small repulsive interaction in the particle-hole channel at momentum transfer $\boldsymbol{\pi}$ already gives rise to an SDW instability at $T = T_s$. If the interaction is attractive in a SC pairing channel, then the SC pairing susceptibility also diverges logarithmically at $T = 0$ and the system becomes a SC at $T = T_c$, unless magnetism interferes.

Previous studies of an itinerant model of small electron and hole FS have found that the *same* interaction, inter-band Josephson-type pair hopping, gives rise to an SDW order and to superconductivity with extended s -wave (s^+) symmetry of the SC order parameter ($\Delta(k) \propto \cos k_x + \cos k_y$ in the folded Brillouin zone),^{13,14} leading to competition between the two orders. The full interactions in SDW and s^+ channels also involve inter-band forward scattering and intra-band Hubbard interaction, respectively, and the full interaction is larger in the SDW channel.¹³ Then at zero doping, which we as-

sociate with near-perfect nesting, the highest instability temperature is that of an SDW state. At a nonzero doping x , nesting is destroyed (either hole or electron pocket gets relatively larger), and SDW susceptibility no longer diverges. Magnetic $T_s(x)$ then goes down with doping and above a particular value of x , the first instability upon cooling is into s^+ SC state. The superconducting state is only weakly affected by doping.

The goal of the present work is to understand how the system evolves from an SDW antiferromagnet to an s^+ superconductor. For this we derive and solve a set of coupled non-linear BCS-type equations for SC and SDW order parameters. We assume that the interactions in the two channels are comparable in strength and that $T_c \lesssim T_s$, where T_s is the transition temperature at zero doping, $T_s = T_s(x=0)$.

We report two results. First, when T_s/T_c is close to unity, the system displays second order SDW and SC transitions at $T_s(x)$ and T_c , whichever is larger. The SDW state is commensurate, with momentum $\boldsymbol{\pi}$. At smaller T , the transition between SDW and SC upon changing x is first order, and there is no stable co-existence region (Fig. 1). This is similar to the phase diagram reported for LaFeAsO_{1-x}F_x in Ref.1. Second, when T_s/T_c gets larger, SDW order becomes incommensurate with momentum $\mathbf{Q} = \boldsymbol{\pi} + \mathbf{q}$ below some $T_s^* = 0.56T_s > T_c$ (an SDW_q phase.^{15,16}) We argue that in this situation SDW_q and SC states co-exist. The co-existence region is initially confined to a small region below T_c , while at lower T the system still displays a first order transition between a commensurate SDW and SC states. As the ratio T_s/T_c increases, the co-existence region extends down to lower T and eventually reaches $T = 0$ (Figs. 2,3).

An incommensurate SDW_q state at finite dopings has been studied in connection with theoretical models for chromium and its alloys by Rice¹⁵ and others,¹² and in connection to pnictides by Cvetkovic and Tesanovic.¹⁶ Such state is a magnetic analog of Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) state,¹⁷ for which doping plays the same role as a magnetic field in a paramagnetically limited superconductor.¹⁵ We found that at $T < T_c$ SDW_q phase exists *only* in combination with s^+ superconductivity.

Model and equations. We neglect double degeneracy of hole and electron states, which does not seem to be essential for the pnictides,^{14,18} and consider a weak-coupling model with two families of fermions, near one hole and one electron FSs of small and near-equal sizes. The free electron part of the Hamiltonian is $\mathcal{H}_0 = \sum_{\mathbf{k}} \left(\xi_c(\mathbf{k}) c_{\mathbf{k}\alpha}^\dagger c_{\mathbf{k}\alpha} + \xi_f(\mathbf{k}) f_{\mathbf{k}\alpha}^\dagger f_{\mathbf{k}\alpha} \right)$, where operators $c_{\mathbf{k}\alpha}$ and $f_{\mathbf{k}\alpha}$ describe fermions near $(0,0)$ and (π,π) , respectively (the momentum \mathbf{k} in $f_{\mathbf{k}\alpha}$ is a deviation from $\boldsymbol{\pi}$).

The dispersions $\xi^{f,c}(\mathbf{k}) = \pm \xi_{\mathbf{k}} + \delta$, where $\xi_{\mathbf{k}} = v_F(k - k_F)$, and δ measures a deviation from a perfect nesting and may be tuned by doping (x) or pressure. The four-fermion part contains interactions in SDW and SC channels and in mean-field (BCS) approximation reduces to the effective quadratic form $\mathcal{H} = \frac{1}{2} \sum_{\mathbf{k}\alpha\beta} \bar{\Psi}_{\mathbf{k}\alpha} \hat{\mathcal{H}} \Psi_{\mathbf{k}\beta}$, with $\bar{\Psi}_{\mathbf{k}\alpha} = (c_{\mathbf{k}\alpha}^\dagger, c_{-\mathbf{k}\alpha}, f_{\mathbf{k}+\mathbf{q}\alpha}^\dagger, f_{-\mathbf{k}-\mathbf{q}\alpha})$, (Ψ - is a conjugated column):

$$\hat{\mathcal{H}} = \left(\begin{array}{cc|cc} \xi^c(\mathbf{k}) & \Delta^c i\sigma_{\alpha\beta}^y & m_{\mathbf{q}} \sigma_{\alpha\beta}^z & 0 \\ -\Delta^{*c} i\sigma_{\alpha\beta}^y & -\xi^c(-\mathbf{k}) & 0 & -m_{\mathbf{q}} \sigma_{\alpha\beta}^z \\ \hline m_{\mathbf{q}}^* \sigma_{\alpha\beta}^z & 0 & \xi^f(\mathbf{k} + \mathbf{q}) & \Delta^f i\sigma_{\alpha\beta}^y \\ 0 & -m_{\mathbf{q}}^* \sigma_{\alpha\beta}^z & -\Delta^{*f} i\sigma_{\alpha\beta}^y & -\xi^f(-\mathbf{k} - \mathbf{q}) \end{array} \right). \quad (1)$$

The two diagonal blocks of the matrix $\hat{\mathcal{H}}$ include the s^+ SC order parameter $\Delta^c = -\Delta^f = \Delta$ for two FS pockets and two off-diagonal blocks contain SDW parameter $m_{\mathbf{q}}$; $\xi^f(\mathbf{k} + \mathbf{q}) = \xi_{\mathbf{k}} + \delta + \mathbf{v}_F \mathbf{q}$ for $q \ll k_F$. The values of $m_{\mathbf{q}}$ and Δ are determined by conventional self-consistency equations

$$m_{\mathbf{q}} = V^{sdw} \sum_{\mathbf{k}} \sigma_{\alpha\beta}^z \langle f_{\mathbf{k}+\mathbf{q}\alpha}^\dagger c_{\mathbf{k}\beta} \rangle, \quad (2a)$$

$$\Delta = V^{sc} \sum_{\mathbf{k}} (-i\sigma^y)_{\alpha\beta} \langle c_{-\mathbf{k}\alpha} c_{\mathbf{k}\beta} \rangle, \quad (2b)$$

where the sums are confined to only $(0,0)$ FS pocket, and V^{sdw} and V^{sc} are the couplings in the particle-hole SDW channel and in the particle-particle SC s^+ channel.¹³ Taken alone, V^{sc} leads to an s^+ SC state with critical temperature T_c , while V^{sdw} leads to an SDW state with transition temperature T_s at $\delta = 0$. The SDW order yields a real magnetization $M(R) \sim \cos \mathbf{Q}\mathbf{R}$ at wave vector $\mathbf{Q} = \boldsymbol{\pi} + \mathbf{q}$. The couplings V^{sdw} and V^{sc} undergo logarithmic renormalizations from fermions with energies between ϵ_F and much larger bandwidth W and flow to the same value when $W/\epsilon_F \rightarrow \infty$ (Ref. 13). For any finite W/ϵ_F , V^{sdw} is the largest of the two.

The correlators in Eqs. (2) are related to components of the Green's function $\hat{G}(\mathbf{k}, \tau)_{\alpha\beta} = -\langle T_\tau \Psi(\tau)_{\mathbf{k}\alpha} \bar{\Psi}(0)_{\mathbf{k}\beta} \rangle$, defined as the inverse of $\hat{G}^{-1} = i\epsilon_n - \hat{\mathcal{H}}$, where $\epsilon_n = \pi T(2n + 1)$ are the Matsubara frequencies. The Green's functions in Eqs. (2) can be explicitly integrated over $\xi_- = [\xi^f(\mathbf{k} + \mathbf{q}) - \xi^c(\mathbf{k})]/2 = \xi_{\mathbf{k}} + \frac{1}{2} \mathbf{v}_F \mathbf{q}$. Removing the coupling constants $2N_f |V^{sc}|$ and $2N_f |V^{sdw}|$ (N_f is the density of states at the Fermi surface per spin) and the upper cutoffs of the frequency sums in favor of the

transition temperatures $T_{c,s}$, we obtain from Eqs. (2)

$$\ln \frac{T}{T_c} = 2\pi T \sum_{n>0} \text{Re} \left\langle \frac{(E_n + i\delta_{\mathbf{q}})/E_n}{\sqrt{(E_n + i\delta_{\mathbf{q}})^2 + m_q^2}} - \frac{1}{|\epsilon_n|} \right\rangle \quad (3a)$$

and

$$\ln \frac{T}{T_s} = 2\pi T \sum_{n>0} \text{Re} \left\langle \frac{1}{\sqrt{(E_n + i\delta_{\mathbf{q}})^2 + m_q^2}} - \frac{1}{|\epsilon_n|} \right\rangle. \quad (3b)$$

where angle brackets denote Fermi surface average, $E_n = \sqrt{\epsilon_n^2 + |\Delta|^2}$, $\delta_{\mathbf{q}} = [\xi^c(\mathbf{k}) + \xi^f(\mathbf{k} + \mathbf{q})]/2 = \delta + \frac{1}{2} \mathbf{v}_F \mathbf{q}$, and T_s and T_c are solutions of the linearized equations, respectively for SDW ($\Delta = \delta = 0$) and SC ($m_q = 0$). This system of equations is solved to find all possible uniform SC and (generally) incommensurate SDW states.

Note that for $\Delta = 0$, Eq. (3b) for SDW order coincides with the gap equation for a paramagnetically limited superconductor with m_q instead of superconducting order parameter, δ instead of magnetic field H , and incommensurateness vector \mathbf{q} instead of the total momentum of a Cooper pair.^{12,15,16}

We will also need the free energy $F(\Delta, m_q)$ for these mean-field order parameters to pick out the state with the lowest F . We find $F(\Delta, m_q)$ in two complementary approaches: by differentiating with respect to interaction parameters,¹⁹ and by using Luttinger-Ward functional and extending to a finite m_q the derivation of the condensation energy for a BCS SC.^{20,21} Both methods yield,

$$\frac{\Delta F(\Delta, m_q)}{4N_f} = \frac{|\Delta|^2}{2} \ln \frac{T}{T_c} + \frac{m_q^2}{2} \ln \frac{T}{T_s} - \pi T \sum_{\epsilon_m} \text{Re} \left\langle \sqrt{(E + i\delta_q)^2 + m_q^2} - |\epsilon_m| - \frac{|\Delta|^2}{2|\epsilon_m|} - \frac{m_q^2}{2|\epsilon_m|} \right\rangle, \quad (4)$$




FIG. 1: (color online) (a) Phase diagram for $T_s/T_c = 1.5$ as a function of δ controlled by doping. Thick solid and dashed lines are second-order SDW and SC transitions at $T_s(\delta)$ and T_c , and dotted line - first order transition between commensurate SDW_0 and s^+ SC. Thin lines - physically inaccessible transitions. The pure magnetic $T_s(\delta)$ line follows the curve of ‘paramagnetically limited superconductivity’. The superconducting T_c is independent of δ in our model. Light lines denoted δ_Δ and δ_m are instability lines of SC and SDW states. (b) SDW and SC order parameters, and (c) free energy for SDW, SC and unstable mixed phases at $T/T_c = 0.1$.

where $\Delta F(\Delta, m_q) = F(\Delta, m_q) - F(0, 0)$. We solve self-consistency equations (3) for Δ and m_q at finite δ (i.e., doping) and arbitrary q and select the solution with q minimizing the free energy.

Results. The results of our calculations are shown in Figs. 1 - 3. We find that the system behavior depends on the ratio $T_s/T_c > 1$. When this ratio is close to unity, the system only develops a commensurate SDW order with $m_{q=0} = m$ (see Fig.1). The SDW and SC transitions at $T_s(\delta)$ and T_c (which is independent of δ , see Eq. (3a) with $m_q = 0$) are of second order. Below tricritical point at which $T_s(\delta) = T_c$, the transition between the states ($m \neq 0, \Delta = 0$) and ($\Delta \neq 0, m = 0$), is first order, and there is no region where m and Δ co-exist.

The first-order transition at $T = 0$ can be understood analytically. Setting $q = 0$ and subtracting Eq. (3a) from (3b), we obtain for small δ :

$$\ln \frac{T_s}{T_c} = \frac{\delta^2}{m^2 + \Delta^2}. \quad (5)$$

Setting $\Delta^2 = 0$ yields a linearized SC gap equations (3). We see that in the presence of a nonzero $m_0 = m(T = 0)$, Δ first appears at $\delta_\Delta^2 = m_0^2 \ln(T_s/T_c)$. Similarly, for nonzero Δ_0 , the SDW order grows from $\delta_m^2 = \Delta_0^2 \ln(T_s/T_c)$. Their ratio, $\delta_\Delta/\delta_m = m_0/\Delta_0 = T_s/T_c > 1$, implying that Δ nucleates in the SDW phase at a higher doping whereas m develops in SC state at lower δ (see Fig.1b). This contradicts the very fact that SDW state is stable at smaller dopings than the SC state. The solution with $\Delta, m \neq 0$ then grows in “wrong” direction




FIG. 2: (color online) Same as in Fig. 1, but for $T_s/T_c = 5$. An incommensurate SDW order appears below $T_s(\delta)$ once it becomes smaller than $T_s^* = 0.56T_s > T_c$. Below T_c , a new mixed phase appears in which incommensurate SDW_q order co-exists with SC. At small T , there is no SDW_q state without superconductivity. The transitions into the mixed state are second order from a SC state and first order from a commensurate SDW state. The free energy now shows that near $\delta/2\pi T_s = 0.2$ a mixed state has lower energy than the two pure states.

of δ , and we explicitly verified that it has a higher energy than pure states and therefore is unstable, see Fig. 1c. As both $q = 0$ SDW and SC gaps cover the entire FS, the absence of the state where the two co-exist implies that fully gapped SC and SDW orders cannot co-exist, and only one of these two states is present at a given δ . First order transition between the SDW and SC states occurs at $\delta = \delta_{cr}$, when their free energies coincide. This happens when $-m_0^2/2 + \delta_{cr}^2 = -\Delta_0^2/2$, hence

$$\delta_{cr}^2 = \frac{m_0^2}{2} \left[1 - \left(\frac{T_c}{T_s} \right)^2 \right], \quad (6)$$

which is in between the two second order instability points δ_m and δ_Δ .

Situation changes when T_s/T_c gets larger, and there appears a wider range of dopings where $T_s(\delta) > T_c$. If only commensurate magnetic order SDW_0 was possible, magnetic transition would become first-order below a certain $T_s^* = 0.56T_s$, which at large enough T_s/T_c becomes greater than T_c . In reality, the system avoids a first-order transition and extends the region of magnetic order by forming an incommensurate SDW_q state below T_s^* (see Figs. 2, 3).^{12,15,16} The transition from the normal state to SDW_q state is second order, the subsequent transition to the commensurate SDW_0 state is first order. Once m_q is developed, the actual solution is more complex and includes harmonics with multiple q (Ref. 12), but we ignore this for now.

Our main result is the discovery of a new phase below T_c , in which incommensurate SDW_q order co-exists with s^+ SC order. Physically, the key reason for appearance of




FIG. 3: (color online) Same as in previous two figures, but for intermediate $T_s/T_c = 3$. Mixed phase appears only in a tiny region near the point where T_c and $T_s(\delta, q)$ cross. At smaller T , the system still displays a first order transition (dotted line) between a commensurate SDW_0 state and a SC state.

such phase is that incommensurate SDW_q order does not gap the excitations on entire FS – the system remains a metal albeit with a modified, smaller FS.^{16,22,23} Once the Fermi surface survives, an attractive pairing interaction gives rise to SC below T_c . Alternatively speaking, for SDW_q order, some parts of the FS become inaccessible to magnetic ‘pairing’, and SC order takes advantage of this, c.f. Refs. 24,25.

At large enough T_s/T_c , the co-existence phase extends to $T = 0$ (see Fig. 2 for $T_s/T_c = 5$). In Fig. 2b we show order parameters at $T = T_c/2$. The transition from a commensurate SDW state into a mixed state is first order with both Δ and the amount of incommensuratness q jumping to finite values. The transition from a SDW_q state into a mixed state, as well as the transition from a mixed state into a pure SC state are of second order with m_q gradually vanishing. Fig. 2c shows the corresponding free energies of all states. Comparing it with Fig. 1c we see that now the SC state becomes unstable at a higher δ and the mixed state now has lower energy than pure SC or SDW states.

The behavior at somewhat smaller T_s/T_c is intermediate between those in Figs. 1 and 2. In figure 3 we show the phase diagram for intermediate $T_s/T_c = 3$. We still have $T^* > T_c$, and the mixed phase still exists, but it

now appears only as a small pocket near the point where $T_s(\delta) = T_c$. At smaller T , the system shows the same behavior as at $q = 0$, i.e., a first-order transition between commensurate SDW and SC states.

To conclude, in this paper we considered SDW and s^+ SC instabilities in a mean-field approximation for a two-band model for the pnictides. We assumed that there are ‘attractive’ interactions in both SDW and extended s-wave SC channels, leading to the nesting-driven SDW magnetism at $T_s(\delta)$, decreasing with δ controlled by doping or pressure, and to s^+ SC at T_c . At zero doping, $T_s(0) = T_s > T_c$, but T_c becomes the first instability at large dopings. The issue we consider is how SDW ordered state transforms into a superconducting s^+ state. We found that the transition is first order, with no intermediate mixed phase, if T_s and T_c are close enough so that SDW order above T_c is commensurate. At larger T_s/T_c , the system develops an incommensurate SDW_q order above T_c . This incommensurate SDW order does not gap the whole FS and allows for a co-existence of magnetism and superconductivity. We found that the mixed phase first appears in a small pocket near $T_s(\delta) = T_c$, but extends as T_s/T_c increases eventually reaching down to $T = 0$. At even larger T_s/T_c , zero temperature phases include a commensurate SDW phase, an s^+ SC phase, and an intermediate mixed phase where s^+ SC co-exists with an incommensurate SDW order, but there is no purely incommensurate SDW at $T = 0$. The transition from the mixed phase to a commensurate SDW phase is first order, and to the SC phase is second order.

Finally, in this paper, we only considered a mixed state with a uniform SC order (zero total momentum of a pair). In principle, an inhomogeneous SC state (a true FFLO state) is possible because incommensurate SDW_q order breaks the symmetry between FS points with \mathbf{k} and $-\mathbf{k}$. This, however, should not change the phase diagram as a non-uniform SC state may only exist at large enough m_q , i.e., near a first order transition into a commensurate SDW phase, possibly extending a mixed state into SDW_0 region. For large m_q , an approximation by a single q is not sufficient. A more sophisticated analysis in this region is called for.

We acknowledge with thanks useful discussions with D. Maslov, V. Cvetkovic, I. Eremin, I. Mazin, and Z. Tesanovic. This work was supported by nsf-dmr 0604406 (A.V.Ch).

¹ H. Luetkens, H. H. Klauss, M. Kraken, F. J. Litterst, T. Dellmann, R. Klingeler, C. Hess, R. Khasanov, A. Amato, C. Baines, et al., arXiv.org:0806.3533 (2008).

² J. Zhao, Q. Huang, C. de la Cruz, S. Li, J. W. Lynn, Y. Chen, M. A. Green, G. F. Chen, G. Li, Z. Li, et al., arXiv:0806.2528 (2008).

³ H. Chen, Y. Ren, Y. Qiu, W. Bao, R. H. Liu, G. Wu, T. Wu, Y. L. Xie, X. F. Wang, Q. Huang, et al., arXiv:0807.3950 (2008).

⁴ Y. Kamihara, T. Watanabe, M. Hirano, and H. Hosono, J. Amer. Chem. Soc. **130**, 3296 (2008).

⁵ R. H. Liu, G. Wu, T. Wu, D. F. Fang, H. Chen, S. Y. Li, K. Liu, Y. L. Xie, X. F. Wang, R. L. Yang, et al., Phys. Rev. Lett. **101**, 087001 (2008).

⁶ D. J. Singh and M.-H. Du, Phys. Rev. Lett. **100**, 237003 (2008).

⁷ C. Liu, G. D. Samolyuk, Y. Lee, N. Ni, T. Kondo, A. F. Santander-Syro, S. L. Bud’ko, J. L. McChesney, E. Roten-

- berg, T. Valla, et al., Phys. Rev. Lett. **101**, 177005 (2008).
- ⁸ C. Liu, T. Kondo, M. E. Tillman, R. Gordon, G. D. Samolyuk, Y. Lee, C. Martin, J. L. McChesney, S. Bud'ko, M. A. Tanatar, et al., arXiv.org:0806.2147 (2008).
- ⁹ S. E. Sebastian, J. Gillett, N. Harrison, P. H. C. Lau, D. J. Singh, C. H. Mielke, and G. G. Lonzarich, J. Phys.: Cond. Matter **20**, 422203 (2008).
- ¹⁰ T. Kondo, A. F. Santander-Syro, O. Copie, C. Liu, M. E. Tillman, E. D. Mun, J. Schmalian, S. L. Bud'ko, M. A. Tanatar, P. C. Canfield, et al., Phys. Rev. Lett. **101**, 147003 (2008).
- ¹¹ H. Ding, K. Nakayama, P. Richard, S. Souma, T. Sato, T. Takahashi, M. Neupane, Y.-M. Xu, Z.-H. Pan, A. Federov, et al., arXiv:0812.0534 (2008).
- ¹² N. Kulikov and V.V.Tugushev, Sov. Phys. Usp. **27**, 954 (1984), [Usp.Fiz.Nauk **144** 643 (1984)].
- ¹³ A. V. Chubukov, D. Efremov, and I. Eremin, Phys. Rev. B **78**, 134512 (2008).
- ¹⁴ F. Wang, H. Zhai, Y. Ran, A. Vishwanath, and D.-H. Lee, arXiv:0807.0498 (2008).
- ¹⁵ T. Rice, Phys. Rev. B **2**, 3619 (1970).
- ¹⁶ V. Cvetkovic and Z. Tesanovic, arXiv.org:0808.3742 (2008).
- ¹⁷ P.Fulde and R.Ferrell, Phys. Rev. **135** A550 (1964); A. I. Larkin and Y.N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47** 1136 (1964) [Sov. Phys. JETP **20**, 762 (1965)].
- ¹⁸ M. M. Korshunov and I. Eremin, Phys. Rev. B **78**, 140509 (2008).
- ¹⁹ K.Machida and T.Matsubara, J.Phys.Soc.Jpn. **50**, 3231 (1981).
- ²⁰ J. Bardeen and M. Stephen, Phys. Rev. **136**, A1485 (1964).
- ²¹ D. Rainer and J. W. Serene, Phys. Rev. B **13**, 4745 (1976).
- ²² I. Mazin, D. Singh, M. Johannes, and M. Du, Phys. Rev. Lett. **101**, 057003 (2008).
- ²³ I. I. Mazin and M. D.Johannes, arXiv.org:0807.3737 (2008).
- ²⁴ K.Machida, J.Phys.Soc.Jpn. **50**, 2195 (1981).
- ²⁵ M. Kato and K. Machida, J.Phys.Soc.Jpn. **56**, 2136 (1987).