# Jordan \*-homomorphisms on $C^*$ -algebras

#### M. Eshaghi Gordji

Department of Mathematics, Semnan University, P. O. Box 35195-363, Semnan, Iran e-mail: madjid.eshaghi@gmail.com

## N. Ghobadipour

Department of Mathematics, Semnan University, P. O. Box 35195-363, Semnan, Iran e-mail: ghobadipour.n@gmail.com

### Choonkil Park

Department of Mathematics, Hanyang University, Seoul 133-791, South Korea e-mail: baak@hanyang.ac.kr

**Abstract.** In this paper, we investigate Jordan \*-homomorphisms on  $C^*$ -algebras associated with the following functional inequality  $\|f(\frac{b-a}{3}) + f(\frac{a-3c}{3}) + f(\frac{3a+3c-b}{3})\| \le \|f(a)\|$ . We moreover prove the superstability and the generalized Hyers-Ulam stability of Jordan \*-homomorphisms on  $C^*$ -algebras associated with the following functional equation

$$f(\frac{b-a}{3}) + f(\frac{a-3c}{3}) + f(\frac{3a+3c-b}{3}) = f(a).$$

#### 1. Introduction

The stability of functional equations was first introduced by Ulam [21] in 1940. More precisely, he proposed the following problem: Given a group  $G_1$ , a metric group  $(G_2, d)$  and a positive number  $\epsilon$ , does there exist a  $\delta > 0$  such that if a function  $f: G_1 \longrightarrow G_2$  satisfies the inequality  $d(f(xy), f(x)f(y)) < \delta$  for all  $x, y \in G_1$ , then there exists a homomorphism  $T: G_1 \to G_2$  such that  $d(f(x), T(x)) < \epsilon$  for all  $x \in G_1$ ? As mentioned above, when this problem has a solution, we say that the homomorphisms from  $G_1$  to  $G_2$  are stable. In 1941, Hyers [5] gave a partial solution of Ulam's problem for the case of approximate additive mappings under the assumption that  $G_1$  and  $G_2$  are Banach spaces. In 1978, Th. M. Rassias [17] generalized the theorem of Hyers by considering the stability problem with unbounded Cauchy differences. This phenomenon of stability that was introduced by Th. M. Rassias [17] is called generalized Hyers-Ulam stability or Hyers-Ulam-Rassias stability. **Theorem 1.1.** Let  $f: E \longrightarrow E'$  be a mapping from a norm vector space E into a Banach space E' subject to the inequality

$$||f(x+y) - f(x) - f(y)|| \le \epsilon(||x||^p + ||y||^p)$$
(1.1)

 $<sup>^{0}</sup>$  2000 Mathematics Subject Classification: Primary 17C65; 39B82; 46L05; 47Jxx; 47B48; 39B72.

 $<sup>^{0}</sup>$  Keywords: Jordan \*-homomorphism,  $C^{\ast}\text{-algebra},$  generalized Hyers-Ulam stability, functional equation and inequality.

for all  $x, y \in E$ , where  $\epsilon$  and p are constants with  $\epsilon > 0$  and p < 1. Then there exists a unique additive mapping  $T : E \longrightarrow E'$  such that

$$||f(x) - T(x)|| \le \frac{2\epsilon}{2 - 2^p} ||x||^p$$
 (1.2)

for all  $x \in E$ . If p < 0 then inequality (1.1) holds for all  $x, y \neq 0$ , and (1.2) for  $x \neq 0$ . Also, if the function  $t \mapsto f(tx)$  from  $\mathbb{R}$  into E' is continuous for each fixed  $x \in E$ , then T is  $\mathbb{R}$ -linear.

During the last decades several stability problems of functional equations have been investigated by many mathematicians. A large list of references concerning the stability of functional equations can be found in [1]–[20].

**Definition 1.2.** Let A, B be two  $C^*$ -algebras. A  $\mathbb{C}$ -linear mapping  $f: A \to B$  is called a Jordan \*-homomorphism if

 $\begin{cases} f(a^2) = f(a)^2, \\ f(a^*) = f(a)^* \end{cases}$ 

for all  $a \in A$ .

C. Park [15] introduced and investigated Jordan \*-derivations on  $C^*$ - algebras associated with the following functional inequality

$$||f(a) + f(b) + kf(c)|| \le ||kf(\frac{a+b}{k} + c)||$$

for some integer k greater than 1 and proved the generalized Hyers-Ulam stability of Jordan \*-derivations on  $C^*$ -algebras associated with the following functional equation

$$f(\frac{a+b}{k}+c) = \frac{f(a)+f(b)}{k} + f(c)$$

for some integer k greater than 1 (see also [16, 12, 13]).

In this paper, we investigate Jordan \*-homomorphisms on  $C^*$ -algebras associated with the following functional inequality

$$||f(\frac{b-a}{3}) + f(\frac{a-3c}{3}) + f(\frac{3a+3c-b}{3})|| \le ||f(a)||.$$

We moreover prove the generalized Hyers-Ulam stability of Jordan \*-homomorphisms on  $C^*$ -algebras associated with the following functional equation

$$f(\frac{b-a}{3}) + f(\frac{a-3c}{3}) + f(\frac{3a+3c-b}{3}) = f(a).$$

# 2. Jordan \*-Homomorphisms

In this section, we investigate Jordan \*-homomorphisms on  $C^*$ - algebras. Throughout this section, assume that A, B are two  $C^*$ -algebras.

**Lemma 2.1.** Let  $f: A \to B$  be a mapping such that

$$\|f(\frac{b-a}{3}) + f(\frac{a-3c}{3}) + f(\frac{3a+3c-b}{3})\|_{B} \le \|f(a)\|_{B}$$
 (2.1)

for all  $a, b, c \in A$ . Then f is additive.

*Proof.* Letting a = b = c = 0 in (2.1), we get

$$||3f(0)||_B \le ||f(0)||_B$$
.

So f(0) = 0. Letting a = b = 0 in (2.1), we get

$$||f(-c) + f(c)||_B \le ||f(0)||_B = 0$$

for all  $c \in A$ . Hence f(-c) = -f(c) for all  $c \in A$ . Letting a = 0 and b = 6c in (2.1), we get

$$||f(2c) - 2f(c)||_B \le ||f(0)||_B = 0$$

for all  $c \in A$ . Hence

$$f(2c) = 2f(c)$$

for all  $c \in A$ . Letting a = 0 and b = 9c in (2.1), we get

$$||f(3c) - f(c) - 2f(c)||_B \le ||f(0)||_B = 0$$

for all  $c \in A$ . Hence

$$f(3c) = 3f(c)$$

for all  $c \in A$ . Letting a = 0 in (2.1), we get

$$||f(\frac{b}{3}) + f(-c) + f(c - \frac{b}{3})||_B \le ||f(0)||_B = 0$$

for all  $a, b, c \in A$ . So

$$f(\frac{b}{3}) + f(-c) + f(c - \frac{b}{3}) = 0 (2.2)$$

for all  $a, b, c \in A$ . Let  $t_1 = c - \frac{b}{3}$  and  $t_2 = \frac{b}{3}$  in (2.2). Then

$$f(t_2) - f(t_1 + t_2) + f(t_1) = 0$$

for all  $t_1, t_2 \in A$ . This means that f is additive.

Now we prove the superstability problem for Jordan \*-homomorphisms as follows.

**Theorem 2.2.** Let p < 1 and  $\theta$  be nonnegative real numbers, and let  $f : A \to B$  be a mapping such that

$$||f(\frac{b-a}{3}) + f(\frac{a-3\mu c}{3}) + \mu f(\frac{3a+3c-b}{3})||_{B} \le ||f(a)||_{B},$$
(2.2)

$$||f(a^2) - f(a)^2||_B \le \theta ||a||^{2p},$$
 (2.3)

$$||f(a^*) - f(a)^*||_B \le \theta ||a^*||^p \tag{2.4}$$

for all  $\mu \in \mathbb{T}^1 := \{\lambda \in \mathbb{C} : |\lambda| = 1\}$  and all  $a, b, c \in A$ . Then the mapping  $f : A \to B$  is a Jordan \*-homomorphism.

*Proof.* Let  $\mu=1$  in (2.2). By Lemma 2.1, the mapping  $f:A\to B$  is additive. Letting a=b=0 in (2.2), we get

$$||f(-\mu c) + \mu f(c)||_B \le ||f(0)||_B = 0$$

for all  $c \in A$  and all  $\mu \in \mathbb{T}^1$ . So

$$-f(\mu c) + \mu f(c) = f(-\mu c) + \mu f(c) = 0$$

for all  $c \in A$  and all  $\mu \in \mathbb{T}^1$ . Hence  $f(\mu c) = \mu f(c)$  for all  $c \in A$  and all  $\mu \in \mathbb{T}^1$ . By Theorem 2.1 of [14], the mapping  $f: A \to B$  is  $\mathbb{C}$ -linear. It follows from (2.3) that

$$||f(a^{2}) - f(a)^{2}||_{B} = ||\frac{1}{n^{2}}f(n^{2}a^{2}) - (\frac{1}{n}f(na))^{2}||_{B}$$

$$= \frac{1}{n^{2}}||f(n^{2}a^{2}) - f(na)^{2}||_{B}$$

$$\leq \frac{\theta}{n^{2}}n^{2p}||a||^{2p}$$

for all  $a \in A$ . Thus, since p < 1, by letting n tend to  $\infty$  in last inequality, we obtain  $f(a^2) = f(a)^2$  for all  $a \in A$ . On the other hand, it follows from (2.4) that

$$||f(a^*) - f(a)^*||_B = ||\frac{1}{n}f(na^*) - (\frac{1}{n}f(na))^*||_B$$
$$= \frac{1}{n}||f(na^*) - f(na)^*||_B$$
$$\leq \frac{\theta}{n}n^p||a^*||^p$$

for all  $a \in A$ . Thus, since p < 1, by letting n tend to  $\infty$  in last inequality, we obtain  $f(a^*) = f(a)^*$  for all  $a \in A$ . Hence the mapping  $f : A \to B$  is a Jordan \*-homomorphism.  $\square$ 

**Theorem 2.3.** Let p > 1 and  $\theta$  be a nonnegative real number, and let  $f: A \to B$  be a mapping satisfying (2.2), (2.3) and (2.4). Then the mapping  $f: A \to B$  is a Jordan \*-homomorphism.

*Proof.* The proof is similar to the proof of Theorem 2.2.

We prove the generalized Hyers-Ulam stability of Jordan \*-homomorphisms on  $C^*$ -algebras.

**Theorem 2.4.** Suppose that  $f: A \to B$  is an odd mapping for which there exists a function  $\varphi: A \times A \times A \to \mathbb{R}^+$  such that

$$\sum_{i=0}^{\infty} 3^i \varphi(\frac{a}{3^i}, \frac{b}{3^i}, \frac{c}{3^i}) < \infty, \tag{2.5}$$

$$\lim_{n \to \infty} 3^{2n} \varphi(\frac{a}{3^n}, \frac{b}{3^n}, \frac{c}{3^n}) = 0, \tag{2.6}$$

$$||f(a^*) - f(a)^*||_B \le \varphi(a, a, a),$$
 (2.7)

$$||f(\frac{\mu b - a}{3}) + f(\frac{a - 3c}{3}) + \mu f(\frac{3a - b}{3} + c) - f(a) + f(c^{2}) - f(c)^{2}||_{B} \le \varphi(a, b, c)$$
 (2.8)

for all  $a,b,c\in A$  and all  $\mu\in\mathbb{T}^1$ . Then there exists a unique Jordan \*-homomorphism  $h:A\to B$  such that

$$||h(a) - f(a)||_B \le \sum_{i=0}^{\infty} 3^i \varphi(\frac{a}{3^i}, \frac{2a}{3^i}, 0)$$
 (2.9)

for all  $a \in A$ .

*Proof.* Letting  $\mu = 1$ , b = 2a and c = 0 in (2.8), we get

$$||3f(\frac{a}{3}) - f(a)||_B \le \varphi(a, 2a, 0)$$

for all  $a \in A$ . Using the induction method, we have

$$||3^n f(\frac{a}{3^n}) - f(a)|| \le \sum_{i=0}^{n-1} 3^i \varphi(\frac{a}{3^i}, \frac{2a}{3^i}, 0)$$
(2.10)

for all  $a \in A$ . In order to show the functions  $h_n(a) = 3^n f(\frac{a}{3^n})$  form a convergent sequence, we use the Cauchy convergence criterion. Indeed, replace a by  $\frac{a}{3^m}$  and multiply by  $3^m$  in (2.10), where m is an arbitrary positive integer. We find that

$$||3^{m+n}f(\frac{a}{3^{m+n}}) - 3^m f(\frac{a}{3^m})|| \le \sum_{i=m}^{m+n-1} 3^i \varphi(\frac{a}{3^i}, \frac{2a}{3^i}, 0)$$
(2.11)

for all positive integers. Hence by the Cauchy criterion the limit  $h(a) = \lim_{n \to \infty} h_n(a)$  exists for each  $a \in A$ . By taking the limit as  $n \to \infty$  in (2.10) we see that

$$||h(a) - f(a)|| \le \sum_{i=0}^{\infty} 3^{i} \varphi(\frac{a}{3^{i}}, \frac{2a}{3^{i}}, 0)$$

and (2.9) holds for all  $a \in A$ . Let  $\mu = 1$  and c = 0 in (2.8), we get

$$||f(\frac{b-a}{3}) + f(\frac{a}{3}) + f(\frac{3a-b}{3}) - f(a)||_{B} \le \varphi(a,b,0)$$
 (2.12)

for all  $a, b, c \in A$ . Multiplying both sides (2.12) by  $3^n$  and Replacing a, b by  $\frac{a}{3^n}, \frac{b}{3^n}$ , respectively, we get

$$||3^{n}f(\frac{b-a}{3^{n+1}}) + 3^{n}f(\frac{a}{3^{n+1}}) + 3^{n}f(\frac{3a-b}{3^{n+1}}) - 3^{n}f(\frac{a}{3^{n}})||_{B} \le 3^{n}\varphi(\frac{a}{3^{n}}, \frac{b}{3^{n}}, 0)$$
(2.13)

for all  $a, b, c \in A$ . Taking the limit as  $n \to \infty$ , we obtain

$$h(\frac{b-a}{3}) + h(\frac{a}{3}) + h(\frac{3a-b}{3}) - h(a) = 0$$
 (2.14)

for all  $a, b, c \in A$ . Putting b = 2a in (2.14), we get  $3h(\frac{a}{3}) = h(a)$  for all  $a \in A$ . Replacing a by 2a in (2.14), we get

$$h(b-2a) + h(6a-b) = 2h(2a)$$
(2.15)

for all  $a, b \in A$ . Letting b = 2a in (2.15), we get h(4a) = 2h(2a) for all  $a \in A$ . So h(2a) = 2h(a) for all  $a \in A$ . Letting 3a - b = s and b - a = t in (2.14), we get

$$h(\frac{t}{3}) + h(\frac{s+t}{6}) + h(\frac{t}{3}) = h(\frac{s+t}{2})$$

for all  $s,t\in A$ . Hence h(s)+h(t)=h(s+t) for all  $s,t\in A$ . So, h is additive. Letting a=c=0 in (2.12) and using the above method, we have  $h(\mu b)=\mu h(b)$  for all  $b\in A$  and all  $\mu\in\mathbb{T}$ . Hence by the Theorem 2.1 of [14], the mapping  $f:A\to B$  is  $\mathbb{C}$ -linear.

Now, let  $h': A \to B$  be another  $\mathbb{C}$ -linear mapping satisfying (2.9). Then we have

$$||h(a) - h'(a)||_{B} = 3^{n} ||h(\frac{a}{3^{n}}) - h'(\frac{a}{3^{n}})||_{B}$$

$$\leq 3^{n} [||h(\frac{a}{3^{n}}) - f(\frac{a}{3^{n}})||_{B}$$

$$+ ||h'(\frac{a}{3^{n}}) - f(\frac{a}{3^{n}})||_{B}]$$

$$\leq 2 \sum_{i=n}^{\infty} 3^{i} \varphi(\frac{a}{3^{i}}, \frac{2a}{3^{i}}, 0) = 0$$

for all  $a \in A$ . Letting  $\mu = 1$  and a = b = 0 in (2.8), we get  $||f(c^2) - f(c)||_B \le \varphi(0, 0, c)$  for all  $c \in A$ . So

$$||h(c^2) - h(c)^2||_B = \lim_{n \to \infty} 3^{2n} ||f(\frac{c^2}{3^{2n}}) - f(\frac{c}{3^n})^2||_B \le \lim_{n \to \infty} 3^{2n} \varphi(0, 0, \frac{c}{3^n}) = 0$$

for all  $c \in A$ . Hence  $h(c^2) = h(c)^2$  for all  $c \in A$ . On the other hand we have

$$||h(c^*) - h(c)^*||_B = \lim_{n \to \infty} 3^n ||f(\frac{c^*}{3^n}) - f(\frac{c}{3^n})^*||_B \le \lim_{n \to \infty} 3^n \varphi(\frac{c}{3^n}, \frac{c}{3^n}, \frac{c}{3^n}) = 0$$

for all  $c \in A$ . Hence  $h(c^*) = h(c)^*$  for all  $c \in A$ . Hence  $h: A \to B$  is a unique Jordan \*-homomorphism.

**Corollary 2.5.** Suppose that  $f: A \to B$  is a mapping with f(0) = 0 for which there exists constant  $\theta \ge 0$  and  $p_1, p_2, p_3 > 1$  such that

$$||f(\frac{\mu b - a}{3}) + f(\frac{a - 3c}{3}) + \mu f(\frac{3a - b}{3} + c) - f(a) + f(c^{2}) - f(c)^{2}||_{B} \le \theta(||a||^{p_{1}} + ||b||^{p_{2}} + ||c||^{p_{3}}),$$

$$||f(a^*) - f(a)^*||_B \le \theta(||a||^{p_1} + ||a||^{p_2} + ||a||^{p_3})$$

for all  $a,b,c\in A$  and all  $\mu\in\mathbb{T}$ . Then there exists a unique Jordan \*-homomorphism  $h:A\to B$  such that

$$||f(a) - h(a)||_B \le \frac{\theta ||a||^{p_1}}{1 - 3^{(1-p_1)}} + \frac{\theta 2^{p_2} ||a||^{p_2}}{1 - 3^{(1-p_2)}}$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a,b,c) := \theta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3})$  in Theorem 2.4, we obtain the result.  $\square$ 

The following corollary is the Isac-Rassias stability.

Corollary 2.6. Let  $\psi : \mathbb{R}^+ \cup \{0\} \to \mathbb{R}^+ \cup \{0\}$  be a function with  $\psi(0) = 0$  such that

$$\lim_{t \to 0} \frac{\psi(t)}{t} = 0,$$

$$\psi(st) \le \psi(s)\psi(t) \ s, t \in \mathbb{R}^+$$

$$3\psi(\frac{1}{3}) < 1.$$

Suppose that  $f: A \to B$  is a mapping with f(0) = 0 satisfying (2.7) and (2.8) such that

$$\|f(\frac{\mu b-a}{3})+f(\frac{a-3c}{3})+\mu f(\frac{3a-b}{3}+c)-f(a)+f(c^2)-f(c)^2\|_B \leq \theta[\psi(\|a\|)+\psi(\|b\|)+\psi(\|c\|)]$$

for all  $a,b,c \in A$  where  $\theta > 0$  is a constant. Then there exists a unique Jordan \*-homomorphism  $h:A \to B$  such that

$$||h(a) - f(a)||_B \le \frac{\theta(1 + \psi(2))\psi(||a||)}{1 - 3\psi(\frac{1}{3})}$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a,b,c):=\theta[\psi(\|a\|)+\psi(\|b\|)+\psi(\|c\|)]$  in Theorem 2.4, we obtain the result.  $\Box$ 

**Theorem 2.7.** Suppose that  $f: A \to B$  is a mapping with f(0) = 0 for which there exists a function  $\varphi: A \times A \times A \to B$  satisfying (2.7), (2.8) and (2.8) such that

$$\sum_{i=1}^{\infty} 3^{-i} \varphi(3^i a, 3^i b, 3^i c) < \infty, \tag{2.16}$$

$$\lim_{n \to \infty} 3^{-2n} \varphi(3^i a, 3^i b, 3^i c) = 0 \tag{2.17}$$

for all  $a, b, c \in A$ . Then there exists a unique Jordan \*-homomorphism  $h: A \to B$  such that

$$||h(a) - f(a)||_B \le \sum_{i=1}^{\infty} 3^{-i} \varphi(3^i a, 3^i 2a, 0)$$
 (2.18)

for all  $a \in A$ .

*Proof.* Letting  $\mu = 1$ , b = 2a and c = 0 in (2.8), we get

$$||3f(\frac{a}{3}) - f(a)||_B \le \varphi(a, 2a, 0) \tag{2.19}$$

for all  $a \in A$ . Replacing a by 3a in (2.19), we get

$$||3^{-1}f(3a) - f(a)||_B \le 3^{-1}\varphi(3a, 2(3a), 0)$$

for all  $a \in A$ . On can apply the induction method to prove that

$$||3^{-n}f(3^na) - f(a)||_B \le \sum_{i=1}^n 3^{-i}\varphi(3^ia, 2(3^ia), 0)$$
(2.20)

for all  $a \in A$ . In order to show the functions  $h_n(a) = 3^{-n} f(3^n a)$  form a convergent sequence, we use the Cauchy convergence criterion. Indeed, replace a by  $3^m a$  and multiply by  $3^{-m}$  in (2.20), where m is an arbitrary positive integer. We find that

$$\|3^{-(m+n)}f(3^{m+n}a) - 3^{-m}f(3^ma)\| \le \sum_{i=m+1}^{m+n} 3^{-i}\varphi(3^ia, 2(3^ia), 0)$$
 (2.21)

for all positive integers. Hence by the Cauchy criterion the limit  $h(a) = \lim_{n \to \infty} h_n(a)$  exists for each  $a \in A$ . By taking the limit as  $n \to \infty$  in (2.20) we see that

$$||h(a) - f(a)|| \le \sum_{i=1}^{\infty} 3^{-i} \varphi(3^i a, 2(3^i a), 0)$$

and (2.18) holds for all  $a \in A$ .

The rest of the proof is similar to the proof of Theorem 2.4.

**Corollary 2.8.** Suppose that  $f: A \to B$  is a mapping with f(0) = 0 for which there exists constant  $\theta \ge 0$  and  $p_1, p_2, p_3 < 1$  such that

$$||f(\frac{\mu b - a}{3}) + f(\frac{a - 3c}{3}) + \mu f(\frac{3a - b}{3} + c) - f(a) + f(c^{2}) - f(c)^{2}||_{B} \le \theta(||a||^{p_{1}} + ||b||^{p_{2}} + ||c||^{p_{3}}),$$

$$||f(a^{*}) - f(a)^{*}||_{B} \le \theta(||a||^{p_{1}} + ||a||^{p_{2}} + ||a||^{p_{3}})$$

for all  $a,b,c\in A$  and all  $\mu\in\mathbb{T}$ . Then there exists a unique Jordan \*-homomorphism  $h:A\to B$  such that

$$||f(a) - h(a)||_B \le \frac{\theta ||a||^{p_1}}{3^{(1-p_1)} - 1} + \frac{\theta 2^{p_2} ||a||^{p_2}}{3^{(1-p_2)} - 1}$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a,b,c) := \theta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3})$  in Theorem 2.7, we obtain the result.  $\square$ 

The following corollary is the Isac-Rassias stability.

Corollary 2.9. Let  $\psi: \mathbb{R}^+ \cup \{0\} \to \mathbb{R}^+ \cup \{0\}$  be a function with  $\psi(0) = 0$  such that

$$\lim_{t \to 0} \frac{\psi(t)}{t} = 0,$$

$$\psi(st) \le \psi(s)\psi(t) \ s, t \in \mathbb{R}^+,$$

$$3^{-1}\psi(3) < 1.$$

Suppose that  $f: A \to B$  is a mapping with f(0) = 0 satisfying (2.7) and (2.8) such that

$$\|f(\frac{\mu b - a}{3}) + f(\frac{a - 3c}{3}) + \mu f(\frac{3a - b}{3} + c) - f(a) + f(c^2) - f(c)^2\|_B \le \theta[\psi(\|a\|) + \psi(\|b\|) + \psi(\|c\|)]$$

for all  $a,b,c \in A$  where  $\theta > 0$  is a constant. Then there exists a unique Jordan \*-homomorphism  $h:A \to B$  such that

$$||h(a) - f(a)||_B \le \frac{\theta(1 + \psi(2))\psi(||a||)}{1 - 3^{-1}\psi(3)}$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a,b,c) := \theta[\psi(\|a\|) + \psi(\|b\|) + \psi(\|c\|)]$  in Theorem 2.7, we obtain the result.

One can get easily the stability of Hyers-Ulam by the following corollary.

**Corollary 2.10.** Suppose that  $f: A \to B$  is a mapping with f(0) = 0 for which there exists constant  $\theta > 0$  such that

$$||f(\frac{\mu b - a}{3}) + f(\frac{a - 3c}{3}) + \mu f(\frac{3a - b}{3} + c) - f(a) + f(c^{2}) - f(c)^{2}||_{B} \le \theta,$$

$$||f(a^{*}) - f(a)^{*}||_{B} \le \theta$$

for all  $a,b,c\in A$  and all  $\mu\in\mathbb{T}$ . Then there exists a unique Jordan \*-homomorphism  $h:A\to B$  such that

$$||f(a) - h(a)||_B \le \theta$$

for all  $a \in A$ .

*Proof.* Letting  $p_1 = p_2 = p_3 = 0$  in Corollary 2.8, we obtain the result.

#### References

- [1] B. Baak, D. Boo and Th. M. Rassias, Generalized additive mapping in Banach modules and isomorphisms between C\*-algebras, J. Math. Anal. Appl. 314 (2006), 150–161.
- [2] P. W. Cholewa, Remarks on the stability of functional equations, Aequationes Math. 27 (1984), 76–86.
- [3] Z. Gajda, On stability of additive mappings, Internat. J. Math. Math. Sci. 14 (1991), 431–434.
- [4] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, J. Math. Anal. Appl. 184 (1994), 431–436.
- [5] D. H. Hyers, On the stability of the linear functional equation, Proc. Nat. Acad. Sci. USA 27 (1941), 222–224.
- [6] D. H. Hyers, G. Isac and Th. M. Rassias, Stability of Functional Equations in Several Variables, Birkhauser, Basel, 1998.
- [7] D. H. Hyers and Th. M. Rassias, Approximate homomorphisms, Aequationes Math. 44 (1992), 125–153.
- [8] G. Isac and Th. M. Rassias, On the Hyers-Ulam stability of  $\psi$ -additive mappings, J. Approx. Theory **72** (1993), 131–137.
- [9] G. Isac and Th. M. Rassias, Stability of ψ-additive mappings: Applications to nonlinear analysis, Internat. J. Math. Math. Sci. 19 (1996), 219–228.
- [10] K. Jun and Y. Lee, A generalization of the Hyers-Ulam-Rassias stability of Jensen's equation, J. Math. Anal. Appl. 238 (1999), 305–315.
- [11] S. Jung, Hyers-Ulam-Rassias stability of Jensen's equation and its application, Proc. Amer. Math. Soc. 126 (1998), 3137–3143.
- [12] Hark-Mahn Kim, Stability for generalized Jensen functional equations and isomorphisms between C\*-algebras, Bulletin of the Belgian Mathematical Society Simon Stevin, Volume 14, Number 1 (2007), 1-14.

- [13] Mohammad Sal Moslehian, Almost Derivations on C\*-Ternary Rings, Bulletin of the Belgian Mathematical Society - Simon Stevin, Volume 14, Number 1 (2007), 135-142.
- [14] C. Park, Homomorphisms between Poisson JC\*-algebras, Bull. Braz. Math. Soc. 36 (2005), 79–97.
- [15] C. Park, J. An and J. Cui, Jordan \*-derivations on C\*-algebras and JC\*-algebras, Abstact and Applied Analasis (in press).
- [16] Chun-Gil Park, Hyers-Ulam-Rassias stability of a generalized Euler-Lagrange type additive mapping and isomorphisms between C\*-algebras, Bulletin of the Belgian Mathematical Society Simon Stevin, Volume 13, Number 4 (2006), 619-631.
- [17] Th. M. Rassias, On the stability of the linear mapping in Banach spaces, Proc. Amer. Math. Soc. 72 (1978), 297–300.
- [18] Th. M. Rassias, Approximate homomorphisms, Aequationes Math. 44 (1992), 125–153.
- [19] Th. M. Rassias, On the stability of functional equations and a problem of Ulam, Acta Math. Appl. 62 (2000), 23–130. MR1778016 (2001j:39042)
- [20] Th. M. Rassias, On the stability of functional equations in Banach spaces, J. Math. Anal. Appl. 251 (2000), 264–284.
- [21] S. M. Ulam, Problems in Modern Mathematics, Chapter VI, Science ed. Wiley, New York, 1940.