

Generalizations of The Chung-Feller Theorem

Jun Ma^{a,*} Yeong-Nan Yeh^{b,†}

^{a,b} Institute of Mathematics, Academia Sinica, Taipei, Taiwan

Abstract

The classical Chung-Feller theorem [2] tells us that the number of Dyck paths of length n with flaws m is the n -th Catalan number and independent on m . L. Shapiro [7] found the Chung-Feller properties for the Motzkin paths. In this paper, we find the connections between these two Chung-Feller theorems. We focus on the weighted versions of three classes of lattice paths and give the generalizations of the above two theorems. We prove the Chung-Feller theorems of Dyck type for these three classes of lattice paths and the Chung-Feller theorems of Motzkin type for two of these three classes. From the obtained results, we find an interesting fact that many lattice paths have the Chung-Feller properties of both Dyck type and Motzkin type.

Keywords: Chung-Feller Theorem; Dyck path; Lattice path; Motzkin path; Pointed path

1 Introduction

Let \mathcal{S} be a subset of the set $\mathbb{Z} \times \mathbb{Z} \setminus \{(0, 0)\}$, where \mathbb{Z} is the set of the integers. We call \mathcal{S} the *step set*. Let k be an integer.

Definition 1.1 *An (\mathcal{S}, k) -lattice path is a path in $\mathbb{Z} \times \mathbb{Z}$ which:*

- (a) *is made only of steps in \mathcal{S} ;*
- (b) *starts at $(0, 0)$ and ends on the line $y = k$.*

If it is made of m steps and ends at (r, k) , we say that it is of order m and size r .

*Email address of the corresponding author: majun@math.sinica.edu.tw

†Partially supported by NSC 96-2115-M-001-005

Let \mathcal{L}^k be the set of all the (\mathcal{S}, k) -lattice path. For short we call $(\mathcal{S}, 0)$ -lattice paths \mathcal{S} -paths and write \mathcal{L}^0 as \mathcal{L} . Let w and l be two mappings from \mathcal{S} to \mathbb{R} , where \mathbb{R} is the set of the real number. We say that w and l are the weight function and the length function of \mathcal{S} respectively. For any $s \in \mathcal{S}$, $w(s)$ and $l(s)$ are called the *weight* and the *length* of the step s respectively. We can view an (\mathcal{S}, k) -lattice path L of order m as a word $s_1 s_2 \dots s_m$, where $s_j \in \mathcal{S}$. In this word, let s_j denote the j -th letter from the left. Define the weight $w(L)$ and the length $l(L)$ of the path L as

$$w(L) = \prod_{j=1}^m w(s_j) \text{ and } l(L) = \sum_{j=1}^m l(s_j).$$

Moreover, we can consider the path L as a sequence of the points

$$(0, 0) = (x_0, x_0), (x_1, y_1), (x_2, y_2), \dots, (x_m, y_m),$$

where (x_j, y_j) is the end point of the step s_j in the lattice path L for $j \geq 1$. Let $h(s_j) = h_L(s_j) = y_j$ for all $j \geq 1$. We say that $h(s_j)$ is the *height* of the step s_j in L . Define

$$\bar{l}(L) = \sum_{s_j \in L, h(s_j) \leq 0} l(s_j).$$

$\bar{l}(L)$ is called the *non-positive length* of L . A *minimum point* is a point (x_i, y_i) in the path L such that $y_i \leq y_j$ for all $j \neq i$, let $m(L) = y_i$, we call $m(L)$ the *minimum value* of L . A *absolute minimum point* is a minimum point (x_i, y_i) such that the point is the rightmost one among all the minimum points, and the index i is called the *absolute minimum position*, denoted by $mp(L)$. Finally, we define the *absolute minimum length* $ml(L)$ of the path L as

$$ml(L) = \sum_{1 \leq j \leq mp(L)} l(s_j).$$

Definition 1.2 An (\mathcal{S}, k) -nonnegative path is an (\mathcal{S}, k) -lattice path which never goes below the line $y = k$.

Let \mathcal{N}^k be the set of all the (\mathcal{S}, k) -nonnegative path. For short we call $(\mathcal{S}, 0)$ -nonnegative path \mathcal{S} -nonnegative path and write \mathcal{N}^0 as \mathcal{N} .

Now, we set $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$. In this situation, an $(\mathcal{S}, 0)$ -nonnegative path is called Dyck path as well. We may state the classical Chung-Feller Theorem [2] as follows:

The number of $(\mathcal{S}, 0)$ -lattice paths with length n and non-positive length m is equal to the number of the Dyck paths with length n and independent on m .

It is well known that the number of Dyck paths with length n is the n -th Catalan number $c_n = \frac{1}{n+1} \binom{2n}{n}$. The generating function $C(z) := \sum_{n \geq 0} c_n z^n$ satisfies the functional equation $C(z) = 1 + zC(z)^2$ and $C(z) = \frac{1 - \sqrt{1-4z}}{2z}$ explicitly.

The Chung-Feller Theorem were proved by using analytic method in [2]. T.V.Narayana [6] showed the Chung-Feller Theorem by combinatorial methods. S.P.Eu et al. [3] proved the Chung-Feller Theorem by using the Taylor expansions of generating functions and gave a refinement of this theorem. In [4], they gave a strengthening of the Chung-Feller Theorem and a weighted version for Schröder paths. Y.M. Chen [1] revisited the Chung-Feller Theorem by establishing a bijection.

Moreover, if we set $\mathcal{S} = \{(1, 1), (1, -1), (1, 0)\}$, $w(s) = 1$ and $l(s) = 1$ for $s \in \mathcal{S}$, then $(\mathcal{S}, 0)$ -nonnegative paths are the famous Motzkin paths. L. Shapiro [7] found the following Chung-Feller phenomenons for the Motzkin paths.

The number of $(\mathcal{S}, 1)$ -lattice paths with length $n + 1$ and absolute minimum length m is equal to the number of the Motzkin paths with length n and independent on m .

It is well known that the number of Motzkin paths with length n is the n -th Motzkin number m_n . The generating function $M(z) := \sum_{n \geq 0} m_n z^n$ satisfies $M(z) = 1 + zM(z) + z^2M(z)^2$ and explicitly $M(z) = \frac{1-z-\sqrt{1-2z-3z^2}}{2z^2}$. Recently, Shu-Chung Liu et al. [5] use an unify algebra approach to prove chung-Feller theorems for Dyck path and Motzkin path and develop a new method to find some combinatorial structures which have the Chung-Feller property.

The direct motivations of this paper come from the following two problems:

- (1) When $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$, is the number of \mathcal{S} -paths with length n and absolute minimum length m independent on m ?
- (2) When $\mathcal{S} = \{(1, 1), (1, -1), (1, 0)\}$, $w(s) = 1$ and $l(s) = 1$ for $s \in \mathcal{S}$, is the number of \mathcal{S} -paths with length n and non-positive length m independent on m ?

We find that the answers of these two problems are yes. In fact, in this paper, let A and B be two finite subsets of the set \mathbb{P} , where \mathbb{P} is the set of the positive integers. We consider the weighted versions of the following three classes of lattice paths.

Class 1. $\mathcal{S}_1 = \mathcal{S}_A \cup \mathcal{S}_B \cup \{(1, 1)\}$, where $\mathcal{S}_A = \{(2i - 1, -1) \mid i \in A\}$ and $\mathcal{S}_B = \{(2i, 0) \mid i \in B\}$.

For any step $s \in \mathcal{S}_1$, let

$$l(s) = \begin{cases} i & \text{if } s = (2i, 0), \\ i - 1 & \text{if } s = (2i - 1, -1), \\ 1 & \text{if } s = (1, 1), \end{cases} \quad w(s) = \begin{cases} b_i & \text{if } s = (2i, 0), \\ a_i & \text{if } s = (2i - 1, -1), \\ 1 & \text{if } s = (1, 1). \end{cases}$$

Class 2. $\mathcal{S}_2 = \mathcal{S}_A \cup \mathcal{S}_B \cup \{(1, 1)\}$, where $\mathcal{S}_A = \{(i, -1) \mid i \in A\}$ and $\mathcal{S}_B = \{(i, 0) \mid i \in B\}$.

For any step $s \in \mathcal{S}_2$, let

$$l(s) = \begin{cases} i & \text{if } s = (i, 0) \text{ and } (i, -1), \\ 1 & \text{if } s = (1, 1), \end{cases} \quad w(s) = \begin{cases} b_i & \text{if } s = (i, 0), \\ a_i & \text{if } s = (i, -1), \\ 1 & \text{if } s = (1, 1). \end{cases}$$

Class 3. $\mathcal{S}_3 = \mathcal{S}_A \cup \mathcal{S}_B \cup \{(1, 1)\}$, where $\mathcal{S}_A = \{(1, -2i + 1) \mid i \in A\}$ and $\mathcal{S}_B = \{(2i, 0) \mid i \in B\}$.

For any step $s \in \mathcal{S}_3$, let

$$l(s) = \begin{cases} i & \text{if } s = (2i, 0), \\ 0 & \text{if } s = (1, -2i + 1), \\ 1 & \text{if } s = (1, 1), \end{cases} \quad w(s) = \begin{cases} b_i & \text{if } s = (2i, 0), \\ a_i & \text{if } s = (1, -2i + 1), \\ 1 & \text{if } s = (1, 1). \end{cases}$$

First, we give the definition of the pointed lattice paths. Then we define two parameters on the pointed lattice paths: non-positive pointed length and absolute minimum pointed length. So, for any step set \mathcal{S} , we say that the pointed (\mathcal{S}, k) -lattice paths have the *Chung-Feller properties of Dyck type* (resp. *Motzkin type*) if the sum of the weights of all the pointed (\mathcal{S}, k) -lattice paths with length n and non-positive pointed length (resp. absolute minimum pointed length) m are independent on m . Finally, we prove the Chung-Feller theorem of Dyck type for the above three classes of lattice paths and the Chung-Feller theorem of Motzkin type for Classes 1 and 2. From the obtained results, we find an interesting fact that many lattice paths have the Chung-Feller properties of both Dyck type and Motzkin type. These results tell us that there are closed relations between two parameters of lattice paths: non-positive pointed length and absolute minimum pointed length.

This paper is organized as follows. In Section 2, we give the definition of the pointed lattice path and the definitions of two parameters on the pointed lattice path: non-positive pointed length and absolute minimum pointed length. In Section 3, we prove the Chung-Feller Theorem of Dyck type for Classes 1,2,3. In Section 4, we prove the Chung-Feller Theorem of Motzkin type for Classes 1,2. In Section 5, we give some interesting facts and problems.

2 The pointed path

Throughout the paper, we let the step set \mathcal{S}_i as well as the corresponding weight function w and length function l be defined as that in Classes $i = 1, 2, 3$. In this section, we will give the definition of the pointed lattice path and the definitions of two parameters of the pointed lattice path: non-positive pointed length and absolute minimum pointed length.

Let $L = s_1 s_2 \dots s_m$ be an (\mathcal{S}_i, k) -lattice path with $l(s_m) \geq 1$. Recall that L can be viewed as a sequence of the points

$$(0, 0) = (x_0, y_0), (x_1, y_1), (x_2, y_2), \dots, (x_m, y_m),$$

where (x_j, y_j) is the end point of the step s_j in the lattice path L for $j \geq 1$. For Classes 1 and 3 let $\dot{L} = [L, (x_m - 2j, 0)]$ and for Classes 2 let $\dot{L} = [L, (x_m - j, 0)]$ for some $0 \leq j \leq l(s_m) - 1$, moreover, let $p(\dot{L}_i) = j$. \dot{L} is called the *pointed path* since it denote the the path L marked by a point on the x -axis and $p(\dot{L})$ is called the *pointed length* of \dot{L}_i . Let \mathcal{M}_i^k denote the set of all the pointed (\mathcal{S}_i, k) -lattice path in which the length of the final step is no less than 1. For short we write \mathcal{M}_i^0 as \mathcal{M}_i .

Definition 2.1 Given a path $\dot{L} \in \mathcal{M}_i^1$, let $\bar{l}p(\dot{L}) = \bar{l}(L) + p(\dot{L})$. $\bar{l}p(\dot{L})$ is called the *non-positive pointed length* of \dot{L} . Let $mlp(\dot{L}) = ml(L) + p(\dot{L})$. $mlp(\dot{L})$ is called the *absolute minimum pointed length* of \dot{L} .

Example 2.2 Let $\mathcal{S} = \{(1, 1), (1, 0), (5, -1)\}$, $l((1, 1)) = 1$, $l((1, 0)) = 1$ and $l((5, -1)) = 5$. We draw a pointed \mathcal{S} -path \dot{L} of length 14 as follows, where $*$ denote the marked point.




Fig.1. A pointed \mathcal{S} -path \dot{L} of length 14

Note that the path is in Class 2. So, it is easy to see $l(L) = 14$, $p(\dot{L}) = 2$, $\bar{l}(L) = 8$, and $ml(L) = 6$. Hence, $\bar{l}p(\dot{L}) = 10$ and $mlp(\dot{L}) = 8$.

For Classes $i = 1, 2, 3$, define the generating functions

$$D_i(y, z) = \sum_{\dot{L} \in \mathcal{M}_i^1} w(L) y^{\bar{l}p(\dot{L})} z^{l(L)-1}$$

$$M_i(y, z) = \sum_{\dot{L} \in \mathcal{M}_i^1} w(L) y^{mlp(\dot{L})} z^{l(L)-1}.$$

Let $\bar{f}_{i;n,m}$ (resp. $\bar{g}_{i;n,m}$) be the sum of the weights of the pointed $(\mathcal{S}_i, 1)$ -lattice paths $\dot{L} \in \mathcal{M}_i^1$ with length $n + 1$ and non-positive pointed length (resp. absolute minimum pointed length) m for $(n, m) \neq (0, 0)$ and $\bar{f}_{i;0,0} = \bar{g}_{i;0,0} = 1$. It is easy to see

$$D_i(y, z) = \sum_{n \geq 0} \sum_{m=0}^n \bar{f}_{i;n,m} y^m z^n. \quad (1)$$

and

$$M_i(y, z) = \sum_{n \geq 0} \sum_{m=0}^n \bar{g}_{i;n,m} y^m z^n. \quad (2)$$

Let \mathcal{N}_i be the set of all the \mathcal{S}_i -nonnegative path. Define the generating function

$$F_i(z) = \sum_{L \in \mathcal{N}_i} w(L) z^{l(L)}.$$

Lemma 2.3 *For Classes 1, 2 and 3, we have*

- (1) $F_1(z) = 1 + \left(\sum_{i \in B} b_i z^i \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \right) [F_1(z)]^2$
- (2) $F_2(z) = 1 + \left(\sum_{i \in B} b_i z^i \right) F_2(z) + \left(\sum_{i \in A} a_i z^{i+1} \right) [F_2(z)]^2,$
- (3) $F_3(z) = 1 + \left(\sum_{i \in B} b_i z^i \right) F_3(z) + \left(\sum_{i \in A} a_i z^i \right) [F_3(z)]^{i+1}.$

Proof. (1) Given a path $L \in \mathcal{N}_1$ and $L \neq \emptyset$, we suppose that s is the first step of L and discuss the following two cases:

Case I. $s = (2i, 0)$ for some $i \in B$. We can decompose the path L into sR , where $R \in \mathcal{N}_1$. Note that $l(s) = i$ and $w(s) = b_i$. This provide the term $\left(\sum_{i \in B} b_i z^i \right) F_1(z)$.

Case II. $s = (1, 1)$. Let t be the first step returning to the x -axis. We can decompose the path L into $sRtQ$, where $R, Q \in \mathcal{N}_1$ and $t = (2i - 1, -1)$ for some $i \in A$. Note that $l(s) = 1$ and $w(s) = 1$, $l(t) = i - 1$ and $w(t) = a_i$. This provide the term $\left(\sum_{i \in A} a_i z^i \right) [F_1(z)]^2$.

(2) The proof is similar to that of (1).

(3) Given a path $L \in \mathcal{N}_3$ and $L \neq \emptyset$, we suppose that s is the first step of L and discuss the following two cases:

Case I. $s = (2i, 0)$ for some $i \in B$. Similar to Case I in (1), we can obtain the term $\left(\sum_{i \in B} b_i z^i \right) F_3(z)$.

Case II. $s \notin \mathcal{S}_B$. Let $t = (1, -2i + 1)$ be the first step returning to the x -axis for some $i \in A$. Let s_j be the first step $(1, 1)$ of height j for $1 \leq j \leq i$. We can decompose the path L into $s_1 R_1 s_2 R_2 \dots s_i R_i t Q$, where $R_j \in \mathcal{N}_3$ for all j and $Q \in \mathcal{N}_3$. Note that $l(s_j) = 1$ and $w(s_j) = 1$ for all j , $l(t) = 0$ and $w(t) = a_i$. This provide the term $\left(\sum_{i \in A} a_i z^i \right) [F_3(z)]^{i+1}$. ■

For Classes $i = 1, 2, 3$, let $f_{i;n}$ be the sum of the weights of the \mathcal{S}_i -nonnegative paths with length n for $n \geq 1$ and $f_{i;0} = 1$. It is easy to see

$$F_i(z) = \sum_{n \geq 0} f_{i;n} z^n. \quad (3)$$

3 The Chung-Feller property of Dyck type

In this section, we will prove the Chung-Feller Theorem of Dyck type for Classes 1, 2, 3. For $i = 1, 2, 3$, let \mathcal{P}_i be the set of all the pointed \mathcal{S}_i -nonnegative path in which the length of the final step is no less than 1 and define the generating function

$$P_i(y, z) = \sum_{\dot{L} \in \mathcal{P}_i} w(L) y^{p(\dot{L})} z^{l(L)}.$$

Lemma 3.1 *For Classes 1, 2, 3, we have*

$$\begin{aligned} (1) \quad P_1(y, z) &= 1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2 \\ (2) \quad P_2(y, z) &= 1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_2(z) + \left(\sum_{i \in A} a_i z^{i+1} \sum_{j=0}^{i-1} y^j \right) [F_2(z)]^2 \\ (3) \quad P_3(y, z) &= 1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_3(z). \end{aligned}$$

Proof. (1) Given a path $\dot{L} \in \mathcal{P}_1$ and $\dot{L} \neq \emptyset$, we suppose that s is the final step of \dot{L} and $(x, 0)$ is the final point which the path \dot{L} reach. Then $\dot{L} = [L, (x - 2j, 0)]$ for some $0 \leq j \leq l(s) - 1$. We discuss the following two cases:

Case I. $s = (2i, 0)$ for some $i \in B$. We can decompose the path L into Rs , where $R \in \mathcal{N}_1$. Note that $l(s) = i$, $w(s) = b_i$ and $j \in \{0, 1, \dots, i - 1\}$. This provide the term $\left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z)$.

Case II. $s = (2i - 1, -1)$ for some $i \in A$ and $i \geq 2$. Let t be the right-most step leaving the x -axis. We can decompose the path L into $QtRs$, where $R, Q \in \mathcal{N}_1$ and $t = (1, 1)$. Note

that $l(t) = 1$, $w(t) = 1$, $l(s) = i - 1$, $w(s) = a_i$ and $j \in \{0, 1, \dots, i - 2\}$. This provide the term $\left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2$.

(2) The proof is similar to that of (1).

(3) For any $\dot{L} \in \mathcal{P}_3$, suppose s is the final step of \dot{L} . Clearly, $s \neq (1, 1)$. Furthermore, we have $s = (2i, 0)$ for some $i \in B$ since $l(s) \geq 1$. Using the similar method as Case I in (1), we can obtain the identity as desired. ■

Now, we turn to \mathcal{S}_i -path for Classes $i = 1, 2, 3$. Let \mathcal{L}_i be the set of all the \mathcal{S}_i -paths. Define the generating functions

$$G_i(y, z) = \sum_{L \in \mathcal{L}_i} w(L) y^{\bar{l}(L)} z^{l(L)}.$$

Lemma 3.2 *For Classes 1, 2, 3, we have*

$$\begin{aligned} (1) \quad G_1(y, z) &= 1 + \left(\sum_{i \in B} b_i y^i z^i \right) G_1(y, z) + \left(\sum_{i \in A} a_i y^i z^i \right) F_1(yz) G_1(y, z) \\ &\quad + \left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z) G_1(y, z), \end{aligned}$$

$$\text{Equivalently, } G_1(y, z) = \frac{1}{1 - \sum_{i \in B} b_i y^i z^i - \left(\sum_{i \in A} a_i y^i z^i \right) F_1(yz) - \left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z)}$$

$$\begin{aligned} (2) \quad G_2(y, z) &= 1 + \left(\sum_{i \in B} b_i y^i z^i \right) G_2(y, z) + \left(\sum_{i \in A} a_i y^{i+1} z^{i+1} \right) F_2(yz) G_2(y, z) \\ &\quad + \left(\sum_{i \in A} a_i y^i z^{i+1} \right) F_2(z) G_2(y, z), \end{aligned}$$

$$\text{Equivalently, } G_2(y, z) = \frac{1}{1 - \sum_{i \in B} b_i y^i z^i - \left(\sum_{i \in A} a_i y^{i+1} z^{i+1} \right) F_2(yz) - \left(\sum_{i \in A} a_i y^i z^{i+1} \right) F_2(z)}$$

$$\begin{aligned} (3) \quad G_3(y, z) &= 1 + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^i y^{i-j} [F_3(yz)]^{i-j} [F_3(z)]^j \right) G_3(y, z) \\ &\quad + \left(\sum_{i \in B} b_i y^i z^i \right) G_3(y, z). \end{aligned}$$

$$\text{Equivalently, } G_3(y, z) = \frac{1}{1 - \sum_{i \in B} b_i y^i z^i - \sum_{i \in A} a_i z^i \sum_{j=0}^i y^{i-j} [F_3(yz)]^{i-j} [F_3(z)]^j}$$

Proof. (1) Given a path $L \in \mathcal{L}_1$ and $L \neq \emptyset$, we suppose that s is the first step of L . We discuss the following three cases:

Case I. $s = (2i, 0)$ for some $i \in B$. We can decompose the path L into sR , where $R \in \mathcal{L}_1$. Note that $l(s) = i$, $w(s) = b_i$ and $h(s) = 0$. This provide the term $\left(\sum_{i \in B} b_i y^i z^i \right) G_1(y, z)$.

Case II. $s = (2i - 1, -1)$ for some $i \in A$. Let $t = (1, 1)$ be the left-most step returning to the x -axis. We can decompose the path L into $s\overline{Q}tR$, where $R \in \mathcal{L}_1$, and if we view \overline{Q} as a word $s_1 s_2 \dots s_r$ of the steps in \mathcal{S}_1 , then $Q = s_r s_{r-1} \dots s_1 \in \mathcal{N}_1$. Note that $l(t) = 1$, $w(t) = 1$, $l(s) = i - 1$, $w(s) = a_i$, $h(s) = -1$ and $h(t) = 0$. This provide the term $\left(\sum_{i \in A} a_i y^i z^i \right) F_1(yz) G_1(y, z)$.

Case III. $s = (1, 1)$. Let $t = (2i - 1, -1)$ be the first step returning to the x -axis. We can decompose the path L into $sQtR$, where $R \in \mathcal{L}_1$ and $Q \in \mathcal{N}_1$. Note that $l(t) = i - 1$, $w(t) = a_i$, $l(s) = 1$, $w(s) = 1$, $h(s) = 1$ and $h(t) = 0$. This provide the term $\left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z) G_1(y, z)$.

(2) The proof is similar to that of (1).

(3) Given a path $L \in \mathcal{L}_3$ and $L \neq \emptyset$, we suppose that s is the first step of L . We discuss the following two cases:

Case I. $s = (2i, 0)$ for some $i \in B$. Similar to Case I in (1), we can obtain the term $\left(\sum_{i \in B} b_i y^i z^i \right) G_3(y, z)$.

Case II. $s \notin \mathcal{S}_B$. Let $t = (1, -2i + 1)$ be the left-most step which passes the x -axis and has height $-i + j$ for some $i \in A$ and $0 \leq j \leq i$. Furthermore, let s_r be the first step $(1, 1)$ with height r for any $1 \leq r \leq j$. Let u_r be the first step $(1, 1)$ at the right of t with height $-r + 1$ for any $1 \leq r \leq i - j$. So, we can decompose the path L into $s_1 R_1 s_2 R_2 \dots s_j R_j t \overline{Q}_1 u_{i-j} \overline{Q}_2 u_{i-j-1} \dots \overline{Q}_{i-j} u_1 T$, where $R_r \in \mathcal{N}_3$ for all $1 \leq r \leq j$, $T \in \mathcal{L}_3$, and if we view \overline{Q}_r as a word $s'_{r,1} s'_{r,2} \dots s'_{r,k}$ of the step in \mathcal{S}_3 , then $Q_r = s'_{r,k} s'_{r,k-1} \dots s'_{r,1} \in \mathcal{N}_3$ for all $1 \leq r \leq i - j$. Note that $l(s_r) = 1$, $w(s_r) = 1$, $h(s_r) \geq 1$ for all $1 \leq r \leq j$, $l(t) = 0$, $w(t) = a_i$, $h(t) \leq 0$, $l(u_r) = 1$, $w(u_r) = 1$ and $h(u_r) \leq 0$ for all $1 \leq r \leq i - j$. This provide the term $\left(\sum_{i \in A} a_i z^i \sum_{j=0}^i y^{i-j} [F_3(yz)]^{i-j} [F_3(z)]^j \right) G_3(y, z)$. ■

Recall that \mathcal{M}_i^1 is the set of all the pointed $(\mathcal{S}_i, 1)$ -path in which the length of the final step is no less than 1 and $D_i(y, z) = \sum_{\dot{L} \in \mathcal{M}_i^1} w(L) y^{\overline{lp}(\dot{L})} z^{l(L)-1}$ for $i = 1, 2, 3$.

Lemma 3.3 For Classes $i = 1, 2, 3$, we have

$$D_i(y, z) = G_i(y, z)P_i(y, z).$$

Proof. For any $i = 1, 2, 3$, let $\dot{L} \in \mathcal{M}_i^1$. Let s be the right-most step $(1, 1)$ leaving x -axis and reaching the line $y = 1$. We can decompose the path \dot{L} into $Rs\dot{Q}$, where $R \in \mathcal{L}_i$ and $\dot{Q} \in \mathcal{P}_i$. Hence, $D_i(y, z) = G_i(y, z)P_i(y, z)$. \blacksquare

Now, we are in a position to prove the Chung-Feller theorem of Dyck type for Classes 1, 2, 3.

Theorem 3.4 For Classes $i = 1, 2, 3$, let $\bar{f}_{i,n,m}$ be the sum of the weights of the pointed $(\mathcal{S}_i, 1)$ -lattice paths which

- (a.) have length $n + 1$,
- (b.) have non-positive pointed length m ,
- (c.) have the length of the final step no less than 1.

Let $f_{i,n}$ be the sum of the weights of the \mathcal{S}_i -nonnegative paths with length n . Then $\bar{f}_{i,n,m}$ has the Chung-Feller property of Dyck type, i.e., $\bar{f}_{i,n,m} = f_{i,n}$.

Proof. First, we consider Class 1. By Lemmas 3.1, 3.2 and 3.3, we have

$$\begin{aligned} D_1(y, z) &= G_1(y, z)P_1(y, z) \\ &= \frac{1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2}{1 - \sum_{i \in B} b_i y^i z^i - \left(\sum_{i \in A} a_i y^i z^i \right) F_1(yz) - \left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z)} \\ &= \frac{\left[1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2 \right] F_1(yz)}{\left[1 - \sum_{i \in B} b_i y^i z^i - \left(\sum_{i \in A} a_i y^i z^i \right) F_1(yz) - \left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z) \right] F_1(yz)} \\ &= \frac{\left[1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2 \right] F_1(yz)}{1 - \left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z) F_1(yz)} \end{aligned}$$

since Lemma 2.3 tells us $F_1(yz) - \left(\sum_{i \in B} b_i y^i z^i\right) F_1(yz) - \left(\sum_{i \in A} a_i y^i z^i\right) [F_1(yz)]^2 = 1$. Note that

$$\begin{aligned}
& \left[1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2 \right] F_1(yz) \\
&= \left[1 + \left(\sum_{i \in B} b_i z^i \frac{y^i - 1}{y - 1} \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \frac{y^{i-1} - 1}{y - 1} \right) [F_1(z)]^2 \right] F_1(yz) \\
&= \frac{1}{y - 1} \left[y - 1 + \left(\sum_{i \in B} b_i z^i (y^i - 1) \right) F_1(z) + \left(\sum_{i \in A} a_i z^i (y^{i-1} - 1) \right) [F_1(z)]^2 \right] F_1(yz) \\
&= \frac{1}{y - 1} \left[y F_1(yz) + \left(\sum_{i \in B} b_i z^i y^i \right) F_1(z) F_1(yz) + \left(\sum_{i \in A} a_i z^i y^{i-1} \right) [F_1(z)]^2 F_1(yz) - F_1(z) F_1(yz) \right]
\end{aligned}$$

Furthermore, since $\left(\sum_{i \in B} b_i z^i y^i\right) F_1(yz) = F_1(yz) - \left(\sum_{i \in A} a_i y^i z^i\right) [F_1(yz)]^2 - 1$, we get

$$\begin{aligned}
& \left[1 + \left(\sum_{i \in B} b_i z^i \sum_{j=0}^{i-1} y^j \right) F_1(z) + \left(\sum_{i \in A} a_i z^i \sum_{j=0}^{i-2} y^j \right) [F_1(z)]^2 \right] F_1(yz) \\
&= \frac{1}{y - 1} \left[y F_1(yz) - \left(\sum_{i \in A} a_i y^i z^i \right) [F_1(yz)]^2 F_1(z) - F_1(z) + \left(\sum_{i \in A} a_i z^i y^{i-1} \right) [F_1(z)]^2 F_1(yz) \right] \\
&= \frac{[y F_1(yz) - F_1(z)] \left[1 - \left(\sum_{i \in A} a_i y^{i-1} z^i \right) F_1(z) F_1(yz) \right]}{y - 1}.
\end{aligned}$$

Hence,

$$\begin{aligned}
D_1(y, z) &= \frac{y F_1(yz) - F_1(z)}{y - 1} \\
&= \frac{y \sum_{n \geq 0} f_{1;n} y^n z^n - \sum_{n \geq 0} f_{1;n} z^n}{y - 1} \\
&= \sum_{n \geq 0} f_{1;n} z^n \frac{y^{n+1} - 1}{y - 1} \\
&= \sum_{n \geq 0} f_{1;n} z^n \sum_{m=0}^n y^m \\
&= \sum_{n \geq 0} \sum_{m=0}^n f_{1;n} y^m z^n.
\end{aligned}$$

This implies $\bar{f}_{1;n,m} = f_{1;n}$ for all $0 \leq m \leq n$. Similarly, we can prove the theorems for Classes $i = 2, 3$. ■

Corollary 3.5 (Chung-Feller.) Let $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$. Then the number of the $(\mathcal{S}, 0)$ -lattice path with length n and non-positive length m is the n -th Catalan number.

Proof. For any a pointed $(\mathcal{S}, 1)$ -lattice path, we suppose that s is the final step in this path. Then $s = (1, 1)$ since $l((1, -1)) = 0 < 1$. If we delete the final step of this path and erase the marked point, we will obtain an $(\mathcal{S}, 0)$ -lattice path with length n and non-positive length m . By Theorem 3.4, the number of the pointed $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and non-positive pointed length m in which the length of the final step is no less than 1 is equal to the number of the \mathcal{S} -nonnegative paths with length n . By Lemma 2.3, we have $F_1(z) = 1 + z[F_1(z)]^2$ since $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$. Hence, the number of the \mathcal{S}_1 -nonnegative paths with length n is the n -th Catalan number. This complete the proof. ■

Corollary 3.6 Let $\mathcal{S} = \{(1, 1), (1, -1), (1, 0)\}$, $w(s) = 1$ and $l(s) = 1$ for any $s \in \mathcal{S}$. Then the number of the $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and non-positive length m is the n -th Motzkin number.

Proof. For any a pointed $(\mathcal{S}, 1)$ -lattice path, we suppose that the final point in this path is $(x, 1)$. Then the marked point must be $(x, 0)$ since $l(s) = 1$ for any $s \in \mathcal{S}$. So, we can erase the marked point. By Theorem 3.4, the number of the pointed $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and non-positive pointed length m is equal to the number of the \mathcal{S} -nonnegative paths with length n . By Lemma 2.3, we have $F_2(z) = 1 + zF_2(z) + z^2[F_2(z)]^2$ since $\mathcal{S} = \{(1, 1), (1, -1), (1, 0)\}$, $w(s) = 1$ and $l(s) = 1$ for any $s \in \mathcal{S}$. Hence, the number of the \mathcal{S} -nonnegative paths with length n is the n -th Motzkin number. This complete the proof. ■




Fig.2. An example of Corollary 3.6, where $n = 4$

Corollary 3.7 Let $\mathcal{S} = \{(1, 1), (5, -1), (1, -1)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$, $l((5, -1)) = 2$ and $l((1, -1)) = 0$. Then the number of the pointed $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and non-positive pointed length m is equal to the number of the \mathcal{S} -nonnegative path with length n .

We omit the proof of Corollary 3.7. In Fig.3, we show an example of this Corollary with $n = 3$, where $*$ denote the marked point.




Fig.3. An example of Corollary 3.7, where $n = 3$.

Corollary 3.8 Let $\mathcal{S} = \{(1, 1), (2, 0), (1, -3)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$, $l((2, 0)) = 1$ and $l((1, -3)) = 0$. Then the number of the pointed $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and non-positive length m is equal to the number of the \mathcal{S} -nonnegative path with length n .

We omit the proof of Corollary 3.8. In Fig.4, we show an example of this Corollary with $n = 3$.




Fig.4. An example of Corollary 3.8, where $n = 3$.

4 The Chung-Feller property of Motzkin type

In this section, we will prove the Chung-Feller Theorem of Motzkin type for Classes 1,2. For $i = 1, 2$, recall that an $(\mathcal{S}_i, -k)$ -nonnegative path is an $(\mathcal{S}_i, -k)$ -lattice path which never goes below the line $y = -k$, where $k \geq 0$. \mathcal{N}_i^{-k} is the set of all the $(\mathcal{S}_i, -k)$ -nonnegative path. Define the generating functions

$$H_i^k(z) = \sum_{L \in \mathcal{N}_i^{-k}} w(L) z^{l(L)}.$$

Lemma 4.1 *For Classes 1,2, we have*

$$H_1^k(z) = [F_1(z)]^{k+1} \left[\sum_{i \in A} a_i z^{i-1} \right]^k \quad \text{and} \quad H_2^k(z) = [F_2(z)]^{k+1} \left[\sum_{i \in A} a_i z^i \right]^k.$$

Proof. For any a path $L \in \mathcal{N}_i^{-k}$ and $L \neq \emptyset$, we consider the first step s_m with height $-m$, where $1 \leq m \leq k$. Thus we can decompose the path L into $L_0 s_1 L_1 s_2 \dots L_{k-1} s_k L_k$, where $L_r \in \mathcal{N}_i$ for all $0 \leq r \leq k$ and $s_j \in \mathcal{S}_A$ for all j . Thus,

$$H_1^k(z) = [F_1(z)]^{k+1} \left[\sum_{i \in A} a_i z^{i-1} \right]^k \quad \text{and} \quad H_2^k(z) = [F_2(z)]^{k+1} \left[\sum_{i \in A} a_i z^i \right]^k.$$

■

Now we focus on the generating functions $M_i(y, z) = \sum_{\dot{L} \in \mathcal{M}_i^1} w(L) y^{mlp(\dot{L})} z^{l(L)-1}$ for $i = 1, 2$.

Lemma 4.2 *For Classes 1,2, we have*

$$M_1(y, z) = \frac{P_1(y, z) F_1(yz)}{1 - \left[\sum_{i \in A} a_i y^{i-1} z^i \right] [F_1(yz) F_1(z)]},$$

and

$$M_2(y, z) = \frac{P_2(y, z) F_2(yz)}{1 - \left[\sum_{i \in A} a_i y^i z^{i+1} \right] [F_2(yz) F_2(z)]}.$$

Proof. For any $k \geq 0$, let $\mathcal{M}_i^1(k)$ be the set of all the paths L in the set \mathcal{M}_i^1 such that $m(L) = -k$, where $m(L)$ is the minimum value of L . Clearly, $\mathcal{M}_i^1 = \bigcup_{k \geq 0} \mathcal{M}_i^1(k)$. Given a path $\dot{L} \in \mathcal{M}_i^1(k)$ and $L \neq \emptyset$, using the absolute minimum position of L , we can decompose

L into $R(1,1)\dot{T}$, where $R \in \mathcal{N}_i^{-k}$. For the path \dot{T} , we consider the rightmost step $(1,1)$ with height $-m$, where $-1 \leq m \leq k-1$. Thus we can decompose the path T into $L_{k-1}(1,1)L_{k-2}(1,1)\dots L_0(1,1)\dot{Q}$, where L_j is \mathcal{S}_i -nonnegative path for all $0 \leq j \leq k-1$ and $\dot{Q} \in \mathcal{M}_i$, where \mathcal{M}_i is the set of all the pointed \mathcal{S}_i -nonnegative path. Hence, by Lemmas 3.1 and 4.1, we get

$$M_i(y, z) = \sum_{k \geq 0} H_i^k(yz) [F_i(z)]^k z^k P_i(y, z).$$

Hence,

$$\begin{aligned} M_1(y, z) &= P_1(y, z) F_1(yz) \sum_{k \geq 0} [F_1(yz)]^k \left[\sum_{i \in A} a_i y^{i-1} z^{i-1} \right]^k [F_1(z)]^k z^k \\ &= \frac{P_1(y, z) F_1(yz)}{1 - \left[\sum_{i \in A} a_i y^{i-1} z^i \right] [F_1(yz) F_1(z)]}, \end{aligned}$$

and

$$\begin{aligned} M_2(y, z) &= P_2(y, z) F_2(yz) \sum_{k \geq 0} [F_2(yz)]^k \left[\sum_{i \in A} a_i y^i z^i \right]^k [F_2(z)]^k z^k \\ &= \frac{P_2(y, z) F_2(yz)}{1 - \left[\sum_{i \in A} a_i y^i z^{i+1} \right] [F_2(yz) F_2(z)]}. \end{aligned}$$

■

Now, we can prove the following Chung-Feller theorem of Motzkin type for Classes 1,2.

Theorem 4.3 *For Classes $i = 1, 2$, let $\bar{g}_{i;n,m}$ be the sum of the weights of the pointed $(\mathcal{S}_i, 1)$ -lattice paths which*

- (a.) *have length $n + 1$,*
- (b.) *have absolute minimum pointed length m ,*
- (c.) *have the length of the final step no less than 1.*

Let $f_{i;n}$ be the sum of the weights of the \mathcal{S}_i -nonnegative paths with length n . Then $\bar{g}_{i;n,m}$ has the Chung-Feller property of Motzkin type, i.e., $\bar{g}_{i;n,m} = f_{i;n}$.

Proof. In fact, we derived

$$M_i(y, z) = \frac{yF_i(yz) - F_i(z)}{y - 1}.$$

in the proof of Theorem 3.4 for $i = 1, 2$. Hence, the theorems hold. ■

Corollary 4.4 (*L. Shapiro*) Let $\mathcal{S} = \{(1, 1), (1, -1), (1, 0)\}$, $w(s) = 1$ and $l(s) = 1$ for any $s \in \mathcal{S}$. Then the number of the $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and absolute minimum pointed length m is the n -th Motzkin number.

Proof. For any a pointed $(\mathcal{S}, 1)$ -lattice path, we suppose the final point in this path is $(x, 1)$. Then the marked point must be $(x, 0)$ since $l(s) = 1$ for any $s \in \mathcal{S}$. So, we can erase the marked point. By Theorem 4.3, the number of the pointed $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and absolute minimum pointed length m is equal to the number of the \mathcal{S} -nonnegative paths with length n . By Lemma 2.3, we have $F_2(z) = 1 + zF_2(z) + z^2[F_2(z)]^2$ since $\mathcal{S} = \{(1, 1), (1, -1), (1, 0)\}$, $w(s) = 1$ and $l(s) = 1$ for any $s \in \mathcal{S}$. Hence, the number of the \mathcal{S} -nonnegative paths with length n is the n -th Motzkin number. This complete the proof. ■

Corollary 4.5 Let $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$. Then the number of the \mathcal{S} -path with length n and absolute minimum pointed length m in which the length of the final step is no less than 1 is the n -th Catalan number.

Proof. For any a pointed $(\mathcal{S}, 1)$ -lattice path, we suppose the final step in this path is s . Then $s = (1, 1)$ since $l((1, -1)) = 0 < 1$. If we delete the final step of this path and erase the marked point, we will obtain a \mathcal{S} -path with length n and absolute minimum length m . By Theorem 4.3, the number of the pointed $(\mathcal{S}, 1)$ -lattice path with length $n + 1$ and absolute minimum pointed length m in which the length of the final step is no less than 1 is equal to the number of the \mathcal{S} -nonnegative paths with length n . By Lemma 2.3, we have $F_1(z) = 1 + z[F_1(z)]^2$ since $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$. Hence, the number of the \mathcal{S} -nonnegative paths with length n is the n -th Catalan number. This complete the proof. ■




Fig.5. An example of Corollary 4.5, where $n = 2$

5 Conclusions

By Theorems 3.4 and 4.3, the lattice paths in Classes 1 and 2 have the Chung-Feller properties of both Dyck type and Motzkin type. We only prove the Chung-Feller theorem of Dyck type for Class 3. In fact, the lattice paths in Class 3 have the Chung-Feller properties of Motzkin type as well. We don't include this result in this paper since the statements are very complicate.

There are many lattice paths which have the Chung-Feller properties of both Dyck type and Motzkin type. For simplify, we set $\mathcal{S} = \{(1, 1), (1, -1)\}$, $w(s) = 1$ for any $s \in \mathcal{S}$, $l((1, 1)) = 1$ and $l((1, -1)) = 0$. Let θ be a mapping from \mathcal{L} to \mathbb{N} , where \mathcal{L} is the set of all the $(\mathcal{S}, 1)$ -lattice path. θ is called a parameter on $(\mathcal{S}, 1)$ -lattice path. For any $0 \leq m \leq n$, if the number of the $(\mathcal{S}, 1)$ -lattice path L with length n such that $\theta(L) = m$ is independent on m , then we say that θ is a *Chung-Feller parameter* for $(\mathcal{S}, 1)$ -lattice paths. There are two Chung-Feller parameters on $(\mathcal{S}, 1)$ -lattice path: non-positive pointed length and absolute minimum pointed length. To end this paper, we propose a problem: are there the other Chung-Feller parameters on $(\mathcal{S}, 1)$ -lattice paths?

References

- [1] Y.M. Chen, The Chung-Feller theorem revisited, *Discrete Math.* 308 (2008) 1328-1329
- [2] K.L. Chung, W. Feller, On fluctuations in-coin tossing, *Proc. Natl. Acad. Sci. USA* 35 (1949) 605-608
- [3] S. P. Eu, T.S.Fu, Y.N. Yeh, Taylor expansions for Catalan and Motzkin numbers, *Adv. Appl. Math.* 29 (2002) 345-357
- [4] S. P. Eu, T.S.Fu, Y.N. Yeh, Refined Chung-Feller theorems for lattice paths, *J. Combin. Theory Ser. A* 112 (2005) 143-162
- [5] Shu-Chung Liu, Yi Wang and Yeong-Nan Yeh, Chung-Feller property in view of generating functions, submitted
- [6] T.V.Narayana, Cyclic permutation of lattice paths and the Chung-Feller theorem, *Skand. Aktuarietidskr.* (1967) 23-30

- [7] L. Shapiro, Some open questions about random walks, involutions, limiting distributions, and generating functions, *Advances in Applied Math.* 27 (2001), 585-596.