Dephasing of entangled atoms as an improved test of quantum gravity

Mark S. Everitt, Martin L. Jones, and Benjamin T. H. Varcoe

School of Physics and Astronomy, University of Leeds, Leeds, LS2 9JT, United Kingdom

(Dated: April 10, 2019)

In a recent article Wang *et al.* (Class. Quantum Grav. 23 (2006) L59), demonstrated that phase of a particle fluctuated due to interactions with random deviations of a conformal gravitational field. Furthermore they demonstrated that atom interferometers were sensitive to these fluctuations and that sensitivity to Planck scale effects could be achieved with a sufficiently sensitive interferometer. In this paper we demonstrate that a class of entangled states, the N-atom Greenberger-Horne-Zeilinger (GHZ) states, provide a better scaling than atom interferometry and that current experiments are capable of making a significant impact in this field. We outline an experiment which uses atomic beams of rubidium atoms excited to Rydberg states that undergo controlled collisions in high-Q microwave resonators in a sequence that makes the resulting state highly sensitive to conformal field fluctuations. We show that a significant advance in sensitivity is possible.

PACS numbers: 04.80.Cc, 04.60.-m, 03.67.Bg

Unifying physics remains a task of considerable difficulty. For decades physicists have been unable to access the Planck scale where unification is expected to occur and have produced theories of quantum gravity to bridge this gap. Unfortunately extreme energy and length scales make the Planck scale difficult to reach directly. However over the last decade several steps have been taken towards phenomenological tests of quantum gravity. Power and Percival [1] showed that in the weak field and slow motion limit conformal fluctuations couple to massive particles and therefore atom interferometers should be able to detect the background field by making observations of small random relative shifts in phase. The loss of phase leads to a loss of visibility in interference fringes, elevating conformal fluctuations to a measurable scale. There were some approximations made in this paper that were subsequently improved upon by Wang et al. [2, 3] who showed that conformal fluctuations should be a feature in the semiclassical domain. In this letter we concentrate on the semiclassical domain described by Wang *et al.* [2, 3] where quantum gravitational effects are seen as random conformal fluctuations on otherwise flat Minkowski spacetime. We use the analogy that the dephasing seen in matter waves is like Brownian motion in that the origin of the fluctuations causing the dephasing is at the Planck scale but effects can be seen many of orders of magnitude away. The limit of these fluctuations is given as a cut-off length l_{cut} , where l_{cut} is linked to the Planck scale by a factor λ , which is the subject of interest in [1, 2, 3], by $l_{\rm cut} = \lambda l_{\rm Planck}$. λ is a background dependent parameter, which should have a unique prediction from each theory of quantum gravity and it is expected to be in the $10^2 - 10^6$ range [4]. Following the analogy to Brownian motion, λ is like a mean free path; larger values of lambda equate to less frequent kicks to the phase of massive particles, and a lower rate of dephasing.

Wang et al. arrived at the linkage between dephasing

of an atom with the equation [2]

$$\lambda = \left(\frac{8c^4\tau_0\sqrt{2\pi^5}}{9\hbar^2} \cdot \frac{M^2T}{p}\right)^{1/3},$$
 (1)

where τ_0 is the Planck time, M is the mass of the matter wave, T is the time the matter wave is allowed to travel before recombination and p is the parameter which describes dephasing. The parameter p is related to dephasing in the density matrix of an atom (a two state system) by

$$\begin{pmatrix} |\alpha|^2 & \alpha\beta^* \\ \alpha^*\beta & |\beta|^2 \end{pmatrix} \to \begin{pmatrix} |\alpha^2| & 0 \\ 0 & |\beta|^2 \end{pmatrix} + (1-p) \begin{pmatrix} 0 & \alpha\beta^* \\ \alpha^*\beta & 0 \end{pmatrix}$$
(2)

so that when p = 1 the atom is entirely dephased and when it is 0 the atom is in a pure state. For atom interferometry this is all that is needed. The loss of visibility in the interference fringes provides a lower bound on λ by equation (1). Atom interferometry experiments [5] currently put a lower bound on λ of 7600 [2]. Unfortunately the prospects for atom interferometry experiments improving on this are limited as the sensitivity scales unfavorably. Equation (1) shows that for every three orders of magnitude gain in sensitivity, only a single order of magnitude in λ is achieved.

Entanglement has been shown to be a sensitive tool for measuring dephasing and decoherence [6]. This is a problem in the field of quantum information, in which some entangled states which are particularly useful for quantum algorithms are unfortunately very sensitive to decoherence through dephasing. It is therefore interesting to consider the impact of quantum gravity induced dephasing on the correlations between measurements of entangled atoms. Following Sung Jang *et al.* [7], the expectation of the measurement of an *N*-atom Greenberger-Horne-Zeilinger (GHZ) state that has been subjected to p dephasing on each atom is

$$\langle B_1 B_2 \dots B_N \rangle = \frac{1 + (-1)^N}{2} \left(\prod_{i=1}^N \cos\left(\theta_i\right) \right)$$

$$+ (1-p)^N \cos\left(\sum_{i=1}^N \phi_i\right) \left(\prod_{i=1}^N \sin\left(\theta_i\right) \right),$$

$$(3)$$

in an arbitrary basis B_i for each atom i = 1, N. The basis describes the orientation of measurement on the Bloch sphere using the angles θ and ϕ ,

$$B_i = [\sigma_x \cos(\phi_i) + \sigma_y \sin(\phi_i)] \sin(\theta_i) + \sigma_z \cos(\theta_i) , \quad (4)$$

where σ_{α} are the Pauli operators. We choose $\theta = \pi/4$ and $\phi = 0$ to reduce equation (3) to measurements only in σ_x . Physically this would be achieved by applying a resonant $\pi/2$ rotation in the Bloch sphere of an atom and then measuring it, thereby enabling us to detect one of the states $|\pm\rangle = (|e\rangle \pm |g\rangle)/\sqrt{2}$. Under this condition equation (3) reduces to

$$\langle \sigma_x^{\otimes N} \rangle = (1-p)^N \,.$$
 (5)

This can be substituted directly into equation (1) to yield

$$\lambda = \left(\frac{8c^4\tau_0\sqrt{2\pi^5}}{9\hbar^2} \cdot \frac{M^2T}{1-\langle\sigma_x^{\otimes N}\rangle^{1/N}}\right)^{1/3}.$$
 (6)

This is the central result of this letter. In the limit of $\langle \sigma_x^{\otimes N} \rangle \approx 1$ equation (6) reduces to

$$\lambda \approx \left(\frac{8c^4 \tau_0 \sqrt{2\pi^5}}{9\hbar^2} \cdot \frac{NM^2 T}{1 - \langle \sigma_x^{\otimes N} \rangle}\right)^{1/3}, \qquad (7)$$

which corresponds to the sensitivity of a typical experiment. This demonstrates an improvement with respect to atom interferometry of $N^{1/3}$.

We now look at entanglement as a measurement tool and determine the outcome of an experiment. The GHZ state is a pure state in which all parties are 0 or all parties are 1 in equal superposition. For example an N = 3GHZ state is $(|000\rangle + |111\rangle)/\sqrt{2}$. This means that the decoherence of only one particle collapses the entire state leaving no entanglement regardless of the number of particles. These states are used for tests of non-locality.

The fragility of the GHZ class of states distinguishes it from other types of entangled state. The *degree of entan*glement for any GHZ state of length N is always 1, which is in contrast to the Werner state or 'W-state' which has a degree of entanglement proportional to $\log(N)$. The Wstate is simply an equal superposition of all states which contain N - 1 zeros and a one (e.g. for N = 3 the W state is $(|100\rangle + |010\rangle + |001\rangle)/\sqrt{3}$). This kind of entanglement often occurs in systems with a large number of particles such as Bose-Einstein condensates. The sensitivity to dephasing grows with *connectivity* as discussed



FIG. 1: A diagram demonstrating the collisional phase gate with two atoms as the qubits. The atom sources produce synchronized atoms with equal speed. The dashed arrows indicate the trajectory of individual atoms, the solid arrows are used to label components. This arrangement is a controlled phase gate on the atoms input in $(\alpha_1 | g \rangle + \beta_1 | e \rangle) \otimes (\alpha_2 | g \rangle + \beta_2 | e \rangle)$ with arbitrary $\alpha_{1,2}$ and $\beta_{1,2}$.

by Vedral [6]. The connectivity of a GHZ state of N atoms is N, whereas the connectivity of a W state is always 2 [6]. The W-state therefore retains the sensitivity equivalent to a pair of entangled particles and the advantage of entangling a large number of particles is lost. These two states are opposite extremes of entanglement, but they make the point that one needs to be careful in designing the experiment to create the right type of entanglement. These two types of state have been shown to be fundamentally different [8], so it is important to define the type of entanglement that would be created in a specific experiment and show that it has the required properties.

In the experiment that we are proposing an entangled state is constructed between a number of independent atoms. The entanglement is created using the collisional phase gate by Zheng and Guo [9] that was demonstrated by Haroche *et al.* [10]. For this experiment we have in mind a ⁸⁵Rb ladder of Rydberg states; $|e\rangle = 63P$, $|g\rangle = 61D$, and an auxiliary state $|i\rangle = 62P$ which are standard micromaser transitions [11]. Rydberg states are used commonly in the field of quantum optics for their relative longevity and large dipole moment [11]. The atom is effectively a two state system, which is detuned from the cavity resonance by the amount Δ . The Hamiltonian for this system is a two atom Tavis-Cummings Hamiltonian,

$$\frac{H_{\rm TC}}{\hbar} = \sum_{i} \left[\frac{\Delta \sigma_z^{(i)}}{2} + ig \left(\hat{a}^{\dagger} \sigma_-^{(i)} - \hat{a} \sigma_+^{(i)} \right) \right], \quad (8)$$

where the sum runs over the atoms in the interaction, q is the atom-cavity dipole coupling constant, \hat{a}^{\dagger} (\hat{a}) are the bosonic creation (annihilation) operators acting on the single mode cavity field, σ_+ (σ_-) are the atomic raising (lowering) operators and σ_z is the standard Pauli operator. This Hamiltonian is used to describe the evolution of a system with a single mode of the electromagnetic field coupling to a number of atoms simultaneously. In the situation with large detuning and two atoms, the system is reduced to an effective two-state system as the field is adiabatically eliminated. In this system an auxiliary state $|i\rangle$ is sometimes used which is so far detuned from resonance that it may be assumed that it does not interact with the field at all. In this event the Hamiltonian reduces to a detuned Jaynes-Cummings model for the atom not in the auxiliary state. The detuned Jaynes Cummings interaction is given simply as (with the atoms in either order and zero photons in the field)

$$\begin{array}{l} |g,i\rangle \mapsto |g,i\rangle \\ |e,i\rangle \mapsto \mathrm{e}^{-i\gamma t} |e,i\rangle \ , \end{array}$$

$$(9)$$

where γ is the effective coupling constant given by $\gamma = g^2/\Delta$. This interaction requires that the detuning is much larger than the coupling constant $\Delta \gg g$ to eliminate the field as a degree of freedom. The only two cases that they needed to consider for two atom interactions are the trivial case of two ground state atoms entering the cavity, which do not evolve, and the case of an excited atom and a ground state atom which interact by virtual excitation of the field. The case of two excited atoms entering is replaced by the case of one atom in the excited state and one in the auxiliary state. The reason and implementation of this omission will shortly become clear. For this case the Hamiltonian reduces to an effective Hamiltonian in the detuned limit

$$\frac{H_{\text{eff}}}{\hbar} = \gamma \sum_{i=1,2} \left[\sigma_{+}^{(i)} \sigma_{-}^{(i)} \hat{a} \hat{a}^{\dagger} - \sigma_{-}^{(i)} \sigma_{+}^{(i)} \hat{a}^{\dagger} \hat{a} \right] + \gamma \sigma_{+}^{(1)} \sigma_{-}^{(2)} + \gamma \sigma_{+}^{(2)} \sigma_{-}^{(1)}$$
(10)

For a cavity with zero photons this Hamiltonian yields the $|g,g\rangle$ and $|e,g\rangle$ evolutions

$$|g,g\rangle \mapsto |g,g\rangle$$

$$|e,g\rangle \mapsto e^{-i\gamma t} \left[\cos\left(\gamma t\right)|e_1g_2\rangle - i\sin\left(\gamma t\right)|g,e\rangle\right]$$
(11)

In the lab a low thermal photon number $(\langle \hat{a}^{\dagger} \hat{a} \rangle \approx 0)$ is achieved by cooling the cavity following the same procedure as the micromaser experiment [11]. By choosing the interaction time in equation (11) to be $t = \pi/(4\gamma)$ with one atom in $|e\rangle$ and the other in $|g\rangle$, an Einstein-Podolski-Rosen pair is produced. With the rotation of one atom to flip its state this is a GHZ state. Rotation of the atoms is achieved by passing the atoms through the side of a microwave waveguide. Alternatively low Q cavities pumped with microwaves can be used. This allows production of atoms in both states $|e\rangle$ and $|g\rangle$ (although $|g\rangle$ may be produced directly at the cost of more lasers and optics equipment), as well as the final rotation of states into the σ_x basis for measurement.

The experiment outlined is capable of producing only N = 2 GHZ states, and thus is sensitive to $(1 - p)^2$. Extensions to this system allow for larger GHZ states to be prepared. GHZ states with N atoms may be prepared by interacting one atom with the other atoms in turn in a similar scheme to that for 2 atoms. For the more general case, atoms are prepared in $|+\rangle$ states initially. We also invoke the auxiliary level $|i\rangle$, as we are implementing the true phase gate as outlined by Zheng and Guo [9]. As stated above this state does not interact with the field as it is not resonant.

For the general system rotation zones rotate every atom into a $|+\rangle$ state before interaction. N-1 cavities are placed in a line, and a special rotation zone is placed at either end. These rotation zones transfer $|e\rangle$ to $|i\rangle$ and vice versa. One of the atoms passes through every cavity and these rotation zones, interacting with one of the other atoms in each cavity with interaction time $t = \pi/\gamma$. A π pulse is then applied to this atom to flip its state, and a $\pi/2$ pulse to the other atoms. The GHZ state is now prepared and allowed to fly as long as possible for gravitational dephasing to take effect. Finally a $\pi/2$ pulse is applied to every atom for measurement in the σ_x basis. A diagrammatic example based on a current experimental design [12], is given in figure 2 for a 5 atom GHZ state. The experiment is run multiple times to develop the expectation value $\langle \sigma_x^{\otimes N} \rangle$ for substitution into equation (6), placing a new lower bound on λ .



FIG. 2: This example produces a five party GHZ state of atoms. Atoms approach as indicated by arrows to coincide in each cavity. Initially all atoms are in state $|e\rangle$. The rotation zones labelled a are $\theta = 3\pi/2$ pulses and c are $\theta = \pi/2$ pulses. Those zones labelled b act on the first atom (traveling horizontally) to switch $|e\rangle$ components of the state to $|i\rangle$, the auxiliary state, and vice versa. The enclosed region is where the GHZ state exists as it travels for time T related to velocity of atoms v, which is not to scale.

In summary we have shown that N-atom Greenberger-Horne-Zeilinger entangled states provide an avenue of increasing the sensitivity of tests of quantum gravity by a factor of $N^{1/3}$ over single atom interferometry. We have also shown how this state may be produced in a lab and that the effect of gravitational dephasing can be measured using current apparatus. This is therefore within current technological bounds and can be accessed by leveraging existing quantum optics technology.

- W. L. Power and I. C. Percival, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences 456, 955 (2000).
- [2] C. H. Wang, R. Bingham, and T. J. Mendonca, Classical and Quantum Gravity 23, L59 (2006), ISSN 0264-9381.
- [3] C. H. T. Wang, P. M. Bonifacio, R. Bingham, and T. J. Mendonca (2008), 0806.3042.
- [4] R. Bingham, Physica Scripta **T116**, 132 (2005).
- [5] A. Peters, K. Y. Chung, B. Young, J. Hensley, and S. Chu, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences 355, 2223 (1997).
- [6] V. Vedral, Nature 453, 1004 (2008), ISSN 0028-0836.
- [7] S. S. Jang, Y. W. Cheong, J. Kim, and H. W. Lee, Physical Review A (Atomic, Molecular, and Optical Physics) 74 (2006).
- [8] W. Dür, G. Vidal, and J. I. Cirac, Physical Review A 62, 062314 (2000).
- [9] S.-B. Zheng and G.-C. Guo, Physical Review Letters 85, 2392 (2000).
- [10] S. Osnaghi, P. Bertet, A. Auffeves, P. Maioli, M. Brune, J. M. Raimond, and S. Haroche, Physical Review Letters 87, 037902 (2001).
- [11] H. Walther, B. T. H. Varcoe, B. G. Englert, and T. Becker, Reports on Progress in Physics 69, 1325 (2006).
- [12] P. J. Blythe and B. T. H. Varcoe, New J. Phys. 8, 231 (2006).