

COARSE DYNAMICS AND FIXED POINT PROPERTY

TOMOHIRO FUKAYA*

Abstract

We investigate the fixed point property of the group actions on a coarse space and its Higson corona. We deduce the coarse version of Brouwer's fixed point theorem.

1 Introduction

Let X be a proper coarse space and G be a finitely generated semi-group acting on X . Here a coarse structure of G is generated by the 'diagonals'

$$\Delta_g = \{(h, hg) | h \in G\}$$

as g runs over a generating set for G . We will define the 'coarse action' of G on X . In this paper, we always assume that the coarse space is coarsely connected. The notation on the coarse geometry are based on Chapter 2 of the textbook written by John Roe [4].

DEFINITION 1.1. An action of G on X is the *coarse action* if for each element g of G , the map $\Psi_g: X \rightarrow X$ defined by $x \mapsto g \cdot x$ is a coarse map.

DEFINITION 1.2. For a point x_0 of X , we define the orbit map $\Phi_{x_0}: G \rightarrow X$ by $g \mapsto g \cdot x_0$ and also we define:

The orbit of x_0 is *proper* if Φ_{x_0} is a proper map.

The orbit of x_0 is *bornologous* if Φ_{x_0} is a bornologous map.

The orbit of x_0 is *coarse* if Φ_{x_0} is a coarse map.

2000 *Mathematics Subject Classification*. Primary 53C24; Secondary 55C20.

Key words and phrases. Coarse geometry, Higson corona, dynamics, fixed point

*The author is supported by Grant-in-Aid for JSPS Fellows (19-3177) from Japan Society for the Promotion of Science.

The most typical example of the coarse action is mentioned in Lemma 2.1.

THEOREM 1.3. *Assume that $G = \mathbb{N}^k$ or \mathbb{Z}^k and G acts on X as a coarse action. Suppose that there exists a point x_0 of X whose orbit coarse. Then the induced action of G on the Higson corona νX has a fixed point. Namely, there exists a point x of νX such that $g \cdot x = x$ for any element $g \in G$.*

Moreover, Let \overline{X} be a coarse compactification of X . We suppose that the action of G on X is continuous and extends to a continuous action on \overline{X} . Then the action on the boundary $\partial X = \overline{X} \setminus X$ has a fixed point. Namely, there exists a point z of ∂X such that $g \cdot z = z$ for any element $g \in G$.

EXAMPLE 1.4. Let G be a finitely generated group with an element γ of infinite order. Then a group action of \mathbb{Z} on G by $(n, g) \mapsto \gamma^n g$ is a coarse action and the orbit of $e \in G$ is coarse. Thus the action of \mathbb{Z} on the Higson corona νG has a fixed point. Moreover, if G is a hyperbolic group, this action extends to the Gromov boundary $\partial_g G$ (see Proposition 2.5 and Example 2.6). Then this action of \mathbb{Z} on $\partial_g G$ has a fixed point. This is a famous fact on the group action on the boundary of hyperbolic groups. (c.f. Proposition 10 and Théorème 30 in Chapter 8 of [2].)

EXAMPLE 1.5. The wreath product $\mathbb{Z} \wr \mathbb{Z}$ contains \mathbb{Z}^n as a subgroup for any integer n (see page 135 of [4]). Thus for any $n \in \mathbb{N}$, there is a coarse action of \mathbb{Z}^n on $\mathbb{Z} \wr \mathbb{Z}$ and induced action of \mathbb{Z}^n on $\nu(\mathbb{Z} \wr \mathbb{Z})$ has a fixed point.

REMARK 1.6. Let F_2 be a free group of rank 2. By Lemma 2.1, the action of F_2 on F_2 is a coarse action and has a coarse orbit. However, the induced action of F_2 on the Higson corona νF_2 does not have any fixed point (see Proposition 2.9).

Let $f: X \rightarrow X$ be a coarse map. We call a point x of X as a *coarse fixed point* of f if the orbit of x , $\{f^n(x) | n \in \mathbb{N}\}$, is a bounded set.

COROLLARY 1.7 (Coarse version of Brouwer's fixed point theorem). *Let X be a proper metric space. Suppose that $f: X \rightarrow X$ is an isometry map and \overline{X} is a coarse compactification of X such that f extends to a continuous map $\bar{f}: \overline{X} \rightarrow \overline{X}$. If f does not have any coarse fixed points in X , then \bar{f} has a fixed point in ∂X .*

This corollary says that an isometry f always has a coarse fixed point in X or a fixed point in ∂X .

EXAMPLE 1.8. Let M be a compact path metric space. The *cone* CM on M is the quotient space $M \times [0, \infty) / \sim$ where $(x, t) \sim (x', t')$ if and only if either $x = x'$ and $t = t'$ or $t = t' = 0$. We define a continuous function $\tau: [0, \infty) \rightarrow [0, 1)$ by

$$\tau(t) = \frac{t}{1+t}.$$

Set $\overline{CM} = M \times [0, 1] / \sim$. The embedding $CM \rightarrow \overline{CM} : (x, t) \mapsto (x, \tau(t))$ gives us a compactification of CM . We can define an appropriate metric on CM such that CM is a proper metric space. Then \overline{CM} becomes a coarse compactification (see Appendix). Let $f: CM \rightarrow CM$ be an isometry map and we suppose that f extends to a continuous map $\bar{f}: \overline{CM} \rightarrow \overline{CM}$. Then f has a coarse fixed point on X or \bar{f} has a fixed point on the boundary $\partial(CM) \cong M$. If M is ENR, then Lefschetz fixed point theorem implies that \bar{f} has a fixed point since \overline{CM} is contractible. However, Let g be an isometry map on a *punctured cone* $CM^\times = M \times [1/2, \infty)$. We suppose that g extends to a continuous map \bar{g} on a coarse compactification $\overline{CM^\times} = M \times [1/2, 1]$. Then g also has a coarse fixed points on CM^\times or \bar{g} has a fixed point on the boundary $\partial(CM) \cong M$. Since CM^\times is homotopic to M , Lefschetz fixed point theorem does not imply this fact.

EXAMPLE 1.9. Set $M = S^{n-1}$. Then CM is homeomorphic to the n -dimensional Euclidean space \mathbb{R}^n and \overline{CM} is homeomorphic to unit ball $B^n = \{x \in \mathbb{R}^n \mid |x| \leq 1\}$. Any isometry map $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ which can be extends to continuous map $\bar{f}: B^n \rightarrow B^n$ has a coarse fixed point on X or a fixed point on $\partial(\mathbb{R}^n) = S^{n-1}$. This corresponds to the Brouwer's fixed point theorem.

Acknowledgments

We would like to thank the participants in the coarse geometry seminar at Kyoto university, S.Honda, T.Kato, T.Kondo and M.Tsukamoto for several discussions and useful comments.

2 Coarse action

LEMMA 2.1. *Let G be a finitely generated group or $G = \mathbb{N}^k$ with left-invariant word metric for some generating set. The action of G on G by the left-translation $(g, h) \mapsto gh$ is a coarse action. Furthermore, for any point h of G , the orbit of h is coarse. any orbit of $h \in H$ has a coarse orbit.*

PROOF. For given $g \in G$, the map $\Psi_g: G \rightarrow G$ by $h \mapsto gh$ is an isometry map, so it is a coarse map. Let $h \in G$ be given, we consider the orbit map $\Phi_h: G \rightarrow G$ by $g \mapsto gh$. We denote the word length by $|\cdot|$. For any $g, g' \in G$, we have

$$d(\Phi_h(g), \Phi_h(g')) = d(gh, g'h) = |h^{-1}g^{-1}g'h| \leq |g^{-1}g'| + 2|h|.$$

This shows that Φ_h is a large scale Lipschitz map and hence a bornologous map. Let D be a bounded subset of G , which is a finite set. Since Φ_g is injective, $\#\Phi_g^{-1}(D) \leq \#D < \infty$. This shows that Φ_h is a proper map. It follows that Φ_h is a coarse map and the orbit of h is coarse. \square

The coarse action of G on X induces the continuous action on the Higson corona of X . Here we recall the definition and basic properties of the Higson corona. For more details, see Section 2.2 and 2.3 of [4].

Let X be a proper coarse space. Let $f: X \rightarrow \mathbb{C}$ be a bounded continuous function on X . We denote by $\mathbf{d}f$ the function $\mathbf{d}f(x, y) = f(y) - f(x): X \times X \rightarrow \mathbb{C}$. We define that f is a *Higson function* if for any controlled set $E \subset X \times X$, the restriction of $\mathbf{d}f$ to E vanishes at infinity*. We denote $C_h(X)$ as the set of Higson functions on X , which becomes unital C^* -algebra. The compactification hX of X characterized by $C(hX) = C_h(X)$ is called the *Higson compactification*. Its boundary $hX \setminus X$ is denoted νX and is called the *Higson corona* of X . The following are basic properties of the Higson corona.

PROPOSITION 2.2. *Let $f: X \rightarrow Y$ be a coarse map between proper coarse spaces. Then f extends to a continuous map $\nu f: \nu X \rightarrow \nu Y$. If $f, g: X \rightarrow Y$ are close, then $\nu f = \nu g$.*

The Higson corona is a huge space (usually not second countable). So it is useful to find more manageable compactifications. Thus we need to know the relation between the Higson corona and such compactifications.

DEFINITION 2.3. Let X be a proper coarse space and \overline{X} be a compactification of X . We call \overline{X} a *coarse compactification* if the following condition is satisfied: For any controlled set $E \subset X \times X$ of X , its closure \overline{E} in $\overline{X} \times \overline{X}$ meets the boundary $\partial(X \times X) = \overline{X} \times \overline{X} \setminus (X \times X)$ only at the diagonal, that is,

$$\overline{E} \cap \partial(X \times X) \subset \Delta_{\partial X} = \{(\omega, \omega) | \omega \in \partial X\}.$$

PROPOSITION 2.4. *The Higson compactification hX is a coarse compactification. Furthermore, it is a universal coarse compactification, in the sense that, for any coarse compactification \overline{X} of X , the identity map $\text{id}: X \rightarrow X$ extends uniquely to a continuous surjective map $\iota: hX \rightarrow \overline{X}$.*

PROPOSITION 2.5. *Let $f: X \rightarrow X$ be a continuous coarse map and \overline{X} be a coarse compactification. Suppose that f extends to a continuous map $\bar{f}: \overline{X} \rightarrow \overline{X}$. Then $\bar{f}(\partial X) \subset \partial X$ and the following diagram commutes.*

$$\begin{array}{ccc} hX & \xrightarrow{hf=f \cup \nu f} & hX \\ \downarrow \iota & & \downarrow \iota \\ \overline{X} & \xrightarrow{\bar{f}} & \overline{X} \end{array}$$

*That is, for any $\epsilon > 0$, there exists a bounded subset $B \subset X$ such that $|\mathbf{d}f(x, y)| < \epsilon$ for any $(x, y) \in E \setminus B \times B$.

Proof of Proposition 2.5. By the properness of f , we have $\bar{f}(\partial X) \subset \partial X$. Since f is a continuous coarse map, f induces $f^*: C_h(X) \rightarrow C_h(X)$. Because f extends to the continuous map on \bar{X} , f^* maps $C(\bar{X})$ to itself. It is clear that f^* and the inclusion $\iota^*: C(\bar{X}) \hookrightarrow C_h(X)$ commute. \square

EXAMPLE 2.6. Let X be a Gromov hyperbolic space. Then the Gromov compactification is a coarse compactification (Lemma 6.23 of [4]). If G is a hyperbolic group and $\gamma \in G$ has an infinite order, then left-translation $\gamma \cdot: G \rightarrow G$ extends to a continuous map on the Gromov boundary $\partial_g G$ (see Chapter 7 and 8 of [2]).

LEMMA 2.7. Let G be \mathbb{N}^k or \mathbb{Z}^k . A coarse action of G on G by $(g, n) \mapsto g + n$ extends to a trivial action on the Higson corona of G . Namely, for any $g \in G$, the induced map $\nu\Psi_g: \nu G \rightarrow \nu G$ is equal to the identity map.

PROOF. Let $g \in G$ be given. For any $n \in G$, $d(n, \Psi_g(n)) = d(n, g + n) = |g|$. Then Ψ_g and $\text{id}_G: G \rightarrow G$ are close. It follows that $\nu\Psi_g = \text{id}_{\nu G}$. \square

Proof of Theorem 1.3. Since the action is coarsely free, there exists $x_0 \in X$ such that the G -equivariant map $\Phi_{x_0}: G \rightarrow X$ by $g \mapsto g \cdot x_0$ is a coarse map. Thus Φ_{x_0} extends to a continuous map $\nu\Phi_{x_0}$ on νG to νX . For each $g \in G$, the coarse map $\Psi_g: X \rightarrow X$ extends to continuous maps $\nu\Psi_g: \nu X \rightarrow \nu X$ and $\bar{\Psi}_g: \partial X \rightarrow \partial X$. Then we have the following commutative diagram:

$$\begin{array}{ccc} \nu G & \xrightarrow{\nu\Psi_g = \text{id}} & \nu G \\ \nu\Phi_{x_0} \downarrow & & \downarrow \nu\Phi_{x_0} \\ \nu X & \xrightarrow{\nu\Psi_g} & \nu X \end{array}$$

By Lemma 2.7, the upper right arrow is the identity map. We choose any $x' \in \nu G$ and denote $x = \nu\Phi_{x_0}(x')$. Then by the above commutative diagram, we have $\nu\Psi_g(x) = x$.

Moreover, Let \bar{X} be a coarse compactification of X . We suppose that the action of G on X is continuous and extends to a continuous action on \bar{X} . Then we have the following commutative diagram:

$$\begin{array}{ccc} \nu X & \xrightarrow{\nu\Psi_g} & \nu X \\ \iota \downarrow & & \downarrow \iota \\ \partial X & \xrightarrow{\bar{\Psi}_g} & \partial X \end{array}$$

Set $z = \iota(x)$. We have $\bar{\Psi}_g(z) = z$. \square

REMARK 2.8. Suppose that X is a proper metric space and Φ_{x_0} is a continuous map. By the results of Dranishnikov; Keesling; Uspenskij [1], if Φ_{x_0} is a coarse embedding and

$\Phi_{x_0}(G)$ is a closed subset, we can show that the map $\nu\Phi_{x_0}$ in the proof of Theorem 1.3 is an embedding. Indeed, Theorem 1.4 of [1] states that the closure of $\Phi_{x_0}(G)$ in hX is homeomorphic to the Higson corona $\nu(\Phi_{x_0}(G))$ of $\Phi_{x_0}(G)$, whose coarse structure is a bounded coarse structure defined by the induced metric from X . Since Φ_{x_0} is a coarse embedding, νG is homeomorphic to $\nu(\Phi_{x_0}(G))$. Thus $\nu\Phi_{x_0}: \nu G \rightarrow \nu X$ is an embedding. It follows that there are so many fixed points on the Higson corona νX .

PROPOSITION 2.9. *The action of F_2 on νF_2 induced by the left-translation $F_2 \times F_2 \rightarrow F_2$ does not have any fixed point. Namely, there are no point x of νF_2 such that $g \cdot x = x$ for any element g of F_2 .*

PROOF. Gromov compactification is a coarse compactification, and isometry action on X induces the continuous action on the boundary. One can easily see that if the induced action of F_2 on νF_2 has a fixed point, the induced action of F_2 on the Gromov boundary $\partial_g F_2$ of F_2 also has a fixed point. However, we can show that for any point $z \in \partial_g F_2$, there exists an element g of F_2 such that $g \cdot z \neq z$. \square

3 Coarse fixed point

DEFINITION 3.1. Let G be a finitely generated semi-group acting on X . We call a point x of X a *coarse fixed point* if its orbit $G \cdot x = \{g \cdot x | g \in G\} \subset X$ is a bounded set.

REMARK 3.2. If G is an infinite group and x is a coarse fixed point, then the orbit of x is not proper.

We are interested in the problem: When does the converse of Remark 3.2 holds?

PROPOSITION 3.3. *Let X be a coarse space such that any bounded subset $D \subset X$ is a finite set. Suppose that an abelian semi-group \mathbb{N} acts on X . Then a point x_0 of X whose orbit is not proper is a coarse fixed point.*

PROOF. Since the orbit of x_0 is not proper, there exists a bounded set $D \subset X$ such that $\#\{n \in \mathbb{N} | n \cdot x_0 \in D\} = \infty$. Because $\#D < \infty$, there exists $m, n \in \mathbb{N}, m > n$ such that $m \cdot x_0 = n \cdot x_0$. Hence for any $l > m$, there are some $k > 0$ and $r = 0, \dots, m - n - 1$ satisfying $l - n = k(m - n) + r$ and thus $l \cdot x_0 = r \cdot (n \cdot x_0)$. It follows that $\mathbb{N} \cdot x_0 \subset \{x_0, 1 \cdot x_0, 2 \cdot x_0, \dots, (m - 1) \cdot x_0\}$. \square

PROPOSITION 3.4. *Let X be a proper metric space. Suppose that an abelian semi-group \mathbb{N} acts on X as an isometry action. Then a point x_0 of X whose orbit is not proper is a coarse fixed point.*

PROOF. Since the orbit of x_0 is not proper, there exists a bounded set $D \subset X$ such that $\#\{n \in \mathbb{N} | n \cdot x_0 \in D\} = \infty$. We can assume that x_0 lies in D . We notice that for any point x of the orbit $\mathbb{N} \cdot x_0$, there exist $n(x) \in \mathbb{N}$ such that $n(x) \cdot x$ lies in D . We define a bounded subset $K \subset X$ by

$$K = B(D, 1) \cap \mathbb{N} \cdot x_0.$$

Here $B(D, 1) = \{x \in X | \exists y \in D, d(x, y) < 1\}$ is the 1-neighborhood of D . Since \overline{K} is a compact set, there exists $x_1, \dots, x_N \in K$ such that

$$\overline{K} \subset \bigcup_{i=1}^N B(x_i, 1).$$

For $0 \leq i \leq N$, we denote $T_i = n(x_i)$. The point $T_i \cdot x_i$ lies in D . We define a positive number L by

$$L = \max_{0 \leq i \leq N} \max_{0 \leq a \leq T_i} d(x_0, a \cdot x_i).$$

We define inductively a sequence $\{i_k\}_{k=0}^{\infty}$ consisting of integers in $\{1, \dots, N\}$ and an increasing sequence $\{S_j\}_{j=0}^{\infty}$ as follows: Let $S_0 = T_0$. Since $S_0 \cdot x_0 = T_0 \cdot x_0 \in K$, there exists an integer $i_0 \in \{1, \dots, N\}$ such that $S_0 \cdot x_0 \in B(x_{i_0}, 1)$. Assume that we have defined i_0, \dots, i_n and S_0, \dots, S_n such that, for $0 \leq j \leq n$ and $0 \leq a < T_{i_j}$, they satisfy

$$\begin{aligned} S_j &= \sum_{k=0}^{j-1} T_{i_k}, \\ S_j \cdot x_0 &\in B(x_{i_j}, 1), \\ (a + S_j) \cdot x_0 &\in B(x_0, L + 1). \end{aligned}$$

Let $S_{n+1} = T_{i_n} + S_n$. Since $T_{i_n} \cdot x_{i_n} \in D$ and

$$d(S_{n+1} \cdot x_0, T_{i_n} \cdot x_{i_n}) = d(S_n \cdot x_0, x_{i_n}) < 1,$$

we have $S_{n+1} \cdot x_0 \in B(D, 1)$ and thus $S_{n+1} \cdot x_0$ lies in K . Hence there exists i_{n+1} such that $S_{n+1} \cdot x_0$ lies in $B(x_{i_{n+1}}, 1)$. For $0 \leq a < T_{n+1}$, we have $d(x_0, a \cdot x_{i_{n+1}}) \leq L$ and

$$d((a + S_{n+1}) \cdot x_0, a \cdot x_{i_{n+1}}) = d(S_{n+1} \cdot x_0, x_{i_{n+1}}) < 1.$$

It follows that $(a + S_{n+1}) \cdot x_0$ lies in $B(x_0, L + 1)$. This shows that for any integer $l > 0$, $l \cdot x_0$ lies in $B(x_0, L + 1)$ and thus x_0 is a coarse fixed point. \square

REMARK 3.5. Under the assumption of Proposition 3.4, if x_0 is a coarse fixed point, then any point x of X is a coarse fixed point.

If an isometry action does not have any coarse fixed point, the orbits of any points are proper. On the other hand, the orbits by an isometry action are always bornologous.

LEMMA 3.6. *Let X be a proper metric with an isometry action of \mathbb{N} . Then the action is a coarse action and any orbits are bornologous.*

PROOF. An isometry action is a coarse action. For any given point x of X , let $L = d(1 \cdot x, x)$ be the distance between $1 \cdot x$ and x . Then we have $d((i+1) \cdot x, i \cdot x) = L$ for all integers $i > 0$. Hence for any integers $m \geq n > 0$,

$$d(\Phi_x(m), \Phi_x(n)) = d(m \cdot x, n \cdot x) \leq \sum_{i=n}^{m-1} d((i+1) \cdot x, i \cdot x) = L|m-n|.$$

Thus Φ_x is a bornologous map. □

Proof of Corollary 1.7. A coarse action of \mathbb{N} on X is defined by $(n, x) \mapsto f^n(x)$. This is an isometry action. If f does not have any coarse fixed point, then by Lemma 3.4 and Lemma 3.6, any orbit of this action is coarse. Thus Theorem 1.3 implies that \bar{f} has a fixed point. □


4 More example of coarse action

Adding machine

Let $T_2 = (V_2, E_2)$ be a binary tree with the set of vertices V_2 and the set edges E_2 . We identify V_2 with the following sets.

$$V_2 \cong \{*\} \sqcup \bigsqcup_{n \geq 1} \{0, 1\}^n.$$

Here $*$ denotes the base point.



$*$ and 0 (resp. 1) can be joined by the edge $e_{*,0}$ (resp. $e_{*,1}$). Let $x = (i_{n-1}, \dots, i_0)$ and $y = (j_n, \dots, j_0)$ be two vertices. x and y can be joined by the edge $e_{x,y}$ if and only if

$i_0 = j_0, \dots, i_{n-1} = j_{n-1}$. T_2 has the usual metric and is a geodesic space. V_2 has the induced metric from T_2 .

We construct a coarse action of \mathbb{N} on V_2 . Let x be a vertex. If $x = (i_n, \dots, i_0) \neq (1, \dots, 1)$, we define $1 \cdot x = (j_{n-1}, \dots, j_0)$ with

$$\sum_{k=0}^{n-1} j_k 2^k = \sum_{k=0}^{n-1} i_k 2^k + 1.$$

If $x = (\overbrace{1, \dots, 1}^n)$, we define $1 \cdot x = (\overbrace{1, 0, \dots, 0}^{n+1})$. Finally we define $1 \cdot * = (0)$.

The Gromov product of $x = (i_{n-1}, \dots, i_0)$ and $y = (j_{m-1}, \dots, j_0)$ with a base point $*$ is defined to be

$$(x|y) = \frac{1}{2} \{d(x, *) + d(y, *) - d(x, y)\}.$$

We notice that $(x|y) = r$ if and only if $i_0 = j_0, \dots, i_{r-1} = j_{r-1}$ and $i_r \neq j_r$.

LEMMA 4.1. $1 \cdot : V_2 \rightarrow V_2$ is a coarse map.

PROOF. Let $x = (i_{n-1}, \dots, i_0)$ and $y = (j_{m-1}, \dots, j_0)$ be two vertices. Set $(x|y) = r$. Then $i_0 = j_0, \dots, i_{r-1} = j_{r-1}$. It is easy to see that $(1 \cdot x | 1 \cdot y) \geq r$. Hence we have

$$\begin{aligned} d(1 \cdot x, 1 \cdot y) &= d(1 \cdot x, *) + d(1 \cdot y, *) - 2(1 \cdot x | 1 \cdot y) \\ &\leq (n+1) + (m+1) - 2r \\ &= d(x, y) + 2. \end{aligned}$$

Thus $x \mapsto 1 \cdot x$ is a Large scale Lipschitz map. Since the map $1 \cdot$ is injective and every bounded set of V_2 is a finite set, we have the map is proper. \square

This action can be extended to a continuous action on the Gromov boundary $\partial_g V_2$ of V_2 . Here we can identify $\partial_g V_2$ as the Cantor set $\{0, 1\}^{\mathbb{N}}$ with a metric d defined to be

$$d(x, y) = 2^{-(x|y)}.$$

Here $x = (i_k)_{k=0}^{\infty}$, $y = (j_k)_{k=0}^{\infty}$. $(x|y) = r$ if and only if $i_0 = j_0, \dots, i_{r-1} = j_{r-1}$ and $i_r \neq j_r$.

PROPOSITION 4.2. The action of \mathbb{N} on $\partial_g V_2$ is minimal. Indeed, for any point x of $\partial_g V_2$, its orbit $\mathbb{N} \cdot x$ is dense in $\partial_g V_2$.

PROOF. Let $x \in \partial_g V_2$ be given. For any point y of $\partial_g V_2$ and $\epsilon > 0$, we will show that there exists an integer n such that $d(n \cdot x, y) < \epsilon$. Set $N > -\frac{\log \epsilon}{\log 2}$ and $x = (i_k)_{k=0}^{\infty}$. We choose $a = 2^{N+1} - \sum_{k=0}^N i_k 2^k$, then we have

$$a \cdot x = (\dots, \overbrace{0, \dots, 0}^{N+1}).$$

Set $y = (j_k)_{k=0}^\infty$ and $b = \sum_{k=0}^N j_k 2^k$. Then we have

$$(a + b) \cdot x = (\cdots, j_N, \cdots, j_0).$$

It follows that $d((a + b) \cdot x, y) < 2^{-N} < \epsilon$. \square

Especially, the action \mathbb{N} on $\partial_g V_2$ does not have any fixed points. On the other hand, the orbit of $x \in V_2$ is not a coarse fixed point. Thus by Proposition 3.3 and Theorem 1.3, the orbit of $x \in V_2$ can never be bornologous.

Appendix

Metric of the cone and its compactification

Let M be a compact path metric space. The cone CM on M is the quotient space $M \times [0, \infty) / \sim$ where $(x, t) \sim (x', t')$ if and only if either $x = x'$ and $t = t'$ or $t = t' = 0$. According to Roe (section 3.6 of [3]), we construct a metric on CM . Let $\lambda: [0, \infty) \rightarrow [0, \infty)$ be a continuous function with $\lambda(t) = 0$ if and only if $t = 0$. If γ is a path in CM , we define its λ -length, $l_\lambda(\gamma)$, to be

$$\sup \left\{ \sum_{j=0}^{n-1} \left(|t_j - t_{j+1}| + \max\{\lambda(t_j), \lambda(t_{j+1})\} d(x_j, x_{j+1}) \right) \right\}$$

where the supremum is taken over all finite sequence $(x_j, t_j)_{j=0}^n$ of points on the path γ (with (x_0, t_0) and (x_n, t_n) being two end points). We define a metric d_λ on CM by $d_\lambda((x, t), (x', t')) = \inf l_\lambda(\gamma)$ where the infimum is taken over all path joining (x, t) and (x', t') . The metric d_λ is compatible with the topology of CM and it becomes a proper metric space (Proposition 3.47 of [3]).

PROPOSITION 4.3. *Suppose that λ is an increasing, unbounded function, then the compactification $\overline{CM} = M \times [0, 1] / \sim$ is a coarse compactification.*

PROOF. Let $E \subset CM \times CM$ be a controlled set and r_E be a positive number such that

$$\sup\{d_\lambda(\mathbf{x}, \mathbf{y}) \mid (\mathbf{x}, \mathbf{y}) \in E\} < r_E.$$

Let $\{(\mathbf{x}_n, \mathbf{y}_n)\} \subset E$ be a sequence such that \mathbf{x}_n and \mathbf{y}_n tends to x and y in $\partial(CM) \cong M$ respectively. We denote $\mathbf{x}_n = (x_n, t_n)$ and $\mathbf{y}_n = (y_n, s_n)$. It is enough to show that $x = y$ (see Theorem 2.27 of [4]). Suppose that $x \neq y$. Since x_n and y_n converge to x and y respectively, for any $\epsilon > 0$ there exists $N_1 > 0$ such that, for all $n > N_1$, x_n and y_n

satisfy $d(x_n, y_n) > d(x, y) - \epsilon$. Because λ is an increasing, unbounded function, there exists $N_2 > 0$ such that, for all $t > N_2$,

$$\lambda(t) > \frac{r_E}{d(x, y) - \epsilon}.$$

Since \mathbf{x}_n tends to infinity, there exists $N_3 > 0$ such that, for all $n > N_3$, the second component t_n of \mathbf{x}_n satisfies $t_n > N_2$. Thus, if we choose an integer n satisfying $n > \max\{N_1, N_3\}$, then we have $d_\lambda(\mathbf{x}_n, \mathbf{y}_n) > r_E$, this contradicts the definition of r_E . \square

References

- [1] A. N. Dranishnikov, J. Keesling, and V. V. Uspenskij, *On the Higson corona of uniformly contractible spaces*, *Topology* **37** (1998), no. 4, 791–803. MR MR1607744 (99k:57049)
- [2] É. Ghys and P. de la Harpe (eds.), *Sur les groupes hyperboliques d'après Mikhael Gromov*, *Progress in Mathematics*, vol. 83, Birkhäuser Boston Inc., Boston, MA, 1990, Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988. MR MR1086648 (92f:53050)
- [3] John Roe, *Coarse cohomology and index theory on complete Riemannian manifolds*, *Mem. Amer. Math. Soc.* **104** (1993), no. 497, x+90. MR MR1147350 (94a:58193)
- [4] ———, *Lectures on coarse geometry*, *University Lecture Series*, vol. 31, American Mathematical Society, Providence, RI, 2003. MR MR2007488 (2004g:53050)

DEPARTMENT OF MATHEMATICS
KYOTO UNIVERSITY
KYOTO 606-8502
JAPAN

E-mail address: tomo_xi@math.kyoto-u.ac.jp