

# Entanglement of qubits via a nonlinear resonator

M. Kurpas, J. Dajka and E. Zipper

*Institute of Physics, University of Silesia, ul. Uniwersytecka 4, 40-007 Katowice, Poland\**

(Dated: March 15, 2019)

Coherent coupling of two qubits mediated by a nonlinear resonator is studied. It is shown that the amount of entanglement accessible in the evolution depends both on the strength of nonlinearity in the Hamiltonian of the resonator and on the initial preparation of the system. The created entanglement survives in the presence of decoherence.

PACS numbers: 03.67.Lx 03.67.Bg 42.50.Ct 85.25.Dq

## I. INTRODUCTION

Developments in quantum information science rely critically on entanglement - quantum correlations between two or more physical systems e.g. between two qubits. There are various ways of creating entangled states of two qubits. The most natural is to place the qubits close to each other so that they can interact directly by i.e. a mutual inductance or capacitance. There are spectacular experiments performed for solid state qubits using this type of interaction [1, 2, 3, 4, 5]. The interaction of distant qubits requires some 'interaction-carrying' medium: a bus. One of natural candidates is the quantum electromagnetic field i.e. a bosonic bath consisting of finite number of modes. The problem of interaction of qubits with a quantized monochromatic electromagnetic field in a quantum cavity has been discussed in several papers [6, 7, 8, 9, 10, 11]. The field mediating the entanglement (the quantum bus) is usually described by a linear resonator. However, in some coupling schemes [12, 13] an intermediate qubit with one or more Josephson junctions plays a role of a coupler. As the junction is strongly nonlinear, the coupler is treated as a nonlinear resonator.

In this paper we assume two qubits strongly coupled to a resonator and discuss a coherent, nonlocal coupling between qubits via a linear and nonlinear resonator. The goal is to determine conditions which allow to strongly entangle the qubits with each other with the disentangled resonator. The results are applicable to entanglement through the optical and microwave nonlinear cavities under the assumption that the qubit-field interaction is sufficiently weak to be described in the rotating wave approximation [14]. In the first part of the paper we also assume that all effects of dissipation and decoherence are negligible over the studied time scales. We briefly discuss the effect of dissipation in the last section of the paper.

To quantify the entanglement we calculate the negativity  $N$  [15] which is a natural measure of entanglement. Our discussion is general enough to not depend on the specific architecture. However, a candidate for possible

implementation is the flux or charge qubit interacting with microwave radiation.

In Section II we present the model of the system, in Sec. III A and III B we trace the evolution of the originally non-entangled state with a *single excitation*. We consider two initial states which will be shown to result in qualitatively different behaviour of the qubit-qubit entanglement: nonlinearity-assisted entanglement gain or entanglement suppression. In Section IV we investigate the system with *two* and *three excitations*, in Sec. V we verify the results for the negativity by presenting the qubit-qubit density matrix. Finally in Sec. VI we shortly study the influence of decoherence on the qubits entanglement.

## II. THE MODEL

The system under considerations consists of two qubits  $Q_i$ ,  $i = 1, 2$  and a monochromatic resonator  $R$ . We assume that both qubits are coupled to the resonator simultaneously. To simplify the notation we apply units such that  $\hbar = 1$ . In the absence of direct qubit-qubit interaction the Hamiltonian of such a system takes the form

$$H = H_{Q_1} + H_{Q_2} + H_R + H_{Q_1R} + H_{Q_2R}, \quad (1)$$

the qubit Hamiltonian is

$$H_{Q_i} = \frac{\Omega_i}{2} \sigma_z, \quad (2)$$

The interaction terms  $H_{Q_iR}$  read

$$H_{Q_iR} = -\frac{\gamma}{2} (a\sigma^+ + a^\dagger\sigma^-) \quad (3)$$

where  $\sigma^+ = \sigma_x + i\sigma_y$ ,  $\sigma^- = (\sigma^+)^\dagger$  and  $\sigma_z$  are the Pauli matrices and  $\gamma$  is the qubit-field coupling constant depending on the specific architecture. The resonator is described by a non-linear bosonic oscillator of the Hamiltonian

$$H_R = \omega_R \left( a^\dagger a + \frac{1}{2} \right) + V_R, \quad (4)$$

where the term  $V_R$  describes the nonlinearity. We shall discuss two different forms of the nonlinearity. The first

---

\*Electronic address: mkurpas@us.edu.pl

is a polynomial nonlinearity

$$V_R^1 = \alpha (a^2 + (a^\dagger)^2) \quad (5)$$

which is known to be related to the celebrated squeezed states [16, 17]. The second one, of central interest for our considerations, is the cosine-like nonlinearity

$$V_R^2 = \alpha \cos(a + a^\dagger) \quad (6)$$

The cosine term introduces nonlinearities of all orders and in that sense the cosine is 'more nonlinear' than any polynomial. This type of nonlinearity is present e.g. in the Hamiltonian of the microwaves generated by a SQUID [1, 13, 18]. For such a circuit the parameter  $\alpha$  can be changed by introducing into the resonator an adjustable weak link loop with a flux controllable Josephson tunneling energy [18]. A tunable anharmonic  $LC$  circuit is important as it introduces non-uniform level spacing reducing leakage to higher states [12]. In the limit of  $V_R = 0$  one arrives at the Hamiltonian of the Jaynes-Cummings type [19] for which exact solutions are known [8].

The state vector of the system at  $t = 0$  is a tensor product of three constituent states

$$|\psi_{Q_1 Q_2 R}(t=0)\rangle = |\psi_{Q_1}\rangle \otimes |\psi_{Q_2}\rangle \otimes |\psi_R\rangle \quad (7)$$

This choice is certainly justified for weakly interacting systems. The unitary evolution of the initially factorisable state leads in general to the entangled tripartite state. By taking the trace over the  $R$  states one obtains the reduced density operator  $\rho_{Q_1 Q_2}$  which we then use to calculate the negativity [15]

$$N(\rho_{Q_1 Q_2}) = \max(0, -\sum_i \lambda_i), \quad (8)$$

where  $\lambda_i$  are negative eigenvalues of the partially transposed [20] density matrix of the two qubits.  $N$  is 0 for separable states and reaches its maximal value  $N = 1/2$  for maximally entangled states. We look for the conditions under which the qubits get very strongly entangled.

### III. ENTANGLEMENT AND NONLINEARITY

In the following we study the influence of the nonlinear term in the bosonic Hamiltonian (4) on the quantitative and qualitative properties of entanglement of the qubit-qubit system. We consider below only the cosine nonlinearity as we have found that the influence of the nonlinearity (5) is weaker and not qualitatively different. We compare the results with those obtained for the linear single mode resonator, which despite its simplicity is a natural reference system.




FIG. 1: (color online) The  $QQR$  (line A) and  $QQ$  (line B) negativity as a function of dimensionless time  $\omega t$  for the linear case and for the initial state  $|eg0\rangle$ . Lines C and D depict the  $QQ-R$  and  $QQ$  negativity for the initial state  $|gg1\rangle$ ;  $\Omega_i = \omega, \gamma = 0.01\omega$ . Maximal qubit-qubit entanglement is possible when the resonator is not entangled with the qubits (lines C and D at  $\omega t = 110$ ).

#### A. Entanglement gain

The first of the considered initial states is

$$|\psi_{Q_1 Q_2 R}(t=0)\rangle = |eg0\rangle \quad (9)$$

Here  $e$  and  $g$  stand for the excited and ground state of one qubit respectively. The state (9) describes the first qubit in an excited state, the second one in the ground state and there are no photons inside the resonator.

Let us start with an analysis of the system in the absence of the nonlinear term i.e.  $V_R = 0$ . In this case one can perform exact analytical calculations of the wave function and trace the behavior of the system. The wave function of the system at the time  $t$  is [8]

$$\begin{aligned} |\psi_{Q_1 Q_2 R}(t)\rangle &= \frac{1}{2} (1 + \cos(\tilde{\gamma}t)) |eg0\rangle \\ &\quad - \frac{1}{2} (1 - \cos(\tilde{\gamma}t)) |ge0\rangle + \frac{i}{\sqrt{2}} \sin(\tilde{\gamma}t) |gg1\rangle, \end{aligned} \quad (10)$$

where  $\tilde{\gamma} = \sqrt{2}\gamma$ . The time evolution goes in the subspace spanned by the states  $|eg0\rangle, |ge0\rangle, |gg1\rangle$ . For  $\tilde{\gamma}t = n\pi$ ,  $n = 1, 2, \dots$  for which the  $R$  subsystem could be decoupled from the qubits also the coefficient at  $|ge0\rangle$  term becomes zero and the whole system remains disentangled. It is shown in Fig.1 as lines A ( $QQ-R$  entanglement) and B ( $QQ$  entanglement) at  $\omega t \simeq 220$ . For other values of  $\tilde{\gamma}t$  we are not able to obtain the entanglement of the qubits without the intrusion of the resonator states. This results in weak  $QQ$  entanglement with the limiting value  $N_{QQ}^{max} \sim 0.1$ . The inclusion of the nonlinear term ( $V_R \neq 0$ ) allows to go beyond this limit. It is shown in Fig.2 that even for the relatively weak nonlinearity the negativity remains bounded by nothing but its natural

limit i.e.  $N_{QQ}^{max} = 1/2$ .

Increasing the strength of the nonlinearity (Fig.3) the qubits get permanently entangled for a relatively long time which increases with increasing  $\alpha$ . For very strong nonlinearity ( $\alpha = 2$ ) the period is  $25000\omega t$  and the maximal value of  $N_{QQ} \simeq 0.45$  (not shown). Thus we see that for the non-symmetric initial state, when the qubits start from opposite states, the nonlinear resonator can lead, in contrast to the linear one, to the emergence of strongly entangled  $QQ$  states remaining very weakly entangled with the resonator states. In other words we get the coherent quantum state transfer between the qubits through the quantum bus.




FIG. 2: (color online) The unitary evolution of the qubit-qubit entanglement as a function of dimensionless time  $\omega t$  and the nonlinearity coefficient  $\alpha$ . The initial state  $|eg0\rangle$ .  $\Omega_i = \omega, \gamma = 0.01\omega$




FIG. 3: (color online) The qubit-qubit negativity for large values of the nonlinearity coefficient  $\alpha$ . The initial state  $|eg0\rangle$ .  $\Omega_i = \omega, \gamma = 0.01\omega$ .




FIG. 4: (color online) The influence of nonlinearity strength  $\alpha$  of the resonator on the entanglement of the qubits for the initial state (11).  $\Omega_i = \omega, \gamma = 0.01\omega$ .

## B. Entanglement suppression

The situation looks different for the 'symmetric' initial state

$$|\psi_{Q_1 Q_2 R}(t=0)\rangle = |gg1\rangle, \quad (11)$$

when the excitation is placed in the resonator and both qubits are in the ground state. The analytical calculations for the linear resonator ( $V_R = 0$ ) result in the wave function [8]

$$|\psi_{Q_1 Q_2 R}(t)\rangle = \frac{i}{\sqrt{2}} [\sin(\tilde{\gamma}t)|eg0\rangle + \sin(\tilde{\gamma}t)|ge0\rangle] + \cos(\tilde{\gamma}t)|gg1\rangle. \quad (12)$$

We see that at  $\tilde{\gamma}t = \pi/2 + m\pi$ , where  $m$  is an integer, we get maximally entangled qubit-qubit (Bell) state

$$|B_1\rangle = \frac{1}{\sqrt{2}}(|eg\rangle + |ge\rangle). \quad (13)$$

It is important that the qubits can be in the maximally entangled state only, when their common state space is separable from the resonator space. It is shown in Fig.1 (lines  $C$  - the  $QQ$ - $R$  negativity and  $D$  - the  $QQ$  negativity) at  $\omega t \simeq 110$ . As in the case of the initial state  $|eg0\rangle$ , the qubit-qubit negativity  $N_{QQ}$  reaches the value 0.1 (line  $D$ ) when the  $QQ$ - $R$  entanglement is maximal (Fig.1  $\omega t \simeq 50$  line  $C$ ).

The inclusion of the nonlinear term spoils the symmetry (see Eq. (12)) and results in suppression of the qubit-qubit entanglement with the negativity limited by  $N_{QQ}^{max} < 1/2$  depending on the amplitude  $\alpha$  (see Fig.4).

## IV. STATES WITH MANY EXCITATIONS

Finally, let us proceed to a consideration of the evolution of the originally non-entangled states with many

excitations. In Figs 5 and 6 we show the resulting qubit-qubit entanglement for the initial state with two and three excitations respectively

$$|\psi_{Q_1 Q_2 R}(t=0)\rangle = |eg1\rangle \quad (14)$$

$$|\psi_{Q_1 Q_2 R}(t=0)\rangle = |eg2\rangle. \quad (15)$$

If the interaction between the qubits goes via a linear resonator ( $\alpha = 0$ ) the negativity  $N_{QQ}$  (and hence the  $QQ$  entanglement) is very small (solid line). In other words, one is not able to get rid of the resonator degrees of freedom without leakage of the information about the state of the system. The situation changes if the interaction goes via the nonlinear resonator ( $\alpha \neq 0$  lines in Figs 5 and 6) and we obtain strong  $QQ$  entanglement. Thus in this case the influence of the nonlinearity of the "quantum bus" is favorable for the creation of the entanglement of the qubits.

If we start from

$$|\psi_{Q_1 Q_2 R}(0)\rangle = |ee0\rangle \quad (16)$$

the interaction via the linear resonator does not entangle the qubits ( $N_{QQ} = 0$ ). It is in agreement with exact analytical calculations [8]. We have also checked it by calculating the concurrence (not shown) which is another entanglement measure. Surprisingly, switching on the nonlinearity does not improve the situation and the qubits remain disentangled.

For the initial state

$$|\psi(0)\rangle = |gg2\rangle \quad (17)$$

for the interaction via the linear resonator we get the  $QQ$  entanglement with  $N_{QQ}^{max} \simeq 0.18$ . The presence of nonlinearity leads, in general, to a decrease of  $N_{QQ}^{max}$  but for some specific  $\alpha$  ( $\alpha \simeq 0.7$ ) we have found a small increase of the negativity (not shown).

Summarizing, the influence of nonlinearity is favorable for the initial states with one qubit in the excited state. Then the coupling via the quantum bus leads to strong entanglement and a coherent state transfer between the qubits.

## V. QUBIT-QUBIT DENSITY MATRIX

One of the best demonstrations of quantum entanglement is via the state tomography which yields a density matrix of coherently coupled qubits. Such experiments for qubits interacting via a resonant cavity are in progress [10]. As an example we present results of our calculations of the qubit-qubit density matrix for the case discussed in Sec.III A.

In Fig.7 we show the real part of the density matrix for the qubits interacting via the linear resonator. The matrix elements are taken at  $\omega t = 111$  for which  $N_{QQ} = 0.104$  in Fig.2. Looking at the state vector (10), this situation corresponds to equal probability of finding the




FIG. 5: (color online) The influence of the nonlinearity strength  $\alpha$  on the qubit-qubit entanglement for the initial state  $|eg1\rangle$ ,  $\Omega_i = \omega, \gamma = 0.01\omega$ .




FIG. 6: (color online) The influence of the nonlinearity strength  $\alpha$  on the qubit-qubit entanglement for the initial state  $|eg2\rangle$ ,  $\Omega_i = \omega, \gamma = 0.01\omega$ .




FIG. 7: (color online) The real part of the qubit-qubit density matrix for the case of the linear resonator and the initial state (9). The matrix is taken at  $\omega t = 111$  for which  $N_{QQ} = 0.104$  (solid line in Fig.2). The imaginary elements of the matrix are of the order of  $10^{-14}$ .




FIG. 8: (color online) The real part of the qubit-qubit density matrix for the case of the nonlinear resonator ( $\alpha = 0.01$ ) and the initial state (9). The values are taken at  $\omega t = 435$ ,  $N_{QQ} = 0.492$  (dashed line in Fig.2) .




FIG. 9: (color online) The imaginary part of the qubit-qubit density matrix for the case of the nonlinear resonator ( $\alpha = 0.01$ ) and the initial state (9). The values are taken at  $\omega t = 435$ ,  $N_{QQ} = 0.492$  (see dashed line in Fig.2).

excitation in the resonator or in one of the qubits. In Fig.8 and Fig.9 we present the real and imaginary part of the  $QQ$  density matrix for  $\alpha = 0.01$ . We see that in this case the non-diagonal matrix elements being the hallmark of entanglement are much larger as the coupling goes via the nonlinear resonator. The values correspond to dashed line in Fig.2 at  $\omega t = 435$ ,  $N_{QQ} = 0.492$ .

## VI. THE EFFECT OF DECOHERENCE

Design and construction of quantum devices is always affected by the influence of environment. In the following we apply the commonly used Markovian approximation




FIG. 10: (color online) The influence of decoherence on the entanglement of the qubits for the initial state (9).  $\Omega_i = \omega$ ,  $\gamma = 0.01\omega$ ,  $\alpha = 0.01$ ,  $T_R = 5 \cdot 10^{-5}$ ,  $T_{Q_i} = 10^{-5}$  s.

[14] and model the reduced dynamics of the  $QQR$  system in terms of master equation generating complete positive dynamics [22]. We take into account the decoherence of both qubits and the resonator. Following [11, 23] we assume that the effect of environment can be included in terms of two independent Lindblad terms:

$$\dot{\rho}(t) = [L_H - \frac{1}{2}L_D]\rho(t) \quad (18)$$

where the 'conservative part' is given by

$$L_H(\cdot) = -i[H, \cdot] \quad (19)$$

whereas the 'Lindblad dissipator'

$$L_D(\cdot) = L_R(\cdot) + \sum_{i=1} 2L_{Q_i}(\cdot) \quad (20)$$

$$L_R(\cdot) = A^\dagger A(\cdot) + (\cdot)A^\dagger A - 2A(\cdot)A^\dagger \quad (21)$$

$$L_{Q_i}(\cdot) = \Sigma_i^\dagger \Sigma_i(\cdot) + (\cdot)\Sigma_i^\dagger \Sigma_i - 2\Sigma_i(\cdot)\Sigma_i^\dagger \quad (22)$$

is expressed in terms of creation and annihilation operators 'weighted' by suitable lifetimes  $A = a/\sqrt{T_R}$  and  $\Sigma_i = \sigma^-/\sqrt{T_{Q_i}}$ , where  $T_R = 5 \cdot 10^{-5}$  s and  $T_{Q_i} = 10^{-5}$  s are the resonator and the qubits decoherence times respectively. It is shown in Fig.10 that the effect of such a local damping is purely qualitative and the constructive role of the nonlinear term in the oscillator is not obscured provided that the relaxation times are sufficiently large.

## VII. CONCLUSIONS

In this paper we have performed model calculations for the entangling quantum gate where the linear or nonlinear resonator plays the role of the quantum bus mediating the interaction between two, not directly interacting, qubits. The goal was to test the role of nonlinearity and to find conditions, under which the qubits become

strongly entangled. We have considered cosine-like nonlinearity where the discussed effects become most visible. We have found that the influence of the nonlinearity of the resonator, having its own dynamics, depends significantly on the initial state of the investigated system. We have shown two qualitatively different behaviours. The first occurs in the system with non-symmetric initial state which allows for a coherent transition of the excitation. Here the presence of nonlinearity results in the strong enhancement of the entanglement almost up to its maximal value. For sufficiently large amplitude of the nonlinear term, the entanglement becomes (quasi) permanent and does not vanish for a long time. This constructive role of nonlinearity is certainly desired for the applications. For the symmetric initial states the role of the nonlinearity is

no more constructive but rather results in suppression of entanglement. The coherent coupling of qubits survives in the presence of dissipation assuming decoherence times in agreement with recent experiments [9, 10].

### Acknowledgement

The work supported by the Polish Ministry of Science and Higher Education under the grant N 202 131 32/3786, Scientific Network LEPPi Nr 106/E-345/BWSN-0166/2008 and the scholarship from the UP-GOW project co-financed by the European Social Fund.

- 
- [1] J. B. Majer, F. G. Paauw, A. C. J. ter Haar, C. J. P. M. Harmans, J. E. Mooij
  - [2] Yu. A. Pashkin et al. *Nature* **421**, 823 (2003)
  - [3] A. J. Berkley et al. *Science* **300**, 1548 (2003)
  - [4] A. Izmailkov et al. *Phys. Rev. Lett.* **93**, 037003 (2004)
  - [5] S. H. W. van der Ploeg et al. *Phys. Rev. Lett.* **98**, 057004 (2007)
  - [6] A. Blais, R. Huang, A. Wallraff, S. M. Girvin, R. J. Schoelkopf, *Phys. Rev. A* **69**, 062320 (2004).
  - [7] R. Migliore, A. Messina *Phys. Rev. B* **72**, 214508 (2005).
  - [8] A. Yu. Smirnov, A. M. Zagoskin, cond-mat/0207214.
  - [9] J. Majer et al., *Nature* **449**, 443, (2007).
  - [10] M. A. Sillanpaa et al., *Nature* **449**, 438, (2007).
  - [11] E. Zipper, M. Kurpas, J. Dajka and M. Kuś *J. Phys: Condensed Matter*, **20**, 275219 (2008)
  - [12] A. Blais, A. Maassen van den Brink, A. M. Zagoskin, *Phys. Rev. Lett.* **90**, 127901 (2003)
  - [13] A. O. Niskanen et al. *Science* **316**, 723 (2007)
  - [14] C. W. Gardiner, P. Zoller, *Quantum noise*, Springer, Berlin (2000).
  - [15] G. Vidal, R. F. Werner, *Phys. Rev. A* **65**, 032314 (2002)
  - [16] A. Vourdas, *J. Phys. A* **39** R65-R141
  - [17] J. Perina, Z. Hradil, B. Jurco *Quantum optics and fundamentals of physics*, Springer-Verlag New York (1994)
  - [18] M. J. Everitt, T. D. Clark, P. B. Stiffell, A. Vourdas, J. F. Ralph, R. J. Prance, H. Prance, *Phys. Rev. A* **69**, 043804 (2004)
  - [19] E. T. Jaynes, F. W. Cummings, *Proc.IEEE*51, 89, (1963)
  - [20] A. Peres *Phys. Rev. Lett.* **77**, 1413, (1996)
  - [21] M. Steffen, M. Ansmann, R. C. Bialczak, N. Katz, E. Lucero, R. McDermott, M. Neeley, E. M. Weig, A. N. Cleland, J. M. Martinis, *Science* **313**, 5792, 1423 (2006).
  - [22] R. Alicki, K. Lendi, *Quantum dynamical semigroups and applications*, (*Lecture Notes in Physics* **286**), Berlin, Springer (1987)
  - [23] A. Blais, R. Huang, A. Wallraff, S. M. Girvin, R. J. Schoelkopf, *Phys. Rev. A* **69**, 062320 (2004)