

DIMINISHING FUNCTIONALS FOR NONCLASSICAL ENTROPY SOLUTIONS SELECTED BY KINETIC RELATIONS

MARC LAFOREST* AND PHILIPPE G. LEFLOCH†

Abstract. We consider nonclassical entropy solutions to scalar conservation laws with concave-convex flux functions, whose set of left- and right-hand admissible states u_l, u_r across undercompressive shocks is selected by a kinetic function $u_r = \varphi^b(u_l)$. We introduce a new definition for the (generalized) strength of classical and nonclassical shocks, allowing us to propose a generalized notion of total variation functional. Relying only upon the natural assumption that the composite function $\varphi^b \circ \varphi^b$ is uniformly contracting, we prove that the generalized total variation of front-tracking approximations is non-increasing in time, and we conclude with the existence of nonclassical solutions to the initial-value problem. We also propose two definitions of generalized interaction potentials which are adapted to handle nonclassical entropy solutions and we investigate their monotonicity properties. In particular, we exhibit an interaction functional which is globally non-increasing along a splitting-merging interaction pattern.

Key words. hyperbolic conservation law; entropy solution; non-convex flux-function; nonclassical shock; kinetic relation; total variation diminishing; interaction potential.

AMS subject classifications. 35L65; 82C26.

1. Introduction. Consider the following initial-value problem associated with a conservation law in one-space variable :

$$\begin{aligned} u_t + f(u)_x &= 0, \\ u(0, \cdot) &= u_0, \end{aligned} \tag{1.1}$$

where u_0 is a function of bounded variation on \mathbb{R} , and the (smooth) flux $f : \mathbb{R} \rightarrow \mathbb{R}$ is a *concave-convex function*, in the sense that

$$\begin{aligned} u f''(u) &> 0 \quad (u \neq 0), \quad f'''(0) \neq 0, \\ \lim_{|u| \rightarrow +\infty} f'(u) &= +\infty. \end{aligned} \tag{1.2}$$

Following LeFloch [6], we consider nonclassical entropy solutions to this problem. Recall that, in many applications, only a single entropy inequality can be imposed on the solutions, i.e.

$$U(u)_t + F(u)_x \leq 0, \tag{1.3}$$

where the so-called entropy U is a given, strictly convex function and the entropy flux $F(u) := \int^u U'(v) f'(v) dv$ is determined by U . It is not difficult to construct multiple weak solutions to the same initial-value problem (1.1), (1.3), so that one realizes that the single entropy inequality is too lax to determine a unique weak solution. In fact, for initial data restricted to lie in one region of concavity or convexity, the classical theory applies and leads to a unique entropy solution. Non-uniqueness arises when weak solutions contain transitions from positive to negative values, or vice-versa.

*Département de mathématiques et génie industriel, École Polytechnique de Montréal, Montréal, Québec, Canada, H3C 3A7. E-mail: Marc.Laforest@polymtl.ca.

†Laboratoire Jacques-Louis Lions & Centre National de la Recherche Scientifique (CNRS), Université Pierre et Marie Curie (Paris 6), 4 Place Jussieu, 75252 Paris, France. E-mail: pgLeFloch@gmail.com. To appear in: *Portugal. Math. (2009)*.

The above non-uniqueness property is closely related to the fact that discontinuous solutions, in general, depend upon their regularization, that is, different regularizations or approximations to the conservation law (1.1) may well lead to different solutions in the limit. This, in particular, is true for solutions to the Riemann problem, corresponding to the special initial data

$$u_0(x) = \begin{cases} u_l, & x < 0, \\ u_r, & x > 0, \end{cases} \quad (1.4)$$

where u_l, u_r are constant states. Indeed, for a wide class of regularizations, including regularizations by nonlinear diffusion-dispersion terms, nonclassical shocks violating Oleinik's entropy inequalities exist. The selection of a nonclassical solution is (essentially) equivalent to prescribing a *kinetic function* $\varphi^b : \mathbb{R} \rightarrow \mathbb{R}$ which, by definition, provides a characterization of admissible nonclassical shocks connecting states u_-, u_+ , that is,

$$u_+ = \varphi^b(u_-). \quad (1.5)$$

For scalar conservation laws and, more generally, nonlinear hyperbolic systems, LeFloch and co-authors initiated the development of a theory of nonclassical entropy solutions selected by the kinetic relation (1.5); various analytical and numerical aspects have been covered. We refer to [6] for a review of the theory and to [8] for recent results on the numerical issues. On the other hand, the kinetic relation was originally introduced in the context of a hyperbolic-elliptic model describing the dynamics of phase transitions in liquids or solids, for which we refer the reader to Slemrod [12], Truskinovsky [13], and Abeyaratne and Knowles [1]. LeFloch [5] used the Glimm scheme and first investigated mathematically the kinetic relations arising in phase dynamics. The subject has developed extensively since then, and we will not try here to review the literature.

Our objective in the present paper is to discuss the construction of new functionals for the (generalized) total variation and wave interaction potential of nonclassical entropy solutions to (1.1). We consider solutions generated by Dafermos' front-tracking method, when the local Riemann solutions are nonclassical and are determined by a given kinetic relation (1.5). We are interested in deriving uniform estimates for the total variation of solutions and showing that the scheme converges to global-in-time, nonclassical entropy solutions to the initial-value problem (1.1).

The new definition of generalized total variation proposed here is completely natural, and uses in a direct manner a key property satisfied by the kinetic function, that is, the uniform contraction property of the second iterate $\varphi^b \circ \varphi^b$. Indeed, this property (see (2.1), below) motivates our definitions of shock strength and, hence, of total variation. In turn, the proposed construction provides both an improvement of earlier results on the subject and a drastic simplification of earlier arguments (cf., for instance, with Section 8.1 in [6]). Most importantly, it appears that our new arguments are robust and may be generalized to tackle systems of equations.

An outline of this paper follows. In Section 2, we begin with a brief review of the theory of kinetic relations, by emphasizing what we will need in the rest of this paper. We then introduce our new definition of generalized shock strength. In Section 3, we prove that the proposed generalized total variation functional is non-increasing along a sequence of Dafermos' front tracking solutions; cf. Theorem 3.1. In Section 4, we turn to the construction of interaction functionals adapted to nonclassical entropy

solutions. In Theorem 4.2 we show that a rather natural definition leads to an interaction functional which is non-increasing in all but four interaction patterns. Next, in Section 5 we prove that the proposed interaction functional is actually globally non-increasing, at least in the significant case of merging-splitting wave patterns. Such solutions were originally introduced by LeFloch and Shearer [9]. Hence, Theorem 5.1 below demonstrates the relevance of the proposed functional to handle nonclassical entropy solutions. Finally, in Section 6 we investigate a second definition of interaction functional which may be better suited for the extension to systems; see Theorem 4.2.

2. Kinetic function and generalized wave strength.

2.1. Assumptions on the kinetic function. The general theory of kinetic functions φ^b imposes the following conditions on the kinetic function:

- (A1) The map $\varphi^b : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous and one-to-one.
- (A2) $\partial_u \varphi^b(u) < 0$ for all u , and $\varphi^b(0) = 0$.
- (A3) The second iterate $\varphi^b \circ \varphi^b$ is a strict contraction, i.e. for some $K \in (0, 1)$

$$|\varphi^b \circ \varphi^b(u)| \leq K|u|, \quad u \neq 0. \quad (2.1)$$

These are the sole conditions required on the kinetic functions in the present paper. We recall that these assumption are satisfied by quite general regularizations of the conservation law (1.1) and, moreover, natural analogues of the properties (A1), (A2), and (A3) are known to hold for *systems*, when φ^b is properly defined in this more general setting. We refer the reader to the monograph [6] and the references cited therein for further information.

The kinetic function determines, via (1.5), the set of admissible states u_-, u_+ arising on the left- and the right-hand sides of a discontinuity. When $u_- > 0$ ($u_- < 0$, respectively), nonclassical shocks are required when the amplitude of the shock is large enough and the threshold $u_+ < \varphi^\sharp(u_-)$ ($\varphi^\sharp(u_-) < u_+$, resp.) is reached. This threshold function $\varphi^\sharp : \mathbb{R} \rightarrow \mathbb{R}$ is defined as the unique value $\varphi^\sharp(u_-) \notin \{u_-, \varphi^b(u_-)\}$ such that

$$\frac{f(u_-) - f(\varphi^b(u_-))}{u_- - \varphi^b(u_-)} = \frac{f(u_-) - f(\varphi^\sharp(u_-))}{u_- - \varphi^\sharp(u_-)}.$$

Geometrically, $(\varphi^\sharp(u_-), f(\varphi^\sharp(u_-)))$ is the point at which the line between $(u_-, f(u_-))$ and $(\varphi^b(u_-), f(\varphi^b(u_-)))$ crosses the graph of f and therefore, if for instance $u_- > 0$, the following inequalities must hold

$$\varphi^b(u_-) < \varphi^\sharp(u_-) < u_-. \quad (2.2)$$

Under the concave-convex condition (1.2), the nonclassical entropy solution to the Riemann problem with data (1.4), $u_l > 0$, is given by

- i) a shock if $\varphi^\sharp(u_l) \leq u_r$, or
- ii) a nonclassical shock connecting u_l to $\varphi^b(u_l)$ followed by a classical shock connecting $\varphi^b(u_l)$ to u_r if $\varphi^b(u_l) < u_r < \varphi^\sharp(u_l)$, or
- iii) a nonclassical shock connecting u_l to $\varphi^b(u_l)$ followed by a classical rarefaction connecting $\varphi^b(u_l)$ to u_r if $u_r \leq \varphi^b(u_l)$.

When $u_l < 0$ and $u_r > 0$, the nonclassical Riemann solver is similar; see [5] for further details.

Some remarks on our assumptions (A1)–(A3) above are in order. In fact, these properties can be motivated from similar properties satisfied by the so-called *zero dissipation kinetic function* φ_0^b , which is determined from the flux f and the given entropy U and is *formally* associated to a dispersion-only regularization of (1.1). Interestingly, this special kinetic function satisfies a stronger version of (2.1), namely

$$\varphi_0^b \circ \varphi_0^b(u) = u. \quad (2.3)$$

On the other hand, for (non-trivial) regularizations including diffusion terms, the kinetic function satisfies the strict inequality

$$|\varphi^b(u)| < |\varphi_0^b(u)|, \quad u \neq 0. \quad (2.4)$$

In the general theory, (2.1) arises as a consequence of the more fundamental identities (2.3) and (2.4). It is, therefore, natural that our functional involve the function φ_0^b rather than directly φ^b , and its definition makes direct use of the properties (2.3) and (2.4).

Finally, we observe that for any bounded set of values $|u| < M$, properties (A2) and (A3) imply that there exist lower and upper Lipschitz constants $\underline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b) > 0$ and $\overline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b) < 1$ such that for $u \neq 0$

$$\underline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b) |u| < |u - \varphi_0^b \circ \varphi^b(u)| < \overline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b) |u|. \quad (2.5)$$

In the context of the general theory, we note that only the lower bound is non-trivial and that it does hold for large classes of diffusion-dispersion relations. On the other hand, for highly degenerate regularizations, such as the one studied in [2], this condition might be violated.

2.2. New notion of generalized wave strength. We introduce here a new functional, which is equivalent to the total variation and can be expressed entirely in terms of the kinetic function φ^b . In order to hope for an extension to systems, the formulation of the total variation should let the property (2.1) appear as naturally as possible. Intuitively, the following definition of strength is based on the idea of mapping all the states to the convex region of the flux ($u > 0$), using φ_0^b .

DEFINITION 2.1 (Notion of generalized wave strength). *To each classical or nonclassical wave (u_-, u_+) one associates the generalized strength $\sigma(u_-, u_+)$ defined as follows : when $u_- \geq 0$,*

$$\sigma(u_-, u_+) = \begin{cases} |u_- - u_+|, & u_+ \geq 0, \\ |u_- - \varphi_0^b(u_+)|, & u_+ < 0, \end{cases} \quad (2.6)$$

while for $u_- < 0$,

$$\sigma(u_-, u_+) = \begin{cases} |\varphi_0^b(u_-) - \varphi_0^b(u_+)|, & u_+ < 0, \\ |\varphi_0^b(u_-) - u_+|, & u_+ \geq 0. \end{cases} \quad (2.7)$$

The following notation will be helpful for the analysis. Henceforth, classical shocks and rarefactions joining two positive states is denoted by C_+^\downarrow and R_+^\uparrow , respectively. Similarly, when both neighboring states are negative, we write C_-^\uparrow and R_-^\downarrow . When a shock joins a positive state with a negative state, we write C_\pm^\downarrow or N_\pm^\downarrow depending on

whether or not the shock is classical or nonclassical, respectively. When the signs of the neighboring states are reversed, we simply write C_{\mp}^{\uparrow} and N_{\mp}^{\uparrow} . Shocks C_{\pm}^{\downarrow} and C_{\mp}^{\uparrow} are also sometimes called crossing shocks.

The generalized wave strength enjoys several immediate properties.

- The first important property to observe is that the proposed generalized strength is *continuous* as u_+ crosses $\varphi^{\sharp}(u_-)$ and the solution of the Riemann problem goes from a single crossing shock to a nonclassical shock followed by a classical shock.

For $u_- > 0$, this is checked from the inequalities $\varphi^{\flat}(u_-) < \varphi^{\sharp}(u_-) < 0$ and the property (2.1) which yield

$$\begin{aligned} \sigma(u_-, \varphi^{\sharp}(u_-)) &= |u_- - \varphi_0^{\flat} \circ \varphi^{\sharp}(u_-)| \\ &= |u_- - \varphi_0^{\flat} \circ \varphi^{\flat}(u_-)| + |\varphi_0^{\flat} \circ \varphi^{\flat}(u_-) - \varphi_0^{\flat} \circ \varphi^{\sharp}(u_-)| \\ &= \sigma(u_-, \varphi^{\flat}(u_-)) + \sigma(\varphi^{\flat}(u_-), \varphi^{\sharp}(u_-)). \end{aligned}$$

In fact, $\varphi^{\sharp}(u_-)$ could be positive and a similar computation would show that continuity still holds.

- The second important property of the generalized strength is its “equivalence” with the usual notion of strength. For positive non-crossing shocks and rarefactions, the definition is the same as the usual one. When the rarefaction and the non-crossing shocks have two negative neighboring states, then there exists a positive constant C' such that

$$\sigma(u_-, u_+) = |\varphi_0^{\flat}(u_-) - \varphi_0^{\flat}(u_+)| \geq C'|u_- - u_+|, \quad (2.8)$$

as long as u_-, u_+ stay within a bounded neighbourhood of the origin. For the shocks C_{\pm}^{\downarrow} and N_{\pm}^{\downarrow} , it suffices to use (2.5) to show that the definition is equivalent to the usual notion of strength :

$$\sigma(N_{\pm}^{\downarrow}) = |u_- - \varphi_0^{\flat} \circ \varphi^{\flat}(u_-)| > C''2|u_-| > C'''|u_- - \varphi^{\flat}(u_-)|. \quad (2.9)$$

The same argument also applies to shocks C_{\mp}^{\uparrow} and N_{\mp}^{\uparrow} . Interestingly, our definition of shock strength makes clear the intuitive idea that it should be increasingly difficult to measure the strength of nonclassical shocks as the zero-diffusion (dispersion-only) case is approached.

3. Diminishing total variation functional. We now introduce front-tracking approximate solutions to (1.1) based on a nonclassical Riemann solver, as was constructed by Dafermos [7] in the classical situation. These approximations are piecewise constant in space and are determined from the nonclassical Riemann solver described in the previous section.

The first step of their construction is to build a piecewise constant approximation of the initial data u_0 which admits finitely many discontinuities and approaches u_0 in the L^1 norm with an error ϵ , for some small ϵ . The Rankine-Hugoniot condition can be used to propagate, in a conservative manner, the discontinuities of the initial data. When the Riemann solver calls for continuous waves, we replace them by a sequence of small discontinuities (u_-, u_+) whose strength satisfy $\sigma(u_-, u_+) < \epsilon$.

When two discontinuities meet, the nonclassical Riemann solver is used to determine the neighboring states, but we continue to enforce that all outgoing waves be discontinuities. Despite the fact that the number of outgoing waves may be larger

than the number of incoming discontinuities, one can check (see [6] for details) that the total number of discontinuities remains bounded for all times, so that the front-tracking approximation can be defined for all times.

For a front-tracking approximation $u : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$, formed entirely of propagating discontinuities (each denoted by α), the inequalities (2.8) and (2.9), as well as the fact that the kinetic functions are Lipschitz continuous, imply that

$$V(u(t)) := \sum_{\alpha} \sigma(u_{-}^{\alpha}, u_{+}^{\alpha}), \quad (3.1)$$

is equivalent to the total variation norm

$$\text{TV}(u(t)) := \sum_{\alpha} |u_{-}^{\alpha} - u_{+}^{\alpha}|, \quad (3.2)$$

where u_{\pm}^{α} denote the left- and right-hand states of the discontinuity α .

Incidentally, we mention that the kinetic function φ_0^b induces an isometry on the space of $\text{BV}(\mathbb{R})$ functions with respect to the norm V :

$$\begin{aligned} \Phi_0^b : \text{BV}(\mathbb{R}) &\rightarrow \text{BV}(\mathbb{R}) \\ u(\cdot) &\mapsto \varphi_0^b \circ u(\cdot). \end{aligned} \quad (3.3)$$

Using (2.3), it is straightforward to check that Φ_0^b is an isometry. This result will not be used directly in this paper.

THEOREM 3.1 (Diminishing total variation functional). *Let φ^b be a kinetic function satisfying the properties (A1)–(A3). For every front-tracking approximation $u : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ to the conservation law (1.1) based on the nonclassical Riemann solver associated with φ^b , the generalized total variation functional $V(u(t))$ is non-increasing. Precisely, the change in V during an interaction is given by:*

$$[V] \leq \begin{cases} -c_1 \sigma(R^{in}), & \text{Cases RC-1, RC-3, CR-1, CR-2, CR-4,} \\ -c_2 \sigma(R^{in}), & \text{Cases RC-2, RN,} \\ -2(\sigma(R^{in}) - \sigma(R^{out})), & \text{Case CR-3,} \\ 0, & \text{all other cases.} \end{cases} \quad (3.4)$$

Here, $c_1 = \min\{1, \underline{\text{Lip}}(\varphi_0^b)\}$, $c_2 = \underline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b)$, while R^{in} and R^{out} denote the incoming and outgoing rarefactions at an interaction, respectively. (The list of interactions is specified in the proof below.)

From this theorem we immediately deduce the following:

COROLLARY 3.2 (Existence of nonclassical entropy solutions). *For any initial data $u_0 \in L^\infty(\mathbb{R}) \cap \text{BV}(\mathbb{R})$ of (1.1), and any sequence of front-tracking approximations u^ϵ such that $u^\epsilon(\cdot, 0)$ converges to $u_0(\cdot)$ in $L^1(\mathbb{R})$, there exists a subsequence of front-tracking approximations that converge in $\text{Lip}([0, T], L^1(\mathbb{R})) \cap L^\infty([0, T], \text{BV}(\mathbb{R}))$ to a solution of the initial value problem (1.1).*

Proof of Theorem 3.1. Our proof is a generalization of the one given in [6] (see Section 8.1). We need here to compute the variation of our functional V by distinguishing between 16 possible interactions, after assuming $u_l > 0$ for definiteness, as described in Section 4.3 of [6]. We note that we have one more case to consider here, since our assumptions on the kinetic function are general enough to allow for the so-called CC-3 interactions. The subscript ' is used to indicate that a wave is outgoing

and with some abuse of notation for the wave strength we write, for example, $\sigma(N_{\pm}^{\downarrow})$ for the shock strength of a nonclassical wave. During a generic interaction between two waves, we denote the states on both sides of the left-hand wave by u_l and u_m while those associated with the right-hand wave are denoted by u_m and u_r . Finally, the bounds on the change of V depend only on the properties (2.1)–(2.5) which are therefore used freely throughout.

Case RC-1 : $(R_+^{\uparrow} C^{\downarrow})-(C^{\downarrow}')$. This case is determined by the constraints

$$\max(\varphi^{\sharp}(u_l), \varphi^{\sharp}(u_m)) < u_r < u_l, \quad 0 < u_l < u_m.$$

This is further subdivided into two subcases depending on the sign of u_r . When $u_r > 0$, then the interactions are entirely classical $(R_+^{\uparrow} C_+^{\downarrow})-(C_+^{\downarrow}')$ and the inequalities $0 < u_r < u_l < u_m$ suffice to check that

$$[V] = \sigma(C_+^{\downarrow}') - \sigma(R_+^{\uparrow}) - \sigma(C_+^{\downarrow}) = -2|u_m - u_l| = -2\sigma(R_+^{\uparrow}).$$

When $u_r < 0$, then the interaction involves crossing shocks $(R_+^{\uparrow} C_{\pm}^{\downarrow})-(C_{\pm}^{\downarrow}')$ and the states involved in measuring the strengths of the waves are

$$0 < \varphi_0^{\flat}(u_r) < u_l < u_m, \quad (3.5)$$

since $\varphi_0^{\flat}(u_r) < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < u_l$. Then

$$\begin{aligned} [V] &= \sigma(C_{\pm}^{\downarrow}') - \sigma(R_+^{\uparrow}) - \sigma(C_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^{\flat}(u_r)| - |u_m - u_l| - |u_m - \varphi_0^{\flat}(u_r)| = -2\sigma(R_+^{\uparrow}). \end{aligned}$$

Case RC-2 : $(R_+^{\uparrow} C_{\pm}^{\downarrow})-(N_{\pm}^{\downarrow} R_{\pm}^{\downarrow}')$. This case is defined by

$$\varphi^{\sharp}(u_m) < u_r \leq \varphi^{\flat}(u_l) < 0 < u_l < u_m.$$

In the first subcase, we assume $\varphi_0^{\flat}(u_r) < u_l$ and use the previous conditions to deduce

$$\varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < \varphi_0^{\flat}(u_r) < u_l < u_m. \quad (3.6)$$

The analysis when $u_l < \varphi_0^{\flat}(u_r)$ will not be treated since only the order of those two terms changes but the conclusions remain the same.

$$\begin{aligned} [V] &= \sigma(N_{\pm}^{\downarrow}') + \sigma(R_{\pm}^{\downarrow}') - \sigma(R_+^{\uparrow}) - \sigma(C_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| + |\varphi_0^{\flat}(u_r) - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| - |u_m - u_l| - |u_m - \varphi_0^{\flat}(u_r)| \\ &= 2|\varphi_0^{\flat}(u_r) - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| - 2|u_m - u_l|. \end{aligned}$$

Since $\varphi^{\sharp}(u_m) < u_r < 0$, from property (2.1) we deduce $0 < \varphi_0^{\flat}(u_r) < \varphi_0^{\flat} \circ \varphi^{\sharp}(u_m) < \varphi_0^{\flat} \circ \varphi^{\flat}(u_m)$. Combining this with (3.6), we find

$$[V] \leq -2\underline{\text{Lip}}(u - \varphi_0^{\flat} \circ \varphi^{\flat})|u_m - u_l| = -2\underline{\text{Lip}}(u - \varphi_0^{\flat} \circ \varphi^{\flat})\sigma(R_+^{\uparrow}).$$

Case RC-3 : $(R_+^{\uparrow} C^{\downarrow})-(N_{\pm}^{\downarrow} C^{\uparrow}')$. The conditions initially satisfied by the neighboring states of the incoming waves are

$$\max(\varphi^{\flat}(u_l), \varphi^{\sharp}(u_m)) < u_r < \varphi^{\sharp}(u_l), \quad 0 < u_l < u_m.$$

In the first subcase, we assume $u_r < 0$ and the interaction is $(R_+^\uparrow C_\pm^\downarrow) - (N_\pm^\downarrow' C_-^\uparrow')$ with the following states appearing in the strength of the waves

$$\varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_l < u_m. \quad (3.7)$$

Therefore, the change in V is

$$\begin{aligned} [V] &= \sigma(N_\pm^\downarrow') + \sigma(C_-^\uparrow') - \sigma(R_+^\uparrow) - \sigma(C_\pm^\downarrow) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_l)| + |\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)| - |u_m - u_l| - |u_m - \varphi_0^b(u_r)| \\ &= -2|u_m - u_l| = -2\sigma(R_+^\uparrow). \end{aligned}$$

In the second subcase with $0 < u_r$, we have $u_r < \varphi^\#(u_l) < \varphi^b \circ \varphi^b(u_l) < \varphi_0^b \circ \varphi^b(u_l)$ and therefore

$$u_r < \varphi_0^b \circ \varphi^b(u_l) < u_l < u_m. \quad (3.8)$$

The change in V is then

$$\begin{aligned} [V] &= \sigma(N_\pm^\downarrow') + \sigma(C_\mp^\uparrow') - \sigma(R_+^\uparrow) - \sigma(C_\pm^\downarrow) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_l)| + |\varphi_0^b \circ \varphi^b(u_l) - u_r| - |u_m - u_l| - |u_m - u_r| \\ &= -2|u_m - u_l| = -2\sigma(R_+^\uparrow). \end{aligned}$$

Case RN : $(R_+^\uparrow N_\pm^\downarrow) - (N_\pm^\downarrow' R_-^\downarrow')$. The states on both sides of the waves satisfy

$$0 < u_l < u_m \quad \text{and} \quad u_r = \varphi^b(u_m).$$

Two cases again occur depending on the relative order of u_l with respect to $\varphi_0^b(u_r) = \varphi_0^b \circ \varphi^b(u_m)$. We consider only the case where

$$\varphi_0^b \circ \varphi^b(u_l) < \varphi_0^b \circ \varphi^b(u_m) < u_l < u_m, \quad (3.9)$$

the other, $\varphi_0^b \circ \varphi^b(u_l) < u_l < \varphi_0^b \circ \varphi^b(u_m) < u_m$, being similar. In this case, the change in V is

$$\begin{aligned} [V] &= \sigma(N_\pm^\downarrow') + \sigma(R_-^\downarrow') - \sigma(R_+^\uparrow) - \sigma(N_\pm^\downarrow) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_l)| + |\varphi_0^b \circ \varphi^b(u_m) - \varphi_0^b \circ \varphi^b(u_l)| \\ &\quad - |u_m - u_l| - |u_m - \varphi_0^b \circ \varphi^b(u_m)| \\ &= 2|\varphi_0^b \circ \varphi^b(u_m) - \varphi_0^b \circ \varphi^b(u_l)| - 2|u_m - u_l| \\ &\leq -2\underline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b)|u_m - u_l| = -2\underline{\text{Lip}}(u - \varphi_0^b \circ \varphi^b)\sigma(R_+^\uparrow). \end{aligned}$$

Case CR-1 : $(C_\pm^\downarrow R_-^\downarrow) - (C_\pm^\downarrow')$. This is a simple case where the states are initially ordered as

$$\varphi^\#(u_l) < u_r < u_m \leq 0 < u_l.$$

This provides $\varphi_0^b(u_m) < \varphi_0^b(u_r) < \varphi_0^b(\varphi^\#(u_l)) < u_l$. The result is :

$$\begin{aligned} [V] &= \sigma(C_\pm^\downarrow') - \sigma(C_\pm^\downarrow) - \sigma(R_-^\downarrow) \\ &= |u_l - \varphi_0^b(u_r)| - |u_l - \varphi_0^b(u_m)| - |\varphi_0^b(u_r) - \varphi_0^b(u_m)| \\ &= -2|\varphi_0^b(u_r) - \varphi_0^b(u_m)| \leq -2\underline{\text{Lip}}(\varphi_0^b)\sigma(R_-^\downarrow). \end{aligned}$$

Case CR-2 : $(C_{\pm}^{\downarrow} R_{\pm}^{\uparrow})-(C_{\pm}^{\downarrow \prime})$. The waves are entirely classical since $0 \leq u_m < u_r < u_l$. There is nothing new to check and the change in V is immediately found to be

$$[V] = -2|u_r - u_m| = -2\sigma(R_{\pm}^{\uparrow}).$$

Case CR-3 : $(C_{\pm}^{\downarrow} R_{\pm}^{\downarrow})-(N_{\pm}^{\downarrow \prime} R_{\pm}^{\downarrow \prime})$. The states begin in the order

$$u_r \leq \varphi^{\flat}(u_l) < \varphi^{\sharp}(u_l) < u_m \leq 0 < u_l.$$

One subcase is obtained when we assume that $u_l < \varphi_0^{\flat}(u_r)$. The important states appearing in the definition of the strength of waves are then ordered as

$$\varphi_0^{\flat}(u_m) < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < u_l < \varphi_0^{\flat}(u_r). \quad (3.10)$$

The change in V is then found to be

$$\begin{aligned} [V] &= \sigma(N_{\pm}^{\downarrow \prime}) + \sigma(R_{\pm}^{\downarrow \prime}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| + |\varphi_0^{\flat}(u_r) - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| - |u_l - \varphi_0^{\flat}(u_m)| - |\varphi_0^{\flat}(u_r) - \varphi_0^{\flat}(u_m)| \\ &= -2|\varphi_0^{\flat} \circ \varphi^{\flat}(u_l) - \varphi_0^{\flat}(u_m)|. \end{aligned}$$

In the second subcase, when $\varphi_0^{\flat}(u_r) < u_l$, we use

$$\varphi_0^{\flat}(u_m) < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < \varphi_0^{\flat}(u_r) < u_l \quad (3.11)$$

to deduce the same equality :

$$\begin{aligned} [V] &= |u_l - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| + |\varphi_0^{\flat}(u_r) - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| - |u_l - \varphi_0^{\flat}(u_m)| - |\varphi_0^{\flat}(u_r) - \varphi_0^{\flat}(u_m)| \\ &= -2|\varphi_0^{\flat} \circ \varphi^{\flat}(u_l) - \varphi_0^{\flat}(u_m)|. \end{aligned}$$

We now observe that in both subcases, the change is equal to a physically relevant quantity

$$[V] = -2(\sigma(R_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\downarrow \prime})).$$

Case CR-4 : $(C_{\pm}^{\downarrow} R_{\pm}^{\downarrow})-(N_{\pm}^{\downarrow \prime} C_{\pm}^{\uparrow \prime})$. The states of the incoming waves satisfy

$$\varphi^{\flat}(u_l) < u_r < \varphi^{\sharp}(u_l) < u_m \leq 0 < u_l. \quad (3.12)$$

This leads us to the inequalities

$$\varphi_0^{\flat}(u_m) < \varphi_0^{\flat}(u_r) < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < u_l. \quad (3.13)$$

The change in our functional is therefore

$$\begin{aligned} [V] &= \sigma(N_{\pm}^{\downarrow \prime}) + \sigma(C_{\pm}^{\uparrow \prime}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)| + |\varphi_0^{\flat} \circ \varphi^{\flat}(u_l) - \varphi_0^{\flat}(u_r)| - |u_l - \varphi_0^{\flat}(u_m)| - |\varphi_0^{\flat}(u_r) - \varphi_0^{\flat}(u_m)| \\ &= -2|\varphi_0^{\flat}(u_r) - \varphi_0^{\flat}(u_m)| \leq -2\underline{\text{Lip}}(\varphi_0^{\flat})\sigma(R_{\pm}^{\downarrow}). \end{aligned}$$

Case CC-1 : $(C_{\pm}^{\downarrow} C^{\downarrow})-(C^{\downarrow'})$. This is another simple case. We begin with

$$\max(\varphi^{\sharp}(u_l), \varphi^{\sharp}(u_m)) < u_r < u_m < u_l, \quad \text{and} \quad 0 \leq u_m.$$

When $0 \leq u_r$, all the waves are classical and it is easy to show that $[V] = 0$. When $u_r < 0$, we still have $\varphi^{\flat}(u_m) < u_r$ and therefore the important states are ordered

$$\varphi_0^{\flat}(u_r) < u_m < u_l.$$

It then easy to check that even in this case, $[V] = 0$.

Case CC-2 : $(C_{\pm}^{\downarrow} C^{\uparrow})-(C^{\downarrow'})$. This interaction is constrained by the initial states in the following manner

$$\varphi^{\sharp}(u_l) < u_m < u_r < \varphi^{\sharp}(u_m) < u_l, \quad \text{and} \quad u_m < 0.$$

Two subcases appear depending on the sign of u_r . When $u_r > 0$, the interaction is $(C_{\pm}^{\downarrow} C_{\mp}^{\uparrow})-(C_{\mp}^{\downarrow'})$. Since $\varphi^{\flat}(u_l) < u_m$, $\varphi_0^{\flat}(u_m) < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < u_l$. Combining this with the fact that $u_r < \varphi^{\sharp}(u_m) < \varphi_0^{\flat}(u_m)$, we can deduce that the states appearing in $[V]$ are ordered

$$u_r < \varphi_0^{\flat}(u_m) < u_l.$$

The change in V is then

$$\begin{aligned} [V] &= \sigma(C_{\mp}^{\downarrow'}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(C_{\mp}^{\uparrow}) \\ &= |u_l - u_r| - |u_l - \varphi_0^{\flat}(u_m)| - |\varphi_0^{\flat}(u_m) - u_r| = 0. \end{aligned}$$

The second subcase treats $u_r < 0$ and interactions $(C_{\pm}^{\downarrow} C_{\mp}^{\uparrow})-(C_{\pm}^{\downarrow'})$. The states used in our definition of the strength of the waves are

$$\varphi_0^{\flat}(u_r) < \varphi_0^{\flat}(u_m) < u_l.$$

A short computation shows that

$$\begin{aligned} [V] &= \sigma(C_{\pm}^{\downarrow'}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(C_{\mp}^{\uparrow}) \\ &= |u_l - \varphi_0^{\flat}(u_r)| - |u_l - \varphi_0^{\flat}(u_m)| - |\varphi_0^{\flat}(u_m) - \varphi_0^{\flat}(u_r)| = 0. \end{aligned}$$

Case CC-3 : $(C_{\pm}^{\downarrow} C^{\downarrow})-(N_{\pm}^{\downarrow'} C^{\uparrow'})$. This interaction represents a typical transition from one crossing shock to a nonclassical shock. The states are

$$\varphi^{\flat}(u_l) < \varphi^{\sharp}(u_m) < u_r < \varphi^{\sharp}(u_l) < u_m < u_l, \quad 0 \leq u_m.$$

The first (and most common) subcase occurs when $u_r < 0$ and the interaction is $(C_{\pm}^{\downarrow} C_{\pm}^{\downarrow})-(N_{\pm}^{\downarrow'} C_{\mp}^{\uparrow'})$. The first subcase needs to further subdivided into two cases. When $u_m < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l)$, all the important states are ordered

$$\varphi_0^{\flat}(u_r) < u_m < \varphi_0^{\flat} \circ \varphi^{\flat}(u_l) < u_l. \quad (3.14)$$

Under these circumstances, the functional V doesn't change.

$$\begin{aligned} [V] &= \sigma(N_{\pm}^{\downarrow'}) + \sigma(C_{-}^{\uparrow'}) - \sigma(C_{+}^{\downarrow}) - \sigma(C_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_l)| + |\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)| - |u_l - u_m| - |u_m - \varphi_0^b(u_r)| = 0. \end{aligned}$$

On the other hand, when $\varphi_0^b \circ \varphi^b(u_l) < u_m$ in the first subcase, we have

$$\varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_m < u_l. \quad (3.15)$$

It is easy to see that we again have $[V] = 0$.

In the second subcase, $0 < u_r$, we again need to introduce two additional subcases to handle the interactions $(C_{+}^{\downarrow} C_{+}^{\downarrow})$ - $(N_{\pm}^{\downarrow'} C_{\mp}^{\uparrow'})$. When $u_m < \varphi_0^b \circ \varphi^b(u_l)$, then the states used in the definition of wave strengths are

$$u_r < u_m < \varphi_0^b \circ \varphi^b(u_l) < u_l.$$

The change in $[V]$ is

$$\begin{aligned} [V] &= \sigma(N_{\pm}^{\downarrow'}) + \sigma(C_{\mp}^{\uparrow'}) - \sigma(C_{+}^{\downarrow}) - \sigma(C_{+}^{\downarrow}) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_l)| + |\varphi_0^b \circ \varphi^b(u_l) - u_r| - |u_l - u_m| - |u_m - u_r| = 0. \end{aligned}$$

When $0 < u_r$ and $\varphi_0^b \circ \varphi^b(u_l) < u_m$, we obtain the same result.

Case CN-1 : $(C_{+}^{\downarrow} N_{\pm}^{\downarrow})$ - $(C_{\pm}^{\downarrow'})$. The states defining the waves are characterized by the inequalities

$$0 < u_m < u_l \quad \text{and} \quad \varphi^{\#}(u_l) \leq u_r = \varphi^b(u_m).$$

This implies that

$$\varphi_0^b \circ \varphi^b(u_m) < u_m < u_l.$$

We deduce

$$\begin{aligned} [V] &= \sigma(C_{\pm}^{\downarrow'}) - \sigma(C_{+}^{\downarrow}) - \sigma(N_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_m)| - |u_l - u_m| - |u_m - \varphi_0^b \circ \varphi^b(u_m)| = 0. \end{aligned}$$

Case CN-2 : $(C_{\pm}^{\downarrow} N_{\mp}^{\uparrow})$ - $(C_{+}^{\downarrow'})$. We begin with states satisfying

$$\varphi^{\#}(u_l) < u_m < 0 \quad \text{and} \quad u_r = \varphi^b(u_m).$$

To measure the wave strengths, we observe that $\varphi^b(u_l) < u_m$ implies $\varphi_0^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < u_l$ and therefore

$$u_r = \varphi^b(u_m) < \varphi_0^b(u_m) < u_l.$$

The jump in V is now

$$\begin{aligned} [V] &= \sigma(C_{+}^{\downarrow'}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(N_{\mp}^{\uparrow}) \\ &= |u_l - \varphi^b(u_m)| - |u_l - \varphi_0^b(u_m)| - |\varphi_0^b(u_m) - \varphi^b(u_m)| = 0. \end{aligned}$$

Case CN-3 : $(C_{\pm}^{\downarrow} N_{\pm}^{\downarrow})-(N_{\pm}^{\downarrow'} C_{\pm}^{\uparrow'})$. The states are initially ordered as

$$0 < u_m < u_l \quad \text{and} \quad u_r = \varphi^b(u_m) < \varphi^{\sharp}(u_l).$$

A first subcase occurs when $u_m < \varphi_0^b \circ \varphi^b(u_l)$. To compute the change in V , we can then use the inequalities

$$\varphi_0^b \circ \varphi^b(u_m) < u_m < \varphi_0^b \circ \varphi^b(u_l) < u_l, \quad (3.16)$$

to deduce

$$\begin{aligned} [V] &= \sigma(N_{\pm}^{\downarrow'}) + \sigma(C_{\pm}^{\uparrow'}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(N_{\pm}^{\downarrow}) \\ &= |u_l - \varphi_0^b \circ \varphi^b(u_l)| + |\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b \circ \varphi^b(u_m)| \\ &\quad - |u_l - u_m| - |u_m - \varphi_0^b \circ \varphi^b(u_m)| \\ &= 0. \end{aligned}$$

Similarly, if $\varphi_0^b \circ \varphi^b(u_l) < u_m$, then the important inequalities become

$$\varphi_0^b \circ \varphi^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < u_m < u_l, \quad (3.17)$$

and $[V] = 0$.

Case NC : $(N_{\pm}^{\downarrow} C_{\pm}^{\uparrow})-(C_{\pm}^{\downarrow'})$. This interaction is constrained by the states

$$u_m = \varphi^b(u_l) \quad \text{and} \quad \varphi^{\sharp}(u_l) < u_r < \varphi^{\sharp}(u_m) < u_l.$$

The first subcase occurs when $u_r < 0$ and the interaction is precisely $(N_{\pm}^{\downarrow} C_{\pm}^{\uparrow})-(C_{\pm}^{\downarrow'})$. The important states are then ordered as

$$\varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_l.$$

With these observations, we find that

$$\begin{aligned} [V] &= \sigma(C_{\pm}^{\downarrow'}) - \sigma(N_{\pm}^{\downarrow}) - \sigma(C_{\pm}^{\uparrow}) \\ &= |u_l - \varphi_0^b(u_r)| - |u_l - \varphi_0^b \circ \varphi^b(u_l)| - |\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)| = 0. \end{aligned}$$

In the second subcase, $u_r > 0$, it is easy to check that

$$u_r < \varphi_0^b \circ \varphi^b(u_l) < u_l,$$

and $[V] = 0$.

Case NN : $(N_{\pm}^{\downarrow} N_{\pm}^{\uparrow})-(C_{\pm}^{\downarrow'})$. This interaction is the limiting case $u_r \rightarrow \varphi^b(u_m)$ of Case NC. By continuity of wave strengths, we must also have $[V] = 0$. \square

4. Quadratic interaction potential (part 1). In the rest of this paper, we investigate two quadratic interaction potentials, keeping in mind from experience with classical shock waves, that different functionals may be of particular interest in different circumstances. We begin by searching for a functional of the form

$$Q(u(t)) := \sum_{\alpha \text{ approaches } \beta} \sigma(u_l^{\alpha}, u_r^{\alpha}) \sigma(u_l^{\beta}, u_r^{\beta}), \quad (4.1)$$

where the proper definition of “pairs of approaching waves” is essential and is now specified.

In Glimm’s original paper [4] for systems of conservation laws, a definition was proposed which, in the scalar case, amounted to stating that pairs of waves were always approaching unless both are rarefactions. The purpose of this section is to investigate the original definition of Glimm in the context of nonclassical shocks.

DEFINITION 4.1. *A wave α is said to weakly approach a wave β , unless both are rarefaction waves. As far as this definition is concerned, waves $R_+^\uparrow, R_-^\downarrow$ are both to be considered as rarefaction waves and $C_+^\downarrow, C_\pm^\downarrow, N_\pm^\downarrow, C_-^\uparrow, C_\mp^\uparrow, N_\mp^\uparrow$ are all to be considered as shock waves.*

Our main result in the present section is as follows.

THEOREM 4.2 (“Weak interaction” potential for nonclassical shocks). *Let φ^b be a kinetic function satisfying the properties (A1)–(A3). Consider the functional Q_{weak} defined by (4.1) where the summation is made over all weakly interacting waves in the sense of Definition 4.1. Then, when evaluated on a sequence of front-tracking solutions, Q_{weak} is strictly decreasing during all interactions except in the cases RC-3, CR-4, CC-3 and CN-3. In fact, for each of these exceptional interactions, there exist initial data for which $V + C_0 Q_{\text{weak}}$ is increasing during the interaction for every positive C_0 .*

In contrast, in Chapter 8 of [6] in joint work by Baiti, LeFloch, and Piccoli, different definitions of both wave strengths and approaching waves are used and the resulting Glimm functional $V + K Q$ is strictly decreasing for some K . In this sense, the interaction functional Q_{weak} above may appear to be less satisfactory. However, our assumptions on the kinetic function are *completely natural*—a major advantage toward a future extension to systems— and, furthermore, an analysis of “splitting/merging” solutions (in the following section) will show that *globally in time* the functional Q_{weak} does decrease.

Several justifications for our definition of potential are now provided, the strongest argument being the requirement of continuity:

1. Given that V is continuous in $BV(\mathbb{R})$ and that Φ_0^b is already an isometry with respect to this $BV(\mathbb{R})$ norm, it is tempting to assume that any reasonable interaction potential Q should also be continuous in $BV(\mathbb{R})$. We observe that any shock C_\mp^\downarrow can be continuously deformed (as measured by Definition 2.1) into, first, a crossing shock C_\pm^\downarrow and, then, a pair of shocks $N_\pm^\downarrow C_\mp^\uparrow$. It is easy to see that imposing continuity would imply Definition 4.1.
2. Another argument can be found by looking at a class of solutions called splitting/merging solutions, introduced in [9] and discussed further in Section 5. These solutions illustrate that some initial data can go through a nearly periodic process of creation and destruction of nonclassical shocks. In particular, nonclassical shocks can indirectly have non-trivial interactions with shocks on their right-hand side.
3. In [6], nonclassical shocks are precluded from interacting with their right-hand neighbours, and it is argued that nonclassical shocks are (slow) undercompressive and, thus, move away from their right-hand neighbors. However, this definition of approaching waves ignores the above-mentioned possibility of nonclassical shocks having indirect interactions with shocks on their right-hand side. In any case, such an interaction functional then would not be

continuous in $BV(\mathbb{R})$.

For the proof of Theorem 4.2 we proceed as follows. During any isolated interaction between two waves in an approximation, the change in $Q = Q_{\text{weak}}$ is of two types

$$[Q] = [Q]_1 + [Q]_2. \quad (4.2)$$

In this decomposition, $[Q]_1$ denotes the change in the products of the strengths of waves either incoming or exiting the interaction and $[Q]_2$ denotes the change in products of strengths of waves where only *one* of the waves was directly involved in the interaction. Moreover, if a wave C is involved in an interaction, we define

$$W(C) := \sum_{\substack{B \text{ approaches } C \\ B \text{ did not interact}}} \sigma(B).$$

According to Definition 4.1, if the incoming wave C_{in} and the outgoing wave C_{out} are of the same type (i.e. both of rarefaction or shock type), then $W(C_{\text{in}}) = W(C_{\text{out}})$. Theorem 4.2 requires the following lemma which emphasize the (wrong) sign of the terms $[Q]_1$ appearing in the most difficult interactions. The lemma gives a precise characterization of the failings of Q_{weak} for those interactions.

LEMMA 4.3. *The interactions involving a crossing shock, classical or not, and a small wave W_{in} are RC-3, RN, CR-4, CC-3 and CN-3. Suppose that the states neighboring the two consecutive waves are u_l, u_m and u_r . For u_l fixed and as a function of the strength of the incoming small wave, the change in $[Q]_1$ is as follows :*

RN : $[Q]_1 < 0$ for all R_{in} ;

CN-3 : $[Q]_1 < 0$ if C_{in} is sufficiently weak;

RC-3 : $[Q]_1 < 0$ if R_{in} is sufficiently strong;

CR-4, CC-3 : $[Q]_1$ can take on both signs, depending on u_m and u_r .

Moreover, for RC-3 and CR-4 interactions, the largest positive change occurs when $\sigma(R_{\text{in}}) = 0$.

Proof. We distinguish between several cases.

Case RN : $(R_+^\uparrow N_\pm^\downarrow) - (N_\pm^{\downarrow'} R_-^{\downarrow'})$. The states appearing in the shock strengths describe two subcases:

$$\text{either } \varphi_0^b \circ \varphi^b(u_l) < \varphi_0^b \circ \varphi^b(u_m) < u_l < u_m,$$

$$\text{or } \varphi_0^b \circ \varphi^b(u_l) < u_l < \varphi_0^b \circ \varphi^b(u_m) < u_m.$$

For both sets of inequalities $\sigma(R_-^{\downarrow'}) < \sigma(R_+^\uparrow)$ and $\sigma(N_\pm^{\downarrow'}) < \sigma(N_\pm^\downarrow)$. As a result,

$$[Q]_1 = \sigma(N_\pm^{\downarrow'})\sigma(R_-^{\downarrow'}) - \sigma(R_+^\uparrow)\sigma(N_\pm^\downarrow) < 0.$$

Case RC-3 : $(R_+^\uparrow C^\downarrow) - (N_\pm^{\downarrow'} C^{\uparrow'})$. The fundamental states describe two subcases (3.7) and (3.8), here rewritten

$$\text{either } \varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_l < u_m,$$

$$\text{or } u_r < \varphi_0^b \circ \varphi^b(u_l) < u_l < u_m.$$

Focusing on the first case, the change in $[Q]_1$ is

$$\begin{aligned} [Q]_1(u_l, u_m, u_r) &= \sigma(N_{\pm}^{\downarrow'})\sigma(C^{\uparrow'}) - \sigma(R_+^{\uparrow})\sigma(C_{\pm}^{\downarrow}) \\ &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)) - (u_m - u_l) (u_m - \varphi_0^b(u_r)). \end{aligned}$$

Clearly, the RC-3 interaction can still occur even as the strength $\sigma(R_+^{\uparrow}) \rightarrow 0$, therefore the expression above shows that $[Q]_1 > 0$ in that limiting case. In fact, we will show that this is the largest value of $[Q]_1$. We begin by computing

$$\frac{\partial [Q]_1}{\partial u_m} = -(u_m - \varphi_0^b(u_r)) - (u_m - u_l) < 0.$$

Then we evaluate $[Q]_1$ when $u_m = p(u_l)$, where

$$p(u_l) := u_l + (u_l - \varphi_0^b \circ \varphi^b(u_l)) = u_l + \sigma(N_{\pm}^{\downarrow'}),$$

and find

$$\begin{aligned} [Q]_1(u_l, p(u_l), u_r) &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)) \\ &\quad - (u_l - \varphi_0^b \circ \varphi^b(u_l)) (2u_l - \varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)) \\ &= -2(u_l - \varphi_0^b \circ \varphi^b(u_l))^2 = -2\sigma(N_{\pm}^{\downarrow'})^2. \end{aligned}$$

Therefore, for the unbounded set of states satisfying $u_m > p(u_l)$ (i.e. $\sigma(R_+^{\uparrow}) > \sigma(N_{\pm}^{\downarrow'})$), we have $[Q]_1 < 0$.

For fixed u_l , the remaining set of states describing an RC-3 interaction form a bounded set in the (u_m, u_r) plane defined by

$$\begin{aligned} u_m &< p(u_l), \\ \varphi^{\#}(u_m) &< u_r < \varphi^{\#}(u_l). \end{aligned}$$

Along the edge $u_m = p(u_l)$, $[Q]_1$ is negative and since $[Q]_1$ is decreasing with respect to u_m , the largest value must occur along the edge $u_r = \varphi^{\#}(u_m)$. We therefore compute the change in $[Q]_1$ along that edge, using u_m as it's parameter.

$$\begin{aligned} &\frac{d}{du_m} \left([Q]_1(u_l, u_m, \varphi^{\#}(u_m)) \right) \\ &= \frac{d}{du_m} \left((u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b \circ \varphi^{\#}(u_m)) \right. \\ &\quad \left. - (u_m - u_l) (u_m - \varphi_0^b \circ \varphi^{\#}(u_m)) \right) \\ &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) \left(-\frac{d\varphi_0^b \circ \varphi^{\#}}{du}(u_m) \right) \\ &\quad - (u_m - \varphi_0^b \circ \varphi^{\#}(u_m)) - (u_m - u_l) \left(1 - \frac{d\varphi_0^b \circ \varphi^{\#}}{du}(u_m) \right) \end{aligned}$$

Using the inequalities for the states and the fact that $\overline{\text{Lip}}(\varphi_0^b \circ \varphi^{\#}) < \overline{\text{Lip}}(\varphi_0^b \circ \varphi^b) < 1$, we deduce that the terms above are negative and therefore, that $[Q]_1$ must attain it's largest value for the smallest value of u_m , namely when $u_m = u_l = \varphi^{\#}(u_r)$ and $\sigma(R_+^{\uparrow}) = 0$.

Assume now that $\varphi^\sharp(u_l) > 0$ and that u_r could be positive but continue to fix u_l . Then the set of states describing an RC-3 interaction form an infinite strip in the (u_m, u_r) plane defined by

$$u_l < u_m, \quad \text{and} \quad \varphi^b(u_l) < u_r < \varphi^\sharp(u_l),$$

where we have assumed the reasonable fact that $(\varphi^\sharp)' > 0$ at $u_r = \varphi^\sharp(u_l)$ (in fact, since $(\varphi^\sharp)' < 0$ at the origin, we want to assume only one change of sign for $(\varphi^\sharp)'$ and more general cases are similar). The same calculation as above shows that $\partial[Q]_1/\partial u_m < 0$ and therefore that the maximum value of $[Q]_1$ must occur when $u_m = u_l$, i.e. $\sigma(R_{\text{in}}) = 0$.

Case CR-4 : $(C_\pm^\downarrow R_-^\downarrow) - (N_\pm^\downarrow C_-^\uparrow)$. The states satisfy (3.13)

$$\varphi_0^b(u_m) < \varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_l,$$

and therefore

$$\begin{aligned} [Q]_1(u_l, u_m, u_r) &= \sigma(N_\pm^\downarrow C_-^\uparrow) - \sigma(C_\pm^\downarrow) \sigma(R_-^\downarrow) \\ &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)) \\ &\quad - (u_l - \varphi_0^b(u_m)) (\varphi_0^b(u_r) - \varphi_0^b(u_m)). \end{aligned}$$

If we let u_r and u_m approach each other while maintaining the condition

$$u_r < \varphi^\sharp(u_l) < u_m,$$

then $\varphi_0^b(u_r) - \varphi_0^b(u_m) \rightarrow 0$ while $u_l - \varphi_0^b(u_m)$ remains bounded. Upon inspection, it is clear that in this limit, $[Q]_1 \rightarrow \sigma(N_\pm^\downarrow C_-^\uparrow)$. We now demonstrate that this is the largest value of $[Q]_1$. When u_l is fixed, the admissible states belong to a rectangular domain

$$(u_m, u_r) \in [\varphi^\sharp(u_l), 0] \times [\varphi^b(u_l), \varphi^\sharp(u_l)].$$

With respect to u_m and u_r , we find

$$\frac{\partial[Q]_1}{\partial u_m} = \frac{d\varphi_0^b}{du}(u_m) \left((\varphi_0^b(u_r) - \varphi_0^b(u_m)) + (u_l - \varphi_0^b(u_m)) \right) < 0,$$

and

$$\frac{\partial[Q]_1}{\partial u_r} = -\frac{d\varphi_0^b}{du}(u_r) \left((u_l - \varphi_0^b \circ \varphi^b(u_l)) + (u_l - \varphi_0^b(u_m)) \right) > 0.$$

Taken together, these inequalities imply that the largest value of $[Q]_1$ occurs at the corner of the domain where $u_m = u_r = \varphi^\sharp(u_l)$, i.e. when $\sigma(R_-^\downarrow) = 0$. On the other hand, the smallest value would occur along the boundary $u_r = \varphi^b(u_l)$, where one finds a negative quantity

$$\begin{aligned} [Q]_1 &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b \circ \varphi^b(u_l)) \\ &\quad - (u_l - \varphi_0^b(u_m)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_m)). \end{aligned}$$

Case CC-3 : $(C_+^\downarrow C^\downarrow)$ - $(N_\pm^\downarrow C^\uparrow')$. As mentioned in the proof of Theorem 3.1, the set of states are subdivided into two subcases depending on the sign of u_r . When $u_r < 0$, we identify two possibilities (3.14) and (3.15), namely

$$\begin{aligned} &\text{either } \varphi_0^b(u_r) < u_m < \varphi_0^b \circ \varphi^b(u_l) < u_l, \\ &\text{or } \varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_m < u_l, \end{aligned}$$

where the second set of inequalities corresponds to a weak C_+^\downarrow . The change in $[Q]_1$ is then

$$\begin{aligned} [Q]_1(u_l, u_m, u_r) &= \sigma(N_\pm^\downarrow) \sigma(C_-^\uparrow') - \sigma(C_+^\downarrow) \sigma(C_\pm^\downarrow) \\ &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b(u_r)) \\ &\quad - (u_l - u_m) (u_m - \varphi_0^b(u_r)). \end{aligned}$$

Consider the function $B(\lambda) := \lambda(1 - \lambda)$ whose maximum on the interval $[0, 1]$ is attained at $\lambda = 1/2$. There exists constants C, λ and λ' such that the change can be rewritten as

$$[Q]_1 = C (B(\lambda') - B(\lambda)).$$

It is therefore clear that $[Q]_1$ will be negative if and if $|\lambda - 1/2| < |\lambda' - 1/2|$. Unfortunately, this set cannot be described in a simple manner in terms of the strengths of the waves.

Case CN-3 : $(C_+^\downarrow N_\pm^\downarrow)$ - $(N_\pm^\downarrow C_-^\uparrow')$. We identify two subcases

$$\begin{aligned} &\text{either } \varphi_0^b \circ \varphi^b(u_m) < u_m < \varphi_0^b \circ \varphi^b(u_l) < u_l, \\ &\text{or } \varphi_0^b \circ \varphi^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < u_m < u_l, \end{aligned}$$

where the second one occurs when the incoming shock C_+^\downarrow is weak. The change can be written as

$$\begin{aligned} [Q]_1(u_l, u_m, u_r) &= \sigma(N_\pm^\downarrow) \sigma(C_-^\uparrow') - \sigma(C_+^\downarrow) \sigma(N_\pm^\downarrow) \\ &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b \circ \varphi^b(u_m)) \\ &\quad - (u_l - u_m) (u_m - \varphi_0^b \circ \varphi^b(u_m)). \end{aligned}$$

Using a bit of algebra, we rewrite the change as

$$\begin{aligned} [Q]_1 &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b \circ \varphi^b(u_m)) \\ &\quad - (u_l - \varphi_0^b \circ \varphi^b(u_l) + \varphi_0^b \circ \varphi^b(u_l) - u_m) (u_m - \varphi_0^b \circ \varphi^b(u_m)) \\ &= (u_l - \varphi_0^b \circ \varphi^b(u_l)) (\varphi_0^b \circ \varphi^b(u_l) - u_m) \\ &\quad + (u_m - \varphi_0^b \circ \varphi^b(u_l)) (u_m - \varphi_0^b \circ \varphi^b(u_m)) \\ &= - (u_m - \varphi_0^b \circ \varphi^b(u_l)) ((u_l - u_m) - (\varphi_0^b \circ \varphi^b(u_l) - \varphi_0^b \circ \varphi^b(u_m))), \end{aligned} \quad (4.3)$$

thus concluding that $[Q]_1 < 0$ as long as C_+^\downarrow is weak. \square

Proof of Theorem 4.2. Throughout the proof, we use liberally the estimates derived in the proof of Theorem 3.1.

Case RC-1 : $(R_+^\uparrow C^\downarrow)-(C^{\downarrow'})$. After examining (3.5), it is immediate that $\sigma(C^\downarrow) > \sigma(C^{\downarrow'})$ and $W(C^\downarrow) = W(C^{\downarrow'})$. Therefore,

$$[Q]_1 = -\sigma(R_+^\uparrow)\sigma(C^\downarrow) < 0$$

and

$$\begin{aligned} [Q]_2 &= W(C^{\downarrow'})\sigma(C^{\downarrow'}) - W(R_+^\uparrow)\sigma(R_+^\uparrow) - W(C^\downarrow)\sigma(C^\downarrow) \\ &= -W(R_+^\uparrow)\sigma(R_+^\uparrow) - W(C^\downarrow)(\sigma(C^\downarrow) - \sigma(C^{\downarrow'})) < 0. \end{aligned}$$

Case RC-2 : $(R_+^\uparrow C_\pm^\downarrow)-(N_\pm^{\downarrow'} R_-^{\downarrow'})$. The conditions defining this case imply two subcases :

$$\begin{aligned} &\text{either } \varphi_0^b \circ \varphi^b(u_l) < \varphi_0^b(u_r) < u_l < u_m, \\ &\text{or } \varphi_0^b \circ \varphi^b(u_l) < u_l < \varphi_0^b(u_r) < u_m. \end{aligned}$$

Given that $\varphi^\sharp(u_m) < u_r < 0$, then property (2.1) leads to

$$0 < \varphi_0^b(u_r) < \varphi_0^b \circ \varphi^\sharp(u_m) < \varphi_0^b \circ \varphi^b(u_m) < u_m.$$

This means that in the first subcase,

$$\sigma(R_-^{\downarrow'}) = |\varphi_0^b(u_r) - \varphi_0^b \circ \varphi^b(u_l)| < |\varphi_0^b \circ \varphi^b(u_m) - \varphi_0^b \circ \varphi^b(u_l)| < |u_m - u_l| = \sigma(R_+^\uparrow).$$

In that subcase, we also have

$$\begin{aligned} \sigma(C_\pm^\downarrow) &= \sigma(R_+^\uparrow) + |u_l - \varphi_0^b(u_r)|, \\ \sigma(N_\pm^{\downarrow'}) &= \sigma(R_-^{\downarrow'}) + |u_l - \varphi_0^b(u_r)|, \end{aligned}$$

then $\sigma(N_\pm^{\downarrow'}) < \sigma(C_\pm^\downarrow)$. Similar arguments prove the same inequalities in the second case.

These inequalities therefore imply that

$$[Q]_1 = \sigma(N_\pm^{\downarrow'})\sigma(R_-^{\downarrow'}) - \sigma(C_\pm^\downarrow)\sigma(R_+^\uparrow) < 0.$$

Using the proposed definition of *weakly approaching waves*, we also deduce

$$\begin{aligned} [Q]_2 &= W(N_\pm^{\downarrow'})\sigma(N_\pm^{\downarrow'}) + W(R_-^{\downarrow'})\sigma(R_-^{\downarrow'}) - W(R_+^\uparrow)\sigma(R_+^\uparrow) - W(C_\pm^\downarrow)\sigma(C_\pm^\downarrow) \\ &= -W(R_-^{\downarrow'})\sigma(R_+^\uparrow) - W(N_\pm^{\downarrow'})\sigma(C_\pm^\downarrow) < 0. \end{aligned}$$

Case RC-3 : $(R_+^\uparrow C^\downarrow)-(N_\pm^{\downarrow'} C^{\uparrow'})$. Two subcases occur depending on the sign of u_r . If $u_r < 0$, then (3.7) holds and when $u_r > 0$, then (3.8) holds. In both of these cases

$$\sigma(C^\downarrow) - \sigma(R_+^\uparrow) = \sigma(C^{\uparrow'}) + \sigma(N_\pm^{\downarrow'}). \quad (4.4)$$

In Lemma 4.3, we already showed that $[Q]_1$ could be positive. For the other interaction term, we use (4.4) to verify

$$\begin{aligned} [Q]_2 &= W(N_\pm^{\downarrow'})\sigma(N_\pm^{\downarrow'}) + W(C^{\uparrow'})\sigma(C^{\uparrow'}) - W(R_+^\uparrow)\sigma(R_+^\uparrow) - W(C^\downarrow)\sigma(C^\downarrow) \\ &= -W(N_\pm^{\downarrow'})\sigma(C^\downarrow) - W(R_+^\uparrow)\sigma(R_+^\uparrow) < 0. \end{aligned}$$

Case RN : $(R_+^\uparrow N_\pm^\downarrow)-(N_\pm^\downarrow R_-^\downarrow')$. The change $[Q]_1$ was studied in Lemma 4.3. For the second term, using the bounds on the wave strengths given in the previous lemma, we again have a negative contribution

$$\begin{aligned} [Q]_2 &= W(N_\pm^\downarrow')\sigma(N_\pm^\downarrow) + W(R_-^\downarrow')\sigma(R_-^\downarrow') - W(R_+^\uparrow)\sigma(R_+^\uparrow) - W(N_\pm^\downarrow)\sigma(N_\pm^\downarrow) \\ &= -W(N_\pm^\downarrow)(\sigma(N_\pm^\downarrow) - \sigma(N_\pm^\downarrow')) - W(R_+^\uparrow)(\sigma(R_+^\uparrow) - \sigma(R_-^\downarrow')) < 0. \end{aligned}$$

Case CR-1 : $(C_\pm^\downarrow R_-^\downarrow)-(C_\pm^\downarrow')$. In this case, the states satisfy

$\varphi_0^b(u_m) < \varphi_0^b(u_r) < \varphi_0^b(\varphi^\#(u_l)) < u_l$ and the wave strengths satisfy $\sigma(C_\pm^\downarrow') < \sigma(C_\pm^\downarrow)$. Since only one wave is outgoing, $[Q]_1 = -\sigma(C_\pm^\downarrow)\sigma(R_-^\downarrow) < 0$. On the other hand, it is easy to check that

$$\begin{aligned} [Q]_2 &= W(C_\pm^\downarrow')\sigma(C_\pm^\downarrow') - W(C_\pm^\downarrow)\sigma(C_\pm^\downarrow) - W(R_-^\downarrow)\sigma(R_-^\downarrow) \\ &= -W(C_\pm^\downarrow)(\sigma(C_\pm^\downarrow) - \sigma(C_\pm^\downarrow')) - W(R_-^\downarrow)\sigma(R_-^\downarrow) < 0. \end{aligned}$$

Case CR-2 : $(C_+^\downarrow R_+^\uparrow)-(C_+^\downarrow')$. This case is entirely classical so it is easy to check that

$$\begin{aligned} [Q]_1 &= -\sigma(C_+^\downarrow)\sigma(R_+^\uparrow) < 0, \\ [Q]_2 &= -W(C_+^\downarrow)(\sigma(C_+^\downarrow) - \sigma(C_+^\downarrow')) - W(R_+^\uparrow)\sigma(R_+^\uparrow) < 0. \end{aligned}$$

Case CR-3 : $(C_\pm^\downarrow R_-^\downarrow)-(N_\pm^\downarrow' R_-^\downarrow')$. A first subcase is defined by the additional condition $u_l < \varphi^b(u_r)$ which provides (3.10)

$$\varphi_0^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < u_l < \varphi_0^b(u_r). \quad (4.5)$$

The relative strengths of the waves are then seen to be $\sigma(R_-^\downarrow') < \sigma(R_-^\downarrow)$ and $\sigma(N_\pm^\downarrow') < \sigma(C_\pm^\downarrow)$. In the second subcase, given by (3.11), these two inequalities still hold. It is now easy to conclude

$$\begin{aligned} [Q]_1 &= \sigma(N_\pm^\downarrow')\sigma(R_-^\downarrow') - \sigma(C_\pm^\downarrow)\sigma(R_-^\downarrow) < 0, \\ [Q]_2 &= -W(C_\pm^\downarrow)(\sigma(C_\pm^\downarrow) - \sigma(N_\pm^\downarrow')) - W(R_-^\downarrow)(\sigma(R_-^\downarrow) - \sigma(R_-^\downarrow')) < 0. \end{aligned}$$

Case CR-4 : $(C_\pm^\downarrow R_-^\downarrow)-(N_\pm^\downarrow' C_-^\uparrow')$. The inequalities (3.13) defining this case suffice to show that $\sigma(N_\pm^\downarrow') + \sigma(C_-^\uparrow') \leq \sigma(C_\pm^\downarrow)$. In Lemma 4.3, we showed that $[Q]_1 > 0$ but the second term is negative

$$[Q]_2 = -W(C_\pm^\downarrow)(\sigma(C_\pm^\downarrow) - \sigma(N_\pm^\downarrow') - \sigma(C_-^\uparrow')) - W(R_-^\downarrow)\sigma(R_-^\downarrow) < 0.$$

Case CC-1 : $(C_+^\downarrow C_+^\downarrow)-(C_+^\downarrow')$. When u_r is positive, all waves are classical and it is easy to show that $[Q]_1 < 0$ and $[Q]_2 = 0$. It is an exercise to see that when $u_r < 0$, then $[Q]_1 < 0$ is still negative and $[Q]_2$ vanishes.

Case CC-2 : $(C_\pm^\downarrow C^\uparrow)-(C_\pm^\downarrow')$. Two subcases appear depending on the sign of u_r but each time,

$$[V] = \sigma(C_\pm^\downarrow') - \sigma(C_\pm^\downarrow) - \sigma(C^\uparrow) = 0.$$

Only one wave is outgoing, so $[Q]_1 < 0$ and the previous identity implies that $[Q]_2 = 0$.

Case CC-3 : $(C_+^\downarrow C^\downarrow)-(N_\pm^\downarrow C^\uparrow)$. Since all waves involved are shocks and $[V] = 0$, then $[Q]_2 = 0$.

Case CN-1 : $(C_+^\downarrow N_\pm^\downarrow)-(C_\pm^\downarrow)$. This is another simple case where the fact that $[V] = 0$ and that $W(\cdot)$ is equal for all waves involved, implies $[Q]_2 = 0$.

Case CN-2 : $(C_+^\downarrow N_\mp^\uparrow)-(C_+^\downarrow)$. Same as Case CN-1.

Case CN-3 : $(C_+^\downarrow N_\pm^\downarrow)-(N_\pm^\downarrow C^\uparrow)$. Two subcases occur but each time $[V] = 0$. All waves are shocks so $[V] = 0$ implies that $[Q]_2 = 0$.

Case NC : $(N_\pm^\downarrow C^\uparrow)-(C^\downarrow)$. Same as Case CN-1.

Case NN : $(N_\pm^\downarrow N_\mp^\uparrow)-(C_+^\downarrow)$. Same as Case CN-1. □

5. Global diminishing property for splitting-merging patterns. In this section we show that, despite the fact that the quadratic interaction potential Q_{weak} increases during interactions of type CR-3, RC-4, CN-3 and CC-3, this potential is indeed *strictly decreasing* globally in time for a large class of perturbations of crossing shocks. Hence, with the local bound we describe below, we provide the first steps towards an analysis of the global-in-time change of Q_{weak} for arbitrary nonclassical entropy solutions. This section, therefore, provides a strong justification for the potential proposed in the previous section.

The *splitting-merging solutions* considered now were introduced in LeFloch and Shearer [9], where a modification of the total variation functional [6] was shown to be strictly decreasing along the evolution of such splitting-merging solutions. The total variation functional V presented in Section 3 also accomplishes this, but here we improve on those results by establishing a similar monotonicity result for the quadratic functional Q_{weak} . Our analysis also brings to light some interesting aspects of splitting-merging solutions that were not seen in [9].

Splitting-merging solutions are, roughly speaking, perturbations of crossing shocks that lead to the creation and destruction of a nonclassical shock. Such solutions contain two (classical and nonclassical) big waves that may merge together (as a classical shock) and also interact with (classical) small waves. A typical initial data for splitting-merging patterns is formed of

- (i) an isolated crossing shock with left- and right-hand states u_-, u_+ satisfying $u_- > 0$ and $\varphi^\sharp(u_-) < u_+$, but $u_+ - \varphi^\sharp(u_-)$ small,
- (ii) followed, on the right-hand side, by a small rarefaction and a small shock.

The rarefaction is sufficiently strong that it has an interaction of type CR-4 with the crossing shock, thereby leading to the creation of a pair of shock waves $N_\pm^\downarrow, C^\uparrow$. If the right-most shock is sufficiently strong, then when it eventually interacts with C^\uparrow and the resulting shock will begin to approach N_\pm^\downarrow . The final interaction of type NC will involve N_\pm^\downarrow and the shock just described, thereby eliminating the nonclassical N_\pm^\downarrow . By adding more waves to the left and the right, this process of creation and destruction of N_\pm^\downarrow can be repeated indefinitely.

We consider a slightly more general configuration in the sense that we do not explicitly demand that a small shock on the right be responsible for the penultimate

NC interaction. Fix some value $u^* > 0$ and define

$$u_0^*(x) = \begin{cases} u^*, & x < 0, \\ \varphi^\sharp(u^*), & x > 0. \end{cases} \quad (5.1)$$

Let θ_ϵ be some function of locally bounded total variation and of oscillation bounded by some small positive ϵ , i.e.

$$\sigma(\theta_\epsilon(x), 0) < \epsilon, \quad x \in \mathbb{R}.$$

Without loss of generality, we may assume that θ_ϵ is piecewise constant. Let u_ϵ^* be the nonclassical solution to the conservation law (1.1) with initial data $u_0^* + \theta_\epsilon$, as generated by the front-tracking method. Assuming the solution initially possesses a single isolated crossing shock located at the origin $x = 0$, that is, assuming that $u_\epsilon^*(0-) > 0$, we see that the crossing shock will be adjacent to many small classical shocks and rarefactions. After an interaction of type RC-3, CR-4, or CC-3, the small waves neighboring C_\pm^\downarrow may lead to the creation of a pair of waves, a nonclassical shock N_\pm^\downarrow and a classical shock C^\uparrow . After the creation of N_\pm^\downarrow , the only types of interaction involving small waves incoming from the left of N_\pm^\downarrow are RN and CN-3. The only types of interaction involving small waves and the shock C^\uparrow , coming from either the left or the right, are entirely classical (CC-1, RC-1 or CR-1). Moreover, no waves can cross C^\uparrow from the right or the left although the small waves that reach N_\pm^\downarrow from the left, will cross and eventually reach C^\uparrow . Therefore, the only way that the nonclassical shock N_\pm^\downarrow can be destroyed is if the shocks N_\pm^\downarrow and C^\uparrow change their speeds and eventually interact back together, leading us back to (a perturbation of) the original crossing shock.

These observations, in particular, imply that *no waves can exit the domain Ω* bounded by the trajectories of N_\pm^\downarrow and C^\uparrow . Our goal, in the present section, is to obtain a local bound on the change of the potential Q_{weak} relative only to the waves entering the domain Ω . It should already be clear that the key here is comparing the total strength of the waves crossing N_\pm^\downarrow to the total strength of the waves terminating at C^\uparrow .

Before stating our main result, we introduce some further notation. Let t_0 be a time of creation of a nonclassical wave N_\pm^\downarrow and denote by t_1, t_2, \dots, t_m the times of the next m interactions between N_\pm^\downarrow and small waves W_i on the right, and let t_{m+1} be the time at which N_\pm^\downarrow is destroyed from an interaction with the shock C^\uparrow . Similarly, let \tilde{t}_i and \bar{t}_i be the times at which an interaction occurs between C^\uparrow and the left incoming waves \widetilde{W}_i or the right incoming waves \overline{W}_i , respectively. We define the *total variation along the trajectory N_\pm^\downarrow* to be

$$\text{TV}(N_\pm^\downarrow) := \sum_{i=1}^m \left| \sigma(N_\pm^\downarrow(t_i+)) - \sigma(N_\pm^\downarrow(t_i-)) \right|, \quad (5.2)$$

and its *signed variation*

$$\text{SV}(N_\pm^\downarrow) := \sigma(N_\pm^\downarrow(t_{m+1}-)) - \sigma(N_\pm^\downarrow(t_0+)). \quad (5.3)$$

Completely similar definitions also apply to the wave C^\uparrow , but an additional decompo-

sition exists by separating the contributions from the left- and the right-hand sides:

$$\begin{aligned} \text{SV}(C^\dagger) &= \left(\sum_{i=1}^{\tilde{m}} \tilde{s}_i \sigma(\tilde{W}_i) \right) + \left(\sum_{i=1}^{\bar{m}} \bar{s}_i \sigma(\bar{W}_i) \right) \\ &=: \text{SV}_L(C^\dagger) + \text{SV}_R(C^\dagger), \end{aligned}$$

where \tilde{s}_i (\bar{s}_i) is $+1$ if \tilde{W}_i (\bar{W}_i) is a shock and -1 otherwise. For convenience, the strengths of the small wave W_i , before and after it has crossed N_\pm^\dagger at some time t_i , are denoted by W_i^- and W_i^+ , respectively.

THEOREM 5.1 (Global diminishing property for splitting-merging patterns). *Let φ^b be a kinetic function satisfying the properties (A1)–(A3), together with the mild requirement at some given reference state $u^* > 0$.*

$$L^* := (\varphi_0^b \circ \varphi^b)'(u^*) \in (1/2, 1).$$

Let u_ϵ^ be the nonclassical solution to the conservation law (1.1) with initial data $u_0^* + \theta_\epsilon$, where u_0^* is defined in (5.1) and the perturbation θ_ϵ is of locally bounded variation and of small amplitude, that is, $\|\theta_\epsilon\|_{L^\infty(\mathbb{R})} < \epsilon$. Suppose that u_ϵ^* exhibits a splitting-merging pattern on the time interval $[t_0, t_{m+1}]$ along successive interactions with small waves (cf. the notation above). Provided ϵ is sufficiently small and the total effect of all waves on the classical shock C^\dagger increases its total strength, that is,*

$$\text{SV}(C^\dagger) \geq 0,$$

then the variation of the potential $Q = Q_{\text{weak}}$ within the domain Ω bounded by the trajectories of N_\pm^\dagger and C^\dagger is negative

$$[Q]_1|_\Omega < 0.$$

Recall that the expression $[Q]_1$ was defined earlier in Section 4. In the above theorem, only contributions within the domain Ω are considered. Two important remarks about our assumptions are in order:

1. The condition on L^* is satisfied for a wide range of fluxes, entropies and nonlinear diffusion-dispersion models, see Theorems 2.3 and 3.3 in [6]. In particular, it is true for the cubic flux $f(u) = u^3 - au$, the entropy $U(u) = u^2/2$ and all ratios of diffusion over dispersion. The non-trivial aspect of this condition is the lower bound $L^* > 1/2$ since the upper bound is always satisfied.
2. Our main assumption $\text{SV}(C^\dagger) \geq 0$ requires that the total effect of the interaction of all waves on C^\dagger is to increase its strength. In fact, this is always the case for the perturbations of splitting-merging solutions within the setting [9]. In our slightly more general setting though, waves crossing through N_\pm^\dagger will change the critical state $\varphi^\sharp(u_l)$ and therefore could conceivably lead to an NC interaction even if C^\dagger interacted only with rarefactions on the right ($\text{SV}(C^\dagger) < 0$). Intuitively, one would like to show that the existence of an NC interaction at time t_{m+1} implies that the wave C became strong enough to change course and therefore that $\text{SV}(C^\dagger) \geq 0$ is satisfied. This is the subject of ongoing research.

It is interesting to check that the condition $\text{SV}(C^\uparrow) \geq 0$ of Theorem 5.1 is in fact necessary, by studying a simple situation involving a single rarefaction R crossing the nonclassical shock N_\pm^\downarrow , and one classical shock C' interacting from the right with C^\uparrow (assuming that φ^\sharp is monotonically decreasing, say). Suppose that the strength of the rarefaction wave after crossing N_\pm^\downarrow is ν_R and that the change in N_\pm^\downarrow during this interaction is ν_L . In Lemma 5.2 below, we will check that $\nu_L < \nu_R$ and that $\nu_L + \nu_R$ is the strength of the original rarefaction. So, assuming that C' interacts with C^\uparrow before the rarefaction does, then the total change in Q is given by :

$$\begin{aligned}
[Q]_1|_\Omega = & + \sigma(N_\pm^\downarrow)\sigma(C^\uparrow) && \text{(due to CR-4),} \\
& - \nu_L(\sigma(N_\pm^\downarrow) + \nu_R) && \text{(due to RN),} \\
& - \sigma(C^\uparrow)\sigma(C') && \text{(due to CC-1),} \\
& - (\sigma(C^\uparrow) + \sigma(C'))\nu_R && \text{(due to RC-1),} \\
& - (\sigma(N_\pm^\downarrow) - \nu_L)(\sigma(C^\uparrow) + \sigma(C') - \nu_R) && \text{(due to NC).}
\end{aligned}$$

The last term can be rewritten as

$$-\sigma(N_\pm^\downarrow)\sigma(C^\uparrow) - \sigma(N_\pm^\downarrow)(\sigma(C') - \nu_R) + \nu_L(\sigma(C^\uparrow) + \sigma(C')) - \nu_L\nu_R.$$

The contribution from the RC-1 interaction can control the third term. Unfortunately, if ν_R is large with respect to $\sigma(C')$, i.e. $\text{SV}(C') < 0$, then the CC-1 term cannot be used to control the second term.

For the proof of Theorem 5.1 we will need the following two lemmas. The interest of the first lemma is to make more precise the (mainly linear) dependence of the change $[Q]_1$ in terms of the incoming wave $\sigma(W_i^-)$. The second lemma provides an estimate which closely relates the signed variations of C^\uparrow and N_\pm^\downarrow .

LEMMA 5.2 (Interactions with the nonclassical shock). *Consider interactions RN and CN-3 at the time t_i involving a left-incoming weak wave W_i and the nonclassical shock N_\pm^\downarrow . Then, there exists a positive constant $L_i < 1$ such that*

$$\begin{aligned}
\sigma(W_i^+) &= L_i\sigma(W_i^-), \\
\sigma(N_\pm^\downarrow(t_i+)) &= \sigma(N_\pm^\downarrow(t_i-)) + s_i(1 - L_i)\sigma(W_i^-), \\
[Q]_1 &= -(1 - L_i)\sigma(W_i^-) \left(\sigma(N_\pm^\downarrow(t_i-)) - s_i L_i \sigma(W_i^-) \right) < 0,
\end{aligned}$$

where s_i is $+1$ if W_i is a shock and -1 otherwise. Moreover, one has

$$L_i = L^* + \mathcal{O}(\epsilon). \quad (5.4)$$

Proof. We consider only a RC-3 interaction, since the calculations for an incoming rarefaction are similar and have been essentially treated in Lemma 4.3, equation (4.3). When W_i is small, then the states are ordered as in (3.17), namely

$$\varphi_0^b \circ \varphi^b(u_m^i) < \varphi_0^b \circ \varphi^b(u_i^i) < u_m^i < u_i^i.$$

Then, we have

$$\sigma(W_i^+) = \left| \varphi_0^b \circ \varphi^b(u_i^i) - \varphi_0^b \circ \varphi^b(u_m^i) \right| =: L_i |u_i^i - u_m^i| = L_i \sigma(W_i^-),$$

with $L_i < 1$ since $\varphi_0^b \circ \varphi^b$ is a contraction. On the other hand, relation (5.4) is obvious since the u_m^i, u_i^i belong to a neighborhood of u^* of size ϵ .

Finally, the outgoing nonclassical shock has strength

$$\begin{aligned} \sigma(N_{\pm}^{\downarrow}(t_i+)) &= |u_i^i - \varphi_0^b \circ \varphi^b(u_i^i)| \\ &= |u_i^i - u_m^i| + |u_m^i - \varphi_0^b \circ \varphi^b(u_m^i)| - \left| \varphi_0^b \circ \varphi^b(u_i^i) - \varphi_0^b \circ \varphi^b(u_m^i) \right| \\ &= \sigma(N_{\pm}^{\downarrow}(t_i-)) + (1 - L_i)\sigma(W_i^-), \end{aligned}$$

while the change $[Q]_1$ takes the form

$$\begin{aligned} [Q]_1 &= \sigma(N_{\pm}^{\downarrow}(t_i+))\sigma(W_i^+) - \sigma(N_{\pm}^{\downarrow}(t_i-))\sigma(W_i^-) \\ &= \left(\sigma(N_{\pm}^{\downarrow}(t_i-)) + (1 - L_i)\sigma(W_i^-) \right) L_i\sigma(W_i^-) - \sigma(N_{\pm}^{\downarrow}(t_i-))\sigma(W_i^-) \\ &= -(1 - L_i)\sigma(W_i^-) \left(\sigma(N_{\pm}^{\downarrow}(t_i-)) - L_i\sigma(W_i^-) \right) < 0. \end{aligned}$$

□

LEMMA 5.3 (Property of the signed variations). *Setting $\lambda^* := (1 - L^*)/L^*$, one has*

$$\left| \lambda^* \text{SV}_L(C^\uparrow) - \text{SV}(N_{\pm}^{\downarrow}) \right| \leq \mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow}). \quad (5.5)$$

Proof. Using Lemma 5.2, we compute

$$\begin{aligned} \text{SV}(N_{\pm}^{\downarrow}) &= \sum_{i=1}^m s_i(1 - L_i)\sigma(W_i^-) \\ &= \sum_{i=1}^m s_i(1 - L^*)\sigma(W_i^-) + \sum_{i=1}^m s_i(L^* - L_i)\sigma(W_i^-) \\ &= \lambda^* \sum_{i=1}^m s_i L^* \sigma(W_i^-) + \sum_{i=1}^m s_i(L^* - L_i)\sigma(W_i^-), \end{aligned}$$

thus

$$\text{SV}(N_{\pm}^{\downarrow}) = \lambda^* \text{SV}_L(C^\uparrow) + (1 + \lambda^*) \sum_{i=1}^m s_i(L^* - L_i)\sigma(W_i^-).$$

For each index i , $L^* - L_i = \mathcal{O}_i(\epsilon)$, so if we take $\mathcal{O}(\epsilon) = \max_i \mathcal{O}_i(\epsilon)$ and use $\underline{L} := \min_i L_i$, then the difference is

$$\begin{aligned} \left| \lambda^* \text{SV}_L(C^\uparrow) - \text{SV}(N_{\pm}^{\downarrow}) \right| &\leq 2\mathcal{O}(\epsilon) \sum_{i=1}^m |\sigma(W_i^-)| \\ &\leq \frac{2}{\underline{L}} \mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow}) = \mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow}). \end{aligned}$$

□

Proof of Theorem 5.1. To help the reader understand the wave interactions (and cancellations) in this proof, we begin with a few preliminary comments. Simply put, the final NC interaction should provide a quadratic term which is, later, cancelled by a similar quadratic term when the two waves merge. One may naively expect the sum

$$-\sigma(N_{\pm}^{\downarrow}(t_{m+1}-))\sigma(C^{\uparrow}(t_{m+1}-)) + \sigma(N_{\pm}^{\downarrow}(t_0+))\sigma(C^{\uparrow}(t_0+))$$

to be negative. Of course, the cumulative strength of the changes during the interactions with the small waves must also be taken into account. The difference between the strength of the initial and final waves $N_{\pm}^{\downarrow}, C^{\uparrow}$ is measured by the signed variation SV along those two shocks. Our proof below shows that, along the trajectories N_{\pm}^{\downarrow} and C^{\uparrow} , the change $[Q]_1$ is negative and proportional to the total variation $\text{TV}(N_{\pm}^{\downarrow}) + \text{TV}(C^{\uparrow})$. The total variation being larger than the signed variation, after further analysis one can conclude that $[Q]_1|_{\Omega} < 0$.

The key technical information is provided by Lemma 5.3, which implies that, up to a quantity of order $\mathcal{O}(\epsilon)\text{TV}(N_{\pm}^{\downarrow})$, we have $\text{SV}(N_{\pm}^{\downarrow}) = \lambda^* \text{SV}_L(C^{\uparrow})$ and, in particular, that the signed variations have the same sign. We now have all the tools necessary to proceed with the proof of Theorem 5.1.

The perturbation θ_{ϵ} has bounded oscillation and therefore it can only alter the right-hand state of C^{\uparrow} by an amount ϵ . The small waves entering Ω through N_{\pm}^{\downarrow} only alter its strength by $|(1 - L_i)\sigma(W_i^-)| < \epsilon$. In both cases, we expect that everywhere along their trajectories,

$$\begin{aligned}\sigma(N_{\pm}^{\downarrow}(t)) &> \sigma(N_{\pm}^{\downarrow}(t_0+)) - 2\epsilon, \\ \sigma(C^{\uparrow}(t)) &> \sigma(C^{\uparrow}(t_0+)) - 2\epsilon.\end{aligned}$$

Ignoring the negative contribution to $[Q]_1$ coming from the interaction that generated N_{\pm}^{\downarrow} (which, anyway, can be arbitrarily small), and neglecting also all classical interactions inside Ω (for which $[Q]_1 \leq 0$ and possibly 0), we have

$$\begin{aligned}[Q]_1\Big|_{\Omega} &\leq +\sigma(N_{\pm}^{\downarrow}(t_0+))\sigma(C^{\uparrow}(t_0+)) \\ &\quad - \sum_{i=1}^m (1 - L_i)\sigma(W_i^-) \left(\sigma(N_{\pm}^{\downarrow}(t_i-)) - s_i L_i \sigma(W_i^-) \right) \\ &\quad - \sum_{i=1}^{\bar{m}} \sigma(\widetilde{W}_i)\sigma(C^{\uparrow}(\widetilde{t}_i-)) - \sum_{i=1}^{\overline{m}} \sigma(\overline{W}_i)\sigma(C^{\uparrow}(\overline{t}_i-)) \\ &\quad - \sigma(N_{\pm}^{\downarrow}(t_{m+1}-))\sigma(C^{\uparrow}(t_{m+1}-)),\end{aligned}$$

and after simplification

$$\begin{aligned}[Q]_1\Big|_{\Omega} &\leq +\sigma(N_{\pm}^{\downarrow}(t_0+))\sigma(C^{\uparrow}(t_0+)) \\ &\quad - \left(\sigma(N_{\pm}^{\downarrow}(t_0+)) - 3\epsilon \right) \text{TV}(N_{\pm}^{\downarrow}) - \left(\sigma(C^{\uparrow}(t_0+)) - 2\epsilon \right) \text{TV}(C^{\uparrow}) \\ &\quad - \left(\sigma(N_{\pm}^{\downarrow}(t_0+)) + \text{SV}(N_{\pm}^{\downarrow}) \right) \left(\sigma(C^{\uparrow}(t_0+)) + \text{SV}(C^{\uparrow}) \right).\end{aligned}$$

Introduce constants $\eta_N, \eta_C \in [0, 1]$ to be determined later in the proof. The terms

can then be split in the following way.

$$\begin{aligned}
[Q]_1 \Big|_{\Omega} &\leq - \left((1 - \eta_N) \sigma(N_{\pm}^{\downarrow}(t_{0+})) - 3\epsilon \right) \text{TV}(N_{\pm}^{\downarrow}) \\
&\quad - \left((1 - \eta_C) \sigma(C^{\uparrow}(t_{0+})) - 2\epsilon \right) \text{TV}(C^{\uparrow}) \\
&\quad - \sigma(N_{\pm}^{\downarrow}(t_{0+})) \left(\eta_N \text{TV}(N_{\pm}^{\downarrow}) + \text{SV}(C^{\uparrow}) \right) \\
&\quad - \sigma(C^{\uparrow}(t_{0+})) \left(\eta_C \text{TV}(C^{\uparrow}) + \text{SV}(N_{\pm}^{\downarrow}) \right) - \text{SV}(N_{\pm}^{\downarrow}) \text{SV}(C^{\uparrow}).
\end{aligned} \tag{5.6}$$

If η_N, η_C are known a priori, the first two terms are negative (and can be neglected) by taking ϵ sufficiently small. Clearly, the main difficulty now lies in the sign of $\text{SV}(N_{\pm}^{\downarrow})$ and $\text{SV}(C^{\uparrow})$. If both are positive, then all the terms are negative and we have proved our result. Recall that $\text{SV}(C^{\uparrow}) \geq 0$, by assumption, so that it suffices to consider the (only possibly unfavorable) case $\text{SV}(N_{\pm}^{\downarrow}) < 0$.

Lemma 5.3 implies that we can assume that $\text{SV}_L(C^{\uparrow})$ is also negative, provided we accept a small correction term $\mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow})$. From this, we deduce that $\text{SV}_R(C^{\uparrow}) > 0$ and in fact, necessarily

$$|\text{SV}_L(C^{\uparrow})| < |\text{SV}_R(C^{\uparrow})|.$$

We immediately note that any correction terms of order $\mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow})$ can be included into the first term of the decomposition (5.6) and, therefore, taking a smaller ϵ if necessary, we can make the new term negative:

$$-\left((1 - \eta_N) \sigma(N_{\pm}^{\downarrow}(t_{0+})) - \mathcal{O}(\epsilon) \right) \text{TV}(N_{\pm}^{\downarrow}) < 0.$$

This fact is used below without further comment.

Using Lemma 5.3 we have $|\text{SV}(N_{\pm}^{\downarrow})| \approx \lambda^* |\text{SV}_L(C^{\uparrow})| \leq \lambda^* \text{TV}(C^{\uparrow})$, and the fourth term in (5.6) can be written as

$$\begin{aligned}
&-\sigma(C^{\uparrow}(t_{0+})) \left(\eta_C \text{TV}(C^{\uparrow}) + \text{SV}(N_{\pm}^{\downarrow}) \right) \\
&\leq -\sigma(C^{\uparrow}(t_{0+})) \left(\eta_C \text{TV}(C^{\uparrow}) - \lambda^* \text{TV}(C^{\uparrow}) \right) + \mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow}).
\end{aligned}$$

Taking $\eta_C = \lambda^*$ suffices to guarantee that the term vanishes. We now re-organize the third and fifth terms of (5.6) to take advantage of the known signs of the signed variations, as follows:

$$\begin{aligned}
A &:= -\sigma(N_{\pm}^{\downarrow}(t_{0+})) \left(\eta_N \text{TV}(N_{\pm}^{\downarrow}) + \text{SV}(C^{\uparrow}) \right) - \text{SV}(N_{\pm}^{\downarrow}) \text{SV}(C^{\uparrow}) \\
&= -\sigma(N_{\pm}^{\downarrow}(t_{0+})) \left(\eta_N \text{TV}(N_{\pm}^{\downarrow}) + \frac{\eta_N}{2} \text{SV}_L(C^{\uparrow}) \right) \\
&\quad - \sigma(N_{\pm}^{\downarrow}(t_{0+})) \left(\left(1 - \frac{\eta_N}{2} \right) \text{SV}_L(C^{\uparrow}) + \text{SV}_R(C^{\uparrow}) \right) \\
&\quad - \text{SV}(N_{\pm}^{\downarrow}) \text{SV}_L(C^{\uparrow}) - \text{SV}(N_{\pm}^{\downarrow}) \text{SV}_R(C^{\uparrow}) + \mathcal{O}(\epsilon) \text{TV}(N_{\pm}^{\downarrow}).
\end{aligned}$$

Observe that we have the following lower bounds

$$\left(1 - \frac{\eta_N}{2} \right) \text{SV}_L(C^{\uparrow}) + \text{SV}_R(C^{\uparrow}) \geq \frac{\eta_N}{2} \left| \text{SV}_L(C^{\uparrow}) \right|$$

and upper bounds

$$\left| \text{SV}_L(C^\dagger) \right| = \frac{1}{\lambda^*} \left| \text{SV}(N_\pm^\downarrow) \right| + \mathcal{O}(\epsilon) \text{TV}(N_\pm^\downarrow) \leq 2 \text{TV}(N_\pm^\downarrow) + \mathcal{O}(\epsilon) \text{TV}(N_\pm^\downarrow).$$

Using the two previous bounds, we find

$$\begin{aligned} A &\leq -\sigma(N_\pm^\downarrow(t_{0+})) \left(\eta_N \text{TV}(N_\pm^\downarrow) - \eta_N \text{TV}(N_\pm^\downarrow) \right) - \sigma(N_\pm^\downarrow(t_{0+})) \frac{\eta_N}{2} \left| \text{SV}_L(C^\dagger) \right| \\ &\quad - \text{SV}(N_\pm^\downarrow) \text{SV}_L(C^\dagger) - \lambda^* \text{SV}_L(C^\dagger) \text{SV}_R(C^\dagger) + \mathcal{O}(\epsilon) \text{TV}(N_\pm^\downarrow) \\ &\leq - \left| \text{SV}_L(C^\dagger) \right| \left(-\lambda^* \text{SV}_R(C^\dagger) + \frac{\eta_N}{2} \sigma(N_\pm^\downarrow(t_{0+})) \right) \\ &\quad - \text{SV}(N_\pm^\downarrow) \text{SV}_L(C^\dagger) + \mathcal{O}(\epsilon) \text{TV}(N_\pm^\downarrow). \end{aligned}$$

For any fixed value of $\eta_N \in (0, 1)$, the first term is negative because $|\text{SV}_R(C^\dagger)| < \epsilon$ can be assumed a priori small. The second term is negative and the last one can be incorporated into the first term of (5.6). This completes the proof of Theorem 5.1. \square

6. Quadratic interaction potential (part 2). We continue our investigation of interaction functionals adapted to nonclassical shocks. The proposal now is to also involve a weight-function in front of the quadratic terms which is chosen to be proportional to the difference in (normalized) wave speeds. Such a weighting was used successfully by Iguchi and LeFloch [3] as well as by Liu and Yang [11] in their analysis of general hyperbolic systems of conservation laws; see also Liu [10] for earlier pioneering work on systems.

The interaction functional proposed now for nonclassical entropy solutions shares the same monotonicity properties as the functional Q_{weak} , in the sense that it fails for precisely the same interactions. In preparation for the study of systems of conservation laws, a complete study of the scalar case requires an examination of the following functional.

So, we introduce

$$Q_{\text{pos}}(u(t)) := \sum_{x < y} \theta(x, y) \sigma(x) \sigma(y),$$

where $\sigma(x), \sigma(y)$ are the generalized wave strengths defined earlier,

$$\sigma(x) = \begin{cases} \sigma(S), & S \text{ is a wave located at } x, \\ 0, & \text{no wave located at } x. \end{cases}$$

The potential now contains a weight function $\theta(x, y)$ defined by

$$\theta(x, y) := \begin{cases} C_* (\widehat{a}(x) - \widehat{a}(y))^+, & \text{same monotonicity,} \\ 1, & \text{otherwise,} \end{cases}$$

where the constant $C_* > 0$ will be chosen to be sufficiently small and $(k)^+ := \max\{k, 0\}$. More precisely, here we say that two waves have the same monotonicity if after mapping all states to the positive region the waves are either both increasing or both decreasing. The definition of \widehat{a} is as follows:

$$\widehat{a}(x) := \frac{f(\widehat{u}_x^+) - f(\widehat{u}_x^-)}{\widehat{u}_x^+ - \widehat{u}_x^-},$$

where

$$\widehat{u}_x^+ := \begin{cases} u_x^+, & u_x^+ \geq 0, \\ \varphi_0^b(u_x^+), & u_x^+ < 0. \end{cases}$$

This definition allows us to enforce that comparisons only occur between state variables belonging to the same region of convexity, i.e. between positive values. Note that, in the present formalism, rarefaction are possibly interacting with each other.

THEOREM 6.1 (“Positive interaction” potential for nonclassical shocks). *Let φ^b be a kinetic function satisfying the properties (A1)–(A3). Consider the weighted interaction functional Q_{pos} evaluated for a front-tracking approximate solution u to (1.1), whose discontinuities satisfy (1.5), and with initial data $u_0 \in \text{BV}(\mathbb{R})$. If C_* is taken sufficiently small, then Q_{pos} is strictly decreasing during all interactions except in the cases RC-3, CR-4, CC-3 and CN-3 which involve a small wave interacting with a crossing shock.*

Proof. We distinguish between several cases of interactions. For the functional $Q = Q_{pos}$ we introduce a new form of the decomposition (4.2),

$$[Q(u(t))] =: [Q]_1 + \sum_y \sigma(y) B(y),$$

where the sum over y excludes waves involved in the interaction at time t . Given that the waves not involved in the interaction are potentially arbitrary, $[Q]$ will be negative if and only if $[Q]_1$ and $B(y)$ are negative. As usual, we will rely heavily on the inequalities described in the proofs of Theorems 3.1 and 4.2 as well as of Lemma 4.3.

Case RC-1: $(R_+^\uparrow C^\downarrow)$ - $(C^{\downarrow'})$. In this case, a single wave is outgoing and we have $[Q]_1 < 0$. In the proof of Theorem 3.1, we observed that the states of interest satisfy

$$\begin{aligned} &\text{either } u_r < u_l < u_m, \\ &\text{or } \varphi_0^b(u_r) < u_l < u_m. \end{aligned}$$

For both sets of inequalities, $\sigma(C^\downarrow) < \sigma(C^{\downarrow'})$ holds and the normalized speeds are ordered

$$\widehat{a}(C^{\downarrow'}) < \widehat{a}(C^\downarrow) < \widehat{a}(R_+^\uparrow).$$

Suppose there is a shock at some point y with speed Λ , then

$$B(y) = C_* (\widehat{a}(C^{\downarrow'}) - \Lambda)^+ \sigma(C^{\downarrow'}) - C_* (\widehat{a}(C^\downarrow) - \Lambda)^+ \sigma(C^\downarrow) - \sigma(R_+^\uparrow),$$

which by the monotonicity just observed must be negative. On the other hand, if a rarefaction is located at y , then the estimate on the wave strengths shows that

$$B(y) = \sigma(C^{\downarrow'}) - \sigma(C^\downarrow) - C_* (\widehat{a}(R_+^\uparrow) - \Lambda)^+ \sigma(R_+^\uparrow) < 0.$$

Case RC-2: $(R_+^\uparrow C_\pm^\downarrow)$ - $(N_\pm^{\downarrow'} R_\pm^{\downarrow'})$. Our study of this interaction in Theorem 4.2 provides

$$\sigma(N_\pm^{\downarrow'}) < \sigma(C_\pm^\downarrow), \quad \sigma(R_\pm^{\downarrow'}) < \sigma(R_+^\uparrow).$$

In turn, it follows that

$$[Q]_1 = \sigma(N_{\pm}^{\downarrow'}) \sigma(R_{\pm}^{\downarrow'}) - \sigma(C_{\pm}^{\downarrow}) \sigma(R_{\pm}^{\uparrow}) < 0.$$

Next, to control $B(y)$ we first consider the relevant speeds. When $\varphi_0^b(u_r) < u_l$, then inequalities (3.6) are valid and

$$\widehat{a}(R_{\pm}^{\downarrow'}) < \widehat{a}(N_{\pm}^{\downarrow}) < \widehat{a}(C_{\pm}^{\downarrow}) < \widehat{a}(R_{\pm}^{\uparrow}),$$

while when $\varphi_0^b(u_r) > u_l$ we have

$$\widehat{a}(N_{\pm}^{\uparrow'}) < \widehat{a}(R_{\pm}^{\downarrow'}) < \widehat{a}(R_{\pm}^{\uparrow}) < \widehat{a}(C_{\pm}^{\downarrow}).$$

Now, suppose that a shock wave is located at some point y , then

$$\begin{aligned} B(y) = & C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) + \sigma(R_{\pm}^{\downarrow'}) \\ & - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\uparrow}), \end{aligned}$$

which is negative in view of the above inequalities on the speeds and strengths. A similar argument applies if a rarefaction is located at y .

Case RC-3 : $(R_{\pm}^{\uparrow} C^{\downarrow}) - (N_{\pm}^{\downarrow'} C^{\uparrow'})$. This case is more delicate, due to the fact that a nonclassical shock is generated from classical waves. The shock strengths satisfy the relation (4.4) which implies

$$\sigma(N_{\pm}^{\downarrow'}) + \sigma(C^{\downarrow'}) < \sigma(C^{\downarrow}). \quad (6.1)$$

We distinguish between two cases (3.7) and (3.8) depending on the sign of u_r . An inspection of the normalized wave speeds associated to the states in (3.7) and (3.8) demonstrates that

$$\widehat{a}(C^{\uparrow'}) < \widehat{a}(N_{\pm}^{\downarrow'}) < \widehat{a}(R_{\pm}^{\uparrow}) \quad \text{and} \quad \widehat{a}(C^{\uparrow'}) < \widehat{a}(C^{\downarrow}) < \widehat{a}(R_{\pm}^{\uparrow}).$$

We analyzed an interaction term in Lemma 4.3 similar to

$$[Q]_1 = C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \widehat{a}(C^{\uparrow'}))^+ \sigma(N_{\pm}^{\downarrow'}) \sigma(C^{\uparrow'}) - \sigma(R_{\pm}^{\uparrow}) \sigma(C^{\downarrow}),$$

and we saw that it was possible to take $\sigma(R_{\pm}^{\uparrow})$ vanishingly small, while maintaining a non-vanishing product $\sigma(N_{\pm}^{\downarrow'}) \sigma(C^{\uparrow'})$. Therefore, in general we cannot claim that $[Q]_1$ is negative.

We now show that in the limit as $\sigma(R_{\pm}^{\uparrow}) \rightarrow 0$, $B(y)$ is again positive for some speeds Λ . Assuming a shock located at y , the general form of B is

$$\begin{aligned} B(y) = & C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) + C_* (\widehat{a}(C^{\uparrow'}) - \Lambda)^+ \sigma(C^{\uparrow'}) \\ & - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\uparrow}). \end{aligned}$$

If $\varphi^{\sharp}(u_l) < 0$ and inequalities (3.7) hold, then take $u_r \rightarrow \varphi^{\sharp}(u_l)$ and $u_m \rightarrow u_l$. In this limit, $\widehat{a}(C^{\uparrow'}) < \widehat{a}(C^{\downarrow}) < \widehat{a}(N_{\pm}^{\downarrow'})$ and the quantities $\widehat{a}(C^{\uparrow'})$ and $\sigma(N_{\pm}^{\downarrow'})$ are functions only of u_l . Therefore taking $\Lambda \in [\widehat{a}(C^{\downarrow}), \widehat{a}(N_{\pm}^{\downarrow'})]$, we find that for those limiting values

$$B(y) = C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) > 0.$$

Now, if a rarefaction is located at y , then (6.1) suffices to show

$$B(y) = \sigma(N_{\pm}^{\downarrow'}) + \sigma(C^{\uparrow'}) - \sigma(C^{\downarrow}) - C_* (\widehat{a}(R_+^{\uparrow}) - \Lambda)^+ \sigma(R_+^{\uparrow}) < 0.$$

Case RN : $(R_+^{\uparrow} N_{\pm}^{\downarrow}) - (N_{\pm}^{\downarrow'} R_-^{\downarrow'})$. As we saw in the proof of Theorem 3.1, there are two subcases defined by either

$$\begin{aligned} &\text{either } \varphi_0^b \circ \varphi^b(u_l) < \varphi_0^b \circ \varphi^b(u_m) < u_l < u_m, \\ &\text{or } \varphi_0^b \circ \varphi^b(u_l) < u_l < \varphi_0^b \circ \varphi^b(u_m) < u_m. \end{aligned}$$

The normalized speeds satisfy respectively In Case A the speeds satisfy

$$\begin{aligned} &\widehat{a}(R_-^{\downarrow'}) < \widehat{a}(N_{\pm}^{\downarrow'}) < \widehat{a}(N_{\pm}^{\downarrow}) < \widehat{a}(R_+^{\uparrow}), \text{ or} \\ &\widehat{a}(N_{\pm}^{\downarrow'}) < \widehat{a}(R_-^{\downarrow'}) < \widehat{a}(R_+^{\uparrow}) < \widehat{a}(N_{\pm}^{\downarrow}). \end{aligned}$$

In both subcases, we have $\sigma(N_{\pm}^{\downarrow'}) < \sigma(N_{\pm}^{\downarrow})$ and $\sigma(R_-^{\downarrow'}) < \sigma(R_+^{\uparrow})$, and the inequalities on the speeds $\widehat{a}(N_{\pm}^{\downarrow'}) < \widehat{a}(N_{\pm}^{\downarrow})$ and $\widehat{a}(R_-^{\downarrow'}) < \widehat{a}(R_+^{\uparrow})$. Using the observations above, we verify that

$$[Q]_1 = \sigma(N_{\pm}^{\downarrow'}) \sigma(R_-^{\downarrow'}) - \sigma(R_+^{\uparrow}) \sigma(N_{\pm}^{\downarrow}) < 0.$$

If a shock is located at y , then the term

$$B(y) = C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) + \sigma(R_-^{\downarrow'}) + C_* (\widehat{a}(N_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow}) - \sigma(R_+^{\uparrow})$$

is negative. When a rarefaction is located at y , then

$$B(y) = \sigma(N_{\pm}^{\downarrow'}) + C_* (\widehat{a}(R_-^{\downarrow'}) - \Lambda)^+ \sigma(R_-^{\downarrow'}) - \sigma(N_{\pm}^{\downarrow}) - C_* (\widehat{a}(R_-^{\downarrow}) - \Lambda)^+ \sigma(R_-^{\downarrow})$$

is again negative.

Case CR-1 : $(C_{\pm}^{\downarrow} R_-^{\downarrow}) - (C_{\pm}^{\downarrow'})$. Only one wave is outgoing, so we trivially have $[Q]_1 < 0$. The states of interest satisfy

$$\varphi_0^b(u_m) < \varphi_0^b(u_r) < u_l,$$

and therefore the normalized speeds are ordered

$$\widehat{a}(R_-^{\downarrow}) < \widehat{a}(C_{\pm}^{\downarrow}) < \widehat{a}(C_{\pm}^{\downarrow'}).$$

Recall also that $\sigma(C_{\pm}^{\downarrow'}) < \sigma(C_{\pm}^{\downarrow})$. Assume now that a shock is located at y , then

$$B(y) = C_* (\widehat{a}(C_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow'}) - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}) - \sigma(R_-^{\downarrow}).$$

The least favorable case corresponds to the situation where $\Lambda = \widehat{a}(C_{\pm}^{\downarrow})$, since only one negative term remains when $\Lambda > \widehat{a}(C_{\pm}^{\downarrow'})$ and $B(y)$ is increasing with respect to Λ when $\Lambda < \widehat{a}(C_{\pm}^{\downarrow})$. The important term in $B(y)$ when $\Lambda = \widehat{a}(C_{\pm}^{\downarrow})$ is the difference in speeds

$$\begin{aligned} (\widehat{a}(C_{\pm}^{\downarrow'}) - \widehat{a}(C_{\pm}^{\downarrow}))^+ &= \left| \frac{f(u_l) - f(\widehat{u}_r)}{u_l - \widehat{u}_r} - \frac{f(u_l) - f(\widehat{u}_m)}{u_l - \widehat{u}_m} \right| \\ &\leq C |\widehat{u}_r - \widehat{u}_m| = C \sigma(R_-^{\downarrow}). \end{aligned}$$

The constant $C = C(u_l)$ is bounded above since the total variation of the solution u remains bounded. If we therefore assume the a priori bound

$$C_* C \text{TV}(u) < 1, \quad (6.2)$$

then

$$B(y) \leq C_* C \sigma(R_-^\downarrow) \sigma(C_\pm^{\downarrow'}) - \sigma(R_-^\downarrow) = -\sigma(R_-^\downarrow)(1 - C_* C \sigma(C_\pm^{\downarrow'})) < 0.$$

To complete the proof, we assume that a rarefaction is now located at y , and immediately see that

$$B(y) = \sigma(C_\pm^{\downarrow'}) - \sigma(C_\pm^\downarrow) - C_* (\widehat{a}(R_-^\downarrow) - \Lambda)^+ \sigma(R_-^\downarrow) < 0.$$

Case CR-2 : $(C_+^\downarrow R_+^\uparrow)$ - $(C_+^{\downarrow'})$. The waves are entirely classical and the calculations are identical to previous Case CR-1. In particular, we again have to impose condition (6.2) on C_* .

Case CR-3 : $(C_\pm^\downarrow R_-^\downarrow)$ - $(N_\pm^{\downarrow'} R_-^{\downarrow'})$. In the proof of Theorem 3.1, we identified two subcases distinguished by the inequalities (3.10) and (3.11) :

$$\begin{aligned} &\text{either } \varphi_0^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < u_l < \varphi_0^b(u_r), \\ &\text{or } \varphi_0^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < \varphi_0^b(u_r) < u_l. \end{aligned}$$

In both subcases we have

$$\sigma(N_\pm^{\downarrow'}) < \sigma(C_\pm^\downarrow), \quad \sigma(R_-^{\downarrow'}) < \sigma(R_-^\downarrow).$$

In the first subcase, the normalized speeds satisfy

$$\widehat{a}(C_\pm^\downarrow) < \min(\widehat{a}(R_-^\downarrow), \widehat{a}(N_\pm^{\downarrow'})) < \max(\widehat{a}(R_-^\downarrow), \widehat{a}(N_\pm^{\downarrow'})) < \widehat{a}(R_-^{\downarrow'}),$$

while in the second,

$$\widehat{a}(R_-^\downarrow) < \min(\widehat{a}(C_\pm^\downarrow), \widehat{a}(R_-^{\downarrow'})) < \max(\widehat{a}(C_\pm^\downarrow), \widehat{a}(R_-^{\downarrow'})) < \widehat{a}(N_\pm^{\downarrow'}).$$

The key point is that in both subcases, $\widehat{a}(C_\pm^\downarrow) < \widehat{a}(N_\pm^{\downarrow'})$ and $\widehat{a}(R_-^\downarrow) < \widehat{a}(R_-^{\downarrow'})$. Using the inequalities satisfied by only the strengths, we obtain

$$[Q]_1 = \sigma(N_\pm^{\downarrow'})\sigma(R_-^{\downarrow'}) - \sigma(C_\pm^\downarrow)\sigma(R_-^\downarrow) < 0.$$

When a shock is located at y , we can verify that

$$B(y) = C_* (\widehat{a}(N_\pm^{\downarrow'}) - \Lambda)^+ \sigma(N_\pm^{\downarrow'}) + \sigma(R_-^{\downarrow'}) - C_* (\widehat{a}(C_\pm^\downarrow) - \Lambda)^+ \sigma(C_\pm^\downarrow) - \sigma(R_-^\downarrow).$$

The worst case occurs when $\Lambda = \widehat{a}(C_\pm^\downarrow) < \widehat{a}(N_\pm^{\downarrow'})$, and then we find that there exists a constant C such that

$$\begin{aligned} \widehat{a}(N_\pm^{\downarrow'}) - \widehat{a}(C_\pm^\downarrow) &= \left| \frac{f(u_l) - f(\varphi_0^b \circ \varphi^b(u_l))}{u_l - \varphi_0^b \circ \varphi^b(u_l)} - \frac{f(u_l) - f(\widehat{u}_m)}{u_l - \widehat{u}_m} \right| \\ &\leq C(\varphi_0^b \circ \varphi^b(u_l) - \widehat{u}_m) = C(\sigma(R_-^\downarrow) - \sigma(R_-^{\downarrow'})). \end{aligned}$$

We again impose a condition of the form (6.2) on C_* and find

$$\begin{aligned} B(y) &= C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \widehat{a}(C_{\pm}^{\downarrow}))^+ \sigma(N_{\pm}^{\downarrow'}) - (\sigma(R_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\downarrow}')) \\ &= - (1 - CC_* \sigma(N_{\pm}^{\downarrow'})) (\sigma(R_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\downarrow}')) < 0. \end{aligned}$$

Suppose next that there is a rarefaction at y , then

$$B(y) = \sigma(N_{\pm}^{\downarrow'}) + C_* (\widehat{a}(R_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(R_{\pm}^{\downarrow'}) - \sigma(C_{\pm}^{\downarrow}) - C_* (\widehat{a}(R_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(R_{\pm}^{\downarrow}).$$

The largest value of $B(y)$ occurs when $\Lambda = \widehat{a}(R_{\pm}^{\downarrow})$ and then the coefficient of the $\sigma(R_{\pm}^{\downarrow}')$ term satisfies

$$\begin{aligned} \widehat{a}(R_{\pm}^{\downarrow}') - \widehat{a}(R_{\pm}^{\downarrow}) &= \left| \frac{f(\widehat{u}_r) - f(\varphi_0^b \circ \varphi^b(u_l))}{\widehat{u}_r - \varphi_0^b \circ \varphi^b(u_l)} - \frac{f(\widehat{u}_r) - f(\widehat{u}_m)}{\widehat{u}_r - \widehat{u}_m} \right| \\ &\leq C(\varphi_0^b \circ \varphi^b(u_l) - \widehat{u}_m) = C(\sigma(C_{\pm}^{\downarrow}) - \sigma(N_{\pm}^{\downarrow}')). \end{aligned}$$

The same arguments as before apply and again allow us to conclude that $B(y) < 0$.

Case CR-4 : $(C_{\pm}^{\downarrow} R_{\pm}^{\downarrow}) - (N_{\pm}^{\downarrow'} C_{\pm}^{\uparrow'})$. Here, we have

$$\varphi_0^b(u_m) < \varphi_0^b(u_r) < \varphi_0^b \circ \varphi^b(u_l) < u_l.$$

Now, the normalized speed satisfy

$$\widehat{a}(R_{\pm}^{\downarrow}) < \widehat{a}(C_{\pm}^{\uparrow'}) < \widehat{a}(N_{\pm}^{\downarrow'}), \quad \widehat{a}(R_{\pm}^{\downarrow}) < \widehat{a}(C_{\pm}^{\downarrow}) < \widehat{a}(N_{\pm}^{\downarrow}')$$

and for the strengths we have

$$\sigma(N_{\pm}^{\downarrow'}) + \sigma(C_{\pm}^{\uparrow'}) < \sigma(C_{\pm}^{\downarrow}).$$

For the interaction, we obtain a term

$$[Q]_1 = C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \widehat{a}(C_{\pm}^{\uparrow'}))^+ \sigma(N_{\pm}^{\downarrow'}) \sigma(C_{\pm}^{\uparrow'}) - \sigma(C_{\pm}^{\downarrow}) \sigma(R_{\pm}^{\downarrow})$$

similar to the one analyzed in Lemma 4.3. There we saw that it was possible to let $\sigma(R_{\pm}^{\downarrow}) \rightarrow 0$ while keeping both $\sigma(N_{\pm}^{\downarrow'})$ and $\sigma(C_{\pm}^{\uparrow'})$ bounded and non-vanishing.

Therefore, this term is positive for a large class of interactions.

When a shock is located at y , the general term to look at is

$$\begin{aligned} B(y) &= C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) + C_* (\widehat{a}(C_{\pm}^{\uparrow'}) - \Lambda)^+ \sigma(C_{\pm}^{\uparrow'}) \\ &\quad - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}) - \sigma(R_{\pm}^{\downarrow}). \end{aligned}$$

Taking again the limit as $\sigma(R_{\pm}^{\downarrow}) \rightarrow 0$, we find $\widehat{a}(C_{\pm}^{\uparrow'}) < \widehat{a}(C_{\pm}^{\downarrow}) < \widehat{a}(N_{\pm}^{\downarrow}')$. Moreover, when $\Lambda = \widehat{a}(C_{\pm}^{\downarrow})$ we obtain expressions $\sigma(N_{\pm}^{\downarrow'})$ and $\widehat{a}(N_{\pm}^{\downarrow'}) - \widehat{a}(C_{\pm}^{\downarrow})$ that are non-zero and functions only of u_l . Therefore

$$B(y) = C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) > 0.$$

When a rarefaction is located at y , then

$$B(y) = \sigma(N_{\pm}^{\downarrow'}) + \sigma(C_{\pm}^{\uparrow'}) - \sigma(C_{\pm}^{\downarrow}) - C_* (\widehat{a}(R_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(R_{\pm}^{\downarrow}) < 0.$$

Case CC-1 : $(C_{\pm}^{\downarrow} C^{\downarrow})-(C^{\downarrow'})$. There are two subcases determined by the sign of u_r :

$$\varphi_0^b(u_r) < u_m < u_l, \quad \text{or} \quad u_r < u_m < u_l.$$

In both subcases, we have the identities

$$\begin{aligned} \sigma(C^{\downarrow'}) &= \sigma(C_{\pm}^{\downarrow}) + \sigma(C^{\downarrow}), \\ \widehat{a}(C^{\downarrow}) &< \widehat{a}(C^{\downarrow'}) < \widehat{a}(C_{\pm}^{\downarrow}). \end{aligned}$$

Only one wave is outgoing so $[Q]_1 < 0$.

Suppose a shock is located at y , then the largest value of

$$B(y) = C_* (\widehat{a}(C^{\downarrow'}) - \Lambda)^+ \sigma(C^{\downarrow'}) - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}) - C_* (\widehat{a}(C^{\downarrow}) - \Lambda)^+ \sigma(C^{\downarrow})$$

occurs in the interval $\Lambda \in [\widehat{a}(C^{\downarrow}), \widehat{a}(C^{\downarrow'})]$, leaving us with

$$B(y) = C_* (\widehat{a}(C^{\downarrow'}) - \Lambda)^+ \sigma(C^{\downarrow'}) - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}).$$

In the subcase where $u_r > 0$, the definition of σ and \widehat{a} allows us to find

$$\begin{aligned} B(y) &= C_* \left((f(u_l) - f(u_r)) - \Lambda(u_l - u_r) - (f(u_l) - f(u_m)) + \Lambda(u_l - u_m) \right) \\ &= C_* (\widehat{a}(C^{\downarrow}) - \Lambda) \sigma(C^{\downarrow}) \leq 0. \end{aligned}$$

If $u_r < 0$, then similar computations show that $B(y)$ is still negative.

When a rarefaction is located at y , then

$$B(y) = \sigma(C^{\downarrow'}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(C^{\downarrow}) = 0.$$

Case CC-2 : $(C_{\pm}^{\downarrow} C^{\uparrow})-(C^{\downarrow'})$. Two subcases appear depending on the sign of u_r . We will assume throughout that $u_r > 0$, leaving us with the states

$$u_r < \varphi_0^b(u_m) < u_l,$$

and therefore the two identities

$$\begin{aligned} \sigma(C^{\downarrow'}) &= \sigma(C_{\pm}^{\downarrow}) + \sigma(C^{\uparrow}), \\ \widehat{a}(C^{\uparrow}) &< \widehat{a}(C^{\downarrow'}) < \widehat{a}(C_{\pm}^{\downarrow}). \end{aligned}$$

We recognize immediately the same identities as in the Case CC-1. As quick verification shows that the same arguments demonstrate that $[Q]_1 < 0$ and $[Q]_2 < 0$.

Case CC-3 : $(C_{\pm}^{\downarrow} C^{\downarrow})-(N_{\pm}^{\downarrow'} C^{\uparrow'})$. We can distinguish between two subcases :

$$\begin{aligned} \text{either } \varphi_0^b(u_r) &< u_m < \varphi_0^b \circ \varphi^b(u_l) < u_l, \\ \text{or } \varphi_0^b(u_r) &< \varphi_0^b \circ \varphi^b(u_l) < u_m < u_l. \end{aligned}$$

In both subcases, one obtains

$$\widehat{a}(C^{\downarrow}) < \widehat{a}(N_{\pm}^{\downarrow'}) \quad \text{and} \quad \widehat{a}(C^{\uparrow'}) < \widehat{a}(C_{\pm}^{\downarrow}). \quad (6.3)$$

Unfortunately, the conditions above on the shock speeds do not preclude the possibility that $\widehat{a}(C_+^\downarrow) < \widehat{a}(N_\pm^{\downarrow'})$ which would imply that

$$B(y) = C_* (\widehat{a}(N_\pm^{\downarrow'}) - \Lambda)^+ \sigma(N_\pm^{\downarrow'}) + C_* (\widehat{a}(C^{\uparrow'}) - \Lambda)^+ \sigma(C^{\uparrow'}) \\ - C_* (\widehat{a}(C_+^\downarrow) - \Lambda)^+ \sigma(C_+^\downarrow) - C_* (\widehat{a}(C^\downarrow) - \Lambda)^+ \sigma(C^\downarrow)$$

was positive. On the other hand, because $[V] = 0$ we have that $B(y) = 0$ when a rarefaction is located at y .

The sign of $[Q]_1$ is also less than helpful. Using the ideas seen in Lemma 4.3 for the same term, one notices that when u_m equals the transitional state $\varphi_0^b \circ \varphi^b(u_l)$ between the two subcases, then the change

$$[Q]_1 = C_* (\widehat{a}(N_\pm^{\downarrow'}) - \widehat{a}(C^{\uparrow'})) + \sigma(N_\pm^{\downarrow'}) \sigma(C^{\uparrow'}) - C_* (\widehat{a}(C_+^\downarrow) - \widehat{a}(C^\downarrow)) + \sigma(C_+^\downarrow) \sigma(C^\downarrow)$$

vanishes because the quadratic terms and the coefficients are equal. It is clear that for some set of waves close to this transitional value, $[Q]_1$ will also be positive.

Case CN-1 : $(C_+^\downarrow N_\pm^\downarrow) - (C_\pm^{\downarrow'})$. The states are $\varphi_0^b \circ \varphi^b(u_m) < u_m < u_l$ and these provide us with the basic inequalities

$$\sigma(C_\pm^{\downarrow'}) = \sigma(C_+^\downarrow) + \sigma(N_\pm^\downarrow), \\ \widehat{a}(N_\pm^\downarrow) < \widehat{a}(C_\pm^{\downarrow'}) < \widehat{a}(C_+^\downarrow).$$

Under these conditions, the analysis is similar to the CC-1 case.

Case CN-2 : $(C_\pm^\downarrow N_\mp^\uparrow) - (C_+^{\downarrow'})$. The important states are ordered $\varphi^b(u_m) < \varphi_0^b(u_m) < u_l$ and therefore

$$\sigma(C_+^{\downarrow'}) = \sigma(C_\pm^\downarrow) + \sigma(N_\mp^\uparrow), \\ \widehat{a}(N_\mp^\uparrow) < \widehat{a}(C_+^{\downarrow'}) < \widehat{a}(C_\pm^\downarrow).$$

Again, the analysis is similar to the CC-1 case.

Case CN-3 : $(C_+^\downarrow N_\pm^\downarrow) - (N_\pm^{\downarrow'} C_-^{\uparrow'})$. There are two subcases defined by the ordered set of states

$$\text{either } \varphi_0^b \circ \varphi^b(u_m) < u_m < \varphi_0^b \circ \varphi^b(u_l) < u_l, \\ \text{or } \varphi_0^b \circ \varphi^b(u_m) < \varphi_0^b \circ \varphi^b(u_l) < u_m < u_l,$$

where the second case corresponds to a weak incoming shock C_+^\downarrow . It is easy to verify that the shock strengths verify

$$\sigma(N_\pm^\downarrow) < \sigma(N_\pm^{\downarrow'}) \quad \text{and} \quad \sigma(C^{\uparrow'}) < \sigma(C_\pm^\downarrow). \quad (6.4)$$

In both subcases, one obtains the unfortunate inequalities

$$\widehat{a}(N_\pm^\downarrow) < \widehat{a}(N_\pm^{\downarrow'}) \quad \text{and} \quad \widehat{a}(C^{\uparrow'}) < \widehat{a}(C_\pm^\downarrow). \quad (6.5)$$

The analogue to

$$[Q]_1 = C_* (\widehat{a}(N_\pm^{\downarrow'}) - \widehat{a}(C_-^{\uparrow'})) + \sigma(N_\pm^{\downarrow'}) \sigma(C_-^{\uparrow'}) - C_* (\widehat{a}(C_+^\downarrow) - \widehat{a}(N_\pm^\downarrow)) + \sigma(C_+^\downarrow) \sigma(N_\pm^\downarrow)$$

for Q_{weak} was analyzed in Lemma 4.3 and seen to be positive for sufficiently strong waves C_{\pm}^{\downarrow} . For the functional Q_{pos} , $[Q]_1 = 0$ when $u_m = \varphi_0^b \circ \varphi^b(u_l)$ because at that critical value, all inequalities in (6.4) and (6.5) are equalities. Without going into the details, it is easy to verify that this implies that for a non-empty set of waves, $[Q]_1$ will be positive.

Suppose a shock is located at y , then

$$B(y) = C_* (\widehat{a}(N_{\pm}^{\downarrow'}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow'}) + C_* (\widehat{a}(C_{\pm}^{\uparrow'}) - \Lambda)^+ \sigma(C_{\pm}^{\uparrow'}) \\ - C_* (\widehat{a}(C_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(C_{\pm}^{\downarrow}) - C_* (\widehat{a}(N_{\pm}^{\downarrow}) - \Lambda)^+ \sigma(N_{\pm}^{\downarrow}).$$

When we are in the first subcase (a strong C_{\pm}^{\downarrow}), the change $B(y)$ is clearly positive if $\Lambda \in [\widehat{a}(C_{\pm}^{\downarrow}), \widehat{a}(N_{\pm}^{\downarrow'})]$.

Finally, suppose a rarefaction is located at y , then

$$B(y) = \sigma(N_{\pm}^{\downarrow'}) + \sigma(C_{\pm}^{\uparrow'}) - \sigma(C_{\pm}^{\downarrow}) - \sigma(N_{\pm}^{\downarrow}) = 0.$$

Case NC : $(N_{\pm}^{\downarrow} C^{\uparrow}) - (C^{\downarrow'})$. The defining states are $u_r < \varphi_0^b \circ \varphi^b(u_l) < u_l$ and we have

$$\sigma(C^{\downarrow'}) = \sigma(N_{\pm}^{\downarrow}) + \sigma(C^{\uparrow}), \\ \widehat{a}(C^{\uparrow}) < \widehat{a}(C^{\downarrow'}) < \widehat{a}(N_{\pm}^{\downarrow}).$$

See the CC-1 case.

Case NN : $(N_{\pm}^{\downarrow} N_{\pm}^{\uparrow}) - (C_{\pm}^{\downarrow'})$. The analysis of the NC case also applies here since NN interactions are limiting situations obtained as $u_r \rightarrow \varphi^b(u_m)$. □

Acknowledgments. The first author (M.L.) was partially supported by the Natural Sciences and Engineering Research Council (NSERC) of Canada. The second author (PLF) was partially supported by the Centre National de la Recherche Scientifique (CNRS) and the Agence Nationale de la Recherche (ANR) through the grant 06-2-134423 entitled “*Mathematical Methods in General Relativity*” (MATH-GR).

REFERENCES

- [1] R. ABEYARATNE AND J.K. KNOWLES, Kinetic relations and the propagation of phase boundaries in solids, *Arch. Rational Mech. Anal.* 114 (1991), 119–154.
- [2] N. BEDJAOUI AND P.G. LEFLOCH, Diffusive-dispersive travelling waves and kinetic relations V. Singular diffusion and nonlinear dispersion, *Proc. Royal Soc. Edinburgh* 134A (2004), 815–844.
- [3] T. IGUCHI AND P.G. LEFLOCH, Existence theory for hyperbolic systems of conservation laws with general flux-functions, *Arch. Rational Mech. Anal.* 168 (2003), 165–244.
- [4] J. GLIMM, Solutions in the large for nonlinear hyperbolic systems of equations, *Comm. Pure Appl. Math.* 18 (1965), 697–715.
- [5] P.G. LEFLOCH, Propagating phase boundaries: formulation of the problem and existence via the Glimm scheme, *Arch. Ration. Mech. Anal.* 123 (1993), 153–197.
- [6] P.G. LEFLOCH, *Hyperbolic systems of conservation laws: The theory of classical and nonclassical shock waves*, Lectures in Mathematics, ETH Zürich, Birkhäuser, 2002.
- [7] C.M. DAFERMOS, Polygonal approximation of solutions to the initial-value problem for a conservation law, *J. Math. Anal. Appl.* 38 (1972), 33–41.

- [8] P.G. LEFLOCH AND M. MOHAMMADIAN, Why many shock wave theories are necessary. Fourth-order models, kinetic functions, and equivalent equations, *J. Comput. Phys.* 227 (2008), 4162–4189.
- [9] P.G. LEFLOCH AND M. SHEARER, Nonclassical Riemann solvers with nucleation, *Proc. Royal Soc. Edinburgh* 134A (2004), 941–964.
- [10] T.-P. LIU, *Admissible solutions of hyperbolic conservation laws*, *Memoir Amer. Math. Soc.* 30, 1981.
- [11] T.-P. LIU AND T. YANG, Weak solutions of general systems of hyperbolic conservation laws, *Commun. Math. Phys.* 230 (2002), 289–327.
- [12] M. SLEMROD, Admissibility criteria for propagating phase boundaries in a van der Waals fluid, *Arch. Rational Mech. Anal.* 81 (1983), 301–315.
- [13] L. TRUSKINOVSKY, Kinks versus shocks, in “Shock induced transitions and phase structures in general media”, R. Fosdick, E. Dunn, and M. Slemrod ed., *IMA Vol. Math. Appl.*, Vol. 52, Springer-Verlag, New York (1993), pp. 185–229.