

# A REMARK ABOUT MAHLER'S CONJECTURE AND THE MAXIMUM VALUE OF BOX SPLINES

ZHIQIANG XU

ABSTRACT. In this paper, we recast a special case of Mahler's conjecture by the maximum value of box splines. This is the case of polytopes with at most  $2n + 2$  facets. An asymptotic formula for univariate box splines is given. Based on the formula, Mahler's conjecture is proved in this case provided  $n$  is big enough.

## 1. INTRODUCTION

Let  $K$  be a symmetric convex body in  $\mathbb{R}^n$ , and let  $K^*$  be its polar  $\{x : |\langle y, x \rangle| \leq 1 \text{ for all } y \in K\}$ . An old conjecture of Mahler is

$$(1) \quad \text{vol}(K) \cdot \text{vol}(K^*) \geq \frac{4^n}{n!}.$$

We note that an  $n$ -dimensional parallelepiped, which has  $2n$  facets, gives equality in (1). So, the first non-trivial case of Mahler's conjecture is the symmetric convex body  $K$  with  $2n + 2$  facets. Such  $K$  can be realized, up to affine invariance, as a one-codimensional section of an  $(n + 1)$ -dimensional cube. This case has been raised as a separate problem by Ball [1]. Ball also shows an interesting relation between the special case of Mahler's conjecture and solutions of a scaling equation. For each  $r \in \mathbb{R}$ , we set

$$(2) \quad \varphi_A(r) := \text{vol}((H + rA) \cap Q_n)$$

where  $Q_n := [-\frac{1}{2}, \frac{1}{2}]^n$ ,  $A := (a_1, \dots, a_n)$  is an unit vector and  $H := \langle A \rangle^\perp$ . Throughout this paper, without loss of generality, we suppose  $0 < a_1 \leq a_2 \leq \dots \leq a_n$ . In particular, Ball shows that the special case of Mahler's conjecture is equivalent to

$$(3) \quad \varphi_A(0) \cdot E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right) \geq 1,$$

where  $\varepsilon_k$  is a sequence of mutually independent random variables with distribution  $P\{1\} = P\{-1\} = 1/2$ . In [7], the authors prove (3) for  $n \leq 8$  by reducing the problem to a search over a finite set of polytopes for each fixed dimension.

An interesting observation is that  $\varphi_A(\cdot)$  is a box spline, a popular tool in approximation theory. Therefore, we can recast the special case of Mahler's conjecture by the maximum value of box splines. Using the saddle point approximation in statistics, we give an asymptotic formula of  $\varphi_A(\cdot)$ . Based on the asymptotic formula, we obtain the following result:

**Theorem 1.** *Suppose that there is a constant  $c_0$  so that  $a_n/a_1 < c_0$  for any  $n$ . Then there exists a positive integer  $N_0(c_0)$  so that*

$$\varphi_A(0) \cdot E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right) \geq 1$$

when  $n \geq N_0(c_0)$ .

This paper is organized as follows. In Section 2, we introduce box splines and show the relation between box splines and the special case of Mahler's conjecture. In Section 3, we use the saddle point approximation to give an asymptotic formula of univariate box splines. Section 4 presents the proof of Theorem 1.

## 2. BOX SPLINES

Suppose that  $M$  is a  $s \times n$  matrix. The *box spline*  $B(\cdot|M)$  associated with  $M$  is the distribution given by the rule [2, 3]

$$(4) \quad \int_{\mathbb{R}^s} B(x|M)\phi(x)dx = \int_{[0,1]^n} \phi(Mu)du, \quad \phi \in \mathcal{D}(\mathbb{R}^s).$$

By taking  $\phi = \exp(-i\zeta \cdot)$  in (4), we obtain the Fourier transform of  $B(\cdot|M)$  as

$$(5) \quad \widehat{B}(\zeta|M) = \prod_{j=1}^n \frac{1 - \exp(-i\zeta^T m_j)}{i\zeta^T m_j}, \quad \zeta \in \mathbb{C}^s.$$

The following formula shows the relation between box splines and the volume of the section of unit cube (see [4], page 2):

$$(6) \quad B(x|M) = \frac{\text{vol}_n(P \cap [0, 1]^n)}{\sqrt{|\det(MM^T)|}},$$

where  $P := \{y : My = x, y \in \mathbb{R}_+^n\}$ . Set  $A := (a_1, \dots, a_n)$ . Recall that  $\sum_j a_j^2 = 1$  and  $0 < a_1 \leq a_2 \leq \dots \leq a_n$ . Then combining (2) and (6) we have that

$$\varphi_A(\cdot) = B(\cdot + (a_1 + \dots + a_n)/2|A).$$

Also, noting  $B(\cdot|A)$  reaching the maximum value at  $(a_1+a_2+\cdots+a_n)/2$ , (3) is equivalent to

$$\max_x B(x|A) \geq \frac{1}{E(|\sum_{k=1}^n a_k \varepsilon_k|)}.$$

### 3. AN ASYMPTOTIC FORMULA OF UNIVARIATE BOX SPLINES

In this section, we shall present an asymptotic formula of  $B(\cdot|A)$ . In [8], Unser et. al. proved that  $B(\cdot|A)$  tends to the Gaussian function as  $n$  increase provided  $a_1 = a_2 = \cdots = a_n$ . Here, using the saddle point approximation in statistics, for the general matrix  $A$ , we can show the box spline  $B(\cdot|A)$  also converges to the Gaussian function as  $n$  increases:

#### Theorem 2.

$$\lim_{n \rightarrow \infty} B(x|A) = \sqrt{6/\pi} \exp(-6(x - \sum_j a_j/2)^2),$$

where the limit may be taken pointwise or in  $L^p(\mathbb{R})$ ,  $p \in [2, \infty)$ .

*Proof.* The saddle point approximation of  $B(\cdot|A)$  is (see Theorem 6.1 and 6.2 in [5])

$$(7) \quad \frac{1}{(2\pi)^{1/2} |K''(s_0)|^{1/2}} \exp(K(s_0) - s_0 x).$$

Here,

$$K(s) := \ln \prod_i \frac{\exp(a_i s) - 1}{a_i s}$$

and  $s_0$  satisfies  $K'(s_0) = x$ . We can consider  $s_0$  as a function of  $x$ . So,

$$(8) \quad K'(s_0) - x = 0$$

defines an implicit relationship between  $s_0$  and  $x$ . Noting  $K'(0) = (a_1 + \cdots + a_n)/2$ , the equation (8) implies that  $s_0((a_1 + \cdots + a_n)/2) = 0$ . Also, by (8), we have

$$(9) \quad K''(s_0) s'_0 - 1 = 0,$$

which implies that

$$s'_0((a_1 + \cdots + a_n)/2) = 1/K''(0) = 12/\sum_j a_j^2 = 12.$$

Using the similar method, we have

$$s''_0((a_1 + \cdots + a_n)/2) = 0,$$

and

$$s_0'''((a_1 + \cdots + a_n)/2) = 864 \left( \sum_{j=1}^n a_j^4 \right) / 5 = O(1/n).$$

Using Taylor expansion at  $(a_1 + \cdots + a_n)/2$ , one has

$$(10) \quad s_0(x) = 12 \cdot (x - (a_1 + \cdots + a_n)/2) + O(1/n).$$

Also, by (9), we have

$$K''(s_0) = 1/s_0' = 1/12 + O(1/n).$$

Combining Taylor expansion of  $K(\cdot)$  at 0 and (10), one has

$$K(s_0) = 6 \left( \sum_j a_j \right) (x - (a_1 + \cdots + a_n)/2) + 6(x - (a_1 + \cdots + a_n)/2)^2 + O(1/n).$$

By (10), we have

$$\begin{aligned} s_0 \cdot x &= 12(x - (a_1 + \cdots + a_n)/2)x + O(1/n) \\ &= 12(x - (a_1 + \cdots + a_n)/2)^2 + 6 \left( \sum_j a_j \right) (x - (a_1 + \cdots + a_n)/2) + O(1/n). \end{aligned}$$

From (7), the saddle point approximation of  $B(x|A)$  is

$$\sqrt{6/\pi} \exp(-6(x - \sum_j a_j/2)^2) + O(1/n).$$

The properties of the saddle point approximation imply this theorem.  $\square$

#### 4. PROOF OF THE MAIN RESULT

To prove the main result, we firstly introduce a lemma.

**Lemma 3.** *Put*

$$F(s) := \frac{2}{\pi} \int_0^\infty (1 - |\cos(t/\sqrt{s})|^s) t^{-2} dt, \quad s > 0.$$

*Then*

$$E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right) \geq F(a_n^{-2}).$$

*Proof.* By Lemma 1.3 in [6], we have

$$E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right) \geq \sum_{k=1}^n a_k^2 F(a_k^{-2}).$$

Since  $F$  is an increasing function (Lemma 1.4 [6]), we have

$$E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right) \geq \sum_{k=1}^n a_k^2 F(a_k^{-2}) \geq F(a_n^{-2}).$$

□

*Proof of Theorem 1.* To prove the theorem, we only need to prove that there exists a positive integer  $N_0$  so that

$$\max_x B(x|A) \geq \frac{1}{E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right)}$$

when  $n \geq N_0$ . Theorem 2 implies that

$$(11) \quad \lim_{n \rightarrow \infty} \max_x B(x|A) = \sqrt{6/\pi}.$$

Since  $\sum_j a_j^2 = 1$  and  $a_n/a_1 < c_0$ , we have  $\lim_{n \rightarrow \infty} 1/a_n = \infty$ . By [6], we have  $\lim_{s \rightarrow \infty} F(s) = \sqrt{2/\pi}$ . We choose  $\varepsilon_0$  so that

$$0 < \varepsilon_0 < \frac{\sqrt{6/\pi} - \sqrt{\pi/2}}{2}.$$

By Lemma 3, there is a positive integer  $N_1$  so that

$$(12) \quad \frac{1}{E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right)} \leq \frac{1}{F(a_n^{-2})} \leq \sqrt{\pi/2} + \varepsilon_0$$

provided  $n \geq N_1$ . The equation (11) implies that there is a positive integer  $N_2(c_0)$  so that

$$(13) \quad \max_x B(x|A) \geq \sqrt{6/\pi} - \varepsilon_0$$

provided  $n \geq N_2(c_0)$ . We set  $N_0(c_0) := \max\{N_1, N_2(c_0)\}$ . Noting that

$$\sqrt{6/\pi} - \varepsilon_0 \geq \sqrt{\pi/2} + \varepsilon_0,$$

combining (12) and (13), we have

$$\max B(x|A) \geq \frac{1}{E \left( \left| \sum_{k=1}^n a_k \varepsilon_k \right| \right)}$$

when  $n \geq N_0(c_0)$ . □

**Acknowledgments.** Zhiqiang Xu was supported by National Natural Science Foundation of China (10871196) .

## REFERENCES

1. K. Ball, Mahler's conjecture and wavelets, *Discrete and Computational Geometry*, Vol. 13, No.1(1995).
2. C. de Boor and R. DeVore, Approximation by smooth multivariate splines, *Trans. Amer. Math. Soc.*, 276 (1983) 775-788.
3. C. de Boor and K. Höllig, B-splines from parallelepipeds, *J. Anal. Math.*, 42(1982/83)99-115.
4. C. de Boor, K. Höllig and S. Riemenschneider, *Box Splines*, Springer-Verlag, New York, 1993.
5. Daniels, H. E., Saddlepoint approximations in statistics, *Annals Math. Statist.*, 25(1954) 631-650
6. U. Haagerup, The best constants in the Khintchine inequality, *Studia Mathematica*, T. LXX. (1982).
7. M. A. Lopez and S. Reisner, A special case of Mahler's conjecture, *Discrete Comp. Geom.* 20:163-177(1998).
8. M. Unser, A. Aldroubi and M. Eden, On the asymptotic convergence of B-spline wavelets to Gabor functions, *IEEE Transactions on informations theory*, Vol.38, No.2, 1992.

**Authors' addresses:**

Zhiqiang Xu, LSEC, Inst. Comp. Math., Academy of Mathematics and System Sciences, Chinese Academy of Sciences, Beijing, 100080, China, xuzq@lsec.cc.ac.cn