

**(DISK, ESSENTIAL SURFACE) PAIRS OF HEEGAARD
SPLITTINGS THAT INTERSECT IN ONE POINT**

JUNG HOON LEE

ABSTRACT. We consider a Heegaard splitting $M = H_1 \cup_S H_2$ of a 3-manifold M having an essential disk $D \subset H_1$ and an essential surface $F \subset H_2$ with $|D \cap F| = 1$. (We require that $\partial F \subset S$ when H_2 is a compressionbody with $\partial_- H_2 \neq \emptyset$.)

Let F be a genus g surface with n boundary components. From $M = H_1 \cup_S H_2$, we obtain a genus $g(S) + 2g + n - 2$ Heegaard splitting $M = H'_1 \cup_{S'} H'_2$ by cutting H_2 along F and attaching $F \times I$ to H_1 along $\partial F \times I$. As an application, by using a theorem due to Casson and Gordon [1], we give examples of 3-manifolds having two Heegaard splittings of distinct genera where one of the two Heegaard splittings is a strongly irreducible non-minimal genus splitting and it is obtained from the other by the above construction.

1. INTRODUCTION

Every compact 3-manifold M admits a Heegaard splitting and there are various Heegaard splittings as the genus varies. If g is the minimal genus of Heegaard splittings of M , then for each $g' > g$ there exists at least one Heegaard splitting of genus g' — a splitting obtained by stabilizations.

From a Heegaard splitting, we can obtain another Heegaard splitting of different genus which is not just a stabilization if the original one has certain embedded surfaces that intersect in one point. A stabilized Heegaard splitting $H_1 \cup_S H_2$, which has essential disks $D_1 \subset H_1$ and $D_2 \subset H_2$ with $|D_1 \cap D_2| = 1$, can be destabilized and the genus goes down.

Concerning (Disk, Annulus) pairs, many people [11], [10], [8], [4] considered several notions as in the following and their relations with other notions on Heegaard splittings.

- Essential disk $D \subset H_1$ and essential annulus $A \subset H_2$ with $D \cap A = \emptyset$
- Essential disk $D \subset H_1$ and essential annulus $A \subset H_2$ with ∂D equal to one component of ∂A
- Essential disk $D \subset H_1$ and essential annulus $A \subset H_2$ with $|D \cap A| = 1$
- Essential disk $D \subset H_1$ and spanning annulus A in a compressionbody H_2 with $D \cap A = \emptyset$
- Essential disk $D \subset H_1$ and spanning annulus A in a compressionbody H_2 with $|D \cap A| = 1$

In [4], the author considered a Heegaard splitting $H_1 \cup_S H_2$ having an essential disk $D \subset H_1$ and an essential annulus $A \subset H_2$ with $|D \cap A| = 1$ and it was shown that such a Heegaard splitting has the disjoint curve property, a notion which was

1991 *Mathematics Subject Classification.* Primary 57N10, 57M50.

Key words and phrases. Heegaard splitting, essential surface, strongly irreducible.

introduced by Thompson [14], and another Heegaard splitting $H'_1 \cup_{S'} H'_2$ can be obtained from $H_1 \cup_S H_2$ by removing a neighborhood of A from H_2 and attaching it to H_1 . In this case the genus of Heegaard splitting remains unchanged.

In this paper, we consider a Heegaard splitting $H_1 \cup_S H_2$ of a 3-manifold M having an essential disk $D \subset H_1$ and an essential surface $F \subset H_2$ with $|D \cap F| = 1$. We require that $\partial F \subset S$ when H_2 is a compressionbody with $\partial_- H_2 \neq \emptyset$. We denote it as a *strong* (D, F) pair for consistency of terminology with [8]. First we show that if F has genus g and n boundary components, the distance $d(S)$ of $H_1 \cup_S H_2$ is bounded above by $2g + n$ (Theorem 2.3).

From $H_1 \cup_S H_2$ we can obtain another Heegaard splitting $H'_1 \cup_{S'} H'_2$ by removing a neighborhood of F from H_2 and attaching it to H_1 .

Theorem 1.1. *Let $H_1 \cup_S H_2$ be a Heegaard splitting of a 3-manifold with a strong (D, F) pair. Let H'_1 be obtained from H_1 by attaching $F \times I$ along $\partial F \times I$ and H'_2 be obtained from H_2 by cutting along F . Then $H'_1 \cup_{S'} H'_2$ is a Heegaard splitting of genus $g(S) + 2g + n - 2$.*

The construction of new Heegaard surface in Theorem 1.1 resembles quite a bit the *Haken sum* in Moriah, Schleimer, and Sedgwick's paper [6]. In that paper, they considered the Haken sum of Heegaard surface with copies of an incompressible surface in the manifold and obtained infinitely many distinct Heegaard splittings. Also there are related works by Kobayashi [3], and Lustig and Moriah [5]. However, in our case the essential surface lives in one of the compressionbodies (Fig. 1).




FIGURE 1. Haken sum and compressing along an essential surface in a compressionbody

In Theorem 3.5, it is shown that $H'_1 \cup_{S'} H'_2$ has the disjoint curve property if F is not a disk.

In most part of the paper, we are considering the case $\partial F \subset \partial_+ H_2$ when H_2 is a compressionbody with $\partial_- H_2 \neq \emptyset$. In the last part of section 3, we briefly consider the case when F is a spanning annulus in a compressionbody, and show a corresponding result (Theorem 3.6).

For $g \geq 1$ or $n \geq 3$, the genus of $H'_1 \cup_{S'} H'_2$ is greater than that of $H_1 \cup_S H_2$. We give examples of 3-manifolds having two Heegaard splittings of distinct genera where one of the two Heegaard splittings is a strongly irreducible non-minimal

genus splitting and it is obtained from the other by the method in Theorem 1.1. The examples are constructed by doing $1/q$ -Dehn surgery ($|q| \geq 6$) on certain knots and a theorem due to Casson and Gordon is used to show strong irreducibility.

Theorem 1.2. *Let K be a knot with the following properties.*

- *A minimal genus Seifert surface F (of genus g) for K is free.*
- *Every tunnels of an unknotting tunnel system $\{t_1, t_2, \dots, t_t\}$ can be isotoped to lie on F and mutually disjoint.*
- *$\bigcup_{i=1}^t t_i$ cuts F into a connected subsurface $F' \subset F$.*
- *$t + 1 < 2g$*

Let $K(1/q)$ be the manifold obtained by doing $1/q$ -Dehn surgery ($|q| \geq 6$) on K in S^3 . Then $K(1/q)$ has a genus $2g$ strongly irreducible Heegaard splitting and a genus $t + 1$ Heegaard splitting, and the two are related by the construction in Theorem 1.1.

In particular, if K is a torus knot, $K(1/q)$ is a Seifert fibered space over S^2 with three exceptional fibers [8], [13], and Theorem 1.2 gives some insight to the relation of a vertical splitting and a horizontal splitting of such manifolds.

2. HEEGAARD SPLITTING WITH A STRONG (D, F) PAIR

First we show that for a (D, F) pair with $|D \cap F| = 1$, in fact the essentiality of F follows automatically from incompressibility. In other words, we have the following.

Proposition 2.1. *For a Heegaard splitting $H_1 \cup_S H_2$, let D be an essential disk in H_1 and F ($\partial F \neq \emptyset$, $\partial F \subset S$) be an incompressible surface in H_2 such that $|D \cap F| = 1$. Then F is essential in H_2 .*

Proof. Suppose F is not essential in H_2 . Then F is parallel to a subsurface $F' \subset S$ (rel. ∂F). When we go around ∂D , we pass through $\partial F'$ from $S - F'$ to the interior of F' at some time. We should pass through $\partial F'$ at least once more to go around all of ∂D . This is a contradiction since $|D \cap F| = |D \cap \partial F| = 1$ and $\partial F = \partial F'$. \square

Now we consider the distance, due to Hempel [2], of a Heegaard splitting with a strong (D, F) pair. The *distance* $d(S)$ of a Heegaard splitting $H_1 \cup_S H_2$ is the smallest number $n \geq 0$ so that there is a sequence of essential simple closed curves $\alpha_0, \dots, \alpha_n$ in S with α_0 bounding a disk in H_1 , α_n bounding a disk in H_2 and for each $1 \leq i \leq n$, α_{i-1} and α_i can be isotoped in S to be disjoint.

We need the the following technical lemma on boundary compression by Morimoto to get an upper bound for distance.

Lemma 2.2. *(Lemma 5.1 of [9]) Let W be a compact orientable 3-manifold, and let F be an essential surface properly embedded in W such that $\partial F \neq \emptyset$ and ∂F is contained in a single component of ∂W . Let F' be the 2-manifold obtained from F by a boundary compression. Then F' is incompressible and has a component which is not ∂ -parallel. Hence F' is essential.*

Theorem 2.3. *Let $H_1 \cup_S H_2$ be a genus ≥ 2 Heegaard splitting with a strong (D, F) pair. If F has genus g and n boundary components, then the distance $d(S) \leq 2g + n$.*

Proof. Let $\partial F = \beta_1 \cup \dots \cup \beta_n$ and $|D \cap \beta_1| = 1$. Since F is incompressible and not boundary parallel and $\partial F \subset S$, F intersects a meridian disk system of H_2 . By standard innermost disk and outermost arc arguments, we may assume that there is a boundary compressing disk Δ for F , where the boundary compression occurs

toward S . Let $\partial\Delta = \alpha \cup \beta$ where α is an essential arc in F and β is an arc in S . We construct sequence of essential simple closed curves $\alpha_0, \dots, \alpha_k$ with α_0 bounding a disk in H_1 , α_k bounding a disk in H_2 and for each $1 \leq i \leq k$, α_{i-1} and α_i can be isotoped in S to be disjoint, dividing into two cases according to n .

Case 1. $n = 1$.

Take two parallel copies of D in H_1 and connect them with a band along β_1 and push the band slightly into the interior of H_1 to make a disk $D' \subset H_1$. Since $\partial D'$ bounds a once punctured torus in S and $H_1 \cup_S H_2$ is a genus ≥ 2 Heegaard splitting, D' is an essential disk in H_1 . Note that $\partial D'$ is disjoint from β_1 . Take $\partial D'$ as α_0 and β_1 as α_1 .

Case 2. $n > 1$.

In this case, take ∂D as α_0 and any β_i ($i \neq 1$) as α_1 .

Both in Case 1 and Case 2, boundary compress F along Δ to get an essential surface $F_{(1)}$ by Lemma 2.2. All the boundary components of $F_{(1)}$ can be made disjoint from ∂F . Take any component of $\partial F_{(1)}$ as α_2 . Boundary compress $F_{(1)}$ to get an essential surface $F_{(2)}$ by Lemma 2.2. All the boundary components of $F_{(2)}$ can be made disjoint from $\partial F_{(1)}$. Take any component of $\partial F_{(2)}$ as α_3 . In this way, we successively boundary compress until we get an essential disk in H_2 by Lemma 2.2. We can check that the possible maximum number of boundary compressions is $2g + n - 1$. So the possible maximum length sequence of α_i 's would be $\alpha_0, \alpha_1, \dots, \alpha_{2g+n}$. So we conclude that $d(S) \leq 2g + n$. \square

3. OBTAINING NEW HEEGAARD SPLITTINGS

We consider attaching $F \times I$ to a handlebody along $\partial F \times I$. Let $g(X)$ denote the genus of X .

Lemma 3.1. *Let $\gamma_1, \dots, \gamma_n$ be mutually disjoint loops on the boundary of a handlebody H and D be an essential disk of H such that $|\partial D \cap \gamma_1| = 1$ and $\partial D \cap \gamma_i = \emptyset$ ($i = 2, \dots, n$). Let F be a genus g surface with n ($n \geq 1$) boundary components β_1, \dots, β_n .*

If we attach $F \times I$ to H along $\partial F \times I$ so that $\beta_i \times I$ is attached to $N(\gamma_i; \partial H) \cong \gamma_i \times I$ ($i = 1, 2, \dots, n$), then the resulting manifold is a handlebody of genus $g(H) + 2g + n - 2$.

Proof. Let p be the intersection point $D \cap \gamma_1$. Consider the neighborhood $D \times I$ in H and $\gamma_1 \times I$ in ∂H . We can assume that $\partial(D \times I) \cap (\gamma_1 \times I)$ is a small rectangle R containing p . Let R' be the rectangle in $\beta_1 \times I$ that is attached to R .

Since F is a genus g surface with n boundary components, there are mutually disjoint essential arcs a_1, \dots, a_{2g+n-1} in F such that F cut along $a_1 \cup \dots \cup a_{2g+n-1}$ is a disk. In particular, take such an essential arc system so as to satisfy that one of the two points of ∂a_1 is attached to p . More precisely, we take the rectangular parallelepiped neighborhood $a_1 \times I \times I$ of a_1 in $F \times I$ to be equal to $R' \times I$.

Let H' be $cl(H - (D \times I))$. Since $|D \cap \gamma_1| = 1$, D is a non-separating essential disk in H . So H' is a handlebody of genus $g(H) - 1$. Attach a rectangular parallelepiped neighborhood $a_i \times I \times I$ of a_i taken in $F \times I$ to H' along $\partial a_i \times I \times I$ for each $i = 2, \dots, 2g + n - 1$. Since each $a_i \times I \times I$ ($i = 2, \dots, 2g + n - 1$) can be considered as a 1-handle, the resulting manifold H'' is a genus $g(H) + 2g + n - 3$ handlebody.

Observe that $cl(F \times I - (\bigcup_{i=1}^{2g+n-1} a_i \times I \times I))$ is homeomorphic to a 3-ball B , which is attached to H'' along two subdisks of its boundary. Then $H''' = H'' \cup B$

is a handlebody of genus $g(H) + 2g + n - 2$. Observe also that $(D \times I) \cup (R' \times I)$ is a 3-ball attached to H''' along $(D \times \partial I) \cup$ (three faces of $\partial(R' \times I)$), which is a disk on the boundary of a 3-ball. So the genus remains unchanged after attaching $(D \times I) \cup (R' \times I)$ to H''' . Hence we conclude that the resulting manifold after attaching $F \times I$ to H along $\partial F \times I$ is a genus $g(H) + 2g + n - 2$ handlebody. \square

Now we consider removing a neighborhood of incompressible surface from a compressionbody. The following lemma is well-known. It can be found, for example, as Lemma 2 in Schulten's paper [12].

Lemma 3.2. *Let F ($\partial F \neq \emptyset$) be an incompressible surface properly embedded in a compressionbody H with $\partial F \subset \partial_+ H$. Then F cuts H into compressionbodies (or a compressionbody).*

Using Lemma 3.1 and Lemma 3.2, we give a proof of Theorem 1.1.

Proof. (of Theorem 1.1.) By Lemma 3.1, H'_1 is a handlebody of genus $g(S) + 2g + n - 2$. By Lemma 3.2, H'_2 is union of two compressionbodies or a compressionbody. Since $\partial H'_1$ is same as $\partial_+ H'_2$, H'_2 is connected. Therefore $H'_1 \cup_{S'} H'_2$ is a Heegaard splitting of genus $g(S) + 2g + n - 2$. \square

Corollary 3.3. *Let $H_1 \cup_S H_2$ be a Heegaard splitting of a 3-manifold M with a strong (D, A) pair where A is an annulus. Let H'_1 be obtained from H_1 by attaching $A \times I \subset H_2$ along $\partial A \times I$ and H'_2 be obtained from H_2 by cutting along A . Then $H'_1 \cup_{S'} H'_2$ is a Heegaard splitting of same genus with $H_1 \cup_S H_2$.*

Remark 3.4. Corollary 3.3 is a generalization of Definition 14 of [12].

A Heegaard splitting $H_1 \cup_S H_2$ is said to have the *disjoint curve property* ([14]) if there are essential disks $D_1 \subset H_1$, $D_2 \subset H_2$ and an essential loop $\gamma \subset S$ such that $(\partial D_1 \cup \partial D_2) \cap \gamma = \emptyset$. It is equivalent to that the distance $d(S)$ is less than or equal to two. The newly obtained Heegaard splitting $H'_1 \cup_{S'} H'_2$ of Theorem 1.1 has the disjoint curve property if the switch from $H_1 \cup_S H_2$ to $H'_1 \cup_{S'} H'_2$ is not a destabilization.

Theorem 3.5. *If F is not a disk, the Heegaard splitting $H'_1 \cup_{S'} H'_2$ obtained in Theorem 1.1 has the disjoint curve property.*




FIGURE 2. An essential disk E' in H'_1 satisfying the disjoint curve property

Proof. Recall the proof of Lemma 3.1. In the proof of Lemma 3.1, H''' was obtained from H'' by attaching a 1-handle. Consider a meridian disk (co-core) E of the 1-handle. Also remember that $H'_1 = H_1 \cup (F \times I)$ was obtained from H''' by attaching a 3-ball along a 2-disk on its boundary. Then E is enlarged to an essential disk E' in H'_1 , which can be taken as two parallel copies of D attached by a band in $F \times I$. See Fig. 2. Since the band is equivalent to an $(\text{arc}) \times I$ in $F \times I$ with both endpoints of the arc in the same component of ∂F , we can take an essential loop $\gamma \subset F$ which is disjoint from E' . Since F is incompressible in H_2 , we can see that γ is an essential loop in the new Heegaard surface S' . Take a boundary compressing disk $\Delta \subset H_2$ for F . Let $\partial\Delta = \alpha \cup \beta$ where α is an essential arc in F . Then after cutting H_2 along F , Δ is an essential disk in H'_2 . We may assume that α belongs to $F \times \{0\}$ and β belongs to $F \times \{1\}$. So Δ is disjoint from γ . So we conclude that the triple (E', Δ, γ) satisfies the disjoint curve property. \square

Now we consider the case when F is a spanning annulus in a compressionbody. Given a compressionbody H with $\partial_- H \neq \emptyset$, there exists a meridian disk system $\{D_1, \dots, D_k\}$ of H such that H cut along $\bigcup_{i=1}^k D_i$ is $\partial_- H \times I \cup$ (possibly empty) 3-balls. A *spanning annulus* of a compressionbody H is an annulus which can be expressed as $\gamma \times I$ in the compressionbody structure, where γ is an essential loop in $\partial_- H$. By showing the analogue of Lemma 3.1 and Lemma 3.2, we can state the following.




FIGURE 3. Compressing along a spanning annulus

Theorem 3.6. *Let $H_1 \cup_S H_2$ be a Heegaard splitting of a 3-manifold M with a strong (D, A) pair where A is a spanning annulus. Let $A = \gamma \times I$ in the compressionbody structure and $\gamma \subset \Sigma \subset \partial_- H_2$ and Σ has genus g . Let H'_1 and H'_2 be obtained as follows (Fig. 3).*

- H'_1 is obtained from H_1 by attaching $A \times I$ along $(\partial A - \gamma) \times I$ and attaching $\Sigma \times I$ along $\gamma \times I$.
- H'_2 is obtained from H_2 by cutting along A and shrinking the part adjacent to Σ

Then $H'_1 \cup_{S'} H'_2$ is a Heegaard splitting of genus $g(S) + g - 1$. In particular, if Σ is a torus, it has same genus with $H_1 \cup_S H_2$.

Proof. By following the procedure as in the proof of Lemma 3.1, we can see that H'_1 is a compressionbody of genus $g(S) + g - 1$. One remarkable point is that the partition of components of ∂M is changed for the Heegaard splitting — Σ belongs to $\partial_- H_2$ before the change and $\partial_- H'_1$ after the change.

By cutting H_2 along A , H_2' becomes a genus $g(S) + g - 1$ compressionbody and parts of Σ is connected to $\partial_+ H_2$. We conclude that $H_1' \cup_{S'} H_2'$ is a Heegaard splitting of genus $g(S) + g - 1$. \square

4. EXAMPLES

Let K be a knot admitting a minimal genus free Seifert surface F of genus g . Then F is incompressible in $cl(S^3 - N(F))$. Also F is incompressible in the product neighborhood $N(F) = F \times I$. Since $N(F)$ and $cl(S^3 - N(F))$ are handlebodies, this gives a Heegaard splitting $N(F) \cup_{\Sigma} cl(S^3 - N(F))$ of S^3 .

Now we are going to construct a strongly irreducible Heegaard splitting from Σ by Dehn surgery on K . Remove a neighborhood $N(K)$ from S^3 . Let $K(1/q)$ denote the manifold obtained by $1/q$ -filling on $cl(S^3 - N(K))$. We can assume that the filling solid torus T is attached to $N(F) = F \times I$ along an annulus $\partial F \times I$. Note that if we perform $1/q$ -filling, a meridian curve $(1, 0)$ of the filling solid torus is mapped to $(1, q)$ curve and longitude $(0, 1)$ of filling solid torus is mapped to longitude $(0, 1)$. So $N(F) \cup T$ is a handlebody. Then we get the Heegaard splitting $(N(F) \cup T) \cup_{\Sigma'} cl(S^3 - N(F))$ for $K(1/q)$. (Alternatively, we can regard Σ' is obtained from Σ by Dehn twists on K $|q|$ times.) By a theorem due to Casson and Gordon [1], Σ' is a strongly irreducible Heegaard splitting if $|q| \geq 6$. Here we refer the statements in ([7], Appendix).

Theorem 4.1. (Casson-Gordon) *Suppose $M = H_1 \cup_{\Sigma} H_2$ is a weakly reducible Heegaard splitting for the closed manifold M . Let K be a simple closed curve in Σ such that $\Sigma - N(K)$ is incompressible in both H_1 and H_2 . Then Σ' , for all $|q| \geq 6$, is a strongly irreducible Heegaard splitting for the Dehn filled manifold $M(1/q)$.*

Proof. (of Theorem 1.2.) $K(1/q)$ has a genus $2g$ Heegaard splitting $(N(F) \cup T) \cup_{\Sigma'} cl(S^3 - N(F))$. The strong irreducibility of it is already shown above.

Note that $cl(S^3 - N(K \cup (\bigcup_{i=1}^t t_i)))$ is a genus $t + 1$ handlebody and this handlebody remains untouched during the $1/q$ -surgery on K . So $(T \cup (\bigcup_{i=1}^t N(t_i))) \cup_{S'} cl(S^3 - N(K \cup (\bigcup_{i=1}^t t_i)))$ is a genus $t + 1$ Heegaard splitting for the Dehn filled manifold $K(1/q)$.

By assumption of Theorem 1.2, every tunnels of an unknotting tunnel system $\{t_1, t_2, \dots, t_t\}$ for K can be isotoped to lie on F and mutually disjoint, and $\bigcup_{i=1}^t t_i$ cuts F into a connected subsurface $F' \subset F$. Since F' is a subsurface of F , F' is incompressible in $cl(S^3 - N(K \cup (\bigcup_{i=1}^t t_i)))$. We can see that the meridian of the filling solid torus T intersect F' in one point since the determinant of the matrix $\begin{pmatrix} 1 & 0 \\ q & 1 \end{pmatrix}$ is 1. So the genus $t + 1$ Heegaard splitting $(T \cup (\bigcup_{i=1}^t N(t_i))) \cup_{S'} cl(S^3 - N(K \cup (\bigcup_{i=1}^t t_i)))$ has a strong (D, F') pair. Since $t + 1 < 2g$, we can see that the genus $2g$ strongly irreducible non-minimal genus splitting $(N(F) \cup T) \cup_{\Sigma'} cl(S^3 - N(F))$ and the genus $t + 1$ splitting $(T \cup (\bigcup_{i=1}^t N(t_i))) \cup_{S'} cl(S^3 - N(K \cup (\bigcup_{i=1}^t t_i)))$ is related by the construction as in Theorem 1.1. \square

In particular, if K is a torus knot, $K(1/q)$ is a Seifert fibered space over S^2 with three exceptional fibers. The splitting induced by an unknotting tunnel is the ‘‘vertical’’ splitting and the strongly irreducible non-minimal genus splitting is the ‘‘horizontal’’ splitting [8], [13]. Hence Theorem 1.2 gives some insight to the relation of a vertical splitting and a horizontal splitting of such manifolds (Fig. 4) (Fig. 5).




FIGURE 4. Genus $t + 1$ Heegaard splitting with a strong (D, F') pair




FIGURE 5. Strongly irreducible non-minimal genus Heegaard splitting of genus $2n$

REFERENCES

- [1] A. Casson and C. Gordon, *Manifolds with irreducible Heegaard splittings of arbitrary large genus*, Unpublished.
- [2] J. Hempel, *3-manifolds as viewed from the curve complex*, *Topology* **40** (2001), no. 3, 631–657.
- [3] T. Kobayashi, *A construction of 3-manifolds whose homeomorphism classes of Heegaard splittings have polynomial growth*, *Osaka J. Math.* **29** (1992), no. 4, 653–674.
- [4] J. H. Lee, *On (Disk, Annulus) pairs of Heegaard splittings that intersect in one point*, to appear in *Bull. Korean Math. Soc.*
- [5] M. Lustig and Y. Moriah, *3-manifolds with irreducible Heegaard splittings of high genus*, *Topology* **39** (2000), no. 3, 589–618.
- [6] Y. Moriah, S. Schleimer, and E. Sedgwick, *Heegaard splittings of the form $H + nK$* , *Communications in Analysis and Geometry*, **14** (2006), no. 2, 215–247.

- [7] Y. Moriah and J. Schultens, *Irreducible Heegaard splittings of Seifert fibered spaces are either vertical or horizontal*, *Topology* **37** (1998), no. 5, 1089–1112.
- [8] Y. Moriah and E. Sedgwick, *The Heegaard structure of Dehn-filled manifolds*, arXiv:0706.1927,
- [9] K. Morimoto, *Planar surfaces in a handlebody and a theorem of Gordon-Reid*, *Proceedings of Knots 96* edited by Shin'ichi Suzuki, pp. 123–146.
- [10] T. Saito, *Disjoint pairs of annuli and disks for Heegaard splittings*, *J. Korean Math. Soc.* **42** (2005), no. 4, 773–793.
- [11] S. Schleimer, *The disjoint curve property*, *Geom. Topol.* **8** (2004), 77–113.
- [12] J. Schultens, *Additivity of tunnel number for small knots*, *Comment. Math. Helv.* **75** (2000), no. 3, 353–367.
- [13] E. Sedgwick, *The irreducibility of Heegaard splittings of Seifert fibered spaces*, *Pacific J. Math.* **190** (1999) no. 1, 173–199.
- [14] A. Thompson, *The disjoint curve property and genus 2 manifolds*, *Topology Appl.* **97** (1999), no. 3, 273–279.

SCHOOL OF MATHEMATICS, KIAS, 207-43, CHEONGNYANGNI 2-DONG, DONGDAEMUN-GU, SEOUL, KOREA. TEL: +82-2-958-3736

E-mail address: `jhlee@kias.re.kr`