

STABLE CONCORDANCE OF KNOTS IN 3-MANIFOLDS

ROB SCHNEIDERMAN

ABSTRACT. Knots and links in 3-manifolds are studied by applying intersection invariants to singular concordances. The resulting link invariants generalize the Arf invariant, the mod 2 Sato-Levine invariants, and Milnor's triple linking numbers. Besides fitting into a general theory of Whitney towers, these invariants provide obstructions to the existence of a singular concordance which can be homotoped to an embedding after stabilization by connected sums with $S^2 \times S^2$. Results include classifications of stably slice links in orientable 3-manifolds, stable knot concordance in products of an orientable surface with the circle, and stable link concordance for many links of null-homotopic knots in orientable 3-manifolds.

1. INTRODUCTION

A generic homotopy of a knotted circle in a 3-manifold M traces out a properly immersed annulus in the product $M \times I$ of M with an interval. Knot invariants can be extracted from homotopy invariants of such a singular concordance by factoring out the contributions from self-homotopies of knots. For instance, the (relative) self-linking invariants of [26] are determined by the self-intersection numbers of immersed annuli in the group ring $\mathbb{Z}[\pi_1 M]$, and are characterized as the complete obstruction to finding an “improved” singular concordance in the sense that all of its singularities are paired by Whitney disks. In this setting, we will describe how a generalization of the intersection invariant of [29] can be used to define “higher order” linking invariants which give obstructions to building a “higher order Whitney tower”, meaning that intersections between annuli and the interiors of previously existing Whitney disks can be paired by a second layer of Whitney disks. As a corollary, we can in many cases determine which links can cobound a collection A of properly immersed annuli in $M \times I$ such that A is homotopic (rel boundary) to an embedding after stabilization by taking connected sums with copies of $S^2 \times S^2$. If one end of such a *stable concordance* is the unlink then it can be capped off to form *stable slice* disks.

Besides detecting many infinite families of non-concordant knots and links, the invariants defined here will give:

- A complete characterization of links in orientable 3-manifolds which are stably slice.
- A classification of stable concordance for many links of null-homotopic knots in orientable 3-manifolds with torsion-free fundamental group.
- A classification of stable concordance for knots in products $F \times S^1$ of an orientable surface with the circle.

The approach taken here fits into the developing theory of Whitney towers in 4-manifolds ([5, 6, 27, 28, 29, 30]), but of equal interest in the current setting is how the topology of a 3-manifold interacts with its knot theory. This interaction is reflected in the indeterminacies in the invariants coming from self-homotopies of knots, which can be studied using classical 3-dimensional structure theorems, following the approach of [19, 20, 21]. It turns out that these indeterminacies depend only on lower order invariants which can often be easily computed in the 3-manifold.

The algebraic linking invariants of [26] are defined for all knots in a large family of orientable 3-manifolds. For a link of null-homotopic knots the invariants reduce to the usual equivariant linking form, but for knots in non-trivial (free) homotopy classes these invariants are *relative* invariants, which compare homotopic knots to a chosen “base knot”. One of the main observations of [26] is that although such a choice of base knot is not canonical, there are certain preferred knots which “maximize” the geometric characterization of the invariants by “minimizing” the indeterminacies. These knots are called *spherical knots* in [26] because all their corresponding indeterminacies come from intersections between the knots and 2-spheres in the 3-manifold. (In general, knot self-homotopies project to tori in M which can contribute indeterminacies even when M is irreducible.) It is only with respect to spherical knots that the invariants of [26] classify *order 1 Whitney concordance*, i.e. give the complete obstruction to the existence of a singular concordance having all double points paired by Whitney disks.

In the present paper we take a more streamlined approach by directly prescribing the analogous “indeterminacy-minimizing” base knots (and links) which give the desired classification of stable concordance and equivalent characterizations in terms of Whitney towers. Note that we now have many more equivalence classes to examine; for instance in a non-simply-connected 3-manifold there are infinitely many order 1 Whitney concordance classes of null-homotopic knots. We will describe

how to choose appropriate knots so that the resulting indeterminacies only depend on the given equivalence class. These indeterminacies will be computable in terms of intersections between knots and 2-spheres, disks, and annuli in M .

The rest of this introduction will introduce enough terminology to state the main results, leaving detailed definitions and proofs to the body of the paper. The reader should note that there are some essentially cosmetic differences in notation between this paper and the closely related papers [26] and [29]. The notation used here is chosen to fit with the more recent literature on Whitney towers, and clarifications will be made as terminology is introduced. While a familiarity with [26] and [29] is certainly helpful, the present paper is to a large extent self-contained, including sketches of much relevant background material. In particular, properties of Whitney disks are described in detail, and also well-illustrated in the examples.

We work in the smooth oriented category, with orientations usually suppressed from notation. The unit interval $I = [0, 1]$ will be reparametrized without mention. When possible we follow the convention of knot theory to confuse an immersion with its image.

1.1. Intersection trees and stable embeddings. The homotopy invariant denoted by τ in [29] was defined for immersed 2-spheres in 4-manifolds with vanishing Wall intersection invariant by counting “higher order” intersections between Whitney disks and the 2-spheres. This invariant has since been shown to fit into an obstruction theory for immersed surfaces in 4-manifolds in terms of Whitney towers, where an order n Whitney tower \mathcal{W} has an order n obstruction $\tau_n(\mathcal{W})$ which lives in an abelian group generated by trivalent trees decorated by fundamental group elements ([5, 6, 27, 28, 30]). The vanishing of $\tau_n(\mathcal{W})$ implies the existence of an order $n + 1$ Whitney tower on the underlying immersed surfaces. As discussed next and described in detail in Section 2, in this language $\tau_0(\mathcal{W})$ corresponds to Wall’s intersection form, and $\tau_1(\mathcal{W})$ corresponds to the τ in [29].

Let $A = A_1, A_2, \dots, A_m$ be a collection of properly immersed simply connected compact oriented surfaces in an oriented 4-manifold X . Such an A is called a *Whitney tower of order zero*. Wall’s intersection $\lambda(A_i, A_j)$ and self-intersection $\mu(A_i)$ invariants count signed elements of $\pi_1 X$ associated to sheet-changing paths through the intersection points of A . The equivalent *order zero intersection tree* $\tau_0(A)$ takes values in a group $\mathcal{T}_0(\pi_1 X)$ of decorated order zero trees (oriented edges labelled by elements of $\pi_1 X$), and gives the complete obstruction to the existence of an *order 1 Whitney tower* \mathcal{W} on A , where \mathcal{W} consists of the A_i




FIGURE 1. A Whitney disk $W_{(i,j)}$ pairing intersections between surface sheets A_i and A_j , and the Y-tree t_p associated to an intersection point p between the interior of $W_{(i,j)}$ and A_k .

together with a collection of Whitney disks pairing all the intersections and self-intersections among the A_i .

For A admitting such an order one Whitney tower \mathcal{W} , the order one intersection tree $\tau_1(A) := \tau_1(\mathcal{W})$ takes values in an abelian group $\mathcal{T}_1(\pi_1 X)/\text{INT}(A)$ generated by decorated order one trees (univalent Y-trees having a single trivalent vertex). Here each Y-tree $t_p \in \tau_1(\mathcal{W})$ corresponds to a transverse intersection point p between an A_k and the interior of a Whitney disk $W_{(i,j)}$ pairing intersections between A_i and A_j . The univalent vertices of t_p inherit the labels i , j , and k ; and the oriented edges of t_p are decorated by elements of $\pi_1 X$ determined by sheet-changing paths through $W_{(i,j)}$ (see Figure 1 and Section 2.3 below). The target group includes *intersection relations* $\text{INT}(A)$ which are determined by the order zero invariants $\lambda(A_i, S)$ where S ranges over immersed 2-spheres representing generators of $\pi_2 X$.

As shown in [29], $\tau_1(A)$ does not depend on the choice of \mathcal{W} , and has the following geometric characterizations:

Theorem 1 ([29]). *The following are equivalent:*

- (i) $\tau_1(A)$ vanishes.
- (ii) A is homotopic (rel boundary) to A' admitting an order 2 Whitney tower.
- (iii) A is homotopic (rel boundary) to A' admitting a height 1 Whitney tower.

Here an *order 2 Whitney tower* includes a second layer of Whitney disks pairing all intersections between the A_i and the first layer of Whitney disks; and a *height 1 Whitney tower* is an order 1 Whitney tower such that all the interiors of the Whitney disks are disjoint from the

A_i . This geometric characterization of τ_1 corresponds to the fact that all the relations in the target are realizable by controlled manipulations of the Whitney tower.

Definition 1.1.1. A collection A of properly immersed surfaces A_i in a 4-manifold X can be *stably embedded* if for some n the A_i are homotopic (rel boundary) to pairwise disjoint embeddings in the connected sum of X with n copies of $S^2 \times S^2$.

Here the sums are assumed to be taken along balls which are disjoint from A . Note that in the literature ‘stable embedding’ usually means up to homology, rather than homotopy (see e.g. [1]).

The invariant $\tau_1(A)$ is an obstruction to homotoping A to an embedding, and the following corollary gives a stable converse:

Corollary 1. $\tau_1(A)$ vanishes if and only if A can be stably embedded.

Corollary 1 follows from applying the “Norman trick” ([24]) to the height 1 Whitney tower in statement (iii) of Theorem 1, as explained in subsection 2.5 below.

1.2. Stably slice links. A link L in a 3-manifold M is *stably slice* if the components of $L \subset M \times \{0\}$ bound a collection of properly immersed disks in $M \times I$ which can be stably embedded.

As discussed later in subsection 2.6, for a link L of null-homotopic knots in M the concordance invariant $\tau_0(L) := \tau_0(D) \in \mathcal{T}_0(\pi_1 M)$, where D is any collection of properly immersed disks in $M \times I$ bounded by $L \subset M \times \{0\}$, is the complete obstruction to L bounding an order 1 Whitney tower on immersed disks in $M \times I$. For links with vanishing order 0 obstruction, there is a similarly defined order 1 invariant:

Theorem 2. For $L \subset M$ with vanishing $\tau_0(L)$, the order 1 intersection tree $\tau_1(L) := \tau_1(D) \in \mathcal{T}_1(\pi_1 M)$ is a well-defined concordance invariant of L .

Note that in this setting the intersection relations $INT(D)$ turn out to be trivial so $\tau_1(L)$ takes values in $\mathcal{T}_1(\pi_1 M)$. We get the following characterization of stably slice links.

Corollary 2. L is stably slice if and only if $\tau_1(L) = 0 \in \mathcal{T}_1(\pi_1 M)$.

If $\pi_1 M$ is non-trivial and left-orderable, as many 3-manifold groups are [2], then $\mathcal{T}_1(\pi_1 M)$ is isomorphic to $\mathbb{Z}^\infty \oplus \mathbb{Z}_2^\infty$ (Proposition 2.3.1 below). Each element of $\mathcal{T}_1(\pi_1 M)$ can be realized by a link formed by tying copies of the Borromean rings into a trivial link (Figure 2).




FIGURE 2. Tying the Borromean rings into an unlink along bands representing elements a , b , and c in $\pi_1 M$ yields a link L , with $\tau_1(L)$ represented by a Y-tree t_p as in Figure 1. This is illustrated on the right, which shows t_p in a Whitney disk for the cancelling pair of intersection points in a null homotopy of L .

Remark 1 (Finite type invariants). The reader familiar with finite type invariants will recognize that operation of Figure 2 can be effected by a Y-graph clasper or clover surgery ([14, 12]). The links bounding order 1 Whitney towers are exactly the π -algebraically split links which arise in the setting of finite type 3-manifold invariants proposed by Garoufalidis and Levine in [13]. The equivariant triple $\bar{\mu}$ -invariants defined in [13], which characterize *surgery equivalence* of such links (in the sense of [22]), correspond to the projection of $\tau_1(L)$ to the \mathbb{Z}^∞ factors of $\mathcal{T}_1(\pi_1 M)$. Thus the \mathbb{Z}_2 factors of τ_1 detect infinite families of surgery equivalent links which are not (even stably) concordant.

Remark 2 (The Arf invariant). For a knot K in a simply connected M , $\tau_1(K) \in \mathcal{T}_1(\pi_1 M) \cong \mathbb{Z}_2$ is just the Arf invariant $\text{Arf}(K)$, which is known to be the complete obstruction to stably slicing K ([7, 23]). For K null-homotopic in general M , mapping $\pi_1 M$ to the trivial group induces $\mathcal{T}_1(\pi_1 M) \rightarrow \mathbb{Z}_2$ which takes $\tau_1(K)$ to $\text{Arf}(K)$. It is easy to construct K with trivial $\text{Arf}(K)$ but non-trivial $\tau_1(K)$ by tying an even number of Borromean rings (with varying $\pi_1 M$ decorations) into the unknot. Similar comments apply for the mod 2 reduction of the Sato-Levine invariant of a 2-component link ([25]).

Remark 3. Since $\tau_1(-L) = -\tau_1(L)$, where $-L$ denotes L with reversed orientations on all components (fixing the orientation of M), the \mathbb{Z}^∞ factors in $\tau_1(L)$ represent obstructions to the existence of a (stable) concordance between L and $-L$.




FIGURE 3. On the left, a clasp link, built from the unlink on the right, using the indicated guiding arcs.

1.3. Stable concordance of null-homotopic links. Two links L and L' in M are *stably concordant* if there exists a collection A of properly immersed annuli in $M \times I$, bounded by $L \subset M \times \{0\}$ and $L' \subset M \times \{1\}$, such that A can be stably embedded.

To classify stable concordance we will need to extend the definition of τ_1 to properly immersed *annuli* supporting an order 1 Whitney tower in such a way that Theorem 1 and Corollary 1 still hold. Restricting our attention to links of null-homotopic knots corresponds to inessential annuli, that is, proper immersions of annuli $(A, \partial A) \looparrowright (X, \partial X)$ which take the circle factors of A to the trivial element of $\pi_1 X$. In this case, the new indeterminacies present in $\tau_1(A)$ correspond to the boundaries of the Whitney disks winding around the annulus, and are computable from $\partial A \subset \partial X$. In the setting of links in 3-manifolds, these new INT indeterminacies, and others coming from non-separating 2-spheres, turn out to only depend on the lower order linking invariant $\tau_0(L)$, where the link $L \subset \partial A$ consists of the half of the boundary components of A sitting in either end of $M \times I$.

For links of null-homotopic knots, order 1 Whitney concordance is classified by $\tau_0(L) \in \mathcal{T}_0(\pi_1 M)$ (4.3 below and [27]). Any element $z \in \mathcal{T}_0(\pi_1 M)$ can be realized as $\tau_0(L_z)$ for a link L_z created by adding clasps to the unlink, with the clasps guided by loops determining the group elements in a representative of z (See Figure 3).

Fixing such L_z , for any L with $\tau_0(L) = z$ we define the *relative order 1 intersection tree* $\tau_1(L_z, L)$:

$$\tau_1(L_z, L) := \tau_1(A) \in \frac{\mathcal{T}_1(\pi_1 M)/\text{INT}(z)}{\Phi(z)}$$

where A is any singular concordance between L_z and L admitting an order 1 Whitney tower. Here, the intersection relations $\text{INT}(z) := \text{INT}(A)$ turn out to only depend on z ; and the *indeterminacy subgroup* $\Phi(z) < \mathcal{T}_1(\pi_1 M)/\text{INT}(z)$, which is determined by order 1 intersections between 2-spheres and L_z , also turns out to only depend on z (Lemma 3.2.1, and 3.3).

In irreducible 3-manifolds $\Phi(z)$ is trivial for all z , and in many 3-manifolds the quotient by $\Phi(z)$ is sufficient to account for all indeterminacies in the choice of A , so that $\tau_1(L_z, L)$ is a well-defined link concordance invariant for all z . However, as explained below in subsection 5.2, slightly restricting z will greatly expand the number of cases for which $\tau_1(L_z, L)$ is well-defined. For now, we observe that there is an action of the m -fold cartesian product $\Psi := (\pi_1 M)^m$ on $\mathcal{T}_0(\pi_1 M)$ which corresponds to choices of basings of the m components of L in computing $\tau_0(L)$. The stabilizer $\Psi_z < \Psi$ of z is called *untwisted* if all elements of Ψ_z induce the trivial permutation of the elements of $\pi_1 M$ in a representative of z , and *twisted* otherwise (see 4.5 below). This notion of twisting will be seen to correspond to Möbius bands and singular Seifert fibers, and in general a twisted stabilizer Ψ_z can only occur for certain special forms of z .

As explained in Section 4, by choosing a clasp link L_z in each order 1 Whitney concordance class we get the following classification theorem for many links of null-homotopic knots:

Theorem 3. *For any orientable 3-manifold M with $\pi_1 M$ torsion-free, and any $z \in \mathcal{T}_0(\pi_1 M)$ with Ψ_z untwisted, there exists a link L_z in M such that for any links L and L' with $\tau_0(L) = \tau_0(L') = z \in \mathcal{T}_0(\pi_1 M)$ the following are equivalent:*

- (i) $\tau_1(L_z, L) = \tau_1(L_z, L')$.
- (ii) L and L' are stably concordant.
- (iii) L and L' are order 2 Whitney concordant.
- (iv) L and L' are height 1 Whitney concordant.

Here the equality in statement (i) is understood to be up to an action of Ψ_z by left multiplication. By Proposition 5.2.1, Ψ_z will be untwisted for all z if M contains no circle bundles over non-orientable surfaces with orientable total space, and no Seifert fibered submanifolds with singular fibers. The notions of order n (resp. height n) Whitney

concordance refer to the equivalence relations on links of cobounding immersed annuli supporting an order n (resp. height n) Whitney tower in $M \times I$ (c.f. Theorem 1).

It follows that $\tau_1(L_z, \cdot)$ is a concordance invariant, and the effect of changing the choice of L_z is computed in Section 5. Again, all elements in the target can be realized by tying Borromean rings into L_z , but the $\text{INT}(z)$ and $\Phi(z)$ relations show that many such Borromean surgeries do not change the stable concordance class of L_z .

The presence of twisted stabilizers can lead to additional indeterminacies coming from self-homotopies of L_z as illustrated below in subsection 5.3. A general method for dealing with such additional indeterminacies would be interesting, as would be an extension of τ_1 and Theorem 3 to the case where $\pi_1 M$ has torsion.

1.4. Stable concordance of essential knots. In the case where $M \cong F \times S^1$ is the product of an orientable surface $F \neq S^2$ with the circle, the stable concordance classification of knots is completed by Theorem 5 at the end of Section 6, which extends the results of Theorem 3 to knots in arbitrary non-trivial free homotopy classes. The corresponding extension of τ_1 to essential annuli requires two changes: To account for choices of paths, the decorations on Y-trees are taken in coset spaces of $\pi_1 M$ by a cyclic group; and to account for choices of Whitney disk boundaries, the INT relations must be again modified. The resulting INT relations turn out to correspond to solutions of certain Baumslag-Solitar equations – which are highly restricted in 3-manifold groups – and can be computed in terms of lower order intersections among annuli, as explained in Section 6.

2. INTERSECTION TREES AND STABLE EMBEDDINGS

This section recalls the definitions of the intersection invariants τ_0 and τ_1 for simply connected surfaces immersed in 4-manifolds, and sketches the proof that τ_1 is a well-defined homotopy invariant – emphasizing aspects that will be relevant to the generalization of τ_1 to immersed annuli. Proofs of Corollary 1 and Theorem 2 from the introduction are given in subsections 2.5 and 2.6. Additional background material on immersed surfaces in 4-manifolds can be found in Chapter 1 of [8] (and absorbed from the examples below in Section 5).

2.1. The order zero intersection tree τ_0 . For $A = A_1, A_2, \dots, A_m$ a collection of properly immersed simply connected oriented surfaces in an oriented 4-manifold X , Wall's hermitian intersection form λ, μ is defined as follows. The surfaces are perturbed into general position and

equipped with *whiskers* (basepoints joined by a path to the basepoint of X). To each transverse intersection point $p \in A_i \cap A_j$ is associated a fundamental group element $g_p \in \pi_1 X$ determined by a loop through A_i and A_j which changes sheets at p . Summing over all such intersection points, with the usual notion of the sign $\epsilon_p = \pm 1$, defines the intersection and self-intersection “numbers”:

$$\lambda(A_i, A_j) = \sum_{p \in A_i \cap A_j} \epsilon_p \cdot g_p \in \mathbb{Z}[\pi_1 X]$$

and

$$\mu(A_i) = \sum_{p \in A_i \cap A_i} \epsilon_p \cdot g_p \in \frac{\mathbb{Z}[\pi_1 X]}{\mathbb{Z}[1] \oplus \{z - \bar{z}\}}.$$

Modding out by the action of the involution $z \mapsto \bar{z}$ on $\mathbb{Z}[\pi_1 X]$ induced by the map $g \mapsto g^{-1}$ on $\pi_1 X$ accounts for choices of orientations on the sheet-changing loops through the self-intersections of A_i . The relations killing $\mathbb{Z}[1]$ correspond to indeterminacies in the $\mu(A_i)$ due to local cusp homotopies which create self-intersections in A_i whose double-point loops determine the trivial element $1 \in \pi_1 X$.

Basic singularity theory shows that $\lambda(A_i, A_j)$ and $\mu(A_i)$ are invariant under homotopy of A (rel boundary). Since the indeterminacy relations can be realized (by introducing local cusp homotopies and orienting double-point loops appropriately), the vanishing of λ and μ implies that, after perhaps a homotopy (rel boundary), all intersection points of A occur in *cancelling pairs* having equal group elements and opposite signs. As described in detail below, each such cancelling pair admits a Whitney disk.

If Whitney disks can be found that are all disjoint, with embedded interiors disjoint from A , and satisfying a normal framing condition described below, then such Whitney disks can be used to guide a homotopy of A to an embedding. It can always be arranged that the Whitney disks are disjoint and framed, but in general they will have interior intersections with A . The obstruction theory of Whitney towers attempts to pair up these higher order intersections with higher order Whitney disks, and the resulting invariants associated to such a tower of Whitney disks naturally take values in groups generated by univalent trees. These trees are associated to unpaired intersection points, and sit as subsets bifurcating down through the Whitney tower, with the trivalent vertices corresponding to Whitney disks, and the edges corresponding to sheet-changing paths.

In this language, the proper immersion A is called a *Whitney tower of order zero*, and Wall’s intersection form is an invariant of order zero

(since there are no Whitney disks involved). In the notation of Whitney towers, the information contained in λ and μ is expressed as the order zero *intersection tree*

$$\tau_0(A) := \sum \epsilon_p \cdot t_p \in \mathcal{T}_0(\pi_1 X)$$

where each t_p is an oriented edge (order zero trees have no trivalent vertices) which is decorated by g_p and has endpoint vertices labelled by i and j for $p \in A_i \cap A_j$. We will use the notation $(g)_{ij}$ to denote an order zero tree which is decorated by g and oriented from i to j . The target $\mathcal{T}_0(\pi_1 X)$ is the abelian group additively generated by such decorated edges modulo two kinds of relations: The orientation relations $(g)_{ij} = (g^{-1})_{ji}$ reflect the Hermitian nature of the pairing λ for $i \neq j$, and correspond for $i = j$ to the relations $z = \bar{z}$ in the self-intersection invariants $\mu(A_i)$ of the components of A . The framing relations $(1)_{ii} = 0$ correspond to the relations $\mathbb{Z}[1] = 0$ in the $\mu(A_i)$ by killing any edges decorated by trivial group elements and having both vertices labeled by i . (With these framing relations one has a homotopy invariant; without the framing relations one has a *regular* homotopy invariant.)

Thus, $\tau_0(A)$ is exactly Wall's intersection form rewritten in Whitney tower notation, and the vanishing of $\tau_0(A)$ is equivalent to the existence of Whitney disks pairing all intersection points in A (perhaps after performing some cusp homotopies). In the language of Whitney towers we say that such A admits an *order 1 Whitney tower*. The Whitney disks in an order one Whitney tower are required to have pairwise disjointly embedded boundaries. The interiors of the Whitney disks are allowed to be immersed, and may intersect each other as well as A . All Whitney disks are required to be properly framed as described next.

2.2. Whitney disk framings. Consider a cancelling pair of intersections p and q between A_i and A_j , with $g_p = g_q \in \pi_1 X$ and $\epsilon_p = -\epsilon_q$. The union of any pair of arcs, one in A_i and the other in A_j , each connecting p and q (but avoiding all other singularities), forms a loop contractible in X , and any (immersed) disk W bounded by such a loop is a *Whitney disk*. Since p and q have opposite signs, the restriction $\nu_{\partial W}$ of the normal disk bundle ν_W of W in X to ∂W is the trivial bundle $S^1 \times D^2$ (if the signs of p and q were equal, then we would get the non-orientable disk bundle over the circle).

The trivial D^2 -bundle $\nu_{\partial W}$ has a nowhere-vanishing *Whitney section* constructed as follows (see Figure 4): Denote by $\partial_i W$ (resp. $\partial_j W$) the arc of ∂W that lies in A_i (resp. A_j). Over $\partial_i W$ choose a splitting of




FIGURE 4. The Whitney section over the boundary of a (zero-)framed Whitney disk is indicated by the dotted loop shown on the left for a clean Whitney disk W in a 3-dimensional slice of 4-space, and on the right in the (pulled back) normal disk bundle over ∂W .

$\nu_{\partial W}$ given by a vector field v_i^t tangent to A_i and a vector field v_i^n normal to both W and A_i . Over $\partial_j W$ we have a similar splitting $\{v_j^t, v_j^n\}$ of $\nu_{\partial W}$, and we may arrange that these splittings coincide at p and q with $v_i^t = v_j^n$ and $v_i^n = v_j^t$. The Whitney section of $\nu_{\partial W}$ is constructed by taking v_i^t over $\partial_i W$, and v_j^n over $\partial_j W$. If this section of $\nu_{\partial W}$ can be extended to a nowhere-vanishing section of ν_W , then we say that W is *framed*.

For a chosen orientation, the relative Euler number of the Whitney section gives an integer obstruction to extending the section over W , so “framed” really means “zero-framed”. This obstruction can always be killed at the cost of creating interior intersections between W and A_i (or A_j) by a *boundary twisting* modification of W which changes the obstruction by ± 1 . Also, the framing obstruction can be changed by ± 2 at the cost of creating an interior self-intersection of W by performing a cusp homotopy, called an *interior twist*. Boundary and interior twists are illustrated below in Figure 22 and Figure 19 of Section 5.

2.3. The order 1 intersection tree τ_1 . The invariant denoted τ in [29] corresponds in the general theory of Whitney towers ([5, 6, 30]) to the order 1 intersection tree τ_1 . For a collection A of properly immersed simply connected surfaces with vanishing $\tau_0(A)$, $\tau_1(A)$ is defined as follows. Choose an order 1 Whitney tower \mathcal{W} on A , with fixed whiskers on the A_i and on each of the Whitney disks in \mathcal{W} . To each intersection point $p \in W_{(i,j)} \cap A_k$ between A_k and the interior of a Whitney disk $W_{(i,j)}$ pairing intersections between A_i and A_j is associated a trivalent Y-tree t_p as illustrated in Figure 1. The univalent vertices are labelled

$$\begin{array}{c}
 \text{AS} \\
 \begin{array}{c} k \\ \downarrow c \\ \begin{array}{cc} b \swarrow & \searrow i \end{array} \end{array} + \begin{array}{c} k \\ \downarrow c \\ \begin{array}{cc} i \swarrow & \searrow j \end{array} \end{array} = 0 = \begin{array}{c} i \\ \downarrow a \\ \begin{array}{cc} b \swarrow & \searrow i \end{array} \end{array} + \begin{array}{c} j \\ \downarrow b \\ \begin{array}{cc} j \swarrow & \searrow i \end{array} \end{array} \\
 \text{FR} \\
 \text{OR} \\
 \begin{array}{c} k \\ \downarrow c \\ \begin{array}{cc} b \swarrow & \searrow i \end{array} \end{array} = \begin{array}{c} k \\ \downarrow \bar{c} \\ \begin{array}{cc} b \swarrow & \searrow i \end{array} \end{array} \\
 \text{HOL} \\
 \begin{array}{c} k \\ \downarrow c \\ \begin{array}{cc} b \swarrow & \searrow i \end{array} \end{array} = \begin{array}{c} k \\ \downarrow cg \\ \begin{array}{cc} bg \swarrow & \searrow i \end{array} \end{array}
 \end{array}$$

FIGURE 5. The AS *antisymmetry*, FR *framing*, OR *orientation*, and HOL *holonomy* relations in $\mathcal{T}_1(\pi_1 X)$, assuming a fixed planar orientation on the trivalent vertex of all Y-trees. Here a , b , c , and g are elements of $\pi_1 X$, with $\bar{c} = c^{-1}$. These edge labels can also be extended linearly to elements of $\mathbb{Z}[\pi_1 X]$.

by i , j and k , and the oriented edges are decorated by elements a , b , and c in $\pi_1 X$ determined by sheet-changing loops through $W_{(i,j)}$ and the components of A using the chosen whiskers, with the loop orientations corresponding to the edge orientations. Fixing (arbitrary) orientations on the Whitney disks in \mathcal{W} associates a sign $\epsilon_p = \pm 1$ to each interior intersection point p , and determines a cyclic orientation at the trivalent vertex of the corresponding tree t_p via the following convention: As illustrated in Figure 1, the tree t_p sits as an embedded subset of $W_{(i,j)}$. The two edges of t_p labelled i and j determine a “corner” of $W_{(i,j)}$ which does not contain the k -labelled edge, and the orientation of t_p is taken to be that induced by $W_{(i,j)}$ if and only if this corner contains the *positive* intersection point between A_i and A_j that is paired by $W_{(i,j)}$.

(Due to differing orientation conventions there is a global sign difference between τ of [29] and the τ_n of more recent papers ([5, 30]), including τ_1 here.)

The *order 1 intersection tree* $\tau_1(\mathcal{W})$ is defined by summing the t_p over all unpaired intersection points in \mathcal{W} :

$$\tau_1(\mathcal{W}) := \sum \epsilon_p \cdot t_p \in \frac{\mathcal{T}_1(\pi_1 X)}{INT(A)}$$

The abelian group $\mathcal{T}_1(\pi_1 X)$ is additively generated by the above described decorated Y-trees, modulo the relations shown in Figure 5. The $INT(A)$ relations are shown in Figure 6.




FIGURE 6. The $\text{INT}(A)$ *intersection* relations. Here S_{ij} varies over 2-spheres representing a generating set for $\pi_2 X$, and the order zero invariant $\lambda_0(S_{ij}, A)$ counts only intersections between S_{ij} and A , with the vertex label k assigned to the intersections between S_{ij} and A_k (and edge labels from $\pi_1 X$).

2.3.1. *Notation and normal forms for \mathcal{T}_1 .* Note that the AS relation implies that the generators appearing in the FR relation are 2-torsion. By the OR edge-orientation relation, we can normalize all edges on generators of $\mathcal{T}_1(\pi_1 X)$ to be oriented towards the trivalent vertex. We will use the cyclically oriented triple $(a, b, c)_{ijk}$, to denote such a Y-tree having univalent labels i, j, k , and corresponding group elements a, b , and c .

For fixed triples of univalent vertex labels, it will be convenient to represent the corresponding subgroups of $\mathcal{T}_1(\pi_1 X)$ as quotients of the integral group ring $\mathbb{Z}[\pi_1 X \times \pi_1 X] \cong \mathbb{Z}[\pi_1 X] \times \mathbb{Z}[\pi_1 X]$ by using the HOL relation to normalize one edge decoration to the trivial group element $1 \in \pi_1 X$ – for instance $(a, b, c)_{ijk} = (1, ba^{-1}, ca^{-1})_{ijk}$ – and summing group element decorations inside the parentheses. This corresponds to the notation used in [29] for $\tau = \tau_1$.

Assume that \prec is a *right-order* on $\pi_1 X$ – a strict total ordering of the elements of $\pi_1 X$ such that if $g \prec h$ then $gf \prec hf$ – and that the components of A (the univalent vertex labels) are ordered. Then we have the following normal form for $\mathcal{T}_1(\pi_1 X)$:

First consider a Y-tree whose univalent vertices all have the same label, and whose edge decorations are distinct group elements. Use the HOL relation to trivialize one edge decoration, and denote the resulting Y-tree by the ordered pair $(g, h) := (1, g, h)$, where we suppress the common univalent vertex label.

Such a generator (g, h) is not involved in any FR relation (since $1 \neq g \neq h \neq 1$), and the orbit of (g, h) under the AS and HOL relations in this notation is:

$$(g, h) = -(g^{-1}, hg^{-1}) = (hg^{-1}, g^{-1}) = -(gh^{-1}, h^{-1}) = (h^{-1}, gh^{-1}) = -(h, g).$$

Now one can check that for each of the possible relative orderings of 1 , g , and h given by \prec , there is exactly one generator (a, b) in the orbit of (g, h) such that $1 \prec a \prec b$; for instance if $h \prec 1 \prec g$, then (h^{-1}, gh^{-1}) satisfies $1 \prec h^{-1} \prec gh^{-1}$.

In the case where $g = h$, but still assuming all univalent labels are the same, then the generator is 2-torsion and we have the following orbit under the AS, HOL, and FR relations:

$$(g, g) = (g^{-1}, g^{-1}) = (g^{-1}, 1) = (1, g^{-1}) = (g, 1) = (1, g).$$

From each such orbit we get a unique generator $(1, a)$ such that $1 \preceq a$ (or a unique generator (a, a) with $1 \preceq a$).

This same approach works for other combinations of univalent labels, of course with different possible choices of convention. We state here the rest of a set of generators in normal form, written in ordered triple notation, with non-decreasing univalent labels (from left to right) which correspond to the cyclic orientation of the Y-tree:

$$\begin{aligned} (1, g, h)_{ijk} & \quad \text{for distinct labels } i < j < k, \\ (1, g, h)_{iij} & \quad \text{with } 1 \prec g \quad \text{for labels } i < j, \\ (1, g, h)_{ijj} & \quad \text{with } g \prec h \quad \text{for labels } i < j, \end{aligned}$$

and

$$(1, 1, g)_{iij} \quad \text{for labels } i < j.$$

These last generators are 2-torsion.

This construction of a normal form, together with the observation that a (non-trivial) right-orderable group is infinite (it must be torsion free), gives the following proposition:

Proposition 2.3.1. *If $\pi_1 X$ is non-trivial and right-orderable, then $\mathcal{T}_1(\pi_1 X)$ is isomorphic to $\mathbb{Z}^\infty \oplus \mathbb{Z}_2^\infty$.*

2.4. Homotopy invariance of τ_1 . We summarize here the proof in [29] that $\tau_1(A) := \tau_1(\mathcal{W})$ is a well-defined (regular) homotopy invariant of A , indicating the relevant points that will have to be addressed when we allow A to have annular components. The proof proceeds by first checking that the value of $\tau_1(\mathcal{W})$ in $\mathcal{T}_1(\pi_1 X)/\text{INT}(A)$ does not depend on the choices in constructing the Whitney tower \mathcal{W} . Showing homotopy invariance is then reduced to checking invariance under finger moves.

With an eye towards our interest in the case $X = M \times I$, we will make the simplifying assumptions that $\pi_1 X$ is torsion-free, and that X is spin, i.e., the second Stiefel-Whitney class $\omega_2 X$ is trivial.

Independence of tree decorations and signs: First of all, the group elements decorating the tree edges do not depend on the choices




FIGURE 7. The 2-sphere S formed as the union of two Whitney disks can be pushed off of A near its equator – where the collars of the Whitney disks are joined along their boundaries.

of sheet-changing loops since the A_i and the Whitney disks are simply connected. The OR relations account for the orientation choices on the loops, which we will always assume to be oriented *into* the Whitney disks, so that all tree edges are oriented towards the trivalent vertex. The decorations do not depend on the choices of whiskers on the Whitney disks by the HOL relations.

The sign ϵ_p associated to each cyclically oriented t_p is independent of the chosen Whitney disk orientations (for fixed orientations on the A_i) by the AS antisymmetry relations and the “positive corner” convention (subsection 2.3).

Independence of Whitney disk interiors: For fixed boundaries on the Whitney disks, the INT and FR relations guarantee that changing the Whitney disk interiors does not change $\tau_1(A)$; the idea is that such a change is given by intersections with 2-spheres: Consider any Whitney disk $W = W_{(i,j)}$ in \mathcal{W} pairing p and q in $A_i \cap A_j$. If W' is another Whitney disk with same boundary as W , then the union of W and W' is a topological 2-sphere S , which may not be smooth along $\partial W = \partial W'$. We may arrange (by a small isotopy and after perhaps performing some boundary twists on W) that along their common boundary the collars C and C' of W and W' point in opposite directions (Figure 7) so that $W \cup W'$ is an immersed 2-sphere, still denoted S , which is oriented by the orientation of W together with the opposite orientation of W' . The boundary twists on W (n twists into A_i and m twists into A_j) have changed $\tau_1(\mathcal{W})$ by $n(a, b, a)_{iji} + m(a, b, b)_{ijj} \in \mathcal{T}_1(\pi_1 X)$ – which is 2-torsion by the AS relations – and have changed the framing of W by $n + m \in \mathbb{Z}$. But by extending the Whitney section over the equator to a normal push-off of all of S we see that $n + m$ is equal modulo 2 to $\omega_2(S) \in \mathbb{Z}_2$ which vanishes since X is spin (note that any self-intersections of S do not contribute to $\omega_2(S)$). So by the AS and FR relations $n(a, b, a)_{iji} + m(a, b, b)_{ijj} = 0 \in \mathcal{T}_1(\pi_1 X)$.




FIGURE 8. The signed tree $\epsilon_p \cdot t_p$ assigned to an intersection point $p \in \partial_{\epsilon_j} W_j \cap \partial_{\epsilon_k} W_k$ between Whitney disk boundaries in the case $\epsilon_k = + = \epsilon_j$, with the orientations of A_i and t_p both the same as the plane of the paper.

Since the circle of intersection between S and A can be perturbed away without creating any new intersections (Figure 7), it follows that the change in $\tau_1(\mathcal{W})$ resulting from replacing W by W' is exactly described by an $\text{INT}(A)$ relation with $S = S_{ij}$.

Independence of Whitney disk boundaries: To show that $\tau_1(\mathcal{W})$ does not depend on the choices of boundaries of the Whitney disks, for fixed pairings of the cancelling singularities in A , it is convenient to weaken the definition of an order 1 Whitney tower by allowing transverse intersections among the boundaries of the Whitney disks (following [29] and 10.8 of [8]). The definition of τ_1 is extended to such Whitney towers by assigning trees to the boundary intersections between Whitney disks in the following way (see Figure 8).

Consider two Whitney disks, W_j and W_k , pairing intersections that A_i has with A_j and A_k , respectively. Note that even if W_j and W_k have no interior intersections with any A_l , they still have naturally associated Y-trees, with only the “descending” edges and vertices decorated and labelled, as in the left hand side of Figure 8. Choose whiskers so that the edges labelled by i are decorated by the trivial element $1 \in \pi_1 X$, which then determines elements g_j and g_k , respectively, decorating the edges labeled j and k . The cyclic orientation of a Y-tree sitting in a Whitney disk W corresponds to the orientation of W by the “positive corner” convention described above (2.3), and induces an orientation of ∂W (which corresponds to the convention that $\overrightarrow{\partial W}$ together with a second inward pointing vector give the orientation of W). We will use the notation ∂_+ to indicate a Whitney disk boundary arc that is oriented towards its positive intersection point, and ∂_- for a Whitney disk boundary arc that is oriented towards its negative intersection point.

Now let $p \in \partial_{\epsilon_j} W_j \cap \partial_{\epsilon_k} W_k$ be a point such that the ordered pair of tangent vectors $(\overrightarrow{\partial_{\epsilon_j} W_j}, \overrightarrow{\partial_{\epsilon_k} W_k})_p$ is equal to the orientation of A_i at p . Define the tree t_p associated to such a p by:

$$\epsilon_p \cdot t_p := \epsilon_k \epsilon_j (1, g_k^{\epsilon_k}, g_j^{\epsilon_j})_{ikj}$$

where $\epsilon \in \{+, -\} \hat{=} \{+1, -1\}$.

One can check that this definition of t_p does not depend on the choices made. The extended version of $\tau_1(A)$ is defined by including such t_p in the sum. Since all boundary intersections can be eliminated by finger moves which create interior intersections having the exact same trees (Figure 3 in [29]), this extended definition can always be reduced to the original one.

Note that, properly interpreted, the formula assigning t_p to $p \in \partial W_j \cap \partial W_k$ also works when $W_j = W_k$, including the case $i = j = k$. For instance, for W pairing self-intersections of A_i , and $p \in \partial_- W \cap \partial_+ W$ such that the orientation of A_i is equal to $(\overrightarrow{\partial_- W}, \overrightarrow{\partial_+ W})_p$, then $\epsilon_p \cdot t_p = -(1, g, g^{-1})_{iii}$, where the Y-tree associated to W has decorations 1 and g on the edges dual to the $+$ and $-$ boundary arcs respectively. (This formula will be used in Section 5 below.)

The proof of independence of Whitney disk boundaries now goes as follows. Since the components of A are all simply connected, any configuration of Whitney disk boundaries can be achieved by a regular homotopy of (collars of) the Whitney disk boundaries, fixing the intersection points of A (Clarification: we mean here that this regular homotopy is induced by a regular homotopy of the inverse images of the Whitney disk boundaries in the domain of A , and extends to a regular homotopy of collars of the Whitney disks in X). During such a homotopy, τ_1 does not change since boundary intersections come and go in cancelling pairs, or accompanied by a cancelling interior intersection (when pushing over an intersection point of A , see Figure 5 in [29]).

Note that this step uses the fact that (the domains of) the components of A are simply connected, and will have to be modified when we allow A to have immersed annular components.

Independence of intersection pairings: The independence of $\tau_1(W)$ on the choices of pairings of the intersections of A follows easily from a construction pictured in Figure 6 of [29] (originally from 10.8 of [8]). In the presence of 2-torsion in $\pi_1 X$ there is also a subtle indeterminacy corresponding to the pairing of the inverse images of the

self-intersections of the A_i which was first explained in [31], and is covered by a more general INT relation ([29]) (which also covers the case where X is not spin).

Homotopy invariance: Having established the independence of $\tau_1(A) := \tau_1(\mathcal{W})$ on the choice of Whitney tower \mathcal{W} , homotopy invariance can be seen as follows. Up to isotopy, a generic homotopy (rel ∂) between surfaces in a 4-manifold is a sequence of finger moves (which create a pair of intersections), Whitney moves (which eliminate a pair of intersections), and local cusp homotopies (births and deaths of local self-intersections). In a *regular* homotopy it may be arranged that there are *only* finger moves and Whitney moves, and that the finger moves all occur before the Whitney moves. Since finger moves and Whitney moves are inverse to each other, it follows that if A is regularly homotopic to A' , then there is A'' which differs (up to isotopy) from each of A and A' by only finger moves which create cancelling intersections paired by local clean Whitney disks. Since these finger moves and their local clean Whitney disks can be assumed to be disjoint from all other Whitney disks in any Whitney tower on A or A' , it follows that $\tau_1(A) = \tau_1(A'') = \tau_1(A')$.

Geometric characterization: That the vanishing of $\tau_1(A)$ leads to an order 2 and height 1 Whitney tower follows from the fact that all the relations in $\mathcal{T}_1(\pi_1 X)$ can be realized by controlled manipulations of Whitney towers ([29]). For instance, the FR and INT relations can be realized by creating clean Whitney disks, then twisting and tubing into 2-spheres. The new relations introduced later for the generalization of $\tau_1(A)$ which allows A to have annular components will be similarly realizable.

2.5. Proof of Corollary 1.

Proof. The equivalence of statements (i) and (iii) in Theorem 1 is proved as Theorem 2 of [29] in the case where A consists of a single connected component, but the exact same proof goes through for multiple components. Thus, the vanishing of $\tau_1(A)$ means that all the singularities of A can be paired by (framed) Whitney disks whose interiors are disjoint from A . If the Whitney disk interiors are also disjointly embedded, then A is homotopic to an embedding without any stabilization. Otherwise, it may be arranged by splitting the Whitney disks using finger moves that each Whitney disk is embedded, and has at most a single transverse interior intersection with some other Whitney disk. Each such intersection point $p \in W \cap W'$ can be eliminated after taking a connected sum of X with a copy of $S^2 \times S^2$ by the Norman trick ([24]), which is illustrated schematically in Figure 9: If S and S'




FIGURE 9. The Norman trick.

are dual 2-sphere factors generating $\pi_2(S^2 \times S^2)$, then W can be tubed into S , and p can be eliminated by tubing W' into S' along W . This operation can be done without creating any new singularities, and since S and S' are both 0-framed, the resulting Whitney disks are still framed. After one stabilization for each interior intersection between Whitney disks, the resulting disjointly embedded framed Whitney disks guide a homotopy of A to an embedding in the connected sum of X with (finitely many) copies of $S^2 \times S^2$.

On the other hand, if $A \looparrowright X$ is homotopic to an embedding in the connected sum of X with copies of $S^2 \times S^2$, then since $\tau_1(A)$ only depends on the homotopy class (rel boundary) of A , and since taking connected sums of X with $S^2 \times S^2$ does not change the $\text{INT}(A)$ relations, it follows that $\tau_1(A)$ vanishes. \square

2.6. Proof of Theorem 2. This subsection gives a proof of the concordance invariance of the order 1 intersection tree $\tau_1(L)$ of a link (Theorem 2), which at the same time shows the concordance invariance of the order zero intersection tree $\tau_0(L)$.

Proof. Consider an oriented m -component link L of null-homotopic knots in an oriented 3-manifold M , with D any collection of properly immersed disks in $M \times I$ bounded by $L \subset M \times \{0\}$. For n equal to 0 or 1, if D admits an order n Whitney tower, then we want to show that $\tau_n(L) := \tau_n(D)$ only depends on the concordance class of L .

The key lemma is:

Lemma 2.6.1 ([26]). *Any m elements of $\pi_2(M \times I)$ are represented by m embedded pairwise disjoint 2-spheres.*

Proof. A well known consequence of the Sphere Theorem of 3-manifold theory is that $\pi_2 M$ is generated as a module over $\pi_1 M$ by disjoint embeddings (the 2-spheres that decompose M into prime factors, together with any spherical boundary components and cross-sections of any $S^1 \times S^2$ factors, see Proposition 3.12 of [15]). Tubing these generators together in $M \times I$ does not create any new intersections, so all

elements of $\pi_2(M \times I)$ are represented by embeddings, which can be isotoped to be pairwise disjoint using the product structure. \square

We also need the following general properties of $\tau_n(A)$ for $A \looparrowright X$, which can be checked directly from the definitions: If $-A$ denotes a flip of orientation on (all components of) A , then $\tau_0(-A) = \tau_0(A)$ and $\tau_1(-A) = -\tau_1(A)$, where the orientation of the ambient 4-manifold X is fixed. On the other hand, flipping the orientation of X while fixing the orientation of A has the effect of multiplying $\tau_0(A)$ by -1 , but preserves $\tau_1(A)$.

To see that $\tau_n(L)$ does not depend on the choice of D , let D' be another singular null-concordance of L . Then the union S of D and D' along L , in two copies of $M \times I$ identified along M , determines m elements of $\pi_2(M \times I) \cong \pi_2 M$. The definition of $\tau_n(L) := \tau_n(D)$ depends on fixing a convention for the how the orientation of $L \subset M \times \{0\}$ induces an orientation of $D \looparrowright M \times I$. It follows that, after orienting S in $M \times I$ (which requires reorienting one of the original copies of $M \times I$ and the corresponding singular null-concordance), we have

$$0 = \tau_n(S) = \pm(\tau_n(D) - \tau_n(D'))$$

where the first equality comes from Lemma 2.6.1 and the homotopy invariance of τ_n .

It also follows from Lemma 2.6.1 that the $\text{INT}(D)$ relations are trivial in the case $n = 1$.

The sense in which $\tau_n(L)$ is a concordance invariant requires interpreting its value modulo the effect on $\mathcal{T}_n(\pi_1 M)$ of whisker choices. This will be discussed in detail in Section 4. For now we observe that if A is a concordance from L' to L then, up to the change-of-whisker action, $\tau_n(L') = \tau_n(A \cup_L D) = \tau_n(D) = \tau_n(L)$ since A has no singularities. This same argument shows that $\tau_n(L)$ is invariant under order $n + 1$ Whitney concordance since then all singularities of A of order less than $n + 1$ occur in cancelling pairs. \square

3. τ_1 FOR INESSENTIAL ANNULI

In this section we extend the definition of $\tau_1(A)$ to allow A to have components which are *inessential* properly immersed annuli, meaning that the induced map $\mathbb{Z} \rightarrow \pi_1 X$ on fundamental groups is trivial.

This generalized τ_1 is illustrated by a pairing between links and 2-spheres in M described in 3.1 below, and computed in 3.2. These results are used in 3.3 to define the indeterminacy subgroup $\Phi(z)$ in the definition of the relative order 1 intersection tree $\tau_1(L_z, L)$ of 1.3.

(The reader who is only interested in links in irreducible 3-manifolds can skip these three subsections.)

We continue to assume that X is spin, and $\pi_1 X$ is torsion-free.

Let $A \looparrowright X$ be a collection of properly immersed surfaces admitting an order 1 Whitney tower. We now allow the components of A to include inessential annuli as well as 2-disks and 2-spheres. In order to extend the definition (2.3) of $\tau_1(A)$ to such A it will only be necessary to add some additional $\text{INT}(A)$ relations.

Starting with the original definition 2.3, we proceed by examining the steps in the proof in 2.4 that $\tau_1(A)$ is a well-defined homotopy invariant.

First of all, note that $\tau_0(A)$ is still a well-defined homotopy invariant for such A (with the vanishing of $\tau_0(A)$ equivalent to the existence of an order 1 Whitney tower on A) since in inessential annuli the elements of $\pi_1 X$ associated to sheet-changing paths are still well-defined (because loops in A which do not change sheets must represent $1 \in \pi_1 X$). For the same reason the edge decorations from $\pi_1 X$ on the trees in $\tau_1(A)$ are still well-defined.

The key step in 2.4 which relies on the assumption that the components of A are simply connected is the argument showing independence of the choices of Whitney disk boundaries, which uses the fact that any two configurations of arcs with fixed endpoints in a simply connected surface are related by a homotopy (rel endpoints). This is no longer true when A has annular components, and to account for the indeterminacies which correspond to changing the homotopy classes (rel endpoints) of the Whitney disk boundaries on annular components of A we will include new $\text{INT}(A)$ relations.

Consider any Whitney disk W_{ij} pairing intersections between an annular component A_i of A and any component A_j (including possibly $j = i$). Since A_i is inessential, push-offs of either component of ∂A_i bound immersed disks in X . Let D_{ij} be an immersed disk, with ∂D_{ij} a parallel push-off of a component of ∂A_i sitting in a collar on ∂A_i . The normal disk-bundle of D_{ij} in X has a nowhere vanishing section given by pushing tangentially along A_i , and we require that this section extends to a nowhere vanishing section over D_{ij} ; this can always be arranged by boundary-twisting D_{ij} if necessary. As illustrated in Figure 10, by half-tubing a collar of W_{ij} into a collar of D_{ij} along any embedded arc in A_i we can change the homotopy class of ∂W_{ij} (rel endpoints) in A_i by \pm a generator of $\pi_1 A_i$ (meaning that ∂W_{ij} is band-summed into a loop which pulls back to a generator of the fundamental group of the domain of A_i).




FIGURE 10. Changing the homotopy class in A_i of the boundary of a Whitney disk leads to new $\text{INT}(A)$ relations determined by the order zero intersection invariant $\lambda_0(D_{ij}, A)$, with the corresponding univalent labels taken from the components of A .

The framing requirement on D_{ij} means that this operation preserves the framing on W_{ij} , and the resulting change in the contribution of W_{ij} to $\tau_1(A)$ can be expressed as the relation illustrated in Figure 10, where a and b are the descending group elements for W_{ij} , and the order zero invariant $\lambda_0(D_{ij}, A)$ counts only intersections between D_{ij} and all the A_k (not intersections among the A_k), as was the case for the 2-spheres S_{ij} in the original $\text{INT}(A)$ relations of Figure 6.

By the proof of the independence of τ_1 on the Whitney disk interiors (2.4), using the usual $\text{INT}(A)$ relations, this expression does not depend on the choice of D_{ij} (or on the choice of the component of ∂A_i). Since the complement of any number of embedded arcs in an annulus is path-connected, this operation can be iterated any number of times on any number of Whitney disks, to get any choices of homotopy classes (rel endpoints) for the Whitney disk boundaries, so all the resulting indeterminacies are linear combinations of the expressions in Figure 10. By including these expressions into the $\text{INT}(A)$ relations, and otherwise defining $\tau_1(A)$ as before, we get a well-defined homotopy invariant, with A allowed to have inessential properly immersed annular components.

We remark that the original $\text{INT}(A)$ relations are determined by computing order zero intersection invariants for a generating set for $\pi_2 X$ (as a module over $\pi_1 X$). These extended $\text{INT}(A)$ relations are determined by computing finitely many more order zero intersection invariants, which in our applications to links will essentially correspond to the order zero intersection tree of the link.

These new $\text{INT}(A)$ relations can be realized by performing finger-moves on A to create clean Whitney disks with prescribed group elements,

and then performing the above operation. Since the usual geometric manipulations of Whitney towers – such as “transferring” intersections between Whitney disks (Figure 10 in [29]) – also work for annular components, it follows that the geometric statements in Theorem 1 hold for this generalized version of $\tau_1(A)$.

Summarizing, we have:

Theorem 4. *Theorem 1 and Corollary 1 also hold if any components of A are inessential properly immersed annuli.*

3.1. The order 1 intersection pairing of links and spheres. In this subsection we use Theorem 4 to define an order 1 intersection pairing between links and 2-spheres which determines the indeterminacy subgroup $\Phi(z)$ of Theorem 3.

Let L be any link of null-homotopic knots, and S be any immersed 2-sphere in M . Perturb S into $M \times I$, and extend L to the collection of embedded annuli $A = L \times I \subset M \times I$. By Lemma 2.6.1, S is homotopic to an embedding, and since the components of L are null-homotopic it follows that $\tau_0(A, S) = 0$. So we can define the pairing

$$\tau_1(L, S) := \tau_1(A, S) \in \mathcal{T}_1(\pi_1 M) / \text{INT}(z)$$

where we write $\text{INT}(z)$ for $\text{INT}(A, S)$, since (by Lemma 2.6.1 and the fact that the components of L are null-homotopic) the only INT relations come from the boundary L , and are determined by $z = \tau_0(L)$ as follows. Using the ordering of the link components we can write $\tau_0(L)$ in a *normal form*:

$$\tau_0(L) = z = \sum_{i \leq j} z_{ij} = \sum_{i \leq j} \sum_p e_p \cdot (g_p)_{ij},$$

where the coefficients e_p are non-zero integers, with the understanding that the indices p depend on i, j – this is just collecting together all the intersections between D_i and D_j that have the same group element g_p . The group elements are assumed to be distinct, i.e. $(g_p)_{ij} \neq (g_{p'})_{ij}$ for $p \neq p'$; and the only indeterminacy in this normal form for z is in case $i = j$ we choose only one of $(g_p)_{ii}$ or $(g_p^{-1})_{ii}$.

Then each INT relation corresponding to a 2-disk D_{ij} as in Figure 10, with the group element a normalized to 1 by a HOL relation, can be written:

$$0 = (1, g, \sum_{k \neq i} z_{ki})_{ijk} + (1, g, z_{ii} + \overline{z_{ii}})_{iji}$$

where the first term is determined by order zero linking between L_i and the other components L_k for $k \neq i$; and the second term is determined by order zero linking between L_i and a 0-parallel push-off L'_i . Note that




FIGURE 11. On the left, the clasp knot K and (one hemisphere of) the cross-sectional 2-sphere S in $M = S^1 \times S^2$ (with only $S^1 \times D^2$ shown). On the right, a collar of a 0-framed immersed 2-disk D bounded by a parallel copy of K in $K \times I$.

the second term is independent of the above mentioned indeterminacy in the normal form of z (it corresponds to the relation $\lambda(D, D') = \mu(D) + \overline{\mu(D')}$ in Wall's intersection form for D' a zero-parallel push-off of D – see the next example).

3.1.1. Example. This subsection computes the pairing $\tau_1(L, S)$ in a case where L is a knot, illustrating its relation to the study of L up to stable concordance by defining the indeterminacy subgroup $\Phi(z)$ and target group of Theorem 3 in a simple example. It also serves as motivation and a warm-up for the more complicated computations in the subsequent two subsections.

Consider the case $M = S^1 \times S^2$, with $\pi_1 M \cong \mathbb{Z}$ generated multiplicatively by x . Let $x^n S$ denote a cross-sectional 2-sphere S equipped with a whisker corresponding to $x^n \in \pi_1 M$, and let L be the clasp knot $K \subset M$ as in Figure 11 with $\tau_0(K) = x \in \mathcal{T}_0(\langle x \rangle)$. (In this case K is just a positive Whitehead double of a circle factor representing the generator x ; by the “belt trick” there are two such Whitehead doubles up to isotopy.)

The target for $\tau_1(K, x^n S)$ is the quotient of $\mathcal{T}_1(\langle x \rangle)$, with univalent labels from $\{K, S\}$, by the intersection relations $\text{INT}(K \times I, S) = \text{INT}(x)$ which can be computed explicitly from intersecting a (framed) null-homotopy D of a parallel copy of K with $K \times I$ (crossing changes in the right hand side of Figure 11):

$$\text{INT}(x) = (1, x^r, x + x^{-1} - 2)_{KjK} = (1, x^r, x + x^{-1})_{KjK} = 0$$




FIGURE 12. The order 1 Whitney tower on $K \times I$ and $x^n S$. (Whiskers are not shown.)

for $j \in \{K, S\}$ and all $r \in \mathbb{Z}$. (Note that the contributions $(1, x^r, -2) = -2(1, x^r, 1)$ from the two negative twists which preserve the framing of D are killed by the AS relations.)

As illustrated in Figure 12, $K \times I$ and $x^n S$ support an order 1 Whitney tower \mathcal{W} with a single Whitney disk, and

$$\tau_1(\mathcal{W}) = (1, x^{-1}, x^n)_{KKS} = (x, 1, x^{n+1})_{KKS} = -(1, x, x^{n+1})_{KKS}$$

where the middle equality comes from the HOL relations, and the right-most equality comes from the AS relations.

It follows that $\tau_1(K, x^n S)$ lies in the span of Y-trees having two univalent vertices labeled by K and one by S . Using the HOL relation to trivialize one K -labeled edge decoration, we can write any such Y-tree as

$$(x^l, x^m) := (1, x^l, x^m)_{KKS} \quad l, m \in \mathbb{Z}.$$

By the AS and HOL relations we have

$$(x^l, x^m) = -(x^{-l}, x^{m-l}),$$

so the relevant subgroup of $\mathcal{T}_1(\langle x \rangle)$ is isomorphic to $\mathbb{Z}^\infty \oplus \mathbb{Z}_2^\infty$, with the \mathbb{Z} factors generated by (x^l, x^m) , for $l \in \mathbb{N}$ and $m \in \mathbb{Z}$, and the \mathbb{Z}_2 factors generated by $(1, x^m)$, for $m \in \mathbb{Z}$.

Now, in terms of these generators the relevant INT(x) relations $(x + x^{-1}, x^r) = (x, x^r) + (x^{-1}, x^r) = 0$ give:

$$(x, x^r) = (x, x^{r+1})$$

for all $r \in \mathbb{Z}$ (using $(x^{-1}, x^r) = -(x, x^{r+1})$ by HOL and AS).

So for all n we have

$$\tau_1(K, x^n S) = -(x, x^{n+1}) = -(x, x) \neq 0 \in \mathcal{T}_1(\langle x \rangle) / \text{INT}(x).$$

The discussion so far has focused on computing $\tau_1(K, x^n S)$. Recall that the relevance to the knot theory of K from our point of view is that $x^n S$ can be tubed into any singular concordance A of K , and the corresponding indeterminacy in $\tau_1(A)$ lives in $\mathcal{T}_1(\langle x \rangle) / \text{INT}(x)$ with *all univalent labels corresponding to K* . These indeterminacies are described algebraically by just replacing the S -labels with K -labels in $\tau_1(K, x^n S)$, which gives the indeterminacy subgroup $\Phi(x) < \mathcal{T}_1(\langle x \rangle) / \text{INT}(x)$, as x^n varies over $\pi_1 M$ with the generator S of $\pi_2 M$ fixed.

Specifically, we can describe the quotient of $\mathcal{T}_1(\langle x \rangle) / \text{INT}(x)$ by $\Phi(x)$ as follows:

Using the natural ordering of $\langle x \rangle \cong \mathbb{Z}$ with the chosen generator x , we can write the generators of $\mathcal{T}_1(\langle x \rangle)$ with all univalent vertices labelled by K in normal form (as in 2.3.1 above):

$$\mathcal{T}_1(\langle x \rangle) = \langle (x^m, x^n) \rangle_{1 \leq m < n} \oplus \langle (x^r, x^r) \rangle_{0 \leq r} \cong \mathbb{Z}^\infty \oplus \mathbb{Z}_2^\infty.$$

Expressing the projected $\text{INT}(x)$ relations $(x, x^r) + (x^{-1}, x^r) = 0$ for $r \in \mathbb{Z}$ in the normal form generators gives the relations:

$$(1) \quad (x, x^r) = (x, x^{r+1}) \quad 1 \leq r \quad \text{and} \quad (x^q, x^{q+1}) = (x^{q+1}, x^{q+2}) \quad 0 \leq q$$

which intersect when $r = 2$ and $q = 1$, so the $\text{INT}(x)$ relations identify an infinite family of normal form generators.

Now the projected image $\Phi(x)$ of each of the pairings $\tau_1(K, x^n S) = -(x, x^{n+1}) = -(x, x)$, for all n , is equal to (the inverse of) one of the generators identified by the INT relations, so the quotient of $\mathcal{T}_1(\langle x \rangle) / \text{INT}(x)$ by $\Phi(x)$ is gotten by setting all the terms in equation (1) equal to zero.

For example, by Theorem 3 any combination of Borromean surgeries on K which correspond to linear combinations of terms in equation (1) do not change the stable concordance class of K . On other hand, there still exists $\mathbb{Z}^\infty \oplus \mathbb{Z}_2^\infty$ many distinct stable concordance classes of knots which are order 1 Whitney concordant to K , as detected by the generators (x^m, x^n) with $|m - n| \geq 2$ and $m \neq 1$.

3.2. Computing $\tau_1(L, S)$. The computations of this subsection are used in the following subsection to describe the indeterminacy subgroup $\Phi(z)$ of Theorem 3.

It follows from Theorem 4 that $\tau_1(L, S)$ is invariant under isotopy of L and homotopy of S in M . In (the proof of) Lemma 3.2.1 below we will see how $\tau_1(L, S)$ is invariant under order 1 Whitney concordance of L , and is completely determined by S and $\tau_0(L)$.

In order to express $\tau_1(L, S)$ as a function of S and $\tau_0(L)$ it will be useful to define a map $\sigma_S : \pi_1 M \rightarrow \mathbb{Z}[\pi_1 M]$ for each S representing an element of $\pi_2 M$. This will be done by defining the map on each generator of $\pi_2 M$ as a $\pi_1 M$ -module, then extending linearly. We may ignore any generators which are spherical boundary components of M , since they clearly will have vanishing $\tau_1(L, S)$ for any L . The remaining generators consist of 2-spheres that decompose M into prime factors, and cross-sections of any $S^1 \times S^2$ factors. The maps are defined slightly differently for each of these two types of spheres. (The first-time reader could at this point jump to Lemma 3.2.1, and then begin to absorb the definition of σ_S during the subsequent proof.)

3.2.1. Separating spheres. Let S be an embedded separating sphere giving a connected sum decomposition of M as $M = M' \sharp_S M''$. Any $g \in \pi_1 M$ can be written $g = h_0 \prod_{l=1}^n g_l h_l$, where the g_l are carried by M' and the h_l are carried by M'' . Denote by $\beta_q := h_q \prod_{l=q+1}^n g_l h_l$, with $\beta_n = h_n$, and define $\sigma_S(g)$ by

$$\sigma_S(g) := \sum_{q=1}^n (g_q - 1) \beta_q \in \mathbb{Z}[\pi_1 M].$$

Thus, here $\sigma_S(g)$ is roughly an alternating sum of “tails” of g . For example, in the case $n = 2$ we have:

$$\sigma_S : g = h_0 g_1 h_1 g_2 h_2 \mapsto (g_1 - 1)(h_1 g_2 h_2) + (g_2 - 1)h_2.$$

3.2.2. Non-separating spheres. The map σ_S is similar but slightly more complicated for non-separating spheres. Let S be a spherical section of the $S^1 \times S^2$ factor in a connected sum decomposition of M as $M = (S^1 \times S^2) \sharp M'$. Any $g \in \pi_1 M$ can be written $g = h_0 \prod_{l=1}^n x^{\epsilon_l r_l} h_l$, where x is represented by the circle in the $S^1 \times S^2$ factor, with $\epsilon_l = \pm 1$ and the r_l positive integers, and the h_l are carried by M' . Let $\beta_q := h_q \prod_{l=q+1}^n x^{\epsilon_l r_l} h_l$, with $\beta_n = h_n$, and define $\sigma_S(g)$ by

$$\sigma_S(g) := \sum_{q=1}^n \epsilon_q \left(\sum_{l=1}^{r_q} x^{\epsilon_q(l + \frac{\epsilon_q - 1}{2})} \right) \beta_q \in \mathbb{Z}[\pi_1 M].$$

For example, in the case $n = 2$, $r_1 = 3 = r_2$, $\epsilon_1 = 1 = -\epsilon_2$ we have:

$$\sigma_S : g = h_0 x^3 h_1 x^{-3} h_2 \mapsto (x^3 + x^2 + x) h_1 x^{-3} h_2 - (1 + x^{-1} + x^{-2}) h_2.$$

So for separating spheres $\sigma_S(g)$ is again a sum of tails of g , but multiplied by the indicated unit-coefficient polynomials in x or x^{-1} .




FIGURE 13. If L' is homotopic to L by A' , then $L' \times I \subset M \times I$ is homotopic to $A' + (L \times I) - A' \looparrowright M \times I$.

3.2.3. Having defined σ_S on generators, the definition for arbitrary elements of $\pi_2 M$ is gotten by extending linearly: $\sigma_{(aS \pm a'S')}(g) := a\sigma_S(g) \pm a'\sigma_{S'}(g)$.

Lemma 3.2.1. *Let S be any immersed 2-sphere in M . For any $z \in \mathcal{T}_0(\pi_1 M)$, and any link $L \subset M$ of null-homotopic knots with $\tau_0(L) = z$,*

$$\tau_1(L, S) = \sum_{i \leq j} \sum_p e_p \cdot (g_p, 1, \sigma_S(g_p))_{ijS} \in \mathcal{T}_1(\pi_1 M) / \text{INT}(z)$$

where $z = \sum_{i \leq j} \sum_p e_p \cdot (g_p)_{ij}$.

Here z is written in normal form as in subsection 3.1 above.

Proof. First of all, we claim that if L' is any other link with $\tau_0(L') = z$, then $\tau_1(L', S) = \tau_1(L, S)$:

As mentioned above in subsection 3.1, the $\text{INT}(L')$ and $\text{INT}(L)$ relations coincide, since they only depend z . The equality of τ_0 implies that there exists a homotopy from L' to L whose trace is a collection of properly immersed annuli $A' \looparrowright M \times I$ supporting an order 1 Whitney tower, and connecting the link components in opposite ends of $M \times I$ (see 4.3 below). By running this homotopy partially forward, then waiting, and then backward, in increasing increments, one can construct a homotopy (rel boundary) from $L' \times I \subset M \times I$ to $A = A' + (L \times I) - A' \looparrowright M \times I$, where we write the composition of homotopies additively from left to right, and $-A'$ is the oppositely oriented A' which runs from L back to L' (see Figure 13). Since A' and $-A'$ each support an order 1 Whitney tower, A supports an order 1 Whitney tower such that no Whitney disks intersect the part of $M \times I$ containing $L \times I$. (If $\tau_0(L') \neq \tau_0(L)$ then this construction would still yield A admitting an order 1 Whitney tower, but there would necessarily be cancelling pairs of intersections with one point from each pair in A' and the other in $-A'$, so that the corresponding Whitney disks would have to intersect the part of $M \times I$ containing $L \times I$.) Now S is homotopic into the part of $M \times I$ containing $L \times I$, where Whitney




FIGURE 14. Four parallel Whitney disks pair intersections between a separating sphere S and a clasp link L with $\tau_0(L) = (g)_{ij}$ for $g = h_0 g_1 h_1 g_2 h_2$. The indicated decorations for the associated trees can be computed from the diagram, giving $\tau_1(L, aS) = (h_0, (g_1 h_1 g_2 h_2)^{-1}, a)_{ijS} - (h_0 g_1, (h_1 g_2 h_2)^{-1}, a)_{ijS} + (h_0 g_1 h_1, (g_2 h_2)^{-1}, a)_{ijS} - (h_0 g_1 h_1 g_2, h_2^{-1}, a)_{ijS} = (g, 1, a g_1 h_1 g_2 h_2 - a h_1 g_2 h_2 + a g_2 h_2 - a h_2)_{ijS}$, where the second equality comes from the HOL relations.

disks can be found for all intersections between S and $L \times I$ such that the Whitney disks do not intersect the parts of $M \times I$ containing A' and $-A'$ (since $\tau_0(L \times I, S) = 0$). It follows from the homotopy invariance of τ_1 that $\tau_1(L', S) = \tau_1(L, S)$, proving the claim.

So to prove the lemma we are free to use any link L with $\tau_0(L) = z$. This will not be hard using a clasp link (Figure 3 and subsection 4.4 below) whose intersections with S can be paired by Whitney disks that are contained in neighborhoods of the arcs guiding the clasps.

Consider the case where S is a separating sphere giving a connected sum decomposition $M = M' \#_S M''$, and $z = +(g)_{ij}$, is represented by a single positively signed order zero tree (edge) oriented from i to j , and labeled by $g \in \pi_1 M$. Writing $g = h_0 \prod_{l=1}^n g_l h_l$, where the g_l are carried by M' and the h_l are carried by M'' , we can create L , with $\tau_0(L) = (g)_{ij}$, from the unlink by introducing a clasp guided by an arc from the i th to the j th unlink component, which intersects S in $2n$ points, so that $A_i = L_i \times I$ has $2n$ cancelling pairs of intersections with S in some interior slice of $M \times I$, as illustrated in Figure 14 for the case $n = 2$.




FIGURE 15.

As can be seen in Figure 14, these cancelling pairs admit embedded Whitney disks, each of which winds along the band of the clasp and has a single interior intersection with $A_j = L_j \times I$. Using the product structure of $M \times I$, these Whitney disks can be made pairwise disjoint by a small perturbation. With our usual convention of orienting all edges towards the trivalent vertex, the group element labels on the trees associated to these Whitney disks can be seen to have the following pattern: If the whiskers on the Whitney disks are chosen to respect the factorization of g , then as we move through the Whitney disks from A_i towards A_j (right to left in the figure), the group elements labelling the i -edge read off increasing sub-words of g , while the group elements labelling the j -edge read off decreasing sub-words of g^{-1} , as illustrated in Figure 15. A choice of whisker on S determines an element $a \in \pi_1 M$ labelling all the edges corresponding to S . Now using the right multiplication of the HOL relations (geometrically, changing the whiskers on the Whitney disks) to get the group element g on the i -edge of each tree, simultaneously changes each j -edge label to 1, and changes the S -edges exactly as described by the map $\sigma_S(g)$ (see the example in 3.2.1 above), once the signs are taken into account, as can be checked in Figure 14 using our positive corner convention. Thus in this case, we have $\tau_1(L, aS) = (g, 1, a\sigma_S(g))_{ijS} = (g, 1, \sigma_{aS}(g))_{ijS}$ as desired.

It is easy to see that $\tau_1(L, S)$ behaves linearly under adding parallel copies of S , with each new copy of S having $2n$ more parallel Whitney disks, with appropriate sign changes for orientation-reversed copies of S .




FIGURE 16. The arc from L_i to L_j representing $g = h_0 x^3 h_1 x^{-3} h_2$ guides a clasp which, via the construction of Figure 14, leads to six Whitney disks – one for each intersection between the arc and the non-separating sphere S . Only the trees of the Whitney disks are shown, with edges decorated by group elements as computed from the diagram.

Fixing L and $z = (g)_{ij}$, and taking a different (non-homotopic) separating sphere S' , gives a different factorization of g , but the same construction works simultaneously to confirm the computation for $\pi_1 M$ -linear combinations of separating spheres:

$$\tau_1(L, aS + a'S') = (g, 1, a\sigma_S(g))_{ijS} + (g, 1, a'\sigma_{S'}(g))_{ijS'} = (g, 1, \sigma_{aS+a'S'}(g)).$$

And for $z = -(g)_{ij}$, changing the sign of the clasp gives a global change of sign for $\tau_1(L, S)$.

Since this entire construction takes place in a neighborhood of the guiding arc for the clasp determining g , it can be iterated any number of times giving the result for $z = \sum_{i \leq j} \sum_p e_p \cdot (g_p)_{ij}$ and combinations of separating spheres.

Checking the case where S is non-separating follows the same construction: Noting that the construction only depends on the arc guiding the clasp, Figure 16 shows a more schematic illustration of the computation of $\tau_1(z, S)$ for S a cross-section of an $S^1 \times S^2$ factor of M , in the case $z = +(g)_{ij}$ with $g = h_0 x^3 h_1 x^{-3} h_2$, where x represents a generator of the circle factor in $\pi_1 M$. Collecting terms and using HOL relations

again gives $\tau_1(L, aS) = (g, 1, \sigma_{aS}(g))_{ijS}$ (check with the example in 3.2.2 above). \square

3.3. The indeterminacy subgroup $\Phi(z)$. In the definition (1.3) of the relative order 1 intersection tree $\tau_1(L_z, L) := \tau_1(A)$, there is an obvious indeterminacy coming from changing the components of A by connected sums with 2-spheres. By Lemma 3.2.1 this indeterminacy is characterized by the subgroup $\Phi(z)$ which we define to be the span of the elements

$$\sum_{i \leq j} \sum_p e_p \cdot (g_p, 1, a\sigma_S(g_p))_{ijk} \in \mathcal{T}_1(\pi_1 M) / \text{INT}(z)$$

for $z = \sum_{i \leq j} \sum_p e_p \cdot (g_p)_{ij}$ written in normal form, where a ranges over $\pi_1 M$ and S ranges over a generating set for $\pi_2 M$. Each such element corresponds to the effect on $\tau_1(A)$ of tubing a 2-sphere aS into the k th component A_k .

4. PROOF OF THEOREM 3

This section clarifies the definition of the relative order 1 intersection tree $\tau_1(L_z, L)$ given in 1.3, and introduces notions relevant to the proof of Theorem 3. The proof is based on three lemmas which are left to the end of the section.

We remark that for links in 3-manifolds the equivalence relations of singular concordance (co-bounding immersed annuli in $M \times I$) and homotopy (level-preserving singular concordance) coincide by [10, 11].

4.1. The Ψ whisker-change actions. The invariants $\tau_0(A)$ and $\tau_1(A)$ depend on fixing whiskers on the m components of A . In the present setting, the corresponding indeterminacies are characterized by the following (left) actions of the m -fold cartesian product $\Psi := (\pi_1 M)^m$ on $\mathcal{T}_0(\pi_1 M)$ and $\mathcal{T}_1(\pi_1 M)$:

The action of $\psi = (\psi_1, \psi_2, \dots, \psi_m) \in \Psi$ on $\mathcal{T}_0(\pi_1 M)$ and $\mathcal{T}_1(\pi_1 M)$, respectively, is defined on generators by

$$(g_p)_{ij} \mapsto (\psi_i g_p \psi_j^{-1})_{ij}$$

and

$$(a, b, c)_{ijk} \mapsto (\psi_i a, \psi_j b, \psi_k c)_{ijk}.$$

4.2. Latitudes of singular concordances. Let $K \subset M$ be a knot, and let $A \looparrowright M \times I$ be a singular concordance from K to K' . A path that goes along a whisker on K in $M \times \{0\}$, then along A to K' , and then along a whisker on K' in $M \times \{1\}$ is called a *latitude* of A . Via projection to $M \times \{0\}$, a latitude of A determines an element in $\pi_1 M$.

For a singular concordance A of a link $L \subset M$, latitudes of the components determine an element $\psi \in \Psi$.

We write compositions of singular concordances additively from left to right. Note that if A_ψ and A_ϕ have latitudes determining ψ and ϕ respectively, then $A_\psi + A_\phi$ has latitudes determining the product $\psi\phi$. Also, if $-A_\psi$ denotes the inversion of A_ψ , then $-A_\psi$ has latitudes determining ψ^{-1} . Basepoints are assumed to be taken in $M \times \{0\}$ (unless otherwise stated), so for $n = 0, 1$ we have:

$$\tau_n(-A_\psi) = -\psi^{-1} \cdot \tau_n(A_\psi),$$

as well as:

$$\tau_n(A_\psi + A_\phi) = \tau_n(A_\psi) + \psi \cdot \tau_n(A_\phi)$$

and

$$\tau_n(A_\psi - A_\phi) = \tau_n(A_\psi) - \psi\phi^{-1} \cdot \tau_n(A_\phi)$$

where the actions of Ψ on \mathcal{T}_0 and \mathcal{T}_1 are both denoted by “ \cdot ”. Note that when inverting a singular concordance, the orientation of $M \times I$ is assumed to remain fixed. Specific orientation conventions are suppressed from notation, but consistent throughout the examples.

4.3. Order 1 Whitney concordance. Consider L and L' with $\tau_0(L) = z = \tau_0(L') \in \mathcal{T}_0(\pi_1 M)$, for some choices of whiskers. Placing L and L' in either end of $M \times I$ and connecting the components of singular null-concordances D and D' by disjointly embedded tubes gives a singular concordance A from L to L' . Using the whiskers on $L \subset M \times \{0\}$, we have

$$\tau_0(A) = \tau_0(D) - \psi \cdot \tau_0(D') = z - \psi \cdot z$$

for some $\psi \in \Psi$ which depends on the whiskers and the choices of tubes. Since we can re-choose the tubes in the construction of A to realize any element in Ψ , we can arrange for ψ to be trivial, so that $\tau_0(A) = 0 \in \mathcal{T}_0(\pi_1 M)$, which implies that L and L' are order 1 Whitney concordant.

(Note that A can be constructed to be the trace of a homotopy from L to L' : Take D and D' to be homotopies to the unlink, and realize the tubes by isotopies of the unlink components.)

It follows that $\tau_0(L) \in \mathcal{T}_0(\pi_1 M)$ classifies order 1 Whitney concordance for links of null-homotopic knots, since the converse is given by the last line of the proof of Theorem 2 in 2.6.

Note that the same argument shows that $\tau_1(L) \in \mathcal{T}_1(\pi_1 M)$ classifies order 2 Whitney concordance, hence stable concordance, for L with $\tau_0(L) = 0$. But the classification for $\tau_0(L) \neq 0$ – as per Theorem 3 – is more subtle (as is the classification in the case of essential knots given in Theorem 5 of Section 6).

4.4. Clasp links. For each $z \in \mathcal{T}_0(\pi_1 M)$ the link L_z in Theorem 3 can be taken to be a *clasp link* constructed from an m -component unlink U in the following way. Fix whiskers and orientations on U . For each $e_p \cdot (g_p)_{ij}$ in z , choose $|e_p|$ embedded *guiding arcs* joining U_i and U_j which represent $g_p \in \pi_1 M$. Replace each guiding arc by adding a band to the i th component which “clasps” the j th component as illustrated in Figure 3, with the sign of the clasp taken so that the intersection point in the trace of the homotopy which undoes the clasp has sign equal to the sign of $e_p \in \mathbb{Z}$.

For given z , any two clasp links L_z and L'_z are order 1 Whitney equivalent, since $\tau_0(L_z) = z = \tau_0(L'_z)$. Fixing a chosen clasp link L_z gives the classification of Theorem 3. Up to isotopy, there are three kinds of indeterminacies coming from the choices in the construction of a clasp link: twisting of the clasps (around the guiding arcs), knotting and linking of the guiding arcs, and the configurations of the guiding arc endpoints on U . As illustrated in Section 5 below, for any two choices the difference $\tau_1(L_z, L'_z)$ can be explicitly computed.

4.5. Stabilizers of the Ψ -action on \mathcal{T}_0 . The proof of Theorem 3, as well as its constraints, will depend on understanding the stabilizers of the Ψ -action on $\mathcal{T}_0(\pi_1 M)$.

For $z \in \mathcal{T}_0(\pi_1 M)$ written in normal form (as in subsection 3.1), the action $z \mapsto \psi \cdot z$ of $\psi = (\psi_1, \psi_2, \dots, \psi_m) \in \Psi$ is

$$z = \sum_{i \leq j} \sum_p e_p \cdot (g_p)_{ij} \mapsto \sum_{i \leq j} \sum_p e_p \cdot (\psi_i g_p \psi_j^{-1})_{ij},$$

so ψ is in the stabilizer Ψ_z of z if and only if for all i and j

$$\sum_p e_p \cdot (g_p)_{ij} = \sum_p e_p \cdot (\psi_i g_p \psi_j^{-1})_{ij},$$

which means that each pair of factors ψ_i and ψ_j permutes the elements g_p , and if $i = j$ possibly sends some g_p to g_p^{-1} . (Of course permuting the g_p in z_{ij} is not sufficient for ψ to be in Ψ_z since the coefficients of the permuted elements would also have to be equal.)

A stabilizer Ψ_z is called *untwisted* if for all $\psi \in \Psi_z$, and all i and j ,

$$g_p = \psi_i g_p \psi_j^{-1},$$

and Ψ_z is called *twisted* otherwise. So untwisted stabilizers are exactly those which induce the identity map on the fundamental group elements in z_{ij} for all i and j .

We will also refer to an element ψ of Ψ_z as *twisted* or *untwisted*, depending on whether ψ induces a non-trivial or trivial permutation of the elements in z ; so that Ψ_z is untwisted if every ψ in Ψ_z is untwisted.

As discussed in 5.2 below, in a sense “most” z have untwisted stabilizers.

4.6. $\tau_1(L_z, L)$ is well-defined. As discussed in subsection 4.3 above, it is natural to interpret the values of the “absolute” invariants $\tau_0(L)$ and $\tau_1(L)$ modulo the respective Ψ -actions. This means that these invariants are completely independent of choices of whiskers.

In the definition of $\tau_1(L_z, L)$ we consider L_z to be fixed, including the choice of whiskers. Then, for each L the value of $\tau_1(L_z, L)$ in the quotient of $\mathcal{T}_1(\pi_1 M)$ by $\text{INT}(z)$ and $\Phi(z)$ is taken modulo the action of the stabilizer Ψ_z . The reason for this is essentially due to Lemma 4.8.2 below as will become clear shortly.

Let A and A' be singular concordances from L_z to L supporting order 1 Whitney towers. Then the composition $A - A'$ is a singular self-concordance of L_z supporting an order 1 Whitney tower. By Lemma 4.8.2 below, latitudes of $A - A'$ determine an element $\psi \in \Psi_z$. Since Ψ_z is assumed to be untwisted, by Lemma 4.8.3 below there exists a self-homotopy $H_{\psi^{-1}}^0$ of L_z with latitudes determining ψ^{-1} such that $\tau_1(H_{\psi^{-1}}^0) = 0 \in \mathcal{T}_1(\pi_1 M)$. Now the singular self-concordance $A - A' + H_{\psi^{-1}}^0$ of L_z has latitudes determining the trivial element $\psi\psi^{-1} \in \Psi$, so by Lemma 4.8.1 below, $A - A' + H_{\psi^{-1}}^0$ is regularly homotopic (rel ∂) to the connected sum of $L_z \times I$ with disjointly embedded 2-spheres. By the homotopy invariance of τ_1 and the definition of $\Phi(z)$ (3.3) we have

$$\tau_1(A - A' + H_{\psi^{-1}}^0) = \tau_1(A) - \psi \cdot \tau_1(A') \in \Phi(z),$$

so $\tau_1(L_z, L)$ does not depend on the choice of A .

4.7. The equivalence of statements (i)-(iv) in Theorem 3. We will show the equivalence of (i) and (iii); the equivalence of (ii), (iii), and (iv) follows from Theorem 4 in Section 3.

If L and L' are order 2 Whitney concordant, then they cobound an immersed annulus A' which supports an order 2 Whitney tower. So any singular concordance A used to compute $\tau_1(L_z, L)$ can be extended by A' to compute $\tau_1(L_z, L') = \tau_1(A + A') = \tau_1(L_z, L)$, since $\tau_1(A')$ vanishes.

On the other hand, the equality $\tau_1(L_z, L') = \tau_1(L_z, L)$ means that if A and A' are order 1 Whitney concordances from L_z to L and L' respectively, then $\tau_1(A') = \psi \cdot \tau_1(A)$ modulo $\Phi(z)$, for some $\psi \in \Psi_z$. After realizing $\text{INT}(z)$ relations and taking connected sums with some 2-spheres (if necessary) we may arrange that $\tau_1(A') = \psi \cdot \tau_1(A) \in \mathcal{T}_1(\pi_1 M)$. By Lemma 4.8.3, there exists a self-homotopy $H_{\psi^{-1}}^0$ of L_z

with $\tau_1(H_{\psi^{-1}}^0) = 0$ and latitudes determining ψ^{-1} . Now the composition $-A + H_{\psi^{-1}}^0 + A'$ is an order 1 Whitney concordance from L to L' , and taking the whiskers on L_z at the end of $-A$ we have

$$\tau_1(-A + H_{\psi^{-1}}^0 + A') = -\tau_1(A) + \psi^{-1} \cdot \tau_1(A') = -\tau_1(A) + \psi^{-1} \psi \cdot \tau_1(A) = 0$$

so by Theorem 4 it follows that $-A + H_{\psi^{-1}}^0 + A'$ is homotopic to a singular concordance from L to L' which supports an order 2 Whitney tower.

4.8. Lemmas. The proof of Theorem 3 is completed by the following three lemmas.

Lemma 4.8.1 ([26]). *If a singular self-concordance A of a knot $K \subset M$ has a latitude that represents the trivial element $1 \in \pi_1 M$, then A is homotopic (rel ∂) to the connected sum of an embedded 2-sphere and the embedded annulus $K \times I$ in $M \times I$.*

Furthermore, if A also supports a Whitney tower of order 1 (or greater), and K has trivial order zero intersections with 2-spheres in M , then the above homotopy (rel ∂) can be taken to be a regular homotopy.

Note that K is not assumed to be null-homotopic.

Proof. The properly immersed A projects to a map of a torus $T \rightarrow M = M \times \{0\}$, with the image of a latitude in A projecting to a loop C in the image of T which is contractible in M . The inverse image \overline{C} of C is algebraically dual to the inverse image \overline{K} of K in the domain torus T , so after a homotopy we may assume that \overline{C} is geometrically dual to \overline{K} and embedded. Then C bounds an immersed 2-disk D in M which surgers the map $T \rightarrow M$ to a map of a 2-sphere $S \rightarrow M$. This surgery lifts to a homotopy (rel ∂) of A in $M \times I$ to $K \times I \sharp S'$, where S' is a lift of S . By Lemma 2.6.1, S' can be assumed to be embedded.

The second statement of Lemma 4.8.1 follows from the fact that, up to isotopy, a homotopy of surfaces in a 4-manifold consists of a sequence of finger moves, Whitney moves, and cusp homotopies. If the cusp homotopies occur in cancelling pairs, then they can be replaced (up to isotopy) by finger moves (pairs of births) and Whitney moves (pairs of deaths), so that the homotopy is a regular homotopy. So, if $K \times I \sharp S'$ and A both have all self-intersections occurring in cancelling pairs – as will be the case under the further assumptions – then the above homotopy may be arranged to be a regular homotopy. \square

Lemma 4.8.2. *For any $z \in \mathcal{T}_0(\pi_1 M)$, let $L_z \subset M$ be any link such that $\tau_0(L_z) = z \in \mathcal{T}_0(\pi_1 M)$, and let $A_\psi \looparrowright M \times I$ be any singular self-concordance of L_z , with latitudes determining $\psi \in \Psi$. Then:*




FIGURE 17. The guiding arc traces out a Whitney disk as a clasp is undone then redone.

If A_ψ admits an order one Whitney tower, then $\psi \in \Psi_z$.

Proof. As described in subsection 4.3, there exists a self-homotopy H_ψ of L_z with latitudes determining ψ such that $\tau_0(H_\psi) = z - \psi \cdot z \in \mathcal{T}_0(\pi_1 M)$. Since the composition $A_\psi - H_\psi$ has latitudes representing the trivial element $\psi\psi^{-1} \in \Psi$, it follows from Lemma 4.8.1 that $A_\psi - H_\psi$ is homotopic (rel ∂) to the connected sum of $L_z \times I$ with some 2-spheres. By the homotopy invariance of τ_0 , and since null-homotopic knots have trivial order zero intersections with 2-spheres, we have that $\tau_0(A_\psi - H_\psi) = 0 \in \mathcal{T}_0(\pi_1 M)$. So, computing in $\mathcal{T}_0(\pi_1 M)$ we have

$$0 = \tau_0(A_\psi - H_\psi) = \tau_0(A_\psi) - \tau_0(H_\psi) = 0 - (z - \psi \cdot z)$$

since $\tau_0(A_\psi) = 0$, which shows that $\psi \in \Psi_z$. \square

Lemma 4.8.3. *Let $L_z \subset M$ be a clasp link with $\tau_0(L_z) = z \in \mathcal{T}_0(\pi_1 M)$, such that Ψ_z is untwisted. Then for any $\psi \in \Psi_z$, there exists a self-homotopy H_ψ^0 of L_z with latitudes determining ψ , and an order one Whitney tower \mathcal{W} on H_ψ^0 , such that $\tau_1(\mathcal{W}) = 0 \in \mathcal{T}_1(\pi_1 M)$.*

Proof. The construction of H_ψ^0 has three steps, and is based on the observation that undoing and re-doing clasps describes a self-homotopy supporting an order 1 Whitney tower whose Whitney disks are described by the guiding arcs of the clasps (Figure 17):

First, undoing the clasps by crossing changes describes a homotopy from L_z to the unlink U , with the guiding arcs tracing out corners of embedded Whitney disks near the crossing-change intersection points (Figure 3). The components of U can be assumed to lie in small 3-balls, and the guiding arcs between U_i and U_j represent the elements $(g_p)_{ij}$ in z_{ij} .

Now, the untwistedness of ψ means that $g_p = \psi_i g_p \psi_j^{-1} \in \pi_1 M$ for all i and j (and p which depends on i and j) which is exactly the condition that a self-isotopy of U around loops representing ψ extends to self-homotopies of the guiding arcs and whiskers which are attached to U . Each of these self-homotopies of guiding arcs traces out the




FIGURE 18. From left to right, the first part of the homotopy H_ψ^0 , and from right to left the third part. The guiding arcs trace out Whitney disks, and gray arcs trace out an extension of the Whitney section. The right most picture includes the surface F .

center rectangle of an immersed Whitney disk in \mathcal{W} . The self-isotopy of U extends to a self-isotopy of the small 3-ball neighborhoods of the components of U together with a neighborhood of the whiskers, which forms a thickened wedge of arcs. So the only singularities created in this second step of the construction are possible crossing changes between the guiding arcs which are interior intersections among the Whitney disks of \mathcal{W} .

The third step is to run the first homotopy backwards from U to L_z , recreating the clasps by crossing changes, each of which gives an intersection point which forms a cancelling pair with the corresponding intersection created when undoing the clasps in the first step. In this third step the guiding arcs shrink down and disappear at the intersection points, tracing out the remaining corners of the Whitney disks in \mathcal{W} .

The union H_ψ^0 of these three steps is a self-homotopy of L_z which by construction has latitudes determining ψ . Assuming for the moment that the Whitney disks are framed, it is clear that $\tau_1(\mathcal{W}) = 0 \in \mathcal{T}_1(\pi_1 M)$ because the only singularities are interior intersections among the Whitney disks which do not contribute to τ_1 (they are *order 2* intersections). It remains to show that the Whitney disks can be arranged to be framed, which will follow essentially from the fact that Chernov's affine self-linking numbers for framed knots in 3-manifolds vanish modulo 2 ([4]).

Recall that the normal section which defines the Whitney disk framing obstruction is defined by pushing one Whitney disk boundary arc tangentially along its surface sheet and pushing the other boundary arc in the normal direction to its surface sheet. Observe that after




FIGURE 19. An *interior twist* on a Whitney disk is the local introduction of a self-intersection. Creation of a ± 1 -intersection changes the framing by ∓ 2 , as can be seen in the intersections with a normal push-off of the Whitney disk shown in grey. Note that near the self-intersection of the Whitney disk there are a pair of intersections between the Whitney disk and its push-off which are just artifacts of the immersion of the disk bundle and do not contribute to the framing obstruction.

a small isotopy ($\pi/2$ radians in the normal circle bundle) the part of the section that was pushed normally to a surface sheet will lie in the surface sheet, except for a short arc near each intersection point (in the right-hand side of Figure 4 picture the back half of the dotted loop sliding up to the top of the torus). This isotopy does not change the framing obstruction, and will make it easier to compute the framings in the construction of \mathcal{W} .

Push-offs of the Whitney disks in \mathcal{W} extending the Whitney sections are given by parallel copies of the guiding arcs in the above description of H_ψ^0 (Figure 18). As shown in Figure 18, the parts of the Whitney disks in the first and third homotopies are disjoint from their push-offs, and so the only possible contributions to framing obstructions can occur in the second part of the homotopy, which we will call H .

At the start of H we have guiding arcs and their parallel copies running between components of U . The components of U bound disjointly embedded 2-disks (whose interiors are disjoint from all guiding arcs), and the guiding arc pairs determine bands between these 2-disks. After thickening the whiskers on U to a wedge of bands joined near the base-point of M , the resulting embedded surface F (shown in the right-most picture of Figure 18) gives a normal framing on each of the (embedded) loops representing the $(g_p)_{ij}$.

Now H is a self-homotopy of the union of U together with its whiskers and the guiding arcs. Clearly H extends to a homotopy of F (which lies in a neighborhood of U union whiskers and guiding arcs). We may assume that H restricts to a self-isotopy on the sub-disk of F which is the union of the disks bounded by U and the wedge of bands

containing the whiskers. At the end of H the guiding arcs are back where they started, but the parallel copies might not be, since the twisting of the bands may have changed (by full twists) during H . The framing obstruction on each Whitney disk in \mathcal{W} is equal to this change in framing of the corresponding loop in F at the end of H . By the main results of [4], these relative framings can only differ by an even integer. So, after perhaps performing interior twists (which do not contribute to τ_1) as illustrated in Figure 19, the Whitney disks in the above construction of \mathcal{W} on H_ψ^0 can assumed to be framed. \square

5. EXAMPLES

This section computes change-of-base-link formulas for $\tau_1(L_z, \cdot)$, characterizes twisted stabilizers, and shows how twisted Ψ_z can lead to non-trivial indeterminacies in $\tau_1(L_z, \cdot)$.

5.1. Clasp links. For given $z \in \mathcal{T}_0(\pi_1 M)$, a clasp link L_z is determined up to isotopy by choices at three steps in the construction: the configuration of endpoints of guiding arcs on the unlink, the isotopy classes (rel endpoints) of the guiding arcs, and the (relative) framing of the band corresponding to the thickening of the guiding arc to a clasp. This subsection illustrates the computation of the order 1 relative intersection tree $\tau_1(L_z, L'_z)$ for clasp links L_z and L'_z , giving a “change of base-link” formula for Theorem 3.

5.1.1. Twisting clasps. Figure 20 illustrates the trace A of a homotopy between clasp links L_z and L'_z which differ by one twist of a clasp representing $g \in \pi_1 M$ decorating $(g)_{ji}$ in z . The cancelling pair of intersections of A admit an embedded Whitney disk whose interior is disjoint from A , but which is not framed, as illustrated in Figure 21. Figure 22 shows how an application of the boundary twist procedure corrects the framing on the Whitney disk at the cost of creating an interior intersection between the Whitney disk and A , giving $\tau_1(L_z, L'_z) = (1, 1, g)_{ij}$, which is 2-torsion by the AS relations. In general, twisting such a band n times would give $\tau_1(L_z, L'_z) = n(1, 1, g)_{ij}$.

5.1.2. Choices of guiding arcs. If L_z and L'_z differ only by a homotopy (rel endpoints) of guiding arcs then $\tau_1(L_z, L'_z) = 0$: This can be seen directly from the “band pass” move illustrated in Figure 23. Or, constructing a homotopy between L_z and L'_z supporting Whitney disks described by the guiding arcs, the crossings between guiding arcs correspond to interior intersections between Whitney disks and do not contribute to τ_1 (as in the proof of Lemma 4.8.3).




FIGURE 20. The trace of a homotopy between clasp links L_z and L'_z which differ by a single twist in one clasp. The Whitney disk described by the vertical arcs is not correctly framed, as shown in Figure 21.




FIGURE 21. The grey arcs show an extension of the Whitney section over the interior of the Whitney disk. The single intersection between the extension and the Whitney disk indicates that the Whitney disk is not framed.




FIGURE 22. Changing the Whitney disk of Figures 20 and 21 by the indicated local procedure of introducing a boundary twist creates a correctly framed Whitney disk at the cost of creating a single intersection (center picture) between the interior of the Whitney disk and the sheet that was twisted around.




FIGURE 23. A homotopy which pushes a clasp of the i th component across a clasp of the j th component gives rise to two parallel oppositely oriented Whitney disks (“inside” the dotted j -clasp) whose interior intersections with the k th component will have cancelling trees.




FIGURE 24. For clasp links L_z and L'_z differing by the illustrated single transposition of guiding arc endpoints, $\tau_1(L_z, L'_z) = (1, g, h)_{ijk}$.

5.1.3. *Guiding arc endpoint configurations.* Figure 24 shows the computation $\tau_1(L_z, L'_z) = (1, g, h)_{ijk}$, where L_z and L'_z are related by a single transposition of guiding arc endpoints along the i th component. The reader familiar with finite type theory will recognize this as a decorated STU relation.

The transposition of the two endpoints of the *same* guiding arc is illustrated in Figure 25, which computes the corresponding change in τ_1 using the formula for intersections between Whitney disk boundaries. This computation will be used in Example 5.3 below.




FIGURE 25. This figure describes a homotopy A from the clasp knot K shown in the upper left to the clasp knot K' which is determined by the guiding arc shown in the lower right. The undoing of the positive clasp creates a positive self-intersection in A , and the redoing of the clasp creates a cancelling negative self-intersection. These self-intersections are paired by a framed Whitney disk W (described by the trace of the guiding arc) whose boundary arcs ∂_+W and ∂_-W intersect in a single point p where ∂_-W pushes across ∂_+W just before the lower right picture as indicated by the horizontal dotted arrow. The dotted vectors pointing lower-right point in the direction of increasing time, and the dotted vectors pointing upper-left point into the past. The orientation of A is given by a tangent vector to K , together with a second tangent vector pointing into the future. The Whitney disk boundary arc ∂_+W is oriented towards the positive self-intersection, and the boundary arc ∂_-W is oriented towards the negative self-intersection. At p the orientation of A agrees with $(\partial_-W, \partial_+W)_p$. Using the assignment of trees to intersections between Whitney disk boundaries given by the formula $\epsilon_p \cdot t_p := \epsilon_k \epsilon_j (1, g_k^{\epsilon_k}, g_j^{\epsilon_j})$ in the “independence of Whitney disk boundaries” discussion in 2.4, and computing from the figure gives $\tau_1(K, K') = \tau_1(A) = -(g, g^{-1}) = +(g^{-1}, g)$, in the notation $(g, h) := (1, g, h)$.

5.2. Twisted stabilizers. This subsection points out that twisted stabilizers only occur in the presence of circle bundles over non-orientable surfaces with orientable total space, or Seifert fibered submanifolds containing singular fibers, and illustrates how twisted stabilizers can lead to self-homotopies of knots having non-trivial τ_1 .

To present the essential ideas with a minimum of subscripts we concentrate on the case where $z = \tau_0(K)$ for a knot $K \subset M$. Identifying $\mathcal{T}_0(\pi_1 M)$ with $\mathbb{Z}[\pi_1 M]$ modulo inversion and trivial elements, and writing $z = \sum e_i g_i$ in normal form (with all $e_i \neq 0$, $g_i \neq 1$, and with $g_i \neq g_j^{\pm 1}$ for $i \neq j$), any element ψ in the stabilizer Ψ_z of z satisfies the equation

$$\sum e_i \psi g_i \psi^{-1} = \sum e_i g_i \quad \text{modulo } g = g^{-1}.$$

So the conjugation action of ψ permutes the (finite) set $\{g_i^{\pm 1}\}$, and by definition ψ is twisted if this permutation is non-trivial, and untwisted if this permutation is the identity.

Assuming that ψ is twisted, there is a (least) natural number $2 \leq n \in \mathbb{N}$ such that ψ^n induces the identity permutation on $\{g_i^{\pm 1}\}$. Writing $\zeta(z) := \cap_i \zeta(g_i)$ for the intersection of the centralizers $\zeta(g_i)$ in $\pi_1 M$ of the g_i , we have $\psi^n \in \zeta(z)$, but $\psi \notin \zeta(z)$. From work of Jaco-Shalen, Casson-Jungreis, and Gabai, non-cyclic centralizers of non-trivial elements of $\pi_1 M$ are carried by a codimension zero *characteristic submanifold* of M which is Seifert fibered ([3, 9, 16, 17, 18]). Since n th roots of any element are contained in the centralizer of that element, it can be checked that both $\zeta(z)$ and $\zeta(\psi^n)$ are carried by the same connected component N of the characteristic submanifold of M .

It follows from section 4 of [18] (in particular Theorem 4.4) that there are only two possible ways for $\psi^n \in \zeta(z)$, but $\psi \notin \zeta(z)$: Either ψ and $\{g_i\}$ are all carried by a submanifold $B \widetilde{\times} S^1 \subset N$ which is the orientable circle bundle over the Möbius band, or ψ is carried by a neighborhood of singular fiber in N and $\{g_i\}$ lie in the canonical subgroup of $\pi_1 N < \pi_1 M$ consisting of elements represented by loops which project to orientation-preserving loops in a base surface for N . In both cases the possible z can be described explicitly as follows.

5.2.1. The twisted circle bundle over the Möbius band. In the first case, we use the notation

$$\pi_1(B \widetilde{\times} S^1) = \langle a, f \mid afa^{-1} = f^{-1} \rangle < \pi_1 M$$

with f represented by a circle fiber, a represented by a core of the Möbius band B , and normal form $a^n f^l$ for $n, l \in \mathbb{Z}$. It is easily checked that the only candidates to be twisted stabilizers are $\psi = a^{2r+1} f^s$, with

$\psi^2 \in \langle a^{2k} \rangle$ central. For all $r, s \in \mathbb{Z}$, the orbits under conjugation by $a^{2r+1}f^s$ are:

$$\{a^{2k}\} \quad \{f^l, f^{-l}\} \quad \{a^{2k}f^l, a^{2k}f^{-l}\}$$

for all nonzero integers k and l ; and

$$\{a^{2k+1}f^l, a^{2k+1}f^{2s-l}\} \quad \{a^{2k+1}, a^{2k+1}f^{2s}\} \quad \{a^{2k+1}f^s\}$$

for all integers k and l with $l \neq s$. So in this case Ψ_z is twisted exactly when z is of the form:

$$\begin{aligned} & \sum_{l \neq 0, k \neq 0} e_{kl}(a^{2k}f^l + a^{2k}f^{-l}) + e_l f^l + e'_k a^{2k} \\ & + \sum_{l \neq s} d_{kl}(a^{2k+1}f^l + a^{2k+1}f^{2s-l}) + d_k(a^{2k+1} + a^{2k+1}f^{2s}) + d'_k a^{2k+1}f^s, \end{aligned}$$

where the coefficients $e_{kl}, e_l, e'_k, d_{kl}, d_k, d'_k$ are integers, with some coefficient other than the e'_k non-zero.

We will show below in 5.3 that there is a self-homotopy A of a knot K having a latitude determining $a \in \Psi_z$ with $\tau_0(K) = z = f + f^3$ such that $\tau_1(A) \neq 0 \in \mathcal{T}_1(\pi_1 M)/\text{INT}(z)$. This illustrates how Lemma 4.8.3 (hence the proof of Theorem 3) can fail if Ψ_z is twisted.

On the other hand, related examples will show that for K with $\tau_0(K) = z = e_l f^l$, all self-homotopies A of K have $\tau_1(A) = 0 \in \mathcal{T}_1(\pi_1 M)/\text{INT}(z)$. Thus, a twisted Ψ_z does not guarantee the failure of Lemma 4.8.3, and so Theorem 3 can be extended to allow some twisted stabilizers.

5.2.2. Singular Seifert fibers. In the case where ψ is carried by a neighborhood of singular fiber in N and $\{g_i\}$ lie in the canonical subgroup of $\pi_1 N < \pi_1 M$, we have $\psi = \phi^r f^s$ for $r, s \in \mathbb{Z}$ where f is represented by a regular fiber of N and ϕ is represented by a singular fiber of N . In this case $n \in \mathbb{N}$ is the smallest power such that ϕ^{rn} is contained in the cyclic (normal) subgroup $\langle f \rangle$ of $\pi_1 N$. We get fixed z of the form:

$$\sum e_k(g_k + \phi^r g_k \phi^{-r} + \phi^{2r} g_k \phi^{-2r} + \cdots + \phi^{(n-1)r} g_k \phi^{-(n-1)r}) + e_{pq} \phi^p f^q$$

where the $\phi^p f^q$ lie in $\zeta(\phi^r f^s)$, so at least one e_k must be non-zero.

5.2.3. Twisted Ψ_z for links. Now let L be a link with $\tau_0(L) = z = \sum_{i \leq j} z_{ij}$, with each z_{ij} written in normal form:

$$z_{ij} = \sum_p e_p(g_p)_{ij}.$$

A twisted $\psi \in \Psi_z$ induces a non-trivial permutation of at least one of the sets $\{g_p^{\pm 1}\}$ for $i = j$ or $\{g_p\}$ for $i < j$.

In the cases $i = j$ the above discussion for knots applies to describe restrictions on z_{ii} and ψ_i for twisted $\psi \in \Psi_z$. (Of course these restrictions are not independent of each other in the presence of non-zero z_{ij} with $i < j$.)

Considering the case where $i < j$ and ψ induces a non-trivial permutation of $\{g_p\}$ (twisting z_{ij}), there exists $n \geq 2$ such that $\psi_i^n g_p \psi_j^{-n} = g_p$ for all g_p . Thus $g_p^{-1} \psi_i^n g_p = \psi_j^n$, and $g_p \psi_j^n g_p^{-1} = \psi_i^n$ for all g_p . It follows from Proposition 4.5 of [18] that ψ_i and ψ_j are carried by singular fibers in a characteristic component $N \subset M$.

A more detailed analysis of twisted stabilizers would be interesting, but we stop here and conclude:

Proposition 5.2.1. *If M contains no orientable circle bundles over non-orientable surfaces and no Seifert fibered submanifolds having singular fibers, then Ψ_z is untwisted for all $z \in \mathcal{T}_0(\pi_1 M)$.*

By direct computation one can extend Lemma 4.8.3 (hence Theorem 3) to certain simple examples of z with twisted $\psi \in \Psi_z$ carried by a singular Seifert fiber, and it is possible that twisted Ψ_z which correspond only to singular Seifert fibers do not contribute any non-trivial indeterminacies to $\tau_1(L_z, \cdot)$, but I do not know of a general construction. The next example shows that Lemma 4.8.3 can indeed fail for twisted Ψ_z which correspond to orientable circle bundles over non-orientable surfaces.

5.3. Knots in the twisted circle bundle over a Möbius band.

Using the notation of subsection 5.2.1 above, this example will consider knots K in $B \widetilde{\times} S^1$ with $\tau_0(K) = z = \sum e_l f^l$ carried by the circle factor. Such z has twisted Ψ_z , with the conjugation action of a (carried by the Möbius band core) inverting the f^l . After giving an example of a self-homotopy A of such a K with non-trivial $\tau_1(A)$ (so that Lemma 4.8.3 fails), we will observe that in some cases Lemma 4.8.3 can be extended to K with twisted Ψ_z .

Throughout this subsection $(g, h) := (1, g, h)$ with the common univalent labels suppressed.

Since $B \widetilde{\times} S^1$ is irreducible, the $\Phi(z)$ relations are trivial throughout this subsection.

5.3.1. A self-homotopy with non-trivial τ_1 . Illustrated in Figure 26, using the unknot-plus-guiding-arcs notation, is a self-homotopy A of a clasp knot K in $B \widetilde{\times} S^1$, with $\tau_0(K) = z = f + f^3$. The right-back and left-front faces of the rectangular prism are identified by a half-rotation, and the top and bottom faces are identified by a translation, forming

$B \tilde{\times} S^1$. The clasp knot K is described by the left-most rectangular unknot U together with two guiding arcs, one which wraps once around the circle fiber and a longer one which wraps around three times. The self-homotopy A pushes K along the length of the Möbius band, so A has a latitude representing the element $a \in \pi_1 B \tilde{\times} S^1$. During the description of A , the guiding arcs can be thought of either as representing the ‘bands’ of K , or as tracing out a pair of Whitney disks in an order 1 Whitney tower on A .

As explained in the caption of Figure 26, $\tau_1(A) = (f, f) + (f^{-1}, f) + (f^3, f^3) + (f^{-3}, f^3)$, which we will show is non-zero in $\mathcal{T}_1(\pi_1 B \tilde{\times} S^1)/\text{INT}(z)$.

Noting that the edge decorations in $\tau_1(A)$ are carried by circle fibers, the relevant $\text{INT}(z)$ relations for $z = f + f^3$ are:

$$0 = (z + \bar{z}, f^r) = (f, f^r) + (f^{-1}, f^r) + (f^3, f^r) + (f^{-3}, f^r) \quad \text{for } r \in \mathbb{Z}.$$

Denoting these relations by Δ_r and writing them in normal form (f^n, f^m) with $0 < n < m$ or $0 = n \leq m$, we have

$$\begin{aligned} \Delta_r &= (f, f^r) - (f, f^{r+1}) + (f^3, f^r) - (f^3, f^{r+3}) & \text{for } 3 < r \\ \Delta_{-r} &= (f^r, f^{r+1}) - (f^{r-1}, f^r) + (f^r, f^{r+3}) - (f^{r-3}, f^r) & \text{for } -r \leq -3 \\ \Delta_3 &= (f, f^3) - (f, f^4) + (1, f^3) - (f^3, f^6) \\ \Delta_2 &= (f, f^2) - (f, f^3) - (f^2, f^3) - (f^3, f^5) \\ \Delta_1 &= (1, f) - (f, f^2) - (f, f^3) - (f^3, f^4) \\ \Delta_{-1} &= (f, f^2) + (1, f) + (1, f^4) + (f^2, f^3) \\ \Delta_{-2} &= (f^2, f^3) - (f, f^2) + (f^2, f^5) + (f, f^3) \end{aligned}$$

with Δ_0 already zero by the AS and FR relations.

We will show that $\tau_1(A)$ is not equal to a finite sum $\sum d_i \Delta_i$ in $\mathcal{T}_1(\pi_1 B \tilde{\times} S^1)$:

In normal form we have $\tau_1(A) = (1, f) - (f, f^2) + (1, f^3) - (f^3, f^6)$. Assume that $(1, f) - (f, f^2) + (1, f^3) - (f^3, f^6) = \sum d_i \Delta_i$. Note that for all $5 \leq r$ the coefficient of (f, f^r) must vanish, so $d_{r-1} + d_r = 0$. Thus, if d_4 is odd, then $\sum d_i \Delta_i$ can not be a finite sum. Similarly, for all $5 \leq r$ the coefficient of (f^{r-1}, f^r) must vanish, so $d_{-(r-1)} + d_{-r} = 0$. Thus, if d_{-4} is odd, then $\sum d_i \Delta_i$ can not be a finite sum.

Working modulo 2, the first three terms in $\tau_1(A)$ give the following relations among the coefficients:

$$\begin{aligned} (1, f) : & \quad d_1 + d_{-1} \equiv 1 \\ (f, f^2) : & \quad d_2 + d_{-2} \equiv 0 \\ (1, f^3) : & \quad d_3 + d_{-3} \equiv 1. \end{aligned}$$

The vanishing of the coefficient of (f, f^3) gives:

$$d_1 + d_2 + d_{-2} + d_3 \equiv 0$$




FIGURE 26. From left to right are pictured five stages of the self-homotopy A of K in the twisted circle bundle $B \tilde{\times} S^1$ over the Möbius band. During the first three stages, the unknot U flips over while fixing its edge that contains the endpoints of the guiding arcs. Between the second and third stages U intersects each guiding arc near one endpoint – contributing $(f, f) + (f^3, f^3)$ to $\tau_1(A)$ – and intersects two strands of the longer guiding arc in cancelling pairs of intersections which contribute nothing to $\tau_1(A)$. From the third stage to the fourth stage is just an ambient isotopy of U together with the guiding arcs which is rotation by 180° in the horizontal plane. From the fourth stage to the fifth stage each guiding arc pushes one of its endpoints across the other – contributing $(f^{-1}, f) + (f^{-3}, f^3)$ to $\tau_1(A)$, as illustrated in Figure 25 – and the vertical strands of the longer guiding arc pass through each other contributing nothing to $\tau_1(A)$. From the fifth stage back to the first stage is just an isotopy across the back face and in from the front face. Thus $\tau_1(A) = (f, f) + (f^{-1}, f) + (f^3, f^3) + (f^{-3}, f^3)$, which will be seen to be non-zero in $\mathcal{T}_1(\pi_1 B \tilde{\times} S^1) / \text{INT}(z)$ (even modulo 2, in case you don't want to check signs).

which implies that $d_1 \equiv d_3$, since $d_2 + d_{-2} \equiv 0$. Now the vanishing of the coefficient of (f^3, f^4) , together with the congruence $d_1 \equiv d_3$, give:

$$0 \equiv d_1 + d_{-3} + d_4 + d_{-4} \equiv d_3 + d_{-3} + d_4 + d_{-4} \equiv 1 + d_4 + d_{-4},$$

so one of d_4 or d_{-4} is odd, and $\sum d_i \Delta_i$ can not be a finite sum.

5.3.2. *Extending Lemma 4.8.3 to some twisted stabilizers.* The construction of the self-homotopy A in Figure 26 generalizes to clasp knots K with $\tau_0(K) = \sum e_l f^l$, yielding self-homotopies A with latitudes representing a , and

$$\tau_1(A) = \sum e_l [(f^l, f^l) + (f^{-l}, f^l)].$$

In the case where $\tau_0(K) = z = e_l f^l$, it is clear that $\tau_1(A) = e_l [(f^l, f^l) + (f^{-l}, f^l)] \in \text{INT}(z)$ by taking $r = l$ in the INT relations, and Lemma 4.8.3 can be extended to such K . (In the case $e_l = 1$ so K has just a single clasp then A can be taken to be a self-isotopy of K .) Notice that composing A with itself in the general case gives a self-homotopy $A + A$ of K with a latitude representing a^2 which is untwisted, and the computation

$$\begin{aligned} \tau_1(A + A) &= \tau_1(A) + a \cdot \tau_1(A) \\ &= \sum e_l [(f^l, f^l) + (f^{-l}, f^l) + (af^l a^{-1}, af^l a^{-1}) + (af^{-l} a^{-1}, af^{-l} a^{-1})] \\ &= \sum e_l [(f^l, f^l) + (f^{-l}, f^l) + (f^{-l}, f^{-l}) + (f^l, f^{-l})] \\ &= 0 \in \mathcal{T}_1(\pi_1 B \widetilde{\times} S^1) \end{aligned}$$

confirms Lemma 4.8.3 (using $(f^{-l}, f^{-l}) + (f^l, f^l) = 0$ and $(f^l, f^{-l}) + (f^{-l}, f^l) = 0$, by HOL, FR and AS relations).

General techniques for computing in the infinitely generated groups $\mathcal{T}_1(\pi_1 B \widetilde{\times} S^1)/\text{INT}(z)$ would be useful.

6. ESSENTIAL KNOTS

The goal of this section is Theorem 5 (subsection 6.4.2) which, together with Theorem 3 in the introduction, classifies stable knot concordance in products $F \times S^1$ of an orientable surface $F \neq S^2$ with the circle. Most of the work involves characterizing new INT relations which arise when defining τ_1 for singular concordances of knots which are not null-homotopic.

Throughout this section we continue to assume that M is an oriented 3-manifold, with $\pi_1 M$ torsion-free, and add the further assumption that M is irreducible. We restrict our attention to knots in such M , eventually focusing on the case where $M = F \times S^1$.

Consider an *essential* knot $K \subset M$ with $[K] = \gamma \neq 1$ generating a cyclic subgroup $\langle \gamma \rangle < \pi_1 M$, for some choice of whisker on K . For

any singular concordance $A \looparrowright M \times I$ of K which supports an order 1 Whitney tower \mathcal{W} , we want to define $\tau_1(A)$ as before by associating decorated Y-trees to the intersections between A and the interiors of the Whitney disks in \mathcal{W} . Defining τ_1 for such an *essential* annulus will require two modifications of the previous definition for inessential annuli: The edge decorations will now be cosets by $\langle \gamma \rangle$, to account for choices of paths in A ; and a reformulation of the $\text{INT}(A)$ relations will account for new indeterminacies in the choices of Whitney disk boundaries. The relevant $\text{INT}(A)$ relations will turn out to be determined by order zero intersections between singular self-concordances which have latitudes representing generators of the centralizer $\zeta(\gamma)$ of γ in $\pi_1 M$.

6.1. Order zero double coset decorations. We start by discussing the order zero intersection invariants that will be used to define the new INT relations, and index the equivalence classes of singular concordances which support an order 1 Whitney tower.

Define the abelian group $\mathcal{T}_0(\langle \gamma \rangle \backslash \pi_1 M / \langle \gamma \rangle)$ in the same way as $\mathcal{T}_0(\pi_1 M)$ except that the order zero trees are decorated by (representatives of) double cosets of $\pi_1 M$ by $\langle \gamma \rangle$.

In defining an order zero intersection invariant for annuli, we will only be concerned with two separate cases: either all univalent vertices will have the same label; or the pairs of univalent vertices on all edges (order zero trees) will be labeled distinctly from a set of two labels. So in the latter case $\mathcal{T}_0(\langle \gamma \rangle \backslash \pi_1 M / \langle \gamma \rangle)$ can be identified with the \mathbb{Z} -span of the double coset space, and in the first case one takes a further quotient by inversion of coset representatives and by killing the double coset represented by the trivial element $1 \in \pi_1 M$.

If A and A' are singular concordances of knots in M representing γ , then sheet-changing paths through the intersections between A and A' determine double cosets, and we define the *order zero non-repeating intersection tree*

$$\lambda_0(A, A') \in \mathcal{T}_0(\langle \gamma \rangle \backslash \pi_1 M / \langle \gamma \rangle)$$

by summing as usual over the signed intersections between A and A' , and ignoring any self-intersections. $\lambda_0(A, A')$ is invariant under homotopy (rel ∂) and is the complete obstruction to pairing $A \cap A'$ by Whitney disks (an order 1 *non-repeating Whitney tower*).

Similarly, the *order zero self-intersection tree*

$$\tau_0(A) \in \mathcal{T}_0(\langle \gamma \rangle \backslash \pi_1 M / \langle \gamma \rangle)$$

is defined by summing over all signed self-intersections of A , is invariant under homotopy (rel ∂), and is the complete obstruction to the existence of an order 1 Whitney tower on A .

6.2. Order 1 coset decorations. Define the abelian group $\mathcal{T}_1(\langle\gamma\rangle\backslash\pi_1 M)$ in the same way as $\mathcal{T}_1(\pi_1 M)$ except that the edges of the Y-tree generators are decorated by (representatives of) left cosets $\langle\gamma\rangle\backslash\pi_1 M$ – here we are assuming that *all edges in our Y-trees are oriented towards the trivalent vertex*. In this section we only consider knots, so all univalent vertices in $\mathcal{T}_1(\langle\gamma\rangle\backslash\pi_1 M)$ have the same label which is suppressed from notation.

$\mathcal{T}_1(\langle\gamma\rangle\backslash\pi_1 M)$ is the quotient of $\mathcal{T}_1(\pi_1 M)$ by the relations:

$$(a, b, c) = (\gamma^q a, \gamma^r b, \gamma^s c) \quad \text{for } q, r, s \in \mathbb{Z}.$$

For an order 1 Whitney tower \mathcal{W} on a singular concordance A of K as above, it is clear that these relations are exactly what is needed to account for choices of sheet-changing paths from A into the Whitney disks \mathcal{W} when defining $\tau_1(A)$.

6.3. INT relations for essential knots. Fixing the boundary and changing the interior of any Whitney disk in \mathcal{W} does not change $\tau_1(\mathcal{W})$ since we are assuming that M is irreducible. Also, by the same argument as in subsection 2.4, changing the boundaries of the Whitney disks in \mathcal{W} by a regular homotopy in A does not change $\tau_1(\mathcal{W})$. However, changing the homotopy class in A of Whitney disk boundaries can change $\tau_1(\mathcal{W})$ as we discuss next.

Consider a Whitney disk W in \mathcal{W} . Changing ∂W by winding one arc m times around A and the other arc n times around A creates a closed curve in A representing $\gamma^m \beta \gamma^n \beta^{-1} \in \pi_1 M$, where β is determined by double-point loops through points paired by W , and the signs of the integers m and n correspond to the direction of winding relative to γ . This closed curve bounds a Whitney disk exactly when $\gamma^m \beta \gamma^n \beta^{-1} = 1$. Solutions to such Baumslag-Solitar equations are highly restricted in 3-manifold groups, for instance if M is sufficiently large then $m = \pm n$ by Jaco ([16]). We make the further assumption that if $\gamma^m \beta \gamma^n \beta^{-1} = 1$, then γ and β commute, as will be the case for all γ when $M = F \times S^1$.

Assuming that $\gamma^m \beta \gamma^n \beta^{-1} = 1$ implies $\gamma \beta \gamma^{-1} \beta^{-1} = 1$, it is sufficient to compute the change in $\tau_1(\mathcal{W})$ due to changing a Whitney disk boundary ∂W by winding one arc around A in the positive γ direction and the other arc around A in the direction of γ^{-1} , and replacing W by a Whitney disk W' with this new boundary. Such a change is illustrated in the left hand side of Figure 27, which also indicates how two thin bands (half-tubes) B and C can be added to W so that $W \cup B \cup C \cup W'$ forms a torus T (here the orientation of W' is reversed). Since A can




FIGURE 27.

be assumed to miss B and C , the order zero intersections between A and T measure the change in τ_1 as follows.

Take a whisker on T to be a common whisker for W and W' where their boundaries meet. As in the previous INT relations, the two descending edges on Y-trees for W and W' are decorated by the same pair of fundamental group elements (now coset representatives), which here can be normalized to 1 and β by choosing the whisker on T appropriately. The change in τ_1 is determined by sheet-changing paths from A into T which represent cosets decorating the third edges of the Y-trees. Note that allowing these paths to cross B is irrelevant since we are working modulo multiplication by γ , but we can not assume to allow paths in T to cross C which would correspond to multiplication by β . So the change in τ_1 is truly measured by order zero intersections between A and the annulus which is gotten from T essentially by cutting out C , as described next.

The band C can be extended to an annulus \overline{C} by adding a small embedded band in W , as indicated by the dotted lines in the right-hand side of Figure 27. Remove \overline{C} from T and extend the resulting boundary components by disjointly embedded collars to parallel copies of K in $M \times \{0\}$ to get an annulus A' which has the same order zero intersections with A as T . Now the change in τ_1 can be expressed as $(1, \beta, \lambda_0(A, A'))$, where λ_0 is the order zero non-repeating intersection tree of subsection 6.1 above. Since A has only cancelling pairs of self-intersections, we can add a parallel copy of A to A' along one boundary component to get a singular concordance A'' of a parallel copy of K such that $\lambda_0(A, A'') = \lambda_0(A, A')$. Now we can extend both A and A'' by $-A$ (and a parallel copy) to get pair of singular *self*-concordances A_1

and A_2 of K and a parallel copy of K . Again, the intersections between $-A$ and its parallel copy occur in cancelling pairs, so the change in τ_1 can be expressed as $(1, \beta, \lambda_0(A_1, A_2))$.

By Lemma 4.8.1 and the homotopy invariance of λ_0 , $\lambda_0(A_1, A_2)$ only depends on the elements determined by latitudes of A_1 and A_2 . Since a latitude of $A_1 = A - A$ determines the trivial element $1 \in \pi_1 M$, A_1 is homotopic to the product self-concordance $K \times I$, also by Lemma 4.8.1. Since A' has a latitude representing β , and A_2 was formed by extending A' by $A - A$, a latitude of A_2 determines the element β in the centralizer $\zeta(\gamma)$ of γ in $\pi_1 M$. Thus, the possible changes in $\tau_1(\mathcal{W})$ are measured by $\lambda_0(K \times I, A_\beta)$ where β ranges over $\zeta(\gamma)$, and we have the following INT relations.

For $[K] = \gamma \in \pi_1 M$, define the $\text{INT}(K)$ relations by:

$$\text{INT}(K) = (1, \beta, \lambda_0(K \times I, A_\beta))$$

where β ranges over $\zeta(\gamma)$ and A_β is any singular self-concordance of K having a latitude representing β . These INT relations are completely determined by the values on a set of generators for $\zeta(\gamma)$ as illustrated in the next subsection.

It follows from this discussion that (assuming γ has only trivial Baumslag-Solitar solutions) for any singular concordance A of K supporting an order 1 Whitney tower \mathcal{W}

$$\tau_1(A) := \tau_1(\mathcal{W}) \in \frac{\mathcal{T}_1(\langle \gamma \rangle \backslash \pi_1 M)}{\text{INT}(K)}$$

does not depend on the choice of \mathcal{W} , and is invariant under regular homotopy (rel ∂) of A . Also, the geometric conditions equivalent to the vanishing of previous versions of τ_1 hold in this setting since again the relations can all be realized by geometric constructions.

6.4. Stable concordance of knots in $F \times S^1$. We now restrict attention to essential knots in $M = F \times S^1$, the product of a compact orientable surface $F \neq S^2$ with the circle. The results in this section, together with Theorem 3 for null-homotopic knots, give a complete characterization of stable knot concordance in this setting.

6.4.1. The order zero relative self-linking invariant. Fix a knot K_γ in a non-trivial free homotopy class together with a whisker such that $[K_\gamma] = \gamma \in \pi_1 M$. For any singular concordance A from K_γ to any knot K , define the *order zero relative self-intersection tree* $\tau_0(K_\gamma, K)$ by:

$$\tau_0(K_\gamma, K) := \tau_0(A) \in \mathcal{T}_0(\langle \gamma \rangle \backslash \pi_1 M / \langle \gamma \rangle).$$

By [26], it is possible to choose K_γ in each free homotopy class so that the order zero relative self-linking with K_γ classifies order 1 Whitney concordance, meaning that $\tau_0(K_\gamma, K) = \tau_0(K_\gamma, K')$ if and only if K and K' are order 1 Whitney concordant, for any knots K and K' (freely) homotopic to K_γ . (This invariant was denoted $\mu(K_\gamma, K)$ in [26].) Here equality is modulo conjugation by the centralizer $\zeta(\gamma)$, i.e. modulo the restriction to $\zeta(\gamma)$ of the Ψ -action on $\mathcal{T}_0(\langle\gamma\rangle \backslash \pi_1 M / \langle\gamma\rangle)$. Note that choices of whiskers on K_γ giving $[K_\gamma] = \gamma$ differ by loops representing elements in $\zeta(\gamma)$, and all latitudes of singular self-concordances of K_γ determine elements in $\zeta(\gamma)$.

As explained in [26], the key property required of K_γ in order for $\tau_0(K_\gamma, \cdot)$ to classify order 1 Whitney concordance is that $\tau_0(H) = 0 \in \mathcal{T}_0(\langle\gamma\rangle \backslash \pi_1 M / \langle\gamma\rangle)$ for any singular self-concordance H of K_γ . (In the present case M is irreducible; in general the required vanishing of $\tau_0(H)$ would be modulo intersections with 2-spheres – hence the terminology “ K_γ is spherical” in [26].) This key property can be checked on self-homotopies having latitudes representing a set of generators for $\zeta(\gamma)$, and can be easily satisfied by choosing K_γ as follows.

If γ is carried by (a multiple of) a circle fiber, then take K_γ to lie in a tubular neighborhood of a fiber. In this case, $\zeta(\gamma) = \pi_1 M$, and K_γ can be isotoped around any generating loop in M . If γ is not carried by a circle fiber, then take K_γ in a tubular neighborhood of a knot K_ρ representing a primitive element $\rho \in \pi_1 M$, with $\gamma = \rho^n$ for some natural number n . In this case $\zeta(\gamma)$ will be isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$ generated by ρ and a fiber element, unless F is either a 2-torus or 2-disk, and in all cases K_γ can be isotoped around loops representing generators of $\zeta(\gamma)$.

6.4.2. The order 1 relative self-linking invariant. Fix now K_γ as above, which classifies order 1 Whitney concordance in a non-trivial free homotopy class represented by $[K_\gamma] = \gamma \in \pi_1 M$. For any $z \in \mathcal{T}_0(\langle\gamma\rangle \backslash \pi_1 M / \langle\gamma\rangle)$, construct a clasp knot K_z such that $\tau_0(K_\gamma, K_z) = z$ by introducing clasps into K_γ (just as was done for null-homotopic knots). For any K which is order 1 Whitney concordant to K_z (equivalently $\tau_0(K_\gamma, K) = z$), define the *order 1 relative self-intersection tree* $\tau_1(K_z, K)$ by:

$$\tau_1(K_z, K) := \tau_1(A) \in \frac{\mathcal{T}_1(\langle\gamma\rangle \backslash \pi_1 M)}{\text{INT}(z)}$$

where A is any concordance from K_z to K , and the $\text{INT}(z)$ relations are:

$$0 = (1, \beta^r, (1 - \beta^r)(z + \bar{z}))$$

with β ranging over a generating set for $\zeta(\gamma)$, and $r \in \mathbb{Z}$.




FIGURE 28. A knot K_g together with a parallel copy sitting in a slice $F \times I$ of $F \times S^1$. A self-isotopy A_f of the parallel copy up around the circle direction creates the indicated crossing-change intersections, giving $\lambda_0(K_g \times I, A_f) = g + g^{-1} - fg - fg^{-1}$.

To see that these are the correct INT relations ($\text{INT}(z) = \text{INT}(K_z)$) as defined above in subsection 6.3) start by considering Figure 28, which illustrates the case where γ is primitive and z is represented by a single element $g \in \pi_1(F \times S^1) \cong \pi_1 F \times \langle f \rangle$, with both γ and g contained in the subgroup $\pi_1 F$ carried by a thickened surface slice which in the figure is represented by the plane of the paper cross an interval. Isotoping a parallel push-off of K_g up around the circle direction describes an annulus A_f which has a latitude representing the circle generator f , and the order zero non-repeating intersection tree

$$\lambda_0(K_g \times I, A_f) = g + g^{-1} - fg - fg^{-1} = (1 - f)(g + g^{-1})$$

can be computed from crossing changes in the figure. Note that introducing twists between K_g and its parallel push-off only creates terms of the form $\pm(1 - f)$ via this construction which do not contribute to $\text{INT}(g)$ because the 2-torsion elements $(1, f, (1 - f)) = (1, f, 1) - (1, f, f)$ are zero in \mathcal{T}_1 by the AS and FR relations.

Iterating this construction confirms the relations $(1, f^r, (1 - f^r)(g + g^{-1}))$ for the generator $f \in \zeta(\gamma)$ in this case. It is easy to see that the same computation holds if g replaced by any sum $z = \sum g_i$, even if g_i are allowed to contain powers of f as factors (with corresponding clasps

guided by arcs wrapping around circle fibers). One can also check that allowing γ to contain factors of f does not change the computation.

Under the assumption that γ is primitive, the computation so far verifies the formula for the $\text{INT}(z)$ relations in the “generic” case where $\zeta(\gamma) \cong \mathbb{Z} \oplus \mathbb{Z}$ is generated by f and γ , since in this case self-homotopies with latitudes representing γ give only trivial relations: A parallel push-off of the product annulus $K_z \times I$ is disjoint from $K_z \times I$ and has a latitude representing γ .

Consider now the case where $\gamma = f$, so that K_z is created by introducing clasps into a circle fiber K_f . In this case f is central, and for any $\beta \in \pi_1 M$, a self-homotopy A_β of K_z with a latitude representing β and $\lambda_0(K_z \times I, A_\beta) = (1 - \beta)(z + \bar{z})$ can be constructed as follows. Undoing the clasps on a parallel copy K'_z of K_z describes a homotopy of K'_z to a knot K'_f which may twist around the circle fiber K_f . The homotopy so far creates intersections with $K_z \times I$ of the form $z + \bar{z}$ (half of the crossing changes in Figure 28 for each clasp). Next, K'_f can be isotoped around M so that a basepoint on K'_f traces out a loop representing β ; this self isotopy of K'_f only creates intersections with $K_z \times I$ of the form $\pm(1 - \beta)$ corresponding to any twists between K'_f and K_f which are pulled apart and then recreated; as mentioned above these intersections do not contribute to $\text{INT}(z)$. Now reintroducing the clasps into K'_f describes the last part of the self-homotopy A_β of K_z , and creates intersections of the form $-\beta(z + \bar{z})$, so that $\lambda_0(K_z \times I, A_\beta) = (1 - \beta)(z + \bar{z})$ as desired.

The cases where γ is not primitive are checked by constructing self-homotopies similar to the case $\gamma = f$ in the previous paragraph. For instance, if $\gamma = f^n$ then K_z is created from K_{f^n} which is contained in a tubular neighborhood of a circle fiber and the rest of the construction as for the case $n = 1$ applies. The same construction applies for $\gamma = \rho^n$, with ρ a primitive generator of $\zeta(\gamma)$ and K_z created from K_{ρ^n} which is contained in a tubular neighborhood of K_ρ ; and also to any γ in the non-generic cases where F is a torus or a disk.

Combined with Theorem 3, the following theorem gives a classification of stable knot concordance in $F \times S^1$:

Theorem 5. *For $K_z \subset M = F \times S^1$ as above, the following are equivalent:*

- (i) $\tau_1(K_z, K) = \tau_1(K_z, K')$.
- (ii) K and K' are stably concordant.
- (iii) K and K' are order 2 Whitney concordant.
- (iv) K and K' are height 1 Whitney concordant.

Proof. The proof of Theorem 5 follows the arguments in subsections 4.6 and 4.7 in the proof of Theorem 3. The key point needed is that τ_1 vanishes on all self-homotopies of K_z which support an order 1 Whitney tower (equivalently, have vanishing τ_0). If H_ψ is a self-homotopy of K_z with a latitude representing ψ , then $\psi \in \zeta(\gamma)$, and by Lemma 4.8.1 $\tau_0(H_\psi)$ only depends on ψ . It follows that $\tau_0(H_\psi) = z - \psi z \psi^{-1}$, since such H_ψ can be constructed by undoing the clasps on K_z to get K_γ , then isotoping K_γ around $\psi \in \zeta(\gamma)$, then redoing the clasps to return to K_z . Thus, such an H_ψ admits an order 1 Whitney tower if and only if $\tau_0(H_\psi)$ vanishes if and only if $\psi \in \Psi_z \cap \zeta(\gamma)$. In the current setting, Ψ_z is always untwisted, and it can be easily checked that for $\psi \in \Psi_z \cap \zeta(\gamma)$ the self-isotopies of K_γ realizing ψ (as described above in subsection 6.4.1) extend to self-isotopies of K_z . \square

REFERENCES

- [1] **C. Bohr**, *Stabilisation, bordism and embedded spheres in 4-manifolds*, Algebr. Geom. Topol. 2 (2002) 219-238
- [2] **S. Boyer, D. Rolfsen, B. Wiest**, *Orderable 3-manifold groups*, Ann. Inst. Fourier 55 (2005) 243-288.
- [3] **A. Casson, D. Jungreis**, *Convergence groups and Seifert fibered 3-manifolds*. Inventiones Math. Vol. 118, (1994) 441-456.
- [4] **V. Chernov**, *Framed knots in 3-manifolds and affine self-linking numbers*, J. Knot Theory Ramifications 14 (2005), no. 6, 791-818.
- [5] **J. Conant, R. Schneiderman, P. Teichner**, *Jacobi identities in low-dimensional topology*, Compositio Mathematica 143 Part 3 (2007) pp.780-810
- [6] **J. Conant, R. Schneiderman, P. Teichner**, *Milnor's invariants, Whitney towers, and grope concordance*, In preparation.
- [7] **M. Freedman, R. Kirby**, *A geometric proof of Rochlin's theorem*, Algebraic and geometric topology (Stanford CA 1976), Part 2, Proc. Sympos. Pure Math. Vol 32 AMS Providence RI (1978) 85-97.
- [8] **M. Freedman, F. Quinn**, *The topology of 4-manifolds*, Princeton Math. Series 39 Princeton, NJ, (1990).
- [9] **D. Gabai**, *Convergence groups are Fuchsian groups*. Ann. of Math. Vol. 36, (1992) 447-510.
- [10] **C. H. Giffen**, *Link concordance implies link homotopy*, Math. Scand. 45 (1979) 243-254.
- [11] **D. Goldsmith**, *Concordance implies homotopy for classical links in M^3* , Comment. Math. Helvetici 54 (1979) 347-355.
- [12] **S. Garoufalidis, M. Goussarov, M. Polyak**, *Calculus of clovers and finite type invariants of 3-manifolds*, Geom. Topol. 5 (2001) 75-108.
- [13] **S. Garoufalidis, J. Levine**, *Homology surgery and invariants of 3-manifolds*, Geometry and Topology Vol. 5 (2001) 75-108.
- [14] **K. Habiro**, *Claspers and finite type invariants of links*, Geometry and Topology Vol. 4 (2000) 1-83.

- [15] **A. Hatcher**, *Notes on basic 3-manifold topology*. Posted at www.math.cornell.edu/~hatcher.
- [16] **W. Jaco**, *Roots, relations and centralizers in 3-manifold groups*. Lecture Notes in Math. Vol. 438, (1975) 283–309.
- [17] **W. Jaco, P. Shalen**, *Seifert fibered spaces in 3-manifolds*. Memoirs of Amer. Math. Soc. Vol. 21 No. 220, (1979).
- [18] **W. Jaco, P. Shalen**, *A new decomposition theorem for irreducible sufficiently-large 3-manifolds*. Proceedings of Symposia in Pure Mathematics, Amer. Math. Soc. Vol. 32, (1978) 71–84.
- [19] **E. Kalfagianni**, *Finite type invariants for knots in 3-manifolds*. Topology 37, (1998) 673–707.
- [20] **P. Kirk, C. Livingston**, *Type 1 knot invariants in 3-manifolds*. Pacific Journal of Mathematics Vol. 183 No. 2, (1998) 305–331.
- [21] **P. Kirk, C. Livingston**, *Knot invariants in 3-manifolds and essential tori*. Pacific Journal of Mathematics Vol. 197 No. 1, (2001) 73–96.
- [22] **J. Levine**, *Surgery equivalence of links*, Topology 26 (1987) 45–61.
- [23] **Y. Matsumoto**, *Secondary intersectional properties of 4-manifolds and Whitney's trick*, Proceedings of Symposia in Pure mathematics Vol. 32 Part 2 (1978) 99–107.
- [24] **R. Norman**, *Dehn's lemma for certain 4-manifolds*, Invent. Math. 7 (1969) 143–147.
- [25] **N. Sato**, *Cobordisms of semi-boundary links*, Topology Appl. 18 (1984) 225–234.
- [26] **R. Schneiderman**, *Algebraic linking numbers of knots in 3-manifolds*, Alg. and Geom. Topology 3 (2003) 921–968.
- [27] **R. Schneiderman**, *Whitney towers and Grotes in 4-manifolds*, Trans. Amer. Math. Soc. 358 (2006), 4251–4278.
- [28] **R. Schneiderman**, *Simple Whitney towers, half-grotes and the Arf invariant of a knot*, Pacific Journal of Mathematics, Vol. 222, No. 1, Nov (2005).
- [29] **R. Schneiderman, P. Teichner**, *Higher order intersection numbers of 2-spheres in 4-manifolds*, Alg. and Geom. Topology 1 (2001) 1–29.
- [30] **R. Schneiderman, P. Teichner**, *Whitney towers and the Kontsevich integral*, Proceedings of a conference in honor of Andrew Casson, UT Austin 2003, Geometry and Topology Monograph Series, Vol. 7 (2004), 101–134.
- [31] **R. Stong**, *Existence of π_1 -negligible embeddings in 4-manifolds: A correction to Theorem 10.5 of Freedman and Quinn*, Proc. of the A.M.S. 120 (4) (1994) 1309–1314.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, LEHMAN COLLEGE,, CITY UNIVERSITY OF NEW YORK, NEW YORK NY