ONE MORE PATHOLOGY OF C*-ALGEBRAIC TENSOR PRODUCTS

V. MANUILOV

ABSTRACT. We define a collection of tensor product norms for C^* -algebras and show that a symmetric tensor product functor on the category of separable C^* -algebras need not be associative.

1. INTRODUCTION

Following E. Kirchberg, [3], we call a bifunctor $(A, B) \to A \otimes_{\alpha} B$ a C^* -algebraic tensor product functor if it is obtained by completing of the algebraic tensor product $A \odot B$ of C^* -algebras in a functional way with respect to a suitable C^* -norm $\|\cdot\|_{\alpha}$. We call such a functor symmetric if the standard isomorphism $A \odot B \cong B \odot A$ extends to an isomorphism $A \otimes_{\alpha} B \cong B \otimes_{\alpha} A$. Similarly, we call it associative if the standard isomorphism $A \odot (B \odot C) \cong (A \odot B) \odot C$ extends to an isomorphism $A \otimes_{\alpha} (B \otimes_{\alpha} C) \cong (A \otimes_{\alpha} B) \otimes_{\alpha} C$ for any C^* -algebras A, B, C. It is well known that both the minimal tensor product functor \otimes_{\min} and the maximal tensor product functor \otimes_{\max} are symmetric and associative.

In this paper we construct a collection of symmetric C^* -algebraic tensor product functors related to asymptotic homomorphisms of C^* -algebras. For technical reasons we restrict ourselves to the category of *separable* C^* -algebras. Using C^* -algebras related to property T groups [9] we show that some of these tensor product functors are not associative.

Recall that asymptotic homomorphisms of C^* -algebras were first defined and studied in [2] in relation to topological properties of C^* -algebras. The most important and the best known case is the case of asymptotic homomorphisms from a suspended C^* -algebra SA to the C^* -algebra K of compact operators, since the homotopy classes of those are the K-homology of A, the E-theory. Asymptotic homomorphisms to other C^* -algebras are less known. For example, it is known that any asymptotic homomorphism to the Calkin algebra is homotopic to a genuine homomorphism [4, 6]. Even less is known about asymptotic homomorphisms to $\mathbb{B}(H)$, where there is no topological obstruction (recall that the K-groups of $\mathbb{B}(H)$ are trivial). Such asymptotic homomorphisms are called *asymptotic representations* and were first studied in relation to the asymptotic tensor product C^* -algebras [7] and to semi-invertibility of C^* -algebra extensions [8].

2. Definition of asymptotic C^* -tensor products

Recall [2] that an asymptotic homomorphism φ from a C^* -algebra A to a C^* -algebra D is a family of maps $\varphi = (\varphi_t)_{t \in [0,\infty)} : A \to D$ satisfying the following properties:

- 1. the map $t \mapsto \varphi_t(a)$ is continuous for any $a \in A$;
- 2. $\lim_{t\to\infty}\varphi_t(a+\lambda b) \varphi_t(a) \lambda\varphi_t(b) = \lim_{t\to\infty}\varphi_t(a^*) \varphi_t(a)^* = \lim_{t\to\infty}\varphi_t(ab) \varphi_t(a)\varphi_t(b) = 0 \text{ for any } a, b \in A \text{ and any } \lambda \in \mathbb{C}.$

Let $\mathbb{L}(H)$ be the algebra of bounded operators on a separable Hilbert space H. Our point is that we would like to consider D as a C^* -subalgebra of $\mathbb{L}(H)$: $D \subset \mathbb{L}(H)$. We also

²⁰⁰⁰ Mathematics Subject Classification. Primary 46L06, Secondary 46L05.

The author acknowledges partial support from RFFI, grant No. 07-01-00046.

V. MANUILOV

view asymptotic homomorphisms to D as asymptotic representations on H taking values in D. We are mostly interested in the special case $D = \mathbb{K}^{\infty} = \prod_{n=1}^{\infty} \mathbb{K} = \prod_{n=1}^{\infty} \mathbb{K}(H_n)$, where $H_n = H$ for all $n \in \mathbb{N}$ and $\mathbb{K} = \mathbb{K}(H)$ is the C^* -algebra of compact operators on H.

Let A, B be separable C^{*}-algebras and let $\varphi = (\varphi_t)_{t \in [0,\infty)}, \psi = (\psi_t)_{t \in [0,\infty)}$ be asymptotic representations of A and B respectively, taking values in D.

Let $A \odot B$ be the algebraic tensor product of A and B. For each $a \in A$ and $b \in B$, we can define elements $a^{\varphi \otimes \psi}, b^{\varphi \otimes \psi} \in C_b([0, \infty), \mathbb{L}(H \otimes H))$ by

$$a^{\varphi \otimes \psi}(t) = \varphi_t(a) \otimes 1_H$$
 and $b^{\varphi \otimes \psi}(t) = 1_H \otimes \psi_t(b).$

Note that $a^{\varphi \otimes \psi}(t) \cdot b^{\varphi \otimes \psi}(t) \in C_b([0,\infty), D \otimes_{\min} D))$, where \otimes_{\min} denotes the minimal tensor product of C^* -algebras.

We can then define a *-homomorphism

$$\varphi \otimes \psi : A \odot B \to C_b([0,\infty), D \otimes_{\min} D) / C_0([0,\infty), D \otimes_{\min} D)$$

such that

$$\varphi \otimes \psi \left(\sum_{i} a_{i} \otimes b_{i} \right) = q \left(\sum_{i} a_{i}^{\varphi \otimes \psi} \cdot b_{i}^{\varphi \otimes \psi} \right),$$

where

$$q: C_b\left([0,\infty), D \otimes_{\min} D\right) \to C_b\left([0,\infty), D \otimes_{\min} D\right) / C_0\left([0,\infty), D \otimes_{\min} D\right)$$

is the quotient map. Note that

$$\|\varphi \otimes \psi(c)\| = \limsup_{t \to \infty} \left\| \sum_{i} \varphi(a_i) \otimes \psi(b_i) \right\|$$

for any $c = \sum_i a_i \otimes b_i \in A \odot B$. We can now define a seminorm $\|\cdot\|_{D,0}$ on $A \odot B$ by

$$||c||_{D,0} = \sup_{\varphi,\psi} ||\varphi \otimes \psi(c)||,$$

where we take the supremum over all pairs (φ, ψ) , where φ and ψ are asymptotic representations of A and B, respectively, taking values in D.

Note that a genuine *-homomorphism from A to D can be considered as an asymptotic representation in the obvious way. So, if $D = \mathbb{L}(H)$ then $\|\cdot\|_{D,0} \ge \|\cdot\|_{\min}$, and $\|\cdot\|_{D,0}$ is a norm. This norm coincides with the symmetric asymptotic tensor norm defined in [8]. More generally, the seminorm $\|\cdot\|_{D,0}$ is a norm if there exist *faithful* asymptotic representations of A and B taking values in D. Remark that there are other C*-algebras D, besides $\mathbb{L}(H)$, that admit faithful asymptotic representations of any separable C*algebra. For example, it follows from [6] that one can take the *coarse Roe algebra* of \mathbb{Z} as D.

In general, the seminorm $\|\cdot\|_{D,0}$ may be degenerate (e.g. it may happen that any asymptotic representation of a C^* -algebra A taking values in some D may be asymptotically equivalent to zero, see Lemma 2 below), so let us define the norm $\|\cdot\|_D$ on $A \odot B$ by

$$||c||_D = \max\{||c||_{\min}, ||c||_{D,0}\},\$$

where $c \in A \odot B$. Clearly, $\|\cdot\|_D$ is a C^{*}-norm, hence a cross-norm, and

$$\|\cdot\|_{\min} \le \|\cdot\|_D \le \|\cdot\|_{\max}.$$

We denote by $A \otimes_D B$ the C^* -algebra obtained by completing $A \odot B$ with respect to the norm $\|\cdot\|_D$. Obviously the correspondence $(A, B) \mapsto A \otimes_D B$ is a C^* -algebraic tensor product functor on the category of separable C^* -algebras.

Lemma 1. The functor \otimes_D is symmetric.

Proof. Obvious.

3. Asymptotic representations taking values in \mathbb{K}^{∞}

Let G be a residually finite infinite property T group, let π_n be the sequence of all non-equivalent irreducible unitary representations on finitedimensional Hilbert spaces H_n and let \overline{A} be the C*-algebra generated by operators $\bigoplus_{n=1}^{\infty} \pi_n(g), g \in G$. We denote by Ethe C*-subalgebra in $\mathbb{L}(\bigoplus_{n=1}^{\infty} H_n)$ generated by \overline{A} and by compact operators: $E = \overline{A} + \mathbb{K}$. Put $A = E/\mathbb{K}$. This C*-algebra was first considered by S. Wassermann and we refer to his paper [9] for more details.

Lemma 2. Let $\varphi = (\varphi_t)_{t \in [0,\infty)} : A \to \mathbb{K}^{\infty}$ be an asymptotic homomorphism. Then φ is asymptotically equivalent to zero, i.e. $\lim_{t\to\infty} \varphi_t(a) = 0$ for any $a \in A$.

Proof. Let $q_n : \prod_{n=1}^{\infty} \mathbb{K} \to \mathbb{K}$ is the projection onto the *n*-th copy. Then, for any $\varepsilon > 0$ there exists t_0 and continuous families $p_n(t) \in \mathbb{K}_n$ of finitedimensional projections such that

$$\|q_n \circ \varphi_t(1) - p_n(t)\| < \varepsilon \tag{1}$$

for any $t > t_0$ and any $n \in \mathbb{N}$. Let N_n denote the rank of $p_n(t)$. Since all projections of the same finite rank are unitarily equivalent, without loss of generality (by changing φ by a unitarily equivalent asymptotic homomorphism) we may assume that all $p_n(t)$ are *t*independent, $p_n(t) = p_n$. Then the formula $\psi_t(a) = p_n(q_n \circ \varphi_t(a))p_n$ defines an asymptotic homomorphism from A to the matrix algebra. The group G with the stated properties is known to be finitely generated, so without loss of generality we may assume that $\psi_t(g_i)$ are unitaries, where $g_i \in G$, $i = 1, \ldots, k$, are generators for G.

Since the direct product of k copies of the unitary group U_{N_n} is compact, so the set $\{(\psi_t(g_1), \ldots, \psi_t(g_k)) : t \in [0, \infty)\}$ has an accumulation point $(u_1, \ldots, u_k) \in U_{N_n}^k$. If we put $\sigma(g_i) = u_i$ then this map extends to a genuine representation of G of dimension N_n . Indeed, G is a quotient of the free group \mathbb{F}_k generated by g_1, \ldots, g_k modulo some relations and each ψ_t and σ obviously define representations of \mathbb{F}_k , which we denote by the same characters. If $r \in \mathbb{F}_k$ is a relation then $\lim_{t\to\infty} ||\psi_t(r) - p_n|| = 0$. Therefore, $\sigma(r) = p_n$, hence σ factorizes through a representation of G.

Suppose that $p_n \neq 0$ for some *n*. This implies that the representation σ is non-zero, hence it equals one of π_j . Since

$$||a|| \ge \limsup_{t \to \infty} ||\psi_t(a)|| \ge ||\sigma(a)|| = ||\pi_j(a)||,$$

we have $\limsup_{n\to\infty} \|\pi_n(a)\| \ge \|\pi_j(a)\|$ for any $a \in A$. Then the identity map of G extends to a *-homomorphism $i : A \to C^*_{\pi_j}(G)$, where $C^*_{\pi}(G)$ denotes the C^* -algebra generated by the representation π . Tensoring it by $\operatorname{id}_{C^*_{\pi_j}(G)}$, where $\overline{\pi}$ denotes the contragredient representation for a representation π , we get a *-homomorphism

$$i \otimes \operatorname{id}_{C^*_{\overline{\pi}_j}(G)} : A \otimes C^*_{\overline{\pi}_j}(G) \to C^*_{\pi_j}(G) \otimes C^*_{\overline{\pi}_j}(G).$$
 (2)

We do not specify the tensor product norm here because $C^*_{\overline{\pi}_j}(G)$ is finitedimensional, hence nuclear. It was shown in [9] that the norm on the left hand side of (2) is strictly smaller than the norm on the right hand side, so this *-homomorphism cannot exist. This contradiction shows that $p_n = 0$ for all n, hence (1) implies that $\lim_{t\to\infty} ||q_n \circ \varphi_t(1)|| = 0$ uniformly in n, therefore, $\lim_{t\to\infty} ||q_n \circ \varphi_t(a)|| = 0$ uniformly in n for any $a \in A$.

Corollary 3. For A defined above, one has $A \otimes_{\mathbb{K}^{\infty}} B = A \otimes_{\min} B$ for any C^* -algebra B.

Proof. Since

 $\|\varphi \otimes \psi(a \otimes b)\| = \limsup_{t \to \infty} \|\varphi_t(a) \otimes \psi_t(b)\| = \limsup_{t \to \infty} \|\varphi_t(a)\| \cdot \|\psi_t(b)\| = 0$ for any $a \in A, b \in B$ and for any asymptotic representations φ and ψ , one has

 $\|a \otimes b\|_{\mathbb{K}^{\infty},0} = \sup_{\omega,\psi} \|\varphi \otimes \psi(a \otimes b)\| = 0,$

hence $||c||_{\mathbb{K}^{\infty},0} = 0$ for any $c \in A \odot B$, therefore,

$$||c||_{\mathbb{K}^{\infty}} = \max\{||c||_{\min}, 0\} = ||c||_{\min}$$

4. An example of an asymptotic representation taking values in \mathbb{K}^{∞}

Let $C = C_0(0, 1]$. We are going to construct an asymptotic representation φ^0 of $C \otimes A$ taking values in \mathbb{K}^{∞} . (We do not specify here the tensor norm since C is nuclear.)

Let $\chi : A \to E$ be a continuous homogeneous selfadjoint selection map, cf. [1]. We denote by P_n the projection in $\bigoplus_{n=1}^{\infty} H_n$ onto H_n . For $a \in A$ put $\alpha(a) = \iota \circ \chi(a)$, where $\iota : E \to \mathbb{L}(\bigoplus_{n=1}^{\infty} H_n)$ is the standard inclusion.

Let $\{\tau_n\}_{n\in\mathbb{N}}$ be a dense sequence of points in (0,1).

For $t = k \in \mathbb{N}$ and for $f \in C$ put

$$\beta_k(f) = \sum_{n=k+1}^{\infty} f(\tau_n) P_n.$$

(this sum also is convergent with respect to the *-strong topology). If k < t < k+1 then put

$$\beta_t(f) = f((t-k)\tau_{k+1})P_{k+1} + \beta_{k+1}(f).$$

Let $F \in C \otimes A$. One can consider F as a continuous function on [0, 1] taking values in A such that F(0) = 0. Put

$$\phi_k(F) = \sum_{n=k+1}^{\infty} P_n \alpha(F(\tau_n)) P_n$$

where the sum is *-strongly convergent, and

$$\phi_t(F) = P_{k+1}\alpha(F((t-k)\tau_{k+1}))P_{k+1} + \phi_k(F)$$

for k < t < k + 1.

Lemma 4. The family of maps $(\phi_t)_{t \in [0,\infty)}$ is an asymptotic representation of $C \otimes A$ taking values in $\prod_{n=1}^{\infty} \mathbb{L}(H_n) \subset \mathbb{K}^{\infty}$.

Proof. By the definition, the maps ϕ_t , $t \in [0, \infty)$, take values in $\prod_{n=1}^{\infty} \mathbb{L}(H_n)$, so we only need to check that algebraic properties hold asymptotically. Let us check that for multiplication, as other properties can be checked in the same way. Let $F_1, F_2 \in C \otimes A$, then the operators

$$K_n = \alpha(F_1(\tau_n)F_2(\tau_n)) - \alpha(F_1(\tau_n))\alpha(F_2(\tau_n)) \in \mathbb{K}$$

lie in a compact subset of \mathbb{K} (the image of [0, 1] under the continuous map $\tau \mapsto \alpha(F_1(\tau)F_2(\tau)) - \alpha(F_1(\tau))\alpha(F_2(\tau)))$, hence

$$\lim_{k \to \infty} \phi_k(F_1 F_2) - \phi_k(F_1)\phi_k(F_2) = \lim_{k \to \infty} \sum_{n=k+1}^{\infty} P_n K_n P_n = 0.$$

Finally, we easily pass to the continuous parameter: $\lim_{t\to\infty} \phi_t(F_1F_2) - \phi_t(F_1)\phi_t(F_2) = 0$.

Note that if $F = f \otimes a \in C \otimes A$ then

$$\phi_t(f \otimes a) = \sum_{n=k}^{\infty} P_n \alpha(a) P_n \cdot \beta_t(f).$$

5. Compairing tensor norms

Let B be the C^{*}-algebra generated by operators $\bigoplus_{n=1}^{\infty} \overline{\pi}_n(g)$, $g \in G$, where $\overline{\pi}$ denotes the contragredient representation for π .

Theorem 5. The tensor products $C \otimes_{\mathbb{K}^{\infty}} (A \otimes_{\mathbb{K}^{\infty}} B)$ and $(C \otimes_{\mathbb{K}^{\infty}} A) \otimes_{\mathbb{K}^{\infty}} B$ are not canonically isomorphic.

Proof. Let $f \in C$ be the identity function, $f(\tau) = \tau$, and let $\{g_1, \ldots, g_m\}$ be a symmetric set of generators of the group G as above. We identify the group elements with the corresponding unitaries in C^* -algebras generated by representations of G (like B) and in their quotients (like A). Let $d = \sum_{i=1}^m f \otimes g_i \otimes g_i \in C \odot A \odot B$. Denote by $\|\cdot\|_1$ and by $\|\cdot\|_2$ the norms on $C \odot A \odot B$ inherited from $C \otimes_{\mathbb{K}^\infty} (A \otimes_{\mathbb{K}^\infty} B)$ and $(C \otimes_{\mathbb{K}^\infty} A) \otimes_{\mathbb{K}^\infty} B$ respectively. Our aim is to show that $\|d\|_1 \neq \|d\|_2$.

It follows from Lemma 3 and from amenability of C that $C \otimes_{\mathbb{K}^{\infty}} (A \otimes_{\mathbb{K}^{\infty}} B) = C \otimes_{\min} (A \otimes_{\min} B)$, so

$$||d||_1 = ||f|| \cdot \left\| \sum_{i=1}^m g_i \otimes g_i \right\|_{\min} = \left\| \sum_{i=1}^m g_i \otimes g_i \right\|_{\min}$$

It was shown in [9] that the latter norm is strictly smaller than m, so

$$||d||_1 < m.$$
 (3)

When estimating the norm $\|\cdot\|_2$ from below, we may use two special asymptotic representations instead of taking the supremum over all of them. Let us take ϕ_t for $C \otimes A$ and the identity representation for B. Then

$$\begin{split} \|d\|_{2} &\geq \lim \sup_{t \to \infty} \left\| \sum_{i=1}^{m} \phi_{t}(f \otimes g_{i}) \otimes \sum_{n=1}^{\infty} \overline{\pi}_{n}(g_{i})P_{n} \right\| \\ &= \lim \sup_{t \to \infty} \left\| \sum_{i=1}^{m} \sum_{n=1}^{\infty} P_{n}\beta_{t}(f)\alpha(g_{i})P_{n} \otimes \sum_{n=1}^{\infty} \overline{\pi}_{n}(g_{i})P_{n} \right\| \\ &\geq \lim \sup_{t \to \infty} \sup_{n} \left\| P_{n}\beta_{t}(f) \sum_{i=1}^{m} \alpha(g_{i})P_{n} \otimes \overline{\pi}_{n}(g_{i})P_{n} \right\| \\ &\geq \lim \sup_{n \to \infty} \left\| P_{n}\beta_{n}(f) \sum_{i=1}^{m} \alpha(g_{i})P_{n} \otimes \overline{\pi}_{n}(g_{i})P_{n} \right\| \\ &= \lim \sup_{n \to \infty} f(\tau_{n}) \cdot \left\| \sum_{i=1}^{m} P_{n}\alpha(g_{i})P_{n} \otimes \overline{\pi}_{n}(g_{i})P_{n} \right\| \\ &\geq \lim \sup_{j \to \infty} f(\tau_{n_{j}}) \cdot \left\| \sum_{i=1}^{m} P_{n_{j}}\alpha(g_{i})P_{n_{j}} \otimes \overline{\pi}_{n_{j}}(g_{i}) \right\|, \end{split}$$

where $\{n_j\}$ is any increasing subsequence of integers. Since the sequence $\{\tau_n\}_{n=1}^{\infty}$ is dense in [0, 1], we can find a subsequence $\{n_j\}_{j=1}^{\infty}$ such that $\lim_{j\to\infty} \tau_{n_j} = 1$. Then

$$\|d\|_{2} \geq \limsup_{j \to \infty} f(\tau_{n_{j}}) \cdot \left\| \sum_{i=1}^{m} P_{n_{j}} \alpha(g_{i}) P_{n_{j}} \otimes \overline{\pi}_{n_{j}}(g_{i}) \right\|$$

$$= \limsup_{j \to \infty} \left\| \sum_{i=1}^{m} P_{n_{j}} \alpha(g_{i}) P_{n_{j}} \otimes \overline{\pi}_{n_{j}}(g_{i}) \right\|$$

$$= \limsup_{j \to \infty} \left\| \sum_{i=1}^{m} \pi_{n_{j}}(g_{i}) \otimes \overline{\pi}_{n_{j}}(g_{i}) \right\|$$

$$= \left\| \sum_{i=1}^{m} \pi_{n_{j}}(g_{i}) \otimes \overline{\pi}_{n_{j}}(g_{i}) \right\| = \sum_{i=1}^{m} 1 = m.$$

On the other hand, $||d||_2 \leq \sum_{i=1}^m ||f \otimes g_i \otimes g_i||_2 = m$, so we have

$$\|d\|_2 = m. (4)$$

Compairing (3) and (4), we conclude that these two norms are different.

References

- R. G. Bartle, L. M. Graves, Mappings between function spaces. Trans. Amer. Math. Soc. 72 (1952), 400–413.
- [2] A. Connes, N. Higson, Déformations, morphismes asymptotiques et K-théorie bivariante. C. R. Acad. Sci. Paris Sér. I Math. 311 (1990), 101–106.
- [3] E. Kirchberg. Exact C^{*}-algebrass, tensor products, and the classification of purely infinite algebras. Proc. Internat. Congress Math., Birkhäuser, 1995, 943–954.
- [4] V. Manuilov, Asymptotic homomorphisms into the Calkin algebra. J. Reine Angew. Math. 557 (2003), 159–172.
- [5] V. Manuilov, K. Thomsen, The Connes-Higson construction is an isomorphism. J. Funct. Anal. 213 (2004), 154–175.
- [6] V. Manuilov, K. Thomsen, E-Theory is a Special Case of KK-Theory. Proc. London Math. Soc. 88 (2004), 455–478.
- [7] V. Manuilov, K. Thomsen, On the asymptotic tensor C^{*}-norm. Arch. Math. 86 (2006), 138–144.
- [8] V. Manuilov, K. Thomsen, On the lack of inverses to C^{*}-extensions related to property T groups. Canad. Math. Bull. 50 (2007), 268–283..
- [9] S. Wassermann, C^{*}-algebras associated with groups with Kazhdan's property T. Ann. Math. 134 (1991), 423–431.

DEPARTMENT OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, LENINSKIE GORY, MOSCOW, 119991, RUSSIA and HARBIN INSTITUTE OF TECHNOLOGY, HARBIN, P. R. CHINA *E-mail address:* manuilov@mech.math.msu.su